



Glycosylinositolphosphoceramides in *Aspergillus fumigatus*.

Catherine Simenel, Bernadette Coddeville, Muriel Delepierre, Jean-Paul Latgé, Thierry Fontaine

► To cite this version:

Catherine Simenel, Bernadette Coddeville, Muriel Delepierre, Jean-Paul Latgé, Thierry Fontaine. Glycosylinositolphosphoceramides in *Aspergillus fumigatus*.. *Glycobiology*, 2008, 18 (1), pp.84-96. 10.1093/glycob/cwm122 . pasteur-00366203

HAL Id: pasteur-00366203

<https://pasteur.hal.science/pasteur-00366203>

Submitted on 6 Mar 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Glycosylinositolphosphoceramides in *Aspergillus fumigatus*

Journal:	<i>Glycobiology</i>
Manuscript ID:	GLYCO-2007-00125.R1
Manuscript Type:	Regular Manuscripts
Date Submitted by the Author:	n/a
Complete List of Authors:	Simenel, Catherine; Institut Pasteur, Unité de Résonance Magnétique Nucléaire des Biomolécules Coddeville, Bernadette; Université des Sciences et Technologies de Lille, Unité de Glycobiologie Structurale et Fonctionnelle Delepierre, Muriel; Institut Pasteur, Unité de Résonance Magnétique Nucléaire des Biomolécules Latge, Jean-Paul; Institut Pasteur, parasitology Mycology Thierry, Fontaine; Institut Pasteur, Unite des Aspergillus
Key Words:	aspergillus fumigatus, glycosylinositolphosphoceramide, NMR, galactofuranose, glycosphingolipid




Figure 1




Figure 3




Figure 4





Figure 5



Glycosylinositolphosphoceramides

in *Aspergillus fumigatus*

Catherine Simenel¹, Bernadette Coddeville², Muriel Delepierre¹, Jean-Paul Latgé³ and Thierry Fontaine^{3*}

Unité de Résonance Magnétique Nucléaire des Biomolécules, CNRS URA 2185 Institut Pasteur, 25 rue du Docteur Roux 75724 Paris cedex 15,

²Unité de Glycobiologie Structurale et Fonctionnelle, UMR 8576 CNRS, Université des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq cedex

Unité des *Aspergillus*, Institut Pasteur, 25 rue du Docteur Roux 75724 Paris cedex 15.

*corresponding author: tfontain@pasteur.fr

Summary

Fungal glycosylinositolphosphoceramides (GIPCs) are involved in cell growth and fungal-host interactions. In this study, 6 GIPCs from the mycelium of the human pathogen *Aspergillus fumigatus* were purified and characterized using Q-TOF mass spectrometry and ¹H, ¹³C and ³¹P NMR. All structures have the same inositolphosphoceramide moiety with the presence of a C_{18:0}-phytosphingosine conjugated to a 2-hydroxylated saturated fatty acid, (2-hydroxy-lignoceric acid). The carbohydrate moiety defines 2 types of GIPC. The first, a mannosylated zwitterionic glycosphingolipid contains a glucosamine residue linked in α1-2 to an inositol ring that has been described in only two other fungal pathogens. The second type of GIPC presents a α-Manp-(1→3)-α-Manp-(1→2)-IPC common core. A galactofuranose residue is found in 4 GIPC structures, mainly at the terminal position via a β1-2 linkage. Interestingly, this galactofuranose residue could be substituted by a choline-phosphate group, as observed only in the GIPC of *Acremonium sp.*, a plant pathogen.

Introduction

Glycosphingolipids (GSLs) are membrane glycolipids found in all eukaryotic cells that are composed of a carbohydrate and a ceramide hydrophobic moiety. In yeast and fungi, three types of GSLs have been identified. The first class is represented by neutral β -glucosyl-ceramide and β -galactosyl-ceramide. This type of GSL plays a crucial role in spore germination, hyphal growth and in the cell cycle (Leverly *et al.*, 2002; da Silva *et al.*, 2004; Barreto-Bergter *et al.*, 2004; Rittershaus *et al.*, 2006). The second class of fungal GSLs is composed of a complex neutral moiety and a saturated ceramide, suggesting a different biosynthetic pathway (Maciel *et al.*, 2002; Aoki *et al.*, 2004b; Barreto-Bergter *et al.*, 2004). The third class is composed of glycosylinositolphosphoceramides (GIPCs) which are acidic GSLs containing a phosphodiester linkage between inositol and ceramide. In contrast to cerebrosides, these phosphorylinositol-containing sphingolipids are not present in mammals but have been detected in protozoa, plants, fungi and nematodes. The sphingolipids are essential for fungal growth since the deletion of inositolphosphorylceramide (IPC) synthase that catalyses the transfer of inositol and phosphate to ceramide is lethal in *Saccharomyces* and *Aspergillus* (Nagiec *et al.*, 1997; Cheng *et al.*, 2001; Hu *et al.*, 2007). In filamentous fungi, the ceramide moiety is composed of a phytosphingosine associated with a saturated long chain fatty acid containing 18 to 26 carbon atoms with or without a hydroxyl group in position 2. The carbohydrate moiety is more variable. Three types of GIPCs have been mainly identified based on the monosaccharide and its linkage to the inositol ring: i) α -Man-(1-2)-IPC found in numerous species (Barr *et al.*, 1984; Leverly *et al.*, 1998; 2001; Heise *et al.*, 2002), ii) α -Man-(1-6)-IPC, found in *Sporothrix schenckii*, (Loureiro y Penha *et al.*, 2001; Toledo *et al.*, 2001a) iii) α -GlcN-(1-2)-IPC described in *S. schenckii*, *Acremonium sp.* and *Aspergillus fumigatus* (Toledo *et al.*, 2001b; 2007; Aoki *et al.*, 2004a). To these core structures, other monosaccharides such as fucose, xylose, galactose or choline-phosphate

could be associated (Jennemann *et al.*, 1999; Heise *et al.*, 2002; Arigi *et al.*, 2007; Gutierrez *et al.*, 2007). The presence of an α -Man-(1-4)-IPC sequence has been only reported in Basidiomycetes, outside the 3 families described suggesting a high level of complexity in the structure of the fungal sphingolipids (Jennemann *et al.*, 2001).

A. fumigatus is a saprophytic, filamentous fungus found in most environments where it plays an important role in the recycling of organic materials. *A. fumigatus* is also an opportunistic pathogen responsible for severe pulmonary diseases, particularly in immunocompromised patients (Latgé, 1999). Man₂-IPC and five other GIPC structures containing additional mannose, galactofuranose, glucosamine or N-acetylglucosamine residues have been previously identified in *A. fumigatus* (Leverly *et al.*, 2001; Toledo *et al.*, 2007). This fungus also produces a lipogalactomannan linked to the cellular membrane through a GlcN-IPC (Costachel *et al.*, 2005). As an effort to identify cell surface glycans and antigens, four new structures of GIPC were isolated from membrane preparations of *A. fumigatus* mycelium and chemically characterized using mass spectrometry and NMR analysis.

Results

The crude membrane preparation of *A. fumigatus* mycelium was treated with chloroform/methanol/water, then with a butanol/water partition to recover a glycolipid preparation. Two liquid chromatographic steps on a DEAE-Sephadex column and Silica 60 column were used to separate 5 glycosphingolipid fractions (Fig. 1). The total amount of GSLs represented around 0.02 % +/- 0.005 of the total mycelium dry weight. DEAE anion exchange chromatography yielded an unbound glycosphingolipid (GSL-U) and negatively charged GSLs. Similar amounts of unbound and bound fractions were purified from a 15-L fermentor in Sabouraud medium of *A. fumigatus*. Silica gel column chromatography was used to purify the unbound GSL-U and 4 fractions of negatively charged GSLs (GSL-A, GSL-B,

GSL-C, GSL-D) (Fig. 1). The GSL-A and GSL-C fractions contain two molecules. Only one spot was detected in fractions B and D.

Composition analysis obtained by GLC and GLC-MS revealed the presence of C_{18:0}-phytosphingosine and a 2-monohydroxylated C_{24:0} fatty acid in all GSL fractions. Minor fatty acids such as 2OH-C_{25:0} and 2OH-C_{26:0} were observed in low amounts (data not shown). Mannose and *myo*-inositol were identified in all GSL fractions, whereas glucosamine was detected only in the GSL-U fraction and galactose only in negatively-charged GSL fractions (GSL-A to D) (Table I).

GSL fractions have been analysed by Q-TOF mass spectrometry to identify the molecular weight of isolated glycosphingolipids; pseudomolecular negative ions m/z are presented in Table 1 and Fig. 2. The pseudomolecular ion at $m/z = 1410$ obtained with GSL-U fraction corresponded to a glycosphingolipid containing 2 hexose residues and 1 hexosamine associated with an inositolphosphoceramide (IPC) whereas the ceramide was composed of a C_{18:0}-phytosphingosine associated with a 2-hydroxylated C_{24:0} fatty acid (Costachel *et al.*, 2005). The presence of minor ions that differ by a mass of 14 or 28 reflected the variability in size of fatty acids. Pseudomolecular ions at $m/z = 1249, 1411, 1573$, obtained with GSL-A, B, C fractions were compatible with a similar IPC with the presence of 2, 3 or 4 hexose residues. A fragmentation pseudomolecular ion at $m/z = 1411$ produced one main ion at $m/z = 745$ [Hex₃-inositol-phosphate]⁻, ions at $m/z = 241$ and 259 corresponding to inositol-phosphate and ions at $m/z = 79$ and 97 corresponding to free phosphate (Fig. 2) (Costachel *et al.*, 2005). Fragmentation of pseudomolecular ions at $m/z = 1249, 1411, 1573$ gave similar patterns of fragmentation with the loss of 666 corresponding to the ceramide moiety (data not shown), characterizing a GIPC and indicating the presence of 2, 3 and 4 hexose residues linked to IPC. In contrast, the fragmentation of a pseudomolecular ion at $m/z = 1576$ in the GSL-D fraction did not produce similar patterns of daughter ions, indicating structural modifications (Fig. 2).

The composition of this GSL-D fraction indicated the presence of the same IPC structure with mannose residues. Indeed, a pseudomolecular mass at $m/z = 1576$ did not correspond to a classical mannosylated IPC, but instead, the presence of a choline-phosphate linked to a Hex₃-IPC. Daughter fragments at m/z 1517 [$\text{CH}_2\text{-CH}_2\text{-P-Hex}_3\text{-IPC}$]⁻, 910 [$(\text{CH}_3)_3\text{N-CH}_2\text{-CH}_2\text{-P-Hex}_3\text{-inositol-P}$]⁻, 851 [$\text{CH}_2\text{-CH}_2\text{-P-Hex}_3\text{-inositol-P}$]⁻ were in agreement with the presence of a choline-phosphate substituent. Moreover, GLC-MS analysis of the sugar-phosphate following methanolysis and trimethylsilylation of the GSL-D fraction permitted the identification of a monosaccharide-6-phosphate with the same retention time as the galactose-6-phosphate obtained from the lipophosphoglycan of *Leishmania donovani* (data not shown). These data suggest the presence of phosphocholine linked to the C-6 of a galactose residue in the GSL-D fraction.

Linkages of monosaccharides were analysed after methylation (Table II). Methylation of the GSL-U fraction revealed the presence of 3 methyl ethers corresponding to a terminal mannose, a mannose substituted in position 3, and a glucosamine substituted in position 6. In agreement with Q-TOF mass spectra and to GSL described in fungal species (Toledo *et al.*, 2001b; 2007; Aoki *et al.*, 2004a), these data suggest the following sequence: Man1-3-Man1-6GlcN. In the GSL-A fraction, the presence of 2 terminal monosaccharides, a mannose and a galactofuranose, without a disubstituted monosaccharide confirmed the presence of two GSL structures. GSL-B fraction contained 3 major methyl ethers with a terminal galactofuranose, indicating, in agreement with the Q-TOF mass spectra, a Gal-Man-Man sequence. In the GSL-C fraction, a disubstituted mannose in position 3 and 6 indicated a branched structure with a terminal galactofuranose and/or mannose. In the GSL-D fraction, 2 main methyl ethers corresponding to monosubstituted mannoses have been identified, but no terminal monosaccharide. A stronger acid hydrolysis yielded a 2,3,5-tri-O-methyl hexitol that should correspond to a galactofuranose substituted in position 6 by a phosphate identified after

methanolysis and trimethylsilylation. In agreement with the Q-TOF mass spectra (Fig. 2), these data suggest a choline-phosphate-6-Gal-f-Man-Man sequence.

NMR analysis of the five glycosphingolipid fractions

The GSL structures of *A. fumigatus* mycelium were elucidated by analysis of homonuclear and heteronuclear two-dimensional NMR experiments.

In agreement with GLC and MS data (Table I), NMR analysis confirmed the presence of an identical ceramide structure in all GSL fractions. From signal integration in the 1D spectrum of the highest concentrated GSL-B fraction, the total number of carbons of the two aliphatic chains was estimated to be 42. A methylene/methyl ratio of 17.5 was in agreement with a C_{18:0} phytosphingosine linked to an unbranched 2-hydroxylated C_{24:0} fatty acid identified by GLC-MS. The carbon linked to the nitrogen atom in the sphingosine base shifted at 53.36ppm was located in the ¹H ¹³C HSQC experiment as well as three ¹H/¹³C shifts corresponding to CHOH groups at 3.451/76.01, 3.348/73.65 and 3.835/74.3 (Fig. 4). A phosphorus atom was detected in the ceramide moiety attached to the glycosidic one. Indeed, three correlations were observed in the ¹H,³¹P HSQC spectrum between the phosphorus atom and two methylene protons of the ceramide on one side and the H1 proton of inositol on the other side (Fig. 3, 4, 5) indicating the sequence inositol-P-ceramide.

GSL-U fraction

The chemical shifts and coupling constants of the glycosidic moiety obtained for the GSL-U fraction are shown in Table III. The 1D ¹H and 2D ¹H, ¹³C gHSQC spectra showed three signals in the anomeric region, in equal proportions as deduced from integration in the 1D spectrum, indicating the presence of three monosaccharide residues (Fig. 3). The ¹H and ¹³C chemical shift analysis and the examination of ³J_{H,H} and ¹J_{C1,H1} values indicated the presence

of 2 α -mannopyranose and one α -glucosamine residues (Bock and Pedersen, 1974; 1983). The *myo*-inositol residue was identified from its H2 equatorial proton in the 1,2,3,4,5,6-cyclohexanehexol ring as assessed by the small $^3J_{H1,H2}$ and $^3J_{H2,H3}$ values (2,7Hz)(Table III).

A NOESY experiment demonstrated a strong correlation between the H1 proton of a mannose residue with the H3 proton of the second mannose residue (Fig. 3). Moreover, C1/H3 and H1/C3 correlations between these two mannose residues were observed in the gHMBC experiment (Fig. 3), indicating a branched sequence of α -Manp-(1 $\rightarrow$ 3)- α -Manp. This sequence accounts for the downfield chemical shift of the C3 at 82.24 ppm of the second mannose residue (Table III). Similarly, dipolar interactions between the H1 proton of the second mannose residue and H6/H6' protons of the glucosamine residue were observed in the NOESY experiment suggesting a $\rightarrow$ 3)- α -Manp-(1 $\rightarrow$ 6)- α -GlcNH₂ sequence. This linkage was confirmed by an gHMBC experiment with the observation of the correlations H1/C6 and C1/H6H6' between the second mannose residue and the glucosamine residue. This is in agreement with a C6 downfield chemical shift at 68.29ppm observed for the glucosamine residue. A strong interaction was also observed in the NOESY experiment between the anomeric proton of the glucosamine residue and the H2 proton of the inositol residue, corroborated by the presence in the gHMBC experiment of a correlation between the C1 carbon of the glucosamine residue and the H2 proton of the inositol residue, indicating the sequence motif: $\rightarrow$ 6)- α -GlcNH₂-(1 $\rightarrow$ 2)-Ins, in agreement with the downfield shift at 85.22 ppm of the inositol C2 carbon. Along with the MS data, the NMR experiments established the following structure: α -Manp-(1 $\rightarrow$ 3)- α -Manp-(1 $\rightarrow$ 6)- α -GlcNH₂-(1 $\rightarrow$ 2)-Ins-(1 $\rightarrow$ O)-P-Cer. This structure has previously been characterized by NMR (Toledo *et al.*, 2007). Differences in 1H and ^{13}C chemical shifts are observed with our results that are associated primarily with the following: i) the ^{13}C referencing method inducing a shift of 3.11ppm, ii) the protonation state of the amine group of the glucosamine residue (Bunel *et al.*, 1993).

GSL-B fraction.

Among the acidic GIPCs, fractions B was analysed first because it was the most abundant; it contained only one GSL spot on HPTLC (Fig. 1). The chemical shifts and coupling constants of the glycosidic moiety obtained for the GSL-B fraction are shown in Table IV. The 1D ^1H and 2D ^1H , ^{13}C gHSQC spectra exhibited three signals of equivalent areas in the anomeric region (5.029/103.85ppm, 5.020/101.71ppm and 4.881/108.70ppm) indicating the presence of three monosaccharide residues (Fig. 4, Table IV). For the two first glycosidic residues, protons were assigned from anomeric proton only up to H3 and H4 respectively using relayed COSY experiments. The missing ring protons partly located in the TOCSY experiment were fully identified using the combination of ^1H , ^{13}C edited HSQC and H2BC experiments recently described as a useful method for tracing the proton-bearing carbon skeleton of a molecule (Petersen *et al.*, 2006). Sugar rings stereochemistry deduced from $^3J_{\text{H,H}}$ coupling constants were consistent with Manp for these two glycosidic residues. Their α -configuration was evident from $^1J_{\text{C1,H1}}$ coupling constant values (169.1 and 173.1Hz). According to the GLC composition and methylation analysis (Tables I and II), the galactose residue was identified in a β -furanosic configuration by its characteristic very low field anomeric ^{13}C resonance at 108.70ppm (Ritchie *et al.*, 1975). The *myo*-inositol residue was identified as previously for the GSL-U fraction. The NMR structural analysis of this fraction explicitly indicated the presence of two α -mannose, one β -galactofuranose and one inositol residues (Table IV).

In the NOESY spectrum, the two inter-residue dipolar interactions observed between the H1 proton of the β -galactofuranose residue and the H1 and H2 protons of a mannose residue did not allow the characterization of a branching point. In the gHMBC spectrum, only H1/C2 and C1/H2 inter-residues correlations between the β -galactofuranose residue and this mannose residue were observed indicating the β -Galf-(1 $\rightarrow$ 2)- α -Manp motif (Fig. 4). This linkage

accounts for the downfield chemical shift of the mannose C2 carbon at 77,08 ppm. In the NOESY spectrum, the H1/H3 interaction between the latter mannose residue and the second mannose residue indicates a $\rightarrow 2$)- α -Manp-(1 $\rightarrow 3$)- α -Manp linkage. This glycosidic sequence was confirmed by H1/C3 and C1/H3 correlations between these two mannose residues observed in the gHMBC experiments but also by the downfield chemical shift of C3 of the second mannose residue at 81.70ppm (Table IV). The strong H1/H2 interaction between the second mannose residue and the inositol residue observed in the NOESY spectrum and the double correlation H1/C2 and C1/H2 observed in the gHMBC experiment between these two residues indicate the following sequence motif: $\rightarrow 3$)- α -Manp-(1 $\rightarrow 2$)-Ins (Fig. 4). This is in agreement with the downfield shift of the inositol C2 carbon resonating at 80.91 ppm (Table IV). Together with methylation and MS data (Fig. 2, Table I), these NMR data established the structure of GSL-B as β -Galf-(1 $\rightarrow 2$)- α -Manp-(1 $\rightarrow 3$)- α -Manp-(1 $\rightarrow 2$)-Ins-(1 $\rightarrow$ O)-P-Cer.

GSL-A fraction.

In agreement with the TLC data, two distinct molecules were observed by NMR in the GSL-A fraction. The chemical shifts and coupling constants of the corresponding glycosidic sequences are shown in Table V. The one-dimensional ^1H spectrum (not shown) exhibited three signals in the anomeric region with areas in the ratio 2/1.2/2. The sugar spin systems assignment and the sequential glycosidic analysis revealed that these signals correspond to five protons belonging to two different molecules present in about equivalent amounts. The GSL-A2 molecule was identical to the GSL-B one. The GSL-A1 structure displayed the identical dimannoside core as GSL-A2 without the β -galactofuranose residue at the non-reducing end. Thus, the C2 carbon resonance of the first mannose residue was not downfield shifted for the GSL-A1 molecule (73.47ppm) upon branching of galactofuranose as observed for the GSL-A2 molecule (77.09ppm). Moreover, only three correlations were observed in the

^1H , ^{31}P HSQC (not shown) between a phosphorus atom (3.099ppm) and the two methylene protons of the ceramide on one side (4.039ppm and 3.658ppm) and the H1 proton of inositol on the other side (3.738ppm). These data are in agreement with the fact that the two molecules differ only at the non-reducing end of the glycosidic moiety. These NMR experiments indicated the presence of the two following structures: GSL A1, α -Manp-(1 $\rightarrow$ 3)- α -Manp-(1 $\rightarrow$ 2)-Ins-(1 $\rightarrow$ O)-P-Cer and GSL A2, β -Galf-(1 $\rightarrow$ 2)- α -Manp-(1 $\rightarrow$ 3)- α -Manp-(1 $\rightarrow$ 2)-Ins-(1 $\rightarrow$ O)-P-Cer

GSL-C fraction.

In this fraction, two different molecules were also observed which differ in their glycosidic sequences (Table VI). The presence of two different molecules was also confirmed by the observation of two close sets of three correlations in the ^1H , ^{31}P HSQC spectrum (data not shown) corresponding to two distinct phosphorus atoms resonating at 2.478ppm and 2.370ppm and interacting with the two methylene protons of the ceramide (δ =4.002 –3.667 ppm and δ =3.997 –3.670 ppm respectively) and the H1 proton of *myo*-inositol at 3.720ppm and 3.713ppm, respectively. The one-dimensional ^1H -NMR spectrum (not shown) exhibited eight H1 resonances of about equal intensity. In the anomeric region of the 2D ^1H , ^{13}C gHSQC spectrum (not shown), six resonances among the eight were grouped two by two that resembled the anomeric region of the GSL-B fraction. Indeed, the sugar spin systems assignment and the sequential glycosidic analysis corresponded to the same glycosidic sequence which is β -Galf-(1 $\rightarrow$ 2)- α -Manp-(1 $\rightarrow$ 3)- α -Manp-(1 $\rightarrow$ 2)-Ins. One of the other anomeric signals at 4.799/111.10 ppm corresponded to a second β -galactofuranose residue identified by its characteristic downfield shifted anomeric carbon. The last anomeric signal at 4.640/102.57ppm was assigned to a α -mannopyranose residue. The ^1H , ^{13}C edited gHSQC methylene region of the GSL-C fraction compared to that of the GSL-B fraction showed two

extra downfield shifted C6 at 68.27ppm and 69.79ppm indicating that the two corresponding mannose residues were 6-O-substituted.

Thus, NMR data allowed the identification of the two following molecules :

GSL-C1, β -Galf(1 $\rightarrow$ 2)- α -Manp(1 $\rightarrow$ 3)-[α -Manp(1 $\rightarrow$ 6)]- α -Manp-(1 $\rightarrow$ 2)-Ins(1 $\rightarrow$ O)-P-Cer
and GSL-C2, β -Galf(1 $\rightarrow$ 2)- α -Manp(1 $\rightarrow$ 3)-[β -Galf(1 $\rightarrow$ 6)]- α -Manp-(1 $\rightarrow$ 2)-Ins(1 $\rightarrow$ O)-P-Cer.

GSL-D fraction.

The chemical shifts and coupling constants obtained for this fraction are summarized in Table VII. The 1D ^1H and 2D $^1\text{H},^{13}\text{C}$ gHSQC spectra exhibited three signals of equivalent areas in the anomeric region (5.108/103.47ppm, 5.069/102.24ppm and 4.939/109.11ppm), indicating the presence of three monosaccharides (Fig. 5). The glycosidic ring spin systems assignment and coupling constants examination permitted the identification of two α -mannose residues and one β -galactofuranose residue as in the GSL-B fraction. Moreover, the NOESY and gHMBC spectra showed the same inter-residue connectivities as those observed for the GSL-B fraction (Fig. 4 and 5), emphasizing the monosaccharide sequence identity with GSL-B. However, differences were observed when comparing $^1\text{H},^{13}\text{C}$ HSQC spectra of the two fractions (Fig. 4 and 5). Thus, a large downfield shift at 68.92ppm and a smaller upfield shift at 72.11ppm were observed for the C6 and C5 carbons respectively of the β -Galf residue for the GSL-D fraction, indicating a 6-O substitution for this residue. Furthermore, in a gHSQC experiment, two new methylene carbons and one trimethyl group have been identified at 4.127/61.62ppm, 3.571/68.79ppm and 3.169/56.48ppm respectively. The COSY experiment showed that these methylene carbons were linked together. The characteristic methyl ^{13}C shift is consistent with the MS data (Fig. 2), indicating the presence of an N-trimethyl group (Fig. 5). In the HMBC spectrum, two $^1\text{H},^{13}\text{C}$ long range interactions were observed between this N-trimethyl group and the second methylene group indicating the presence of the $(\text{CH}_3)_3\text{-N-}$

CH₂-CH₂ motif. Two sets of correlations were observed in the ¹H,³¹P HSQC (Fig. 5). A first set of three correlations was detected between a phosphorus atom (2.974ppm) and the two methylene protons of the ceramide on one side (4.094ppm and 3.660ppm) and the H1 proton of inositol on the other side (3.756ppm), corresponding to the IPC. Another set of two correlations was observed between a second phosphorus atom (0.763ppm) and the two extra methylene protons on the one hand (4.127ppm) and the H6 protons of β-Galf on the other hand (3.764ppm). This was in agreement with the chemical shifts observed for C6 and C5 carbons resonances of the β-Galf residue, indicating its substitution by a phosphocholine residue. In addition, the comparison with the ¹H 1D and ¹H,¹³C HSQC spectra of a phosphatidylcholine reference compound (Sigma) confirmed the resonances assignment of the phosphocholine (not shown). Thus, the NMR analysis, in agreement with methanolysis/trimethylsylation, methylation and MS data, showed that the GSL-D fraction corresponded to Cho-P-(O→6)-β-Galf-(1→2)-α-Manp-(1→3)-α-Manp-(1→2)-Ins(1→O)-P-Cer. The substitution of galactofuranose residue by a phosphate group explains the low amount of galactose detected by GLC (Table 1).

Discussion

The results presented here and in previous studies show that *A. fumigatus* produced at least 9 GIPCs of different structure (Table VIII). All GSLs have the same ceramide moiety composed of a 2-hydroxylated lignoceric acid (2-OH C_{24:0}) associated with a C_{18:0}-phytosphingosine base. This lipid moiety is common to most fungi species. The glycan part has more variability and 2 types of GIPC have been isolated from *A. fumigatus* mycelium. First, a zwitterionic GSL that is the major GIPC from *A. fumigatus* mycelium contains a glucosamine residue linked in α1-2 to the inositol ring. This unusual carbohydrate sequence has been recently described in *A. fumigatus* by Toledo and his colleagues (Toledo *et al.*, 2007)

and has been described in only two other fungal pathogens, *S. schenckii* and *Acremonium* sp. (Toledo *et al.*, 2001a; Aoki *et al.*, 2004a). Secondly, the 5 other acidic GSL structures contain a common sequence α -Man-(1-3)- α -Man-(1-2)-Inositol. This sequence has been previously described in *A. fumigatus* GSL (Leverly *et al.*, 2001; Toledo *et al.*, 2007) and other fungal species (Leverly *et al.*, 1998; 2001; Bennion *et al.*, 2003; Barr *et al.*, 1984). No α -Man-(1-6)-Inositol as found in *S. schenckii* (Toledo *et al.*, 2001a; Loureiro y Penha *et al.*, 2001) and no α -Man-(1-4)-Inositol as found in mushrooms (Jennemann *et al.*, 1999) have been observed in *A. fumigatus* GSLs. Four of the five acidic GSLs analysed contained galactofuranose in *A. fumigatus*. The presence of galactofuranose in GSLs has been reported in human pathogens such as *Histoplasma capsulatum*, *Paracoccidioides brasiliensis*, *A. fumigatus* (Barr *et al.*, 1984; Leverly *et al.*, 1998, Toledo *et al.*, 2007). In these later structures, the galactofuranose residue is linked to the first mannose like for the GSL-C2 of *A. fumigatus*. However, the galactofuranose residue is mainly linked to the terminal non-reduced mannose residue through a β 1-2 linkage (GSL-B). Surprisingly, a choline-phosphate group has been localized to the terminal galactofuranose residue (GSL-D). A choline phosphate in a GSL structure has also been found in *Acremonium* sp., however it is linked to a mannose residue instead of a galactofuranose residue (Aoki *et al.*, 2004a). Some discrepancies are seen between our data and the study of Toledo (Toledo *et al.*, 2007). These authors did not observe the β -galactofuranose linked in β 1-2 to the terminal mannose and the choline-phosphate linked to this β -galactofuranose. In contrast, they described the presence of a mannose residue linked in α 1-2 to Man₂-IPC that we did not observe. These differences are not explained and could be due to the use of different growth conditions or to different strains.

GIPCs have been mainly analysed in human fungal pathogens (*A. niger*, *H. capsulatum*, *P. brasiliensis*, *S. schenckii*, *C. neoformans*), and it has been suggested that GIPCs have an immunological function during fungal infection. The presence of

galactofuranose in these GIPCs that is absent in mammals, seems to play an important role in fungal-host interactions. Earlier studies have shown that galactofuranose containing molecules of *A. fumigatus* are extremely antigenic (Latgé *et al.*, 1994; Costachel *et al.*, 2005; Morelle *et al.*, 2005; Leitao *et al.*, 2003). Galactofuranose residues are also an immunodominant in GSLs of *P. brasiliensis* and *H. capsulatum* (Barr and Lester, 1984; Levery *et al.*, 1998). In *Leishmania major*, a monoclonal antibody against a glycolipid containing terminal galactofuranose residue can reduce the macrophage infectivity of this parasite (Suzuki *et al.*, 2002). The expression in infected human tissues of intelectin that recognizes a single terminal galactofuranose residue (Tsuji *et al.*, 2001) is in agreement with the involvement of galactofuranosylated GSLs during infection. Antibodies from patients with aspergillosis recognized the *A. fumigatus* GIPCs isolated by Toledo *et al.* (2007), however, the role of *A. fumigatus* GIPCs in host cellular immunity has not been defined as yet.

The biosynthetic pathway of sphingolipids begins in the endoplasmic reticulum where it proceeds to the formation of the ceramide moiety. A linkage between the carbohydrate and phytosphingosine occurs in the Golgi apparatus (Funato *et al.*, 2002; Dickson *et al.*, 2006). In yeast, ceramide biosynthesis and inositol addition are essential for growth. Indeed, GIPCs are involved in cell regulation, cell polarity, stress response, trafficking and cell wall integrity (Dickson *et al.* 2006). Similarly, in *Aspergillus*, the gene encoding the IPC synthase is essential for fungal growth as well as *basA* gene-encoding a phytosphingosine (Li *et al.*, 2007). Sphingolipids are involved in polarised growth via the control of actin cytoskeleton (Cheng *et al.*, 2001; Hu *et al.*, 2007). This result is in agreement with the susceptibility of *Aspergillus* species to various inhibitors of sphingolipid synthesis pathway (Zhong *et al.*, 2000; Li *et al.*, 2007). In *A. fumigatus*, 3 types of membrane anchored molecules present the same inositol-phosphoceramide lipid moiety: GPI-anchored protein (Fontaine *et al.*, 2003);

the lipogalactomannan, a GPI-anchored polysaccharide (Costachel *et al.*, 2005) and GIPCs. In this study, the inositol ring of IPC structure could be substituted in position 2 by a mannose or a glucosamine residue. In contrast, previous studies suggested that the glucosamine residue is linked in α 1-6 to the inositol ring in the GPI-anchor structures from GPI-protein or from the lipogalactomannan

In *S. cerevisiae*, 2 homologous genes of IPC mannosyltransferases are involved in the addition of the first mannose residue to the inositol ring. The deletion of both genes is not lethal, but mutants are sensitive to the external Ca^{2+} (Beeler *et al.*, 1997), suggesting a role for the GSLs in cellular stress response. In *C. albicans*, the deletion of the MIT1, an IPC mannosyltransferase homologue induced the absence of MIPC, M(IP)2C and the phospholipomannan (β 1-2mannan linked to MIPC) and decreased virulence of the mutant (Trinel *et al.*, 2002; Mille *et al.*, 2004). In *A. fumigatus*, no such mutant has been described, so the relevance of these GIPCs during *A. fumigatus* growth and host-pathogen interactions is still unknown.

Materials and methods

Fungal culture and membrane preparation

A. fumigatus, strain CBS 144-89 was grown in a 15-L fermenter in 2% glucose and 1% mycopeptone (Biokar Diagnostics, Pantin, France) for 24 h at 25°C as described previously (Hartland *et al.* 1996). The mycelium was collected by filtration under vacuum, washed with water, and then disrupted in 200 mM TrisHCl, 20 mM EDTA, pH 8.0, 1 mM PMSF buffer at 4°C with glass beads (1 mm, diameter) in a Dyno mill apparatus (W. A. Bachofen AG, Basel, Switzerland). The cell wall was removed by centrifugation at 10,000 g 10 min at 4°C. Total membranes were then collected by ultracentrifugation at 125,000 g 60 min at 4°C. Membrane pellet was homogenized in the disruption buffer with a Dounce homogenizer and then

centrifuged once more at 125,000 g for 60 min at 4°C. Pellet was resuspended again in 20 mM TrisHCl, 2 mM EDTA, pH 8.0 and store at -80°C.

Extraction and purification of glycosylinositolphosphatidylceramides

A chloroform/methanol mixture was added to the membrane suspension (20 mg of protein/ml) to obtain a chloroform/methanol/membrane ratio of 10/10/3 respectively. The mixture was stirred for 2 h at room temperature, then centrifuged at 10,000 g for 10 min. The pellet was resuspended in a chloroform/methanol/water (10/10/3, v/v/v) mixture, and the extraction was repeated once. Pooled supernatants were concentrated under vacuum with a rotavapor, and the residue was submitted to a butanol/water partition. The water phase containing GIPCs was dialysed against water and freeze-dried. The residue was dissolved in chloroform/methanol/water (10/15/4, v/v/v) and applied onto a DEAE-Sephadex A-25 column (Pharmacia, 2 x 15 cm) equilibrated in the same solvent at the flow rate of 30 ml/h. Unbound products were eluted with 2 column volumes of solvent, then retained products were eluted by a chloroform/methanol/NH₄Ac 1M (10/15/4) solvent. Carbohydrates were detected by spraying with orcinol sulfuric acid on spot of 2 µl of different fractions on silica sheets. Fractions containing sugars were concentrated and then dialysed against water and freeze-dried. GIPCs were then purified on a silica 60 column (Merck, 1.8 x 30 cm) equilibrated in propanol-1/water/NH₄OH 30% (85/15/5, v/v/v) and eluted at 25ml/h. Samples were deposited onto the silica 60 column and eluted by 3 column volumes of propanol-1/water/NH₄OH 30% (85/15/5, v/v/v), then 3 volumes of propanol-1/water/NH₄OH 30% (80/20/5, v/v/v), then 3 volumes of propanol-1/water/NH₄OH 30% (70/30/5, v/v/v). The presence of carbohydrate in the fraction was detected as described above. Fractions were dialysed against water and freeze-dried.

Analytical methods.

Neutral hexoses were identified by GLC as alditol acetates obtained after hydrolysis (4 N trifluoroacetic acid, 100°C, 4h.)(Sawardeker *et al.*, 1967). Glucosamine and *myo*-inositol were quantified by GLC-MS after hydrolysis (6 N HCL, 110°C 20h), N-acetylation and trimethylsilylation, using the *scyllo*-inositol as a standard (Ferguson, 1993). Lipid analysis was performed by GLC-MS on HF-treated GIPC fraction (aqueous 50% HF, 2 days on ice). Fatty acids and sphingosine bases were released by methanolysis (1 N HCl in MeOH, 80°C, 20h). Fatty acids were extracted with heptane and analysed after trimethylsilylation. The methanol phase containing the sphingosine base was N-acetylated, trimethylsilylated and analysed by GLC-MS. Phosphorylated carbohydrates were identified by GLC-MS after acid methanolysis (1 N HCL in methanol, 80°C, 20h) and trimethylsilylation (Ferguson, 1993). The lipophosphoglycan from *Leishmania donovani*, a kind gift from Pascale Pescher (Unité de Virulence Parasitaire, Institut Pasteur) was used as positive control. Methylation of GSL fractions was performed using the sodium hydroxide procedure (Ciucanu and Kerek, 1984). GSL containing a glucosamine residue was peracetylated with a pyridine/acetic anhydride solution (50/200 µl) overnight at room temperature prior to the methylation procedure. Methyl ethers were analysed by GLC-MS as polyolacetates (Björndal *et al.*, 1970).

HPTLC. GIPC fractions were applied to a 10 cm aluminium-backed silica gel 60 (Merck) and developed with chloroform/methanol/1 M ammonium acetate/NH₄OH 30%/water (180/140/9/9/23). Sugars were detected with orcinol-sulfuric acid.

GLC and GLC-Mass spectrometry. GLC was performed on a Delsi 200 instrument with a flame ionisation detector using a capillary column (30 m x 0.25 mm id) filled with a ECTM-1 (Alltech) under the following conditions: gas vector and pressure, helium 0.7 bar; temperature

program 120 to 180°C at 2°C/min and 180 to 240°C at 4°C/min. GLC-MS was performed on an Automass II apparatus (Finigan) coupled to a CarloErba gas chromatograph (model 8000top), using a capillary column (30 m x 0.25 mm id) filled with a ECTM-1 (Alltech) under the following conditions : gas vector and flow rate, helium 1.2 ml/min; temperature program for inositol and monosaccharide analysis : 100 to 200°C at 5°C/min, 200 to 240°C at 15°C/min and 240°C for 5 min; temperature program for sphingosine base and fatty acid analysis : 100 to 200°C at 10°C/min, 200 to 260°C at 15°C/min and 260°C for 13 min.

Nano-electrospray mass spectrometry. Mass spectrometric analyses were performed in the negative mode using a Q-STAR Pulsar quadrupole time-of-flight (Q-TOF) mass spectrometer (AB/MDS Sciex, Toronto, Canada) equipped with a nano-electrospray ion source (Protana, Odense, Denmark).

The samples in propanol-1/H₂O (25/75) dissolved in chloroform/methanol (50/50) were sprayed from gold-coated 'medium length' borosilicate capillaries (Protana, Odense, Denmark). A potential of -800 V was applied to the capillary tip. The declustering potential was set at -120V and the focusing potential was set at -200V. The molecular ions were then selected in the quadrupole analyzer and partially fragmented in the hexapole collision cell, with the pressure of collision gas (N₂) 5.3×10^{-5} Torr (1Torr = 133.3 Pa). The collision energy was varied between -40 and -80 eV depending on the sample. For the recording of conventional mass spectra, TOF data were acquired by accumulation of 10 MCA (multiple channel acquisition) scans over mass ranges of m/z ranges 500-1800 Daltons for MS analyses and over mass ranges of m/z 50-1800 for MS/MS analyses. Data acquisition was optimized to supply the highest possible resolution and the best signal-to-noise ratio, even in the case of low abundance signals. Typically, the FWHM (full width at half maximum) was 7000 in the measured mass ranges. External calibration was performed prior to each measure using a

4 pmol/ μ l solution of taurocholic acid in acetonitrile/water (50/50, v/v) containing 2 mM of ammonium acetate.

NMR Spectroscopy. NMR spectra were acquired at 50°C on a Varian Inova 500 spectrometer equipped with a triple $^1\text{H}\{^{13}\text{C}/^{15}\text{N}\}$ resonance ^1H PFG probe or an indirect PFG probe for ^1H , ^{31}P experiments. For the low concentration samples, complementary experiments were performed at 35°C on a Varian Inova 600 spectrometer equipped with a cryogenically-cooled triple resonance $^1\text{H}\{^{13}\text{C}/^{15}\text{N}\}$ PFG probe. Samples were dissolved in DMSO- d_6 for NMR (99.96% ^2H atoms, Euriso-top, CEA, Saclay, France) and transferred in 5mm Shigemi tubes (Shigemi Inc., Alison Park, United States). D_2O (99.97% ^2H atoms, Euriso-top, CEA, Saclay, France) was added in order to exchange sugars hydroxyl protons. Since the proton chemical shift of the residual signal of DMSO- d_6 depends on temperature and water content, external referencing was applied for ^1H chemical shifts using a capillary containing a freshly prepared solution of 20 mM DSS in DMSO- d_6 containing less than 0,01% of water. The DSS methyl resonance was set to 0ppm. ^{13}C chemical shifts were then calculated from ^1H chemical shift and gamma ratio relative to DSS. $^{13}\text{C}/^1\text{H}$ gamma ratio of 0.251449530 was used (Wishart *et al.*, 1995). ^{31}P chemical shifts were determined with neat phosphoric acid (Wilmad-Labglas, New Jersey, USA) by substitution method.

For all GSL fractions, the same general strategy was adopted for assignment of nuclei. First, the proton resonances were assigned using two-dimensional COSY and RELAY experiments with one to three relays to follow connectivities from the anomeric proton up to the H5 proton of most of glycosidic residues (Rance *et al.*, 1983; Wagner 1983). The intra glycosidic residue spin systems were often completed by mean of a TOCSY experiment with a long mixing time (120ms) (Griesinger *et al.*, 1988). A ^1H - ^{13}C edited gHSQC experiment allowed to achieve ^{13}C chemical shifts assignment from previously identified ^1H resonances (Willker *et al.*, 1993).

Then, ^1H , ^1H coupling constants analysis from 1D and/or COSY spectrum (^1H resolution of 0.1 Hz and 1.6Hz respectively) was used to assess the identity of monosaccharide residues. Moreover, the anomeric configuration of monosaccharide residues was established from knowledge of $^3J_{1,2}$ values and confirmed by the measurement of the $^1J_{\text{C1H1}}$ heteronuclear coupling constants in the ^1H dimension of the uncoupled gHSQC spectrum (^1H resolution of 0.6Hz) or of the gHMBC spectrum (^1H resolution of 1.2Hz) (Willker *et al.*, 1993). Finally, glycosidic linkages were established via through-space dipolar interactions using ^1H - ^1H NOESY experiment (mixing time of 200ms) (Macura *et al.*, 1981) and/or via three-bond interglycosidic ^1H - ^{13}C correlations using ^1H - ^{13}C gHMBC experiment (long range delay of 60ms). In addition, the branching point between the phospholipid and the glycosidic moieties was identified using the ^1H - ^{31}P gHSQC.

Acknowledgments

We thank Professor R. Calderone (Georgetown University Medical center, Washington, USA) for reading of the manuscript.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abbreviations

- COSY: Correlation spectroscopy
- DMSO: Dimethylsulfoxide
- DSS: 2,2-dimethyl-2-silapentane- 5-sulfonate sodium salt
- GIPC: Glycosylinositolphosphoceramide
- GLC: Gas-liquid chromatography
- GLC-MS: Gas-liquid chromatography-mass spectrometry
- GPI: Glycosylphosphatidylinositol
- GSL: Glycosphingolipid
- H2BC: Heteronuclear two-bond correlation
- gHMBC: Gradient selected heteronuclear multiple bond correlation
- gHSQC: Gradient selected heteronuclear single-quantum correlation
- HPTLC: High performance thin layer chromatography
- IPC: Inositolphosphoceramide
- MS: Mass spectrometry
- NMR: Nuclear magnetic resonance
- NOESY: Nuclear overhauser effect spectroscopy
- Q-TOF: Pulsar quadrupole-time of flight
- RELAY 2D: Relayed COSY
- PFG: Pulsed field gradient
- TOCSY: Total correlation spectroscopy

References

- Aoki K, Uchiyama R, Itonori S, Sugita M, Che FS, Isogai A, Hada N, Hada J, Takeda T, Kumagai H, Yamamoto K. 2004a. Structural elucidation of novel phosphocholine-containing glycosylinositol-phosphoceramides in filamentous fungi and their induction of cell death of cultured rice cells. *Biochem J.* 378:461-472.
- Aoki K, Uchiyama R, Yamauchi S, Katayama T, Itomori S, Sugita M, Hada N, Yamada-Hada J, Takeda T, Kumagai H, Yamamoto K. 2004b. Newly discovered neutral glycosphingolipids in aureobasidin A-resistant *Zygomycetes*. *J. Biol. Chem.* 279:32028-32034.
- Arigi E, Singh S, Kahlili AH, Winter H. C, Goldstein IJ, and Lavery SB. 2007. Characterization of neutral and acidic glycosphingolipids from the lectin-producing mushroom, *Polyporus squamosus*. *Glycobiology* 17:754-766.
- Barr K, Lester RL. 1984. Occurrence of novel antigenic phosphoinositol-containing sphingolipids in the pathogenic yeast *Histoplasma capsulatum*. *Biochemistry.* 23:5581-5588.
- Beeler TJ, Fu D, Rivera J, Monaghan E, Gable K, Dunn TM. 1997. *SUR1 (CSG1/BCL21)*, a gene necessary for growth of *Saccharomyces cerevisiae* in the presence of high Ca^{2+} concentrations at 37 degrees C, is required for mannosylation of inositolphosphorylceramide. *Mol Gen Genet.* 255:570-579.
- Bennion B, Park C, Fuller M, Lindsey R, Momany M, Jennemann R, Lavery SB. 2003. Glycosphingolipids of the model fungus *Aspergillus nidulans*: characterization of GIPCs with oligo- α -mannose-type glycans. *J Lipid Res.* 44:2073-2088.

- Björndal H, Hellerqvist CG, Lindberg B, Svensson S. 1970. Gas-liquid chromatography and mass spectrometry in methylation analysis of polysaccharides. *Angew. Chem. Int.* 9:610-619.
- Bock K, Pedersen C. 1974. A study of ^{13}C coupling constants in hexopyranoses. *J Chem Soc Perkin Trans. II*:293-297.
- Bock K, Pedersen C. 1983. Carbon-13 nuclear magnetic resonance spectroscopy of monosaccharides. *Adv. Carbohydr. Chem. Biochem.* 41:27-44.
- Bunel S, Ibarra C, Moraga E, Blasko A, Bunton CA. 1993. Structures of complexes of cobalt(III) and glucosamine. Applications of molecular mechanics and NMR spectroscopy. *Carbohydr. Res.* 244:1-14.
- Cheng J, Park TS, Fischl AS, Ye XS. 2001. Cell cycle progression and cell polarity require sphingolipid biosynthesis in *Aspergillus nidulans*. *Mol Cell Biol.* 21:6198-6209.
- Ciucanu I, Kerek F. 1984. A simple and rapid method for the permethylation of carbohydrates. *Carbohydr. Res.* 131:209-217.
- Costachel C, Coddeville B, Latgé JP, Fontaine T. 2005. Glycosylphosphatidylinositol-anchored fungal polysaccharide in *Aspergillus fumigatus*. *J Biol Chem.* 280:39835-39842.
- da Silva AF, Rodrigues ML, Farias SE, Almeida IC, Pinto MR, Barreto-Bergter E. 2004. Glucosylceramides in *Colletotrichum gloeosporioides* are involved in the differentiation of conidia into mycelial cells. *FEBS Lett.* 561:137-143.
- Dickson RC, Sumanasekera C, Lester RL. 2006. Functions and metabolism of sphingolipids in *Saccharomyces cerevisiae*. *Prog Lipid Res.* 45:447-465.

- 1
2
3 Ferguson MA. 1993. GPI-membrane anchors: isolation and analysis in Glycobiology, a
4 practical approach. ed. M Fukudo, A Kobata, pp. 349-383: Oxford University, IRL
5 Press
6
7
8
9
10
11 Fontaine T, Magnin T, Melhert A, Lamont D, Latgé JP, Ferguson MA. 2003. Structures of the
12 glycosylphosphatidylinositol membrane anchors from *Aspergillus fumigatus* membrane
13 proteins. Glycobiology. 13:169-177.
14
15
16
17
18
19 Funato K, Vallee B, Riezman H. 2002. Biosynthesis and trafficking of sphingolipids in the
20 yeast *Saccharomyces cerevisiae*. Biochemistry. 41:15105-15114.
21
22
23
24
25 Griesinger C, Otting G, Wüthrich K, Ernst RR. 1988. Clean TOCSY for proton spin system
26 identification in macromolecules. J. Am. Chem. Soc. 110:7870-7872.
27
28
29
30
31 Gutierrez A. LS, Farage L, Melo M. N, Mohana-Borges R. S, Guerardel Y, Coddeville B,
32 Wieruszeski J-M, Mendonça-Previato L, Previato J. O. 2007. Characterization of
33 glycoinositolphosphoryl ceramide structure mutant strains of *Cryptococcus neoformans*.
34 Glycobiology 17:1C-11C.
35
36
37
38
39
40
41 Hartland RP, Fontaine T, Debeaupuis JP, Simenel C, Delepierre M, Latgé JP. 1996. A novel
42 beta-(1-3)-glucanosyltransferase from the cell wall of *Aspergillus fumigatus*. J Biol
43 Chem. 271:26843-26849.
44
45
46
47
48
49 Heise N, Gutierrez AL, Mattos KA, Jones C, Wait R, Previato JO, Mendonca-Previato L.
50 2002. Molecular analysis of a novel family of complex glycoinositolphosphoryl
51 ceramides from *Cryptococcus neoformans*: structural differences between encapsulated
52 and acapsular yeast forms. Glycobiology. 12:409-420.
53
54
55
56
57
58
59
60

- Hu W, Sillaots S, Lemieux S, Davison J, Kauffman S, Breton A, Linteau A, Xin C, Bowman J, Becker J, Jiang B, Roemer T. 2007. Essential Gene Identification and Drug Target Prioritization in *Aspergillus fumigatus*. PLoS Pathog. 3:e24.
- Jennemann R, Bauer BL, Bertalanffy H, Geyer R, Gschwind RM, Selmer T, Wiegandt H. 1999. Novel glycoinositolphosphosphingolipids, basidiolipids, from *Agaricus*. Eur J Biochem. 259:331-338.
- Jennemann R, Geyer R, Sandhoff R, Gschwind RM, Lavery SB, Grone HJ, Wiegandt H. 2001. Glycoinositolphosphosphingolipids (basidiolipids) of higher mushrooms. Eur J Biochem. 268:1190-1205.
- Latgé JP. 1999. *Aspergillus fumigatus* and aspergillosis. Clin Microbiol Rev. 12:310-350.
- Latgé JP, Kobayashi H, Debeaupuis JP, Diaquin M, Sarfati J, Wieruszeski JM, Parra E, Bouchara JP, Fournet B. 1994. Chemical and immunological characterization of the extracellular galactomannan of *Aspergillus fumigatus*. Infect Immun. 62:5424-5433.
- Leitao EA, Bittencourt VC, Haido RM, Valente AP, Peter-Katalinic J, Letzel M, de Souza LM, Barreto-Bergter E. 2003. Beta-galactofuranose-containing O-linked oligosaccharides present in the cell wall peptidogalactomannan of *Aspergillus fumigatus* contain immunodominant epitopes. Glycobiology. 13:681-692.
- Lavery SB, Momany M, Lindsey R, Toledo MS, Shayman JA, Fuller M, Brooks K, Doong RL, Straus AH, Takahashi HK. 2002. Disruption of the glucosylceramide biosynthetic pathway in *Aspergillus nidulans* and *Aspergillus fumigatus* by inhibitors of UDP-Glc:ceramide glucosyltransferase strongly affects spore germination, cell cycle, and hyphal growth. FEBS Lett. 525:59-64.

- Leverly SB, Toledo MS, Straus AH, Takahashi HK. 1998. Structure elucidation of sphingolipids from the mycopathogen *Paracoccidioides brasiliensis*: an immunodominant beta-galactofuranose residue is carried by a novel glycosylinositol phosphorylceramide antigen. *Biochemistry*. 37:8764-8775.
- Leverly SB, Toledo MS, Straus AH, Takahashi HK. 2001. Comparative analysis of glycosylinositol phosphorylceramides from fungi by electrospray tandem mass spectrometry with low-energy collision-induced dissociation of Li(+) adduct ions. *Rapid Commun Mass Spectrom*. 15:2240-2258.
- Loureiro y Penha CV, Todeschini AR, Lopes-Bezerra LM, Wait R, Jones C, Mattos KA, Heise N, Mendonca-Previato L, Previato JO. 2001. Characterization of novel structures of mannosylinositolphosphorylceramides from the yeast forms of *Sporothrix schenckii*. *Eur J Biochem*. 268:4243-4250.
- Li S, Bao D, Yuen G, Harris SD, Calvo AM. 2007. *basA* regulates cell wall organization and asexual/sexual sporulation ratio in *Aspergillus nidulans*. *Genetics* 176:243-253.
- Maciel DM, Rodrigues ML, Wait R, Villas Boas MHS, Tischer CA, Barreto-bergter E. 2002. Glycosphingolipids from *Magnaporthe grisea* cells: expression of a ceramide dihexoside presenting phytosphingosine as the long-chain base. *Arch. Biochem. Biophys*. 405:205-213.
- Macura S, Huang Y, Suter D, Ernst RR. 1981. Two-dimensional chemical exchange and cross-relaxation spectroscopy of coupled nuclear spins. *J Magn Reson*. 43:259-281.
- Mille C, Janbon G, Delplace F, Ibata-Ombetta S, Gaillardin C, Strecker G, Jouault T, Trinel PA, Poulain D. 2004. Inactivation of CaMIT1 inhibits *Candida albicans*

- phospholipomannan beta-mannosylation, reduces virulence, and alters cell wall protein beta-mannosylation. *J Biol Chem.* 279:47952-47960.
- Morelle W, Bernard M, Debeaupuis JP, Buitrago M, Tabouret M, Latgé JP. 2005. Galactomannoproteins of *Aspergillus fumigatus*. *Eukaryot Cell.* 4:1308-1316.
- Nagiec MM, Nagiec EE, Baltisberger JA, Wells GB, Lester RL, Dickson RC. 1997. Sphingolipid synthesis as a target for antifungal drugs. Complementation of the inositol phosphorylceramide synthase defect in a mutant strain of *Saccharomyces cerevisiae* by the AUR1 gene. *J Biol Chem.* 272:9809-9817.
- Petersen BO, Vinogradov E, Kay W, Würtz P, Nyberg NT, Duus JO, Sorensen OW. 2006. H2BC: a new technique for NMR analysis of complex carbohydrates. *Carbohydr Res.* 341:550-556.
- Rance M, Sorensen OW, Bodenhausen G, Wagner G, Ernst RR, Wüthrich K. 1983. Improved spectral resolution in COSY 1H NMR spectra of proteins via double quantum filtering. *Biochem Biophys Res Commun.* 117:479-485.
- Ritchie RG, Cyr N, Korsch B, Koch HJ, Perlin A. 1975. Carbon-13 chemical shifts of furanosides and cyclopentanols. Configurational and conformational influences. *Can. J. Chem.* 53:1424-1433.
- Rittershaus P, Kechichian TB, Allegood JC, Merrill Jr AH, Hennig M, Luberto C, Del Poeta M. 2006. Glucosylceramide synthase in an essential regulator of pathogenicity of *Cryptococcus neoformans*. *J. Clin. Investi.* 116:1651-1659.
- Sawardeker JS, Sloneker JH, Jeanes A. 1967. Quantitative determination of monosaccharides as their alditol acetates by gas liquid chromatography. *Anal Chem.* 37:1602-1604.

- 1
2
3 Suzuki E, Tanaka AK, Toledo MS, Takahashi HK, Straus AH. 2002. Role of beta-D-
4 galactofuranose in *Leishmania major* macrophage invasion. Infect Immun. 70:6592-
5 6596.
6
7
8
9
10
11 Toledo MS, Lavery SB, Bennion B, Guimaraes LL, Castle SA, Lindsey R, Momany M, Park
12 C, Straus AH, Takahashi HK. 2007. Sphingolipids of the mycopathogen *Aspergillus*
13 *fumigatus*: characterization of glycosylinositol phosphorylceramide antigens with
14 Manp(alpha 1-2)Ins and GlcpN(alpha 1-2)Ins core motifs. J. Lipid Res. 48:1801-1824.
15
16
17
18
19
20
21
22 Toledo MS, Lavery SB, Glushka J, Straus AH, Takahashi HK. 2001a. Structure elucidation of
23 sphingolipids from the mycopathogen *Sporothrix schenckii*: identification of novel
24 glycosylinositol phosphorylceramides with core manalpha1-->6Ins linkage. Biochem
25 Biophys Res Commun. 280:19-24.
26
27
28
29
30
31
32
33 Toledo MS, Lavery SB, Straus AH, Takahashi HK. 2001b. Sphingolipids of the
34 mycopathogen *Sporothrix schenckii*: identification of a glycosylinositol
35 phosphorylceramide with novel core GlcNH(2)alpha1-->2Ins motif. FEBS Lett. 493:50-
36 56.
37
38
39
40
41
42
43
44 Trinel PA, Maes E, Zanetta JP, Delplace F, Coddeville B, Jouault T, Strecker G, Poulain D.
45 2002. *Candida albicans* phospholipomannan, a new member of the fungal mannose
46 inositol phosphoceramide family. J Biol Chem. 277:37260-37271.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
- Tsuiji S, Uehori J, Matsumoto M, Suzuki Y, Matsuhisa A, Toyoshima K, Seya T. 2001. Human intelectin is a novel soluble lectin that recognizes galactofuranose in carbohydrate chains of bacterial cell wall. J Biol Chem. 276:23456-23463.
- Wagner G. 1983. Two-dimensional relayed coherence transfer spectroscopy of a protein. J. Magn Reson. 55:151-156.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Willker W, Leibfritz D, Kerssebaum R, Bermel W. 1993. Gradient selection in inverse heteronuclear correlation spectroscopy. *Magn Reson Chem.* 31:287-292.

Wishart DS, Bigam CG, Yao J, Abildgaard F, Dyson HJ, Oldfield E, Markley JL, Sykes BD. 1995. ¹H, ¹³C and ¹⁵N chemical shift referencing in biomolecular NMR. *J Biomol NMR.* 6:135-140.

Zhong W, Jeffries MW, Georgopapadakou NH. 2000. Inhibition of inositol phosphorylceramide synthase by aureobasidin A in *Candida* and *Aspergillus* species. *Antimicrob Agents Chemother.* 44:651-653.

Legends to figures

Figure 1: HPTLC of GSL fractions (U, A, B, C, D) isolated from *A. fumigatus* mycelium. TLC was developed on 10-cm aluminium-coated silica gel 60 with chloroform/methanol/1 M ammonium acetate/ NH_4OH 30%/water (180/140/9/9/23). Sugars were detected using orcinol sulfuric acid.

Figure 2: Nano-electrospray mass spectrometry analysis of GSL-B and GSL-D fractions of *A. fumigatus* mycelium. Mass spectrometric analyses were performed in negative mode using a Q-STAR Pulsar quadrupole time-of-flight (Q-TOF) mass spectrometer equipped with a nano-electrospray ion source. A: negative ion mass spectrum; B: daughter ion mass spectrum of molecular ion.

Figure 3: NMR spectra of GSL-U isolated from *A. fumigatus* mycelium. ^1H - ^{13}C HSQC, ^1H - ^1H NOESY, ^1H - ^{13}C HMBC and ^1H - ^{31}P HSQC.

Figure 4: NMR spectra of GSL-B from *A. fumigatus* mycelium. ^1H - ^{13}C HSQC, ^1H - ^1H NOESY, ^1H - ^{13}C HMBC and ^1H - ^{31}P HSQC.

Figure 5: NMR spectra of GSL-D from *A. fumigatus* mycelium. ^1H - ^{13}C HSQC, ^1H - ^1H NOESY, ^1H - ^{13}C HMBC and ^1H - ^{31}P HSQC.

Table I. Carbohydrate composition and Q-TOF mass analysis of GSL fractions isolated from *A. fumigatus* mycelium
(Man : mannose; Gal : galactose; GlcN : glucosamine; IPC : inositolphosphoceramide; Hex : hexose; Cho-P : choline-phosphate)

GSL Fractions	Monosaccharide composition *				Pseudomolecular ions identified by Q-TOF		
	Man	Gal	GlcN	Inositol	Main ions	Corresponding structures	Minor ions
GSL-U	1	-	0.49	+	1409.8	Hex ₂ -GlcN-IPC	1424.1 ; 1437.8
GSL-A	1	0.36	-	+	1249.1; 1411.2	Hex ₂ -IPC ; Hex ₃ -IPC	1263.1 ; 1277.1 1425.2 ; 1439.2
GSL-B	1	0.49	-	+	1411.2	Hex ₃ -IPC	1425.2 ; 1439.2
GSL-C	1	0.42	-	+	1573.3	Hex ₄ -IPC	1587.3 ; 1601.3
GSL-D	1	0.11	-	+	1576.3	Cho-P-Hex ₃ -IPC	1590.3 ; 1604.4

*: obtained by GC after hydrolysis with TFA (4N, 100°C, 4h) for neutral monosaccharides and HCl (6N, 110°C 20h) for glucosamine and inositol residues.

Table II : Molar ratio of ethyl ethers obtained after permethylation of GIPC fractions isolated from *A. fumigatus* mycelium. Molar ratios were calculated by GLC-flame ionisation detection after hydrolysis (TFA, 4N, 6h, 100°C), reduction with NaBD₄, and acetylation).

Methyl ethers ^a	Linkages	Glycosphingolipid fractions				
		GSL-U	GSL-A	GSL-B	GSL-C	GSL-D
2,3,4,6-Man	Man-	1.3	1.1	0.1	0.7	-
2,3,5,6-Gal	Gal-	-	0.8	0.7	1.1	-
3,4,6-Man	-2-Man-	-	1	1	1	1
2,4,6-Man	-3-Man-	1	1.6	0.7	-	1.1
2,4-Man	-3- ⁶ Man-	-	-	-	0.8	-
2,3,5-Gal	-6-Gal-	-	-	-	-	+ ^b
2,3,4-GlcNAc	-6-GlcN-	0.4	-	-	-	-

^a : numbers indicate the position of methyl groups

^b : this methyl ether was observed after hydrolysis by HCl 4N, 4h, 100°C

Table III. ¹H and ¹³C NMR chemical shifts (ppm) and coupling constants ($J_{H,H}$ and $^1J_{C1H1}$, Hz) for the glycan sequence of the GSL-U fraction.

α -Manp-(1-3)- α -Manp-(1-6)- α -GlcNH2-(1-2)-Ins-(1-O)-P-Cer

	H ₁	H ₂	H ₃	H ₄	H ₅	H ₆ -H _{6'}
	$^3J_{1,2}$	$^3J_{2,3}$	$^3J_{3,4}$	$^3J_{4,5}$	$^3J_{5,6}$ - $^3J_{5,6'}$	$^2J_{6,6'}$
	C ₁	C ₂	C ₃	C ₄	C ₅	C ₆
	$^1J_{C1H1}$					
α -Man-(1-3)-	4.925	3.781	3.606	3.440	3.468	3.524-3.695
	1.7	≈ 3	≈ 10	≈ 10	4.9	10.9
	105.26	73.56	74.07	70.57	76.66	64.41
	169.6					
-3)- α -Man-(1-6)-	4.666	3.932	3.656	3.576	3.468	3.518-3.665
	1.8	3.1	9.1	9.7	nm -5.0	12.00
	103.51	72.54	82.24	69.06	76.80	64.41
	167.6					
-6)- α -GlcNH2-(1-2)-	5.222	2.923	3.652	3.320	3.994	3.594-3.799
	3.5	10.2	10.0	10.8	≈ 0 -5.2	11.0
	101.32	57.91	73.83	72.97	74.30	68.29
	nm					
-2)-myo-Ins-(1-O)-P	3.825	4.168	3.278	3.371	3.048	3.776
	2.7	2.7	10.0	8.5	$^3J_{5,6}=8.5$	$^3J_{6,1}=8.8$
	79.86	85.22	73.38	75.18	78.03	75.11
	$J_{H,P}=10.4$					

nm : not measured

Table IV. ^1H and ^{13}C NMR chemical shifts (ppm) and coupling constants ($J_{\text{H,H}}$, $^1J_{\text{C,H}}$ and $J_{\text{H,P}}$ Hz) for the glycan sequence of the GSL-B fraction.

$\beta\text{-Galf-(1-2)-}\alpha\text{-Manp-(1-3)-}\alpha\text{-Manp-(1-2)-Ins-(1-O)-P-Cer}$

	H ₁	H ₂	H ₃	H ₄	H ₅	H ₆ -H _{6'}
	$^3J_{1,2}$	$^3J_{2,3}$	$^3J_{3,4}$	$^3J_{4,5}$	$^3J_{5,6}$ - $^3J_{5,6'}$	$^2J_{6,6'}$
	C ₁	C ₂	C ₃	C ₄	C ₅	C ₆
	$^1J_{\text{C,H}}$					
$\beta\text{-Galf-(1-2)-}$	4.881	3.788	3.808	3.936	3.525	3.377-3.401
	2.1	3.2	5.2	3.2	6.7-6.0	10.9
	108.70	83.78	80.10	87.25	73.90	65.85
	173.6					
$\text{-2)-}\alpha\text{-Man-(1-3)-}$	5.020	3.917	3.607	3.332	3.584	3.508-3.578
	1.5	3.8	8.9	9.4	5.0-5.0	11.2
	101.71	77.08	72.86	71.05	76.30	64.00
	169.1					
$\text{-3)-}\alpha\text{-Man-(1-2)-}$	5.029	3.927	3.659	3.608	3.867	3.488-3.561
	1.5	3.3	9.4	10.2	5.8- 3.5	11.5
	103.85	72.60	81.70	68.81	76.16	64.13
	173.1					
$\text{-2)-myo-Ins-(1-O)-P}$	3.727	3.977	3.216	3.351	2.948	3.485
	2.4	2.6	11.3	8.9	$^3J_{5,6}=8.9$	$^3J_{1,6}=9.4$
	78.99	80.91	73.63	75.29	78.56	75.49
	139.3					
	$J_{\text{H},\text{P}}=9.2$					

Table V. ¹H and ¹³C NMR chemical shifts (ppm) and coupling constants (J_{H,H} and ¹J_{C1H1}, Hz) for the two glycan sequences of the GSL-A fraction.

GSL-A1: α-Manp-(1-3)-α-Manp-(1-2)-Ins(1-O)-P-Cer

GSL-A2: β-Galf- (1-2)-α-Manp-(1-3)-α-Manp-(1-2)-Ins(1-O)-P-Cer

	H ₁ ³ J _{1,2} C ₁ ¹ J _{C1H1}	H ₂ ³ J _{2,3} C ₂	H ₃ ³ J _{3,4} C ₃	H ₄ ³ J _{4,5} C ₄	H ₅ ³ J _{5,6} - ³ J _{5,6'} C ₅	H ₆ -H _{6'} C ₆
GSL-A1						
α-Manp-(1-3)-	4.905	3.733	3.586	3.446	3.605	3.51-3.58
	2.2	4.2	9.4	8.8	3.595*	3.53*-3.60*
	104.67	73.47	74.00	70.55	76.38	64.14
	170.0				76.42*	64.05*
-3)-α-Manp-(1-2)-	5.044	3.932	3.641	3.638	3.889	3.51-3.58
	1.7	4.5		10.2	6.6	
	103.88	72.63	81.5	68.85	76.18	64.14
	173.4					
-2)-myo-Ins-(1-O)-P	3.738	3.977	3.227	3.369	2.953	3.501
	3.5	3.4	9.8	9.6	9.6	³ J _{6,1} ≈ 9.5
	78.94	80.91	73.66	75.24	78.68	75.50
GSL-A2						
β-Galf- (1-2)-	4.901	3.808	3.821	3.956	3.540	3.401-3.422
	1.8	4.6	4.0	4.5		
	108.72	83.79	80.06	87.27	73.90	65.84
	173.8					
-2)-α-Manp-(1-3)-	5.035	3.935	3.624	3.364	3.605	3.53-3.60
	1.6	4.0	9.4	9.5	3.595*	
	101.66	77.09	72.86	71.13	76.38	64.05
	169.2				76.42*	
-3)-α-Manp-(1-2)-	5.044	3.951	3.665	3.638	3.889	3.51-3.58
	1.7	4.5	10.2	10.2	6.6	
	103.88	72.59	81.85	68.85	76.18	64.14
	173.4					
-2)-myo-Ins-(1-O)-P	3.738	3.977	3.227	3.369	2.953	3.501
	3.5	3.4	9.8	9.6	9.6	³ J _{6,1} ≈ 9.5
	78.94	80.91	73.66	75.24	78.68	75.50

* possible chemical shift

Table VI. ^1H and ^{13}C NMR chemical shifts (ppm) and coupling constants ($J_{\text{H,H}}$ and $^1J_{\text{C,H}}$, Hz) for the two glycan sequences of the GSL-C fraction.

GSL-C1: β -Galf(1-2)- α -Manp(1-3)-[α -Manp (1-6)]- α -Manp-(1-2)- Ins(1-O)-P-Cer

GSL-C2: β -Galf(1-2)- α -Manp(1-3)-[β -Galf (1-6)]- α -Manp-(1-2)- Ins(1-O)-P-Cer

	H ₁ $^3J_{1,2}$ C ₁ $^1J_{\text{C,H}}$	H ₂ $^3J_{2,3}$ C ₂	H ₃ $^3J_{3,4}$ C ₃	H ₄ $^3J_{4,5}$ C ₄	H ₅ $^3J_{5,6}$ - $^3J_{5,6'}$ C ₅	H ₆ -H _{6'} C ₆
GSL-C1						
β -Galf- (1-2)-	4.877 1.1 108.47 174.7	3.777 2.9 83.60	3.800 79.93	3.933 87.07	3.518 73.72	3.360-3.392 65.72
-2)- α -Manp-(1-3)-	4.989 3.2 101.72 169.3	3.933 3.5 76.81	3.604 72.77	3.325 70.79	3.565 76.29	3.507-3.599 3.451*-3.595* 63.74 64.17*
α -Manp-(1-6)-	4.640 1.5 2.5 102.57 169.1	3.642 3.1 73.20	3.506 73.72	3.366 70.07	3.380 76.51	3.445-3.602 64.17
-3,6)- α -Manp-(1-2)-	4.982 3.2 103.83 174.6	3.924 2.5 72.43	3.621 82.02	3.632 68.27	4.021 74.31	3.486-3.702 68.27
-2)- <i>myo</i> -Ins-(1-O)-P	3.706 <i>weak</i> 78.92	3.949 4.8 80.70	3.208 8.5 73.43	3.337 9.0 75.15	2.936 $^3J_{5,6}=8.8$ 78.49	3.480 $^3J_{6,1}=9.4$ 75.42
GSL-C2						
β -Galf- (1-2)-	4.877 1.1 108.47 174.7	3.777 2.9 83.60	3.800 79.93	3.933 87.07	3.518 73.72	3.360-3.392 65.72
-2)- α -Manp-(1-3)-	5.003 2.5 101.63 169.3	3.916 3.4 76.81	3.594 72.77	3.325 70.79	3.565 76.29	3.507-3.599 3.451*-3.595* 63.74 64.17*
β -Galf-(1-6)-	4.799 1.9 2.5 111.1 171.3	3.806 3.7 84.25	3.779 79.87	3.759 85.71	3.486 73.78	3.507-3.599 3.451*-3.595* 63.74 64.17*
-3,6)- α -Manp-(1-2)-	4.982 2.6 103.83 174.6	3.921 2.9 72.43	3.643 10.2 81.58	3.581 10.2 68.87	4.013 6.6 74.97	3.429-3.747 69.79
-2)- <i>myo</i> -Ins-(1-O)-P	3.699 <i>weak</i> 78.93	3.967 4.8 81.13	3.203 8.5 73.43	3.337 9.0 75.15	2.936 $^3J_{5,6}=8.8$ 78.49	3.468 $^3J_{6,1}=9.4$ 75.42

* possible chemical shift

Table VII. ¹H, ¹³C and ³¹P NMR chemical shifts (ppm) and coupling constants ($J_{H,H}$, $^1J_{C,H}$ and $J_{H,P}$, Hz) for the glycan sequence and for the phosphocholine sequence of the GSL-D fraction.

Cho-P-(O-6)-β-Galf-(1-2)-α-Manp-(1-3)-α-Manp-(1-2)-Ins(1-O)-P-Cer

Glycan moiety						
	H ₁ ³ J _{1,2} C ₁ P	H ₂ ³ J _{2,3} C ₂	H ₃ ³ J _{3,4} C ₃	H ₄ ³ J _{4,5} C ₄	H ₅ ³ J _{5,6} - ³ J _{5,6'} C ₅	H ₆ -H _{6'} C ₆ P
P-(O-6)-β-Galf-(1-2)-	4.939	3.894	3.898	3.974	3.696	3.764
	<i>1,8</i>					
	109.11	83.74	79.46	86.66	72.11	68.92
						0.763 <i>J_{H1,P}=7.5</i>
-2)-α-Manp-(1-3)-	5.069	3.964	3.672	3.413	3.643	3.651-3.562
	<i>1.6</i>	<i>5.2</i>	<i>9.6</i>	<i>9.6</i>		
	102.24	77.25	73.19	71.07	76.70	64.14
-3)-α-Man-(1-2)-	5.108	3.959	3.745	3.647	3.958	3.566
	<i>1.9</i>	<i>4.7</i>				
	103.47	72.66	82.26	68.87	76.22	64.29
-2)-myo-Ins-(1-O)-P	3.756	3.982	3.237	3.433	2.973	3.538
	<i>2.4</i>	<i>2.5</i>	<i>9.4</i>	<i>9.1</i>	<i>³J_{5,6}=9.1</i>	<i>³J_{1,6}=9.2</i>
	79.07	80.43	74.07	75.25	79.00	75.74
	2.974					
	<i>J_{H1,P}=8.6</i>					
Choline-phosphate substitution						
	¹ H					
	¹³ C					
	³¹ P					
CH ₂ OP	4.127					
	61.62					
	0.763					
CH ₂ N	3.571					
	68.79					
N(CH ₃) ₃	3.169					
	56.48					

Table VIII : GIPC structures described in *Aspergillus fumigatus*

Structures of GIPC	References
α -Man-(1-3)- α -Man-(1-6)- α -GlcN-(1-2)-Ins-P-cer	Toledo <i>et al.</i> , 2007; this study
α -Man-(1-3)- α -Man-(1-2)-Ins-P-cer	Leverly <i>et al.</i> , 2001; Toledo <i>et al.</i> , 2007; this study
α -Man-(1-2)- α -Man-(1-3)- α -Man-(1-2)-Ins-P-cer	Toledo <i>et al.</i> , 2007
α -Man-(1-3)-[β -Galf-(1-6)]- α -Man-(1-2)-Ins-P-cer	Toledo <i>et al.</i> , 2007
α -Man-(1-2)- α -Man-(1-3)-[β -Galf-(1-6)]- α -Man-(1-2)-Ins-P-cer	Toledo <i>et al.</i> , 2007
β -Galf-(1-2)- α -Man-(1-3)- α -Man-(1-2)-Ins-P-cer	this study
β -Galf-(1-2)- α -Man-(1-3)-[α -Man-(1-6)]- α -Man-(1-2)-Ins-P-cer	this study
β -Galf-(1-2)- α -Man-(1-3)-[β -Galf-(1-6)]- α -Man-(1-2)-Ins-P-cer	this study
Choline-P-6- β -Galf-(1-2)- α -Man-(1-3)- α -Man-(1-2)-Ins-P-cer	this study