



## H9N2 avian influenza virus in a Mediterranean gull

Camille Lebarbenchon, Chung-Ming Chang, Michel Gauthier-Clerc, Frédéric Thomas, François Renaud, Sylvie van Der Werf

### ► To cite this version:

Camille Lebarbenchon, Chung-Ming Chang, Michel Gauthier-Clerc, Frédéric Thomas, François Renaud, et al.. H9N2 avian influenza virus in a Mediterranean gull. *Journal of Molecular and Genetic Medicine*, 2009, 3 (1), pp.121-123. 10.4172/1747-0862.1000023 . pasteur-00457884

**HAL Id: pasteur-00457884**

**<https://pasteur.hal.science/pasteur-00457884>**

Submitted on 6 Mar 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## H9N2 avian influenza virus in a Mediterranean gull

Camille Lebarbenchon<sup>1,2,3,\*</sup>, Chung-Ming Chang<sup>2,3,4,\*</sup>, Michel Gauthier-Clerc<sup>1</sup>, Frédéric Thomas<sup>2</sup>, François Renaud<sup>2</sup>, Sylvie van der Werf<sup>3,\*</sup>

<sup>1</sup>Centre de Recherche de la Tour du Valat, Le Sambuc, 13200 Arles, France, <sup>2</sup>GEMI, UMR CNRS/IRD 2724, IRD, 911 avenue Agropolis BP 64501, 34394 Montpellier cedex 5, France, <sup>3</sup>Unité de Génétique Moléculaire des Virus Respiratoires, URA3015 CNRS, EA302 Université Paris 7, Institut Pasteur, 25-28 rue du Dr Roux, 75724 Paris Cedex 15, France, <sup>4</sup>Present address: Research Center for Emerging Viral Infections, Department of Medical Biotechnology and Laboratory Science, Chang Gung University, Taoyuan, Taiwan

\*Correspondence to: Sylvie van der Werf, Email: svdwerf@pasteur.fr, Tel: +33 145688722, Fax: +33 140613241

\* These authors contributed equally to this work

Received 03 July 2008, Revised 18 July 2008, Accepted 18 July 2008, Published online 15 October 2008

J Mol Genet Med (2009), 3(1), 121-123

© Copyright The Authors: This is an open access article, published under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/2.0/uk/>). This license permits non-commercial use, distribution and reproduction of the article, provided the original work is appropriately acknowledged with correct citation details.


Influenza A H9N2 is a low pathogenic virus present worldwide in domestic birds and, since the mid-1990s, it has been found to be endemic in poultry of southern China (Xu et al, 2007). Several cases of bird-to-human transmission have been reported in this area since 1999. Recently, Xu et al (2007) provided clear evidence that in southern China H9N2 and the highly-pathogenic H5N1 avian influenza viruses (AIV) have exchanged gene segments to generate currently circulating reassortants of both subtypes. Such genetic exchanges are likely to enhance the potential of pandemic spread of these viruses.

During investigation of AIV in wild birds in the Camargue (Rhône delta, South of France), one Mediterranean gull (*Larus melanocephalus*) was tested positive for H9N2 AIV infection by reverse transcription-polymerase chain reaction (Lebarbenchon et al, 2007). In gulls, AIV are divided into American and Eurasian lineages, and some specific subtypes (*i.e.*, H13, H16) are thought to be associated with the gull reservoir. In northern America, the presence of H9 AIV has been reported in shorebirds (including gulls) in combination with all neuraminidase (NA) subtypes except N3. In Europe H9 AIV have rarely been reported in wild birds (Munster et al, 2007) and has never been found in gull species. N2 is, however, the most frequent NA subtype in European waterfowl and a wide diversity of subtype combinations have been found in poultry. Here, we investigated the origin of the H9N2 AIV in Mediterranean gulls by sequencing 413 base pairs (bp) of the hemagglutinin (HA) and 453 bp of the neuraminidase (NA). We performed phylogenetic analysis including sequences from viruses isolated worldwide from

wild birds (especially North American shorebirds) and domestic birds (Europe and southern China).

cDNA sequencing was done with the following set of primers: H9-151f: 5'-CTYCACACAGARCACAATGG-3' and H9-638r: 5'-GTCACACTTGTTGTTGTRTC-3' (Lee et al, 2001); NAN2-up3: 5'-CTTGTTGACAGTATTGGTTCATGGTCT-3' and NAN2-dw2: 5'-AAAGTCTCATAACCTGAGCGAGAATC-3'; using Big Dye Terminator v.1.1 DNA sequencing kit in an Applied Biosystem 3700 automated sequencer. Alignments were performed using complete sequences available from the Influenza Sequence Database, using the progressive alignment algorithm implemented in CLC Free Workbench 4.0.2. Phylogenetic trees were reconstructed using maximum parsimony (MP) methods with the dnaps program of the PHYLIP 3.61 package and the maximum likelihood (ML) with the software PhyML 2.4.4. We used the GTR evolutionary model for the HA and the TN93 for the NA, as selected by Model Generator 0.84. Nucleotide heterogeneity and substitution rates were estimated with a gamma distribution ( $\Gamma$ ). Nodal supports were assessed with 100 bootstrap replicates generated for each method.

The H9 sequence (EU333949) from our specimen is more related to that of American viruses isolated in 2003 from shorebirds (H9N1 and H9N5) than to H9N2 viruses present in Eurasian birds (Figure 1). Analysis of the NA sequence (EU333950) reveals however that the N2 sequence is closely-related to that of viruses isolated from European wild and domestic birds (H5N2, H6N2 and H9N2), and



**Figure 1.** ML consensus phylogram trees for the HA partial gene sequences, using the GTR+ $\Gamma$  ( $\alpha=0.42$ ) evolutionary model. Bootstrap proportions calculated after 100 replications are indicated at nodes in MP (up) and ML (down). Only bootstrap values >90 in at least one of the two methods (MP and ML) are indicated.

clearly unrelated to that of viruses isolated from gulls in northern America (Figure 2).


These findings support different origins of the HA and NA segments for this virus. H9 is likely to result from natural exchanges of viruses between North American and European gull species. Mediterranean gulls are restricted to western Europe, making the possibility of a direct introduction of viruses belonging to the American H9 lineage by this species very unlikely. On the other hand, the occasional presence of North American gulls in France has been reported as small groups of the ring-billed gull (*Larus delawarensis*) cross the Atlantic Ocean each year. The black-headed gull (*Larus ridibundus*) is also likely to be the vector of influenza exchanges between western Europe, where it is widely represented, and the eastern coast of Canada where breeding colonies are present each summer. Investigations into AIV infections among these

bird populations may be of particular interest to understand patterns of intercontinental virus exchange.

The NA segment presumably originated from viruses that circulate in European wild or domestic birds, suggesting that the H9N2 virus we identified results from a reassortment event between American and Eurasian lineages of AIV.

Only a few studies have related natural intercontinental exchanges and gene segment reassortment (e.g., Krauss et al, 2007). Our report further emphasizes that such intercontinental exchanges can occur, as reported by Jackwood and Stallknecht (2007), and may result in the generation of new strains of AIV.

In Europe, H9N2 AIV is rarely detected in wild birds (Munster et al, 2007), suggesting that its circulation is



**Figure 2.** ML consensus phylogram trees for the NA partial gene sequences, using the TN93+ $\Gamma$  ( $\alpha=0.34$ ) evolutionary model. Bootstrap proportions calculated after 100 replications are indicated at nodes in MP (up) and ML (down). Only bootstrap values  $>90$  in at least one of the two methods (MP and ML) are indicated.

probably very limited among these birds. This clearly contrasts with the co-circulation of H9N2 lineages in poultry from Southern China that have a rapid and independent evolution (Figures 1 and 2) and raises questions as to the generation and emergence of potentially pandemic strains.

## COMPETING INTERESTS

None declared.

## ACKNOWLEDGMENTS

CL is supported by a “Tour du Valat/Région Languedoc-Roussillon” PhD fellowship, and CMC by a CNRS post-doctoral fellowship. This work was funded by the French (ANR) “Santé-Environnement/Santé au travail”, the “European Union's FP6” and the “Région PACA”.

## REFERENCES

- Jackwood MW and Stallknecht DE. 2007. Molecular epidemiologic studies on North American H9 avian influenza virus isolates from waterfowl and shorebirds. *Avian Dis*, 51, 558-450.
- Krauss S, Obert CA, Franks J et al. 2007. Influenza in migratory birds and evidence of limited intercontinental virus exchange. *PLoS Pathogens*, 3, e167.
- Lebarbenchon C, Chang C-M, van der Werf S et al. 2007. Influenza A virus in birds during spring migration in the Camargue, France. *J Wildlife Dis*, 43, 789-793.
- Lee MS, Chang PC, Shien JH et al. 2001. Identification and subtyping of avian influenza viruses by reverse transcription-PCR. *J Virol Methods*, 97, 13-22.
- Munster VJ, Baas C, Lexmond P et al. 2007. Spatial, temporal, and species variation in prevalence of influenza A viruses in wild migratory birds. *PLoS Pathogens*, 3, e61.
- Xu KM, Li KS, Smith GJD et al. 2007. Evolution and Molecular Epidemiology of H9N2 influenza A viruses from Quail in southern China, 2000 to 2005. *J Virol*, 81, 2635-2645.