



Emergence of IFN-alpha TRAIL-expressing killer pDCs (IKpDCs) as a consequence of a crosstalk with NK cells. Influence of HIV-1 infection and implication of HMGB1

Marlène Bras, Héla Saidi, Pauline Formaglio, Marie-Thérèse Melki, Marie-Lise Gougeon

► To cite this version:

Marlène Bras, Héla Saidi, Pauline Formaglio, Marie-Thérèse Melki, Marie-Lise Gougeon. Emergence of IFN-alpha TRAIL-expressing killer pDCs (IKpDCs) as a consequence of a crosstalk with NK cells. Influence of HIV-1 infection and implication of HMGB1. 17th International Symposium on HIV and Emerging Infectious Diseases (ISHEID), May 2012, Marseille, France. 9 (Suppl 1), pp.P15, 2012, 10.1186/1742-4690-9-S1-P15 . pasteur-00701408


HAL Id: pasteur-00701408

<https://pasteur.hal.science/pasteur-00701408>

Submitted on 25 May 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

Emergence of IFN-alpha TRAIL-expressing killer pDCs (IKpDCs) as a consequence of a crosstalk with NK cells. Influence of HIV-1 infection and implication of HMGB1

Marlène Bras*, Héla Saidi, Pauline Formaglio, Marie-Thérèse Melki, Marie-Lise Gougeon

From 17th International Symposium on HIV and Emerging Infectious Diseases (ISHEID)
Marseille, France. 23-25 May 2012

Background

Plasmacytoid dendritic cells (pDCs) mainly contribute to antiviral immunity through recognition of viral components resulting in the production of type-I interferon (IFN), a powerful innate antiviral cytokine. IFN-alpha production by pDCs is promoted by a cross-talk with NK cells, that triggers in return the cytotoxicity of NK cells. Given the essential role of pDCs and NK cells in viral control, we addressed the question of the impact of HIV on NK-pDC cross-talk, and the consequences on viral innate immunity.

Methods

pDCs and NK cells were negatively sorted from PBMC of healthy donors. NK cells were kept either unstimulated (rNK) or activated with PMA/ionomycin for 2 hrs (aNK). pDC were either uninfected or infected with R5-HIV-1 BAL (pDCHIV) and cocultured with NK cells at different ratios for 24 h. The fate of both cell types was studied by multiparametric flow cytometry combined to Multianalyte Profiling technology.

Results

HIV-1-infection of primary pDCs induced their maturation, characterized by the expression of maturation markers (HLA-DR, CD80, CD83, CD86) and the homing receptor CCR7. In addition, HIV-1 triggered the emergence of IFN-induced TRAIL expressing killer pDCs. The crosstalk of pDCHIV with aNK cells strongly increased the differentiation of pDCs into killer pDCs. Interestingly,

the alarmin HMGB1, secreted by pDC upon HIV-1 infection, seems to be important in this cross-talk, since its modulation altered pDC maturation and the emergence of IKpDCs. At high concentrations of HIV-1, pDCs were able to activate rNK cells, as assessed by CD69 expression, and to induce IFN-gamma and TNF-alpha expression as well as perforin degranulation by aNK cells.

Conclusion

We report for the first time that NK-pDCHIV crosstalk potentiates the emergence of TRAIL-expressing IFN-alpha-producing pDCs, and also triggers beta-chemokines synthesis and NK cell killing activity in a HMGB1 dependent manner. Overall these data suggest that the dialogue of HIV-infected pDCs with NK cells favors the emergence of both killer pDC and cytotoxic NK cells and promotes host innate immunity through the activation of potent antiviral effectors.

Published: 25 May 2012

doi:10.1186/1742-4690-9-S1-P15

Cite this article as: Bras et al.: Emergence of IFN-alpha TRAIL-expressing killer pDCs (IKpDCs) as a consequence of a crosstalk with NK cells. Influence of HIV-1 infection and implication of HMGB1. *Retrovirology* 2012 **9**(Suppl 1):P15.

* Correspondence: marlene.bras@pasteur.fr
Institut Pasteur, Paris, France