



The bacterial pathogen *Listeria monocytogenes* and the interferon family: type I, type II and type III interferons

Olivier Dussurget, Hélène Bierne, Pascale Cossart

► To cite this version:

Olivier Dussurget, Hélène Bierne, Pascale Cossart. The bacterial pathogen *Listeria monocytogenes* and the interferon family: type I, type II and type III interferons. *Frontiers in Cellular and Infection Microbiology*, 2014, 4 (50), pp.50. 10.3389/fcimb.2014.00050 . pasteur-01145465

HAL Id: pasteur-01145465

<https://pasteur.hal.science/pasteur-01145465>

Submitted on 24 Apr 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Elimination of *L. monocytogenes* is mostly based on the capacity of the host to mount an efficient cellular immune response to infection (Mackanness, 1962; Shi and Pamer, 2011). In particular, the fate of infection depends on the level of macrophage activation and on *Listeria* ability to counteract bactericidal mechanisms of host cells (Shaughnessy and Swanson, 2007; Corr and O'Neill, 2009; Stavru et al., 2011). Bacterial escape from the phagosome and avoidance of autophagy for intracytosolic replication and cell-cell spread have been well characterized. They have been shown to depend on five major virulence factors: the secreted pore-forming toxin listeriolysin O (LLO) and two phospholipases C (PlcA and PlcB) for vacuolar escape, the surface protein ActA for actin-based motility and both ActA and a surface protein of the internalin family, InlK, for autophagy evasion (Cossart, 2011; Dortet et al., 2012). Other strategies of immune escape that lead to the modulation of cytokine expression occur through a variety of mechanisms. Modifications of *L. monocytogenes* peptidoglycan by the *N*-deacetylase PgdA and the *O*-acetyl transferase OatA prevent lysozyme-dependent release of microbe-associated molecular patterns (MAMPs), activation of pathogen-recognition receptors (PRRs) and subsequent production of pro-inflammatory cytokines (Boneca et al., 2007; Aubry et al., 2011; Rae et al., 2011). The toxin LLO induces dephosphorylation of histone H3 and deacetylation of histone H4, which correlate with decreased expression of pro-inflammatory genes, such as the chemokine gene *cxcl2* (Hamon et al., 2007). The secreted internalin InlC inhibits inflammation by interacting with IKK- α , a component of the I κ B-kinase complex, which is essential for NF- κ B activation and expression of pro-inflammatory genes (Gouin et al., 2010). Other evasion mechanisms remain to be characterized, such as the control of the expression of IL-6 by the surface internalin InlH (Personnic et al., 2010). *L. monocytogenes* also has the capacity to modulate interferon (IFN) production during infection. Type I IFN production by infected cells can be controlled by *Listeria* multidrug efflux pumps MdrM and MdrT, via the secretion of the second messenger cyclic-di-AMP (Crimmins et al., 2008; Woodward et al., 2010; Schwartz et al., 2012). Synthesis of type III IFN has also recently been shown to be tuned by *Listeria* nucleomodulin LntA (Lebreton et al., 2011). Our knowledge concerning the role of the IFN cytokine family during listeriosis has rapidly expanded in the last few years and will be the focus of this review.

INTERFERONS

THE IFNs FAMILY

IFNs form a family of proteins secreted by many cell types in response to infection. They were originally named for their capacity to interfere with viral proliferation (Isaacs and Lindemann, 1957). This diverse family is composed of three groups of cytokines, namely type I-, type II-, and type III-IFNs, which are important components of innate immune responses (Table 1). Type I-IFNs consist of IFN- α , IFN- β , IFN- δ , IFN- ϵ , IFN- ζ , IFN- κ , IFN- ν , IFN- τ , and IFN- ω (Levy et al., 2011). Type II-IFN is composed of a single cytokine, IFN- γ (Pestka et al., 2004). Type III-IFNs are IFN- λ 1, IFN- λ 2, and IFN- λ 3 (formerly IL-29, IL-28A, and IL-28B) and IFN- λ 4 (Kotenko, 2011; Prokunina-Olsson

et al., 2013). Type I- and type III-IFNs have similar signal transduction systems (see below) and are phylogenetically closer from each other than type II-IFN (Pestka et al., 2004).

Sequence conservation and chromosome location suggest that type I-IFN genes evolved from a single ancestor through duplication. However, the extent of type I-IFN gene diversification varies greatly depending on the species (Pestka et al., 2004). Generally, a single gene encodes the type I-IFN in fish. In contrast, multiple gene duplications and diversification led to the emergence of sub-types of type I-IFNs in mammals (Table 1). Gene duplication varies also within each sub-type. A single IFN- β gene is found in the human and mouse genomes (Decker et al., 2005; Honda et al., 2006; Durbin et al., 2013). In contrast 13 IFN- α genes and one pseudogene and 14 IFN- α genes and three pseudogenes are found in the human and mouse genomes, respectively (van Pesch et al., 2004; Durbin et al., 2013). A single gene encodes type II-IFN and four genes encode type III-IFNs in human (Decker et al., 2005; Levy et al., 2011). Of note, IFN- λ 1 and IFN- λ 4 are pseudogenes in mice, which prevents the study of these cytokines in this animal model (Table 1) (Fox et al., 2009).

IFN ACTIVITIES: GENERAL OVERVIEW

IFNs are important components of the immune system, which generally trigger cellular protective defenses in response to infection or tumor formation. Type-II IFN (IFN- γ) is a paradigm for this, being an important mediator of innate and adaptive immune responses with a key role in clearance of viral and bacterial pathogens and in tumor control. IFN- γ was first described as an antiviral protein (Wheelock and Sibley, 1965), but is now known to exhibit broader biological activities, non-redundant with that of other types of IFNs. The crucial role of IFN- γ in immunity to infection is reflected by the phenotype of mice lacking the IFN- γ receptor or the IFN- γ gene, which are highly susceptible to *Mycobacterium bovis* BCG infection (Dalton et al., 1993; Kamijo et al., 1993). Genetic deficiencies resulting in the loss of IFN- γ production or signaling in mice lead to increased susceptibility to infections by other intracellular pathogens, such as *L. monocytogenes* (see below), *Salmonella typhimurium* and some viruses (Huang et al., 1993; Harty and Bevan, 1995; Jouanguy et al., 1999). These defects also lead to the loss of tumor control (Kaplan et al., 1998). Patients with deficiencies in the IFN- γ pathway, for instance by mutation in the gene for the IFN- γ receptor 1, are characterized by severe infections with viruses and intracellular bacteria including *L. monocytogenes*, *Salmonella* sp. and mycobacteria (Jouanguy et al., 1996; Newport et al., 1996; Roesler et al., 1999; van de Vosse et al., 2009). IFN- γ mediates macrophage activation, i.e., increased phagocytosis and production of pro-inflammatory cytokines, of microbicidal reactive oxygen and nitrogen species, leading to clearance of intracellular pathogens (Schoenborn and Wilson, 2007). In addition, IFN- γ controls differentiation of T cells in Th1 effector cells, antigen processing and presentation by antigen-presenting cells, which participate to cellular immunity against intracellular pathogens (Schroder, 2003; Hu and Ivashkiv, 2009). The immunostimulatory and immunomodulatory properties of IFN- γ have therapeutic implications. Indeed, IFN- γ is used in patients with chronic granulomatous disease to reduce infection and mortality,

Table 1 | The interferon family.

Type	ST*	Human**	Mouse**	Source	Target	Receptor	Signaling
I	α	189aa-G***	189aa-G***	Ubiquitous	Ubiquitous	IFNAR1	JAK1
	β	187aa-GP	182aa-GP			IFNAR2	TYK2
	δ	—	—				STAT1
	ϵ	208aa	192aa				STAT2
	ζ	—	182aa-G				
	κ	207aa	199aa				
	ν	Pseudogene	—				
	τ	—	—				
	ω	195aa-G	—				
II	γ	166aa-G	155aa-G	NK, NKT, T APC, B	Ubiquitous	IFN γ R1	JAK1
						IFN γ R2	JAK2 STAT1
III	λ 1	200aa-G	Pseudogene	Ubiquitous	Epithelial cells, Hepatocytes	IFNLR1	JAK1
	λ 2	200aa	193aa-G			IL10R2	TYK2
	λ 3	196aa	193aa				STAT1
	λ 4	179aa	Pseudogene				STAT2

*Subtypes.

**Protein length in amino-acids and protein modifications (G: glycosylation, P: phosphorylation).

***13 genes and a pseudogene in the human genome, 14 genes and three pseudogenes in the mouse genome.

although the clinical benefit has not been demonstrated in all studies (Holland, 2010).

Type I-IFNs are produced in responses to viruses, many bacteria and parasites. However, in contrast to type II IFNs, these cytokines are not always protective against bacterial infections. Indeed, the role of type I-IFNs in response to bacterial infection is complex and depends on the microorganism (Decker et al., 2005; Monroe et al., 2010; Carrero, 2013). They contribute to resistance of the host against infection by extracellular bacteria, such as *Escherichia coli*, *Helicobacter pylori*, *Streptococcus agalactiae* and *S. pneumoniae* (Mancuso et al., 2007; Watanabe et al., 2010). In contrast, they are associated with suppression of innate immune responses and increased susceptibility of the host to infection by *L. monocytogenes* (see below), *Brucella abortus*, *Chlamydia muridarum*, *Francisella novicida*, *Salmonella enterica*, *Staphylococcus aureus*, and *Yersinia pestis* (Auerbuch et al., 2004; Carrero et al., 2004; O'Connell et al., 2004; Qiu et al., 2008; Martin et al., 2009; Henry et al., 2010; de Almeida et al., 2011; Patel et al., 2012; Robinson et al., 2012; Archer et al., 2014). These different effects on infection are likely linked to the wide range of cellular responses induced by their downstream effectors, the products of IFN-stimulated genes (ISGs) (Schoggins et al., 2011). Although Type I-IFNs have long been known to induce antiviral response in the infected host (Isaacs and Lindemann, 1957), they can also induce apoptosis, autophagy, differentiation and migration, inhibit proliferation as well as angiogenesis and mediate cellular damage, inflammation or autoimmunity (Trinchieri, 2010). As a result, type I-IFNs have a therapeutic potential that can be used to treat tumors and viral infections (Pestka, 2007; Heim, 2013; Wilson and Brooks, 2013), while being detrimental for the host in response to a subset of pathogens.

Type III-IFNs have been discovered in 2003 (Kotenko et al., 2003; Sheppard et al., 2003) and their activities have been less extensively characterized than those of type I- and type II-IFNs. However, several studies suggest that type I and type III-IFNs share common biological activities (Levy et al., 2011; Zheng et al., 2013). Although type III-IFNs respond to different stimuli, use different receptors and are not always expressed by the same cells as type I IFNs (see below), engagement of type I- and type III-IFN receptors leads to similar transcriptional responses. Like type I-IFNs, type III-IFNs have been involved in antiproliferative and antiviral responses (Iversen and Paludan, 2010; Mordstein et al., 2010; Durbin et al., 2013; Hamming et al., 2013). Recently, type III-IFNs have been shown to be induced in response to bacterial pathogens, but their downstream effects are not yet characterized (Pietilä et al., 2010; Lebreton et al., 2011; Bierre et al., 2012).

ACTIVATION OF IFNs

Transcription of IFN genes is induced rapidly in response to microbial infection. Type I-IFNs can be produced by all cells, while type III-IFNs are secreted by specific cell types, including dendritic and epithelial cells. Type I- and type III-IFNs activation is initiated by detection of MAMPs by PRRs such as endosomal transmembrane Toll-like receptors and cytosolic receptors (Stetson and Medzhitov, 2006; Monroe et al., 2010). Upon recognition of MAMPs, PRRs trigger diverse signaling pathways that involve adaptor proteins and cytosolic or organelle-bound protein scaffolds activating kinases converging to phosphorylation of transcription factors and their subsequent translocation into the nucleus (Figure 1). IRF1, IRF3, IRF4, IRF5, IRF7, and IRF8 are important for transcription of the IFN- α genes, with IRF7 considered as the master regulator of IFN- α response (Honda et al.,




FIGURE 1 | Activation of type I- and type III-IFNs by infection. (1) Cells detect pathogens by membrane-bound PRRs of the TLR family, which trigger the IFN response. Infection of cells with extracellular Gram-negative bacteria leads to activation of TLR4/CD14/MD2 by LPS. Following TLR4 ligation, IRF3 is activated via the adaptors TRAM and TRIF, and TBK1. (2) Intracellular bacteria and viruses are recognized by TLRs located in the endosome. TLR2, TLR3, TLR7/8, and TLR9 detect lipoproteins, dsRNA, ssRNA, and unmethylated CpG DNA, respectively. Engagement of the receptors leads to the induction of MyD88- and TRIF-dependent pathways involving IKK, TBK1, and MAPK-mediated activation of NF κ B, IRF3, IRF7, and AP1, ultimately resulting in transcriptional activation of IFN genes. (3) Infection of cells leads to the release of cyclic-di-nucleotides (c-di-XMP) and nucleic acids (RNA in red, DNA in blue) in the cytosol that signal the presence of microbes and induce expression of interferons. Cyclic-di-nucleotides secreted by bacteria, e.g., through multidrug resistance MdrT and MdrM in *Listeria*, can activate

the protein STING directly and via the cytosolic sensor DDX41. Cytosolic DNA is recognized by DDX41, DAI, cGAS, and IFI16. cGAS converts dsDNA to cGAMP, which stimulates STING. RNA polymerase III (PolIII) can convert cytosolic AT-rich dsDNA to uncapped 5'-triphosphate RNA that, as ssRNA, is sensed by RIGI. Another cytosolic RNA sensor, MDA5, and RIGI signal via the mitochondrial adaptor MAVS and STING. Activated STING recruits TBK1, which phosphorylates transcription factors IRF3 and IRF7. Upon activation, IRFs translocate to the nucleus where they bind to the promoter of type I IFN genes. Bacterial RNA and DNA additionally activate the cytosolic sensor LRRFIP1 that stimulates β -catenin phosphorylation and nuclear translocation where it binds to IRF3, recruits the histone acetyltransferase p300 to the enhanceosome leading to IFN β transcription. Gray font and arrows: general infection pathways, bold black font and arrows: *Listeria* infection pathways. Red strands, RNA; blue strands, DNA. AP1, activator

(Continued)

FIGURE 1 | Continued

protein-1; CD14, cluster of differentiation 14; c-di-XMP, cyclic-di-nucleotide; cGAMP, cyclic-GMP-AMP; cGAS, cGAMP synthase; DAI, DNA-dependent activator of IFN-regulatory factors; DDX41, DEAD (aspartate-glutamate-alanine-aspartate)-box polypeptide 41; ER, endoplasmic reticulum; IFI16, IFN-inducible protein 16; IKK, I κ B kinases; IRAK, interleukin-1 receptor-associated kinase; LP, lipoprotein; LPS, lipopolysaccharide; LRRFIP1, leucine-rich repeat flightless-interacting protein 1; MAM, mitochondria

associated ER membranes; MAPK, mitogen-activated protein kinase; MAVS, mitochondrial antiviral signaling; MD2, myeloid differential protein-2; MDA5, melanoma differentiation-associated gene 5; MyD88, myeloid differentiation factor 88; NF- κ B, nuclear factor of kappa light polypeptide gene enhancer in B-cells; PolIII, RNA polymerase III; RIGI, retinoic acid inducible gene I; STING, stimulator of IFN genes; TBK1, TANK-binding kinase-1; TIRAP, TIR domain-containing adaptor protein; TLR, Toll-like receptor; TRAM, TRIF-related adaptor molecule; TRIF, TIR domain-containing adaptor inducing IFN- β .

2005; Tailor et al., 2007; Levy et al., 2011). Regulation of the IFN- β gene is more complex. Activated IRF3, IRF7, AP-1, and NF- κ B bind to the enhancer/promoter regions of the IFN- β gene and participate to the formation of the enhanceosome, which alters chromatin structure and allows transcription (Panne et al., 2007; Panne, 2008). In contrast, IRFs and NF- κ B independently activate transcription of type III-IFN genes (Iversen and Paludan, 2010).

Regulation of the IFN- γ gene expression is different from that of type I and type III-IFN genes. NK cells and NKT cells are effectors of the innate immune response and primary sources of IFN- γ . Mature NK and NKT cells quickly react to infection by inducing IFN- γ secretion. Upon recognition of ligands expressed on infected cells, NK cell activating-receptors trigger signaling cascades involving adaptor proteins and protein tyrosine kinases leading to activation of Ras/Sos, PLC- γ and MAPK pathways and induction of IFN- γ production (Schoenborn and Wilson, 2007). In addition to receptors, IL-2, IL-15, IL-12, IL-18, and type I-IFNs also contribute to induction of IFN- γ production by NK cells (Newman and Riley, 2007; Schoenborn and Wilson, 2007; Marçais et al., 2013). Similarly, IL-12 and IL-18 induce IFN- γ production by NKT cells (Godfrey and Berzins, 2007). In NK and NKT cells, the IFN- γ gene locus is transcriptionally permissive within accessible chromatin and allows rapid IFN- γ expression upon activation of transcription factors, such as AP-1, NF- κ B, STAT4, and T-bet (Glimcher et al., 2004; Schoenborn and Wilson, 2007; Lazarevic et al., 2013). In addition, naive CD4 and CD8 T cells can differentiate into Th1 CD4 effector T cells and CD8 cytotoxic T lymphocytes capable of IFN- γ secretion (Wilson et al., 2009). IFN- γ production by CD4 and CD8 T cells depends on IL-12, IL-18 and IFN- γ itself and share many signaling pathways with NK cells. Multiple transcription factors act at the IFN- γ promoter, e.g., AP-1, ATF-2/c-Jun, C/EBP, Eomes, Ets-1, NFAT, NF- κ B, Runx3, STATs and T-bet (Schoenborn and Wilson, 2007; Samten et al., 2008; Wilson et al., 2009; Lazarevic et al., 2013). Moreover, distal regulatory elements modify the chromatin and remodel the IFN- γ gene locus to facilitate IFN- γ production (Wilson et al., 2009).

IFN RECEPTORS AND SIGNAL TRANSDUCTION

IFNs are rapidly secreted upon infection and then bind to their receptors on the surface of target cells (Table 1). Type I-IFNs bind the ubiquitous IFNAR receptor, which consists of two chains, IFNAR1 and IFNAR2 (Piehler et al., 2012). Type III-IFNs bind and signal through a different receptor complex, made of two chains: IFNLR1 (also known as IL-28R α) and IL10R2. This receptor is expressed primarily by epithelial cells and hepatocytes (Iversen and Paludan, 2010). Thus, the physiological roles of

type I- and type III-IFNs are distinct because of the different distribution of their receptors in tissues, type III-IFNs acting predominantly at mucosal surfaces (Mordstein et al., 2010; Durbin et al., 2013). Type I- and type III-IFNs use different receptors but trigger the same JAK-STAT signal transduction cascade involving TYK2, JAK1, STAT1, and STAT2 albeit with different kinetics (Figure 2) (Marcello et al., 2006). Ultimately, STAT1, STAT2, and IRF9 form a transcription factor complex, referred to as ISGF3, which translocates to the nucleus and binds to IFN-stimulated responsive elements (ISRE) in the promoter of ISGs (Schindler et al., 2007). Type II-IFN, uses a heterodimeric receptor consisting of IFN γ R1 and IFN γ R2 chains, expressed by many cell types (Bach et al., 1997). IFN- γ activates JAK1, JAK2 and STAT1, leading to transcription of genes bearing a γ -activation sequence (GAS) in their promoter (Figure 2) (Schindler et al., 2007).

IFNs AND LISTERIA INFECTION**INDUCTION OF IFNs IN RESPONSE TO LISTERIA INFECTION**

An initial innate immune response is triggered when *Listeria* MAMPs activate PRRs of host cells such as epithelial cells and macrophages (Figure 1). Infection induces a robust type I-IFN response. In mice, macrophages have been identified as the major source of IFN- β (Stockinger et al., 2009). *In vitro*, IFN- β production by bone-marrow-derived murine macrophages has been shown to require bacterial escape from the phagosome and activation of cytosolic surveillance pathways (O'Riordan et al., 2002). Induction of IFN- β depends on the adaptor protein STING and the cytosolic PRR DDX41, which are activated by bacterial secondary messengers c-di-AMP and c-di-GMP and by bacterial DNA (Woodward et al., 2010; Burdette et al., 2011; Sauer et al., 2011; Parvatiyar et al., 2012; Archer et al., 2014). STING is a direct receptor for cyclic-dinucleotides, including the cellular second messenger cyclic GMP-AMP (cGAMP) which is produced by the cytosolic sensor cGAMP synthase (cGAS) upon interaction with microbial DNA (Ablasser et al., 2013; Gao et al., 2013; Wu et al., 2013; Sun et al., 2013; Schoggins et al., 2014). Interestingly, type I-IFN production requires activation of the RIG-I helicase by *Listeria* RNA in non-immune cells lacking a functional STING signaling pathway (Abdullah et al., 2012; Hagmann et al., 2013). Another cytosolic PRR, the leucine-rich repeat-containing protein LRRFIP1, has also been implicated in IFN- β production by mouse primary peritoneal macrophages in response to *Listeria* infection, possibly by sensing double stranded DNA and RNA (Yang et al., 2010). While production of IFN- β in response to *Listeria* infection is independent from TLRs in bone-marrow-derived macrophages (McCaffrey et al., 2004; Stockinger et al., 2004; O'Connell et al., 2005), TLR-2 contributes significantly




FIGURE 2 | Signal transduction by the type I, type II and type III IFN receptors. Engagement of type I and type III IFNs by their respective receptors, IFNAR1/IFNAR2 and IFNLR1/IL10R2, triggers phosphorylation of JAK1/TYK2 kinases that activate STAT1 and STAT2. Phosphorylated STAT1/STAT2 heterodimers bind IRF9 forming the ISGF3 complex, which translocates to the nucleus where it induces expression of genes with ISRE-dependent promoters. Engagement of type II IFN by the IFN γ R1/IFN γ R2 receptor induces phosphorylation of JAK1/JAK2 kinases that activate STAT1. Phosphorylated STAT1 homodimers translocate to the

nucleus and induces expression of genes with GAS-dependent promoters. CXCL, CXC chemokine ligand; DUOX, dual oxidase; GAS, gamma-IFN activated site; GBP, guanylate-binding protein; IDO, indoleamine 2,3-dioxygenase; IFITM, IFN-induced transmembrane protein; IRF9, IFN-regulatory factor 9; ISG, IFN-stimulated gene; ISGF3, IFN-stimulated gene factor 3; ISRE, IFN-stimulated response element; JAK, Janus activated kinase; MX, myxovirus resistance; NOS2, nitric oxide synthase 2; NOX, NADPH oxidase; OAS, oligoadenylate synthetase; STAT, signal transducer and activator of transcription; TRIM, tripartite motif; TYK2, tyrosine kinase 2.

to IFN- β secretion by peritoneal macrophages, suggesting that specific macrophage populations have evolved different recognition strategies in response to *Listeria* infection (Aubry et al., 2012).

Listeria infection has recently been shown to induce type III-IFN gene expression in cells of epithelial origin, such as intestinal and trophoblast cells and hepatocytes (Lebreton et al., 2011; Bierne et al., 2012). Similar to type I-IFN, type III-IFN induction is triggered by intracellular *Listeria* (Bierne et al., 2012).

Listeria infection also triggers a rapid and robust IFN- γ response. After intravenous infection of mice with *L. monocytogenes*, NK and T cells are the main sources of IFN- γ (Thale and Kiderlen, 2005; Bou Ghanem et al., 2009). IFN- γ producing V18 $^{+}$ - $\gamma\delta$ T cells are other murine immune cells induced at an early stage of *Listeria* infection in mice inoculated intraperitoneally (Hamada et al., 2008). Using oral infection of mice, the natural route of infection in permissive hosts, *L. monocytogenes*

has been shown to induce IFN- γ production by intraepithelial lymphocytes of the small intestine (Okamoto, 1994). More recently, human E-cadherin (hEcad) expressing mice, a mouse line permissive for *Listeria* oral infection (Lecuit et al., 2001), were used to study cells involved in intestinal mucosal immunity. Infection induced IFN- γ production in NK cells of the small intestine (Reynders et al., 2011).

ROLE OF IFNs DURING *LISTERIA* INFECTION

The production of IFN- γ by immune cells promotes bacterial clearance and is thus critical in controlling primary *L. monocytogenes* infections (Zenewicz and Shen, 2007). Injection of neutralizing monoclonal anti-IFN- γ antibodies in mice infected intraperitoneally with *L. monocytogenes* inhibits macrophage activation and increases the mortality rate (Buchmeier and Schreiber, 1985). In addition, resistance of IFN- γ gene or IFN- γ receptor knock-out mice infected intravenously with *L. monocytogenes* is

severely impaired (Huang et al., 1993; Harty and Bevan, 1995). Recent work using cell-type specific inactivation of STAT1 in mice elegantly demonstrated the key role of IFN- γ and STAT1 in macrophage activation and clearance of *Listeria* (Kernbauer et al., 2012). Interestingly, the role of STAT1 was extremely different after infection of immunized mice. STAT1 signaling in T cells and dendritic cells was critical for adaptive immunity to *Listeria*, while IFN- γ -activated macrophages were not essential anymore once memory cells were produced. Upon oral infection of hEcad mice with *Listeria*, IFN- γ contributes to the control of bacterial burden in the intestine and of bacterial dissemination to other organs. For instance, blocking IFN- γ with neutralizing antibodies increases *Listeria* load in the small intestine, the mesenteric lymph nodes and in the spleen of mice infected orally (Reynders et al., 2011).

In contrast to IFN- γ , type I-IFN is beneficial to *L. monocytogenes*. Mice lacking type I-IFN receptor or IRF3 are more resistant to *Listeria* intraperitoneal or intravenous infection (Auerbuch et al., 2004; Carrero et al., 2004; O'Connell et al., 2004; Garifulin et al., 2007; Jia et al., 2009). The role of type I-IFNs in increasing host susceptibility could be explained by modulation of components of the immune response involved in controlling bacterial growth such as induction of T cell apoptosis, resulting in greater IL-10 secretion by phagocytic cells, in turn dampening the innate immune response (Carrero and Unanue, 2006), the downregulation of IFN- γ R (Rayamajhi et al., 2010; Kearney et al., 2013), or neutrophil recruitment (Brzoza-Lewis et al., 2012). As shown recently, STING-dependent activation of type I-IFN reduces the adaptive immune response to *L. monocytogenes* (Archer et al., 2014). In contrast, recent studies showed that type I-IFNs can also play a beneficial role for the host during *Listeria* infection, pointing to the infection route and the timing of type I-IFN production as determinative factors (Pontioli et al., 2012; Kernbauer et al., 2013).

Interestingly, different strains of *L. monocytogenes* have been shown to vary greatly in their capacity to induce IFN- β (Reutterer et al., 2008; Schwartz et al., 2012). The LO28 strain hyperinduces IFN- β (Reutterer et al., 2008). This strain bears a non-functional BrtA (also named TetR), the transcriptional repressor of the multidrug efflux pump MdrT (Schwartz et al., 2012; Yamamoto et al., 2012). In *Listeria*, MdrT allows secretion of c-di-AMP, which triggers IFN- β . Thus, derepression of MdrT in the LO28 strain promotes IFN- β production. Of note, high expression of MdrT in LO28 correlates with both induction of IFN- β and lower virulence. Another *Listeria* multidrug resistance transporter, MdrM, has been involved in the stimulation of IFN- β production, possibly by secreting c-di-AMP (Crimmins et al., 2008; Woodward et al., 2010; Witte et al., 2012).

The role of type III-IFNs during listeriosis remains to be determined. Since *Listeria* colonizes several tissues of epithelial origins, such as the liver, intestine and placenta, it is tempting to speculate that IFN- λ s play a role in the interaction of *Listeria* with epithelia. However, a prerequisite to address this question is the establishment of a new animal model, i.e., a mouse line expressing a human E-cadherin, thus permissive for *Listeria* infection of epithelia (Lecuit et al., 2001) and impaired in type III-IFN responses, such as IL28R α knockout mice (Mordstein et al., 2010).

One should keep in mind that the mouse model is not optimal to address the role of type III-IFN in human listeriosis. Indeed, *IFN- λ 1* is a pseudogene in mice, while human cells produce this cytokine upon infection with *L. monocytogenes*. In addition, the type III-IFN receptor is expressed at very low levels in the mouse liver and the IFN- λ response of the mouse liver is very weak (Mordstein et al., 2010). In line with this, it has been recently shown that mouse hepatocytes, in contrast to human hepatocytes, are not responsive to IFN- λ (Hermant et al., 2014).

ROLE OF IFN-STIMULATED GENES DURING *LISTERIA* INFECTION

The beneficial or detrimental effects of IFNs on *Listeria* infection rely on the functional properties of their downstream effectors. Indeed, IFNs elicit expression of hundreds of interferon-stimulated genes (ISGs), which encode proteins involved in a broad range of cellular functions (reviewed in MacMicking, 2004). However, while about 2,000 ISGs have been identified so far (Rusinova et al., 2012), their functions in immunomodulation remain to be characterized. To date, the contribution of interferon-induced proteins on *Listeria* infection has mostly been studied in the context of the IFN- γ pathway.

The antilisterial activity of IFN- γ in phagocytic cells involves induction of oxidative and nitrosative defences, via increased expression of enzymes that control production of reactive oxygen and nitrogen species, such as NOX2/CYBB, DUOX2, and iNOS/NOS2 (MacMicking, 2012). These enzymes play an important role in protecting infected cells against *Listeria* cytoinvasion (Myers et al., 2003; Lipinski et al., 2009). The assembly of these enzymes requires IFN- γ -inducible guanosine triphosphatases (GTPases) of the Gbp (guanylate binding protein) family (Boehm et al., 1998), which not only participate to oxidative pathways but also regulate autophagy (Kim et al., 2011). Several Gbps have been shown to protect cells from *Listeria* infection by coordinating a potent oxidative and vesicular trafficking program (Kim et al., 2011). IFN- γ also induces the expression of many nuclear genes encoding mitochondrial respiratory chain machinery, via activation of the nuclear receptor ERR α (estrogen-related receptor α). ERR α contributes to mitochondrial ROS production and efficient clearance of *L. monocytogenes* (Sonoda et al., 2007). A family of IFN- γ -induced chemokines (CXCL9, CXCL10, CXCL11) displays direct antimicrobial activity against *L. monocytogenes* (Cole et al., 2001). In dendritic cells, one of the IFN- γ -associated ISGs is the immunoregulatory enzyme indoleamine 2,3-dioxygenase (IDO), a key enzyme of the tryptophan metabolism. IDO is proposed to play a role in the containment of *Listeria* within granulomatous structures, thus avoiding massive T cell activation (Popov et al., 2006).

The function of type I IFN-associated ISGs in *Listeria* infection is less documented. Zwaferink et al. have observed that upregulation of iNOS/NOS2 by IFN- β promotes necrotic death of macrophages (Zwaferink et al., 2007). Additionally, several interferon-inducible proteins belong to inflammasomes; thus, type I IFN may potentiate inflammasome activation and cell death by pyroptosis (Malireddi and Kanneganti, 2013). Yet, the link between these effectors and the observed harmful effects of type I IFNs on the host is still unclear. Likewise, the role of

effectors induced by type III IFNs in *Listeria* infection of epithelia is not understood.

Of interest, a subset of ISGs is amongst the most induced genes in the intestinal tissue of gnotobiotic humanized mice infected orally with *L. monocytogenes* (Archambaud et al., 2012). However, which type of IFNs triggers this response and for which function on the intestinal mucosa remain to be explored. In addition, IFN-independent pathways may contribute to expression of these ISGs.

SUBVERSION OF IFN RESPONSES BY *LISTERIA*

Listeria has evolved several mechanisms to avoid immune detection and evade IFN responses. It has been demonstrated that deacetylation of *Listeria* peptidoglycan by the deacetylase PgdA confers resistance to host lysozyme, thus preventing release of MAMPs, such as DNA, RNA and lipopeptides, that trigger IFN- β production (Boneca et al., 2007). *Listeria pgdA* mutants are rapidly killed in murine macrophages, which produce lysozyme, and induce a strong secretion of IFN- β compared to wild-type *Listeria*. The role of PgdA is not limited to the control of type I-IFN production as a *pgdA* mutant hyperinduces pro-inflammatory cytokines as well. Modification of peptidoglycan by PgdA is an extremely efficient mechanism of immune escape used by *Listeria*, which correlates with its critical role in virulence.

Remarkably, *Listeria* has evolved a sophisticated strategy to modulate, either negatively or positively, the expression of ISGs in epithelial cells, by targeting a chromatin-repressive complex, BAHD1 (Bierne et al., 2009; Lebreton et al., 2011, 2012). Indeed, *Listeria* infection promotes, albeit via an unknown mechanism, the targeting of BAHD1 at the promoter of a set of ISGs, thereby downregulating type I- and type III-IFN responses. On the other hand, *Listeria* can produce a nucleomodulin, LntA, which when secreted by intracellular bacteria, enters the nucleus of infected cells, binds BAHD1 and inhibits its function (Lebreton et al., 2012, 2014). Thus, LntA stimulates IFN responses. Consistent with the presence of HDAC1/2 in the BAHD1-associated complex, the level of acetylation of lysine 9 on histone H3, which is a mark of active chromatin, increases at the promoters of ISGs in the presence of LntA. When, in which host conditions, and how LntA targets BAHD1 specifically at ISGs remains an open question. The LntA-mediated stimulation of type III-IFN responses might support localized pro-bacterial conditions, as was proposed for IFN-I responses.

CONCLUDING REMARKS

We have an extensive knowledge of the molecular and cellular mechanisms involved in *Listeria*-host interactions. Yet, our understanding of the immune response to *Listeria*, and more specifically the role IFNs and of their downstream effectors, is far from complete and often relies on studies performed in cultured cells or in mice. However, murine and human listeriosis differ in many aspects (Lecuit, 2007; Hoelzer et al., 2012). For instance, E-cadherin, the major receptor for *Listeria* in epithelial cells, is not functional for *Listeria* uptake in the mouse. Thus, the route of entry of *Listeria* is not strictly the same in mice and humans. Moreover, ISGs induced in response to infection are not identical in mice and humans. Additionally, murine hepatocytes do

not respond to type III-IFNs (Hermant et al., 2014), precluding the study of these IFNs during infection by human hepatotropic pathogens, such as *L. monocytogenes*. Altogether, species-specific differences provide limits to the use of mouse models in characterizing IFN pathways engaged during *Listeria* infection in humans, especially in key epithelial organs such as the gut, liver and placenta. It will be important to perform future studies using adapted animal models, such as humanized mice permissive to oral infection or transgenic mice with human xenografts (Walters et al., 2006), since the effect of type I-IFN on *Listeria* infection depends on the route and time of infection (Pontiroli et al., 2012; Kernbauer et al., 2013) and type III-IFN requires bacterial interaction with epithelia (Bierne et al., 2012). Finally, numerous ISGs are induced upon *Listeria* infection *in vitro*, but the relevant ISGs and their cellular functions remain to be identified. Validation of ISGs identified in cultured cells in adequate *in vivo* models or deduced from analyses of patient samples, will be required to address the complex role of IFNs and bacterial subversion strategies and provide new insights into *Listeria* pathogenesis.

ACKNOWLEDGMENTS

The authors declare no conflict of interest. The authors' work has been supported in part by the Institut Pasteur, Inserm, INRA, Université Paris-Diderot, LabEx IBEID, ANR (Grant EPILIS), French Ligue Nationale contre le Cancer, Fondation Louis-Jeantet, Fondation Le Roch, and European Research Council (Advanced Grant Award 233348 to Pascale Cossart). Pascale Cossart is an international research scholar of the Howard Hughes Medical Institute.

REFERENCES

- Abdullah, Z., Schlee, M., Roth, S., Mraheil, M. A., Barchet, W., Böttcher, J. B. O., et al. (2012). RIG-I detects infection with live *Listeria* by sensing secreted bacterial nucleic acids. *EMBO J.* 31, 4153–4164. doi: 10.1038/emboj.2012.274
- Ablasser, A., Goldeck, M., Cavlar, T., Deimling, T., Witte, G., Röhl, I., et al. (2013). cGAS produces a 2'-5'-linked cyclic dinucleotide second messenger that activates STING. *Nature* 498, 380–384. doi: 10.1038/nature12306
- Archambaud, C., Nahori, M.-A., Soubigou, G., Bécavin, C., Laval, L., Lechat, P., et al. (2012). Impact of lactobacilli on orally acquired listeriosis. *Proc. Nat. Acad. Sci. U.S.A.* 109, 16684–16689. doi: 10.1073/pnas.1212809109
- Archer, K. A., Durack, J., and Portnoy, D. A. (2014). STING-dependent type I IFN production inhibits cell-mediated immunity to *Listeria monocytogenes*. *PLoS Pathog.* 10:e1003861. doi: 10.1371/journal.ppat.1003861
- Aubry, C., Corr, S. C., Wienerroither, S., Goulard, C., Jones, R., Jamieson, A. M., et al. (2012). Both TLR2 and TRIF contribute to interferon- β production during *Listeria* infection. *PLoS ONE* 7:e33299. doi: 10.1371/journal.pone.0033299
- Aubry, C., Goulard, C., Nahori, M.-A., Cayet, N., Decalf, J., Sachse, M., et al. (2011). OatA, a peptidoglycan O-acetyltransferase involved in *Listeria monocytogenes* immune escape, is critical for virulence. *J. Infect. Dis.* 204, 731–740. doi: 10.1093/infdis/jir396
- Auerbuch, V., Brockstedt, D., Meyer-Morse, N., O'Riordan, M., and Portnoy, D. A. (2004). Mice lacking the type I interferon receptor are resistant to *Listeria monocytogenes*. *J. Exp. Med.* 200, 527–533. doi: 10.1084/jem.20040976
- Bach, E. A., Aguet, M., and Schreiber, R. D. (1997). The IFN gamma receptor: a paradigm for cytokine receptor signaling. *Annu. Rev. Immunol.* 15, 563–591. doi: 10.1146/annurev.immunol.15.1.563
- Bierne, H., Tham, T. N., Batsche, E., Dumay, A., Leguillou, M., Kerneis-Golsteyn, S., et al. (2009). Human BAHD1 promotes heterochromatic gene silencing. *Proc. Natl. Acad. Sci. U.S.A.* 106, 13826–13831. doi: 10.1073/pnas.0901259106
- Bierne, H., Travier, L., Mahlaköiv, T., Tailleux, L., Subtil, A., Lebreton, A., et al. (2012). Activation of type III interferon genes by pathogenic bacteria in infected epithelial cells and mouse placenta. *PLoS ONE* 7:e39080. doi: 10.1371/journal.pone.0039080

- Boehm, U., Guethlein, L., Klamp, T., Ozbek, K., Schaub, A., Fütterer, A., et al. (1998). Two families of GTPases dominate the complex cellular response to IFN- γ . *J. Immunol.* 161, 6715–6723.
- Boneca, I. G., Dussurget, O., Cabanes, D., Nahori, M.-A., Sousa, S., Lecuit, M., et al. (2007). A critical role for peptidoglycan N-deacetylation in *Listeria* evasion from the host innate immune system. *Proc. Natl. Acad. Sci. U.S.A.* 104, 997–1002. doi: 10.1073/pnas.0609672104
- Bou Ghanem, E. N., McElroy, D. S., and D'Orazio, S. E. F. (2009). Multiple mechanisms contribute to the robust rapid gamma interferon response by CD8+ T cells during *Listeria monocytogenes* infection. *Infect. Immun.* 77, 1492–1501. doi: 10.1128/IAI.01207-08
- Brzoza-Lewis, K. L., Hoth, J. J., and Hiltbold, E. M. (2012). Type I interferon signaling regulates the composition of inflammatory infiltrates upon infection with *Listeria monocytogenes*. *Cell. Immunol.* 273, 41–51. doi: 10.1016/j.cellimm.2011.11.008
- Buchmeier, N. A., and Schreiber, R. D. (1985). Requirement of endogenous interferon- γ production for resolution of *Listeria monocytogenes* infection. *Proc. Natl. Acad. Sci. U.S.A.* 82, 7404–7408. doi: 10.1073/pnas.82.21.7404
- Burdette, D. L., Monroe, K. M., Sotelo-Troha, K., Iwig, J. S., Eckert, B., Hyodo, M., et al. (2011). STING is a direct innate immune sensor of cyclic di-GMP. *Nature* 478, 515–518. doi: 10.1038/nature10429
- Camejo, A., Carvalho, F., Reis, O., Leitão, E., Sousa, S., and Cabanes, D. (2011). The arsenal of virulence factors deployed by *Listeria monocytogenes* to promote its cell infection cycle. *Virulence* 2, 379–394. doi: 10.4161/viru.2.5.17703
- Carrero, J., and Unanue, E. (2006). Lymphocyte apoptosis as an immune subversion strategy of microbial pathogens. *Trends Immunol.* 27, 497–503. doi: 10.1016/j.it.2006.09.005
- Carrero, J. A. (2013). Confounding roles for type I interferons during bacterial and viral pathogenesis. *Int. Immunol.* 25, 663–669. doi: 10.1093/intimm/dxt050
- Carrero, J. A., Calderon, B., and Unanue, E. R. (2004). Type I interferon sensitizes lymphocytes to apoptosis and reduces resistance to *Listeria* infection. *J. Exp. Med.* 200, 535–540. doi: 10.1084/jem.20040769
- Cartwright, E. J., Jackson, K. A., Johnson, S. D., Graves, L. M., Silk, B. J., and Mahon, B. E. (2013). Listeriosis outbreaks and associated food vehicles, United States, 1998–2008. *Emerging Infect. Dis.* 19, 1–9. doi: 10.3201/eid1901.120393
- Chiba, S., Nagai, T., Hayashi, T., Baba, Y., Nagai, S., and Koyasu, S. (2011). Listerial invasion protein internalin B promotes entry into ileal Peyer's patches *in vivo*. *Microbiol. Immunol.* 55, 123–129. doi: 10.1111/j.1348-0421.2010.00292.x
- Cole, A. M., Ganz, T., Liese, A. M., Burdick, M. D., Liu, L., and Strieter, R. M. (2001). Cutting edge: IFN-inducible ELR⁺ CXC chemokines display defensin-like antimicrobial activity. *J. Immunol.* 167, 623–627.
- Corr, S. C., and O'Neill, L. A. J. (2009). *Listeria monocytogenes* infection in the face of innate immunity. *Cell. Microbiol.* 11, 703–709. doi: 10.1111/j.1462-5822.2009.01294.x
- Cossart, P. (2011). Illuminating the landscape of host-pathogen interactions with the bacterium *Listeria monocytogenes*. *Proc. Natl. Acad. Sci. U.S.A.* 108, 19484–19491. doi: 10.1073/pnas.1112371108
- Crimmins, G. T., Herskovits, A. A., Rehder, K., Sivick, K. E., Lauer, P., Dubensky, T. W., et al. (2008). *Listeria monocytogenes* multidrug resistance transporters activate a cytosolic surveillance pathway of innate immunity. *Proc. Natl. Acad. Sci. U.S.A.* 105, 10191–10196. doi: 10.1073/pnas.0804170105
- Dalton, D. K., Pitts-Meek, S., Keshav, S., Figari, I. S., Bradley, A., and Stewart, T. A. (1993). Multiple defects of immune cell function in mice with disrupted interferon- γ genes. *Science* 259, 1739–1742. doi: 10.1126/science.8456300
- de Almeida, L. A., Carvalho, N. B., Oliveira, F. S., Lacerda, T. L. S., Vasconcelos, A. C., Nogueira, L., et al. (2011). MyD88 and STING signaling pathways are required for IRF3-mediated IFN- β induction in response to *Brucella abortus* infection. *PLoS ONE* 6:e23135. doi: 10.1371/journal.pone.0023135
- Decker, T., Müller, M., and Stockinger, S. (2005). The Yin and Yang of type I interferon activity in bacterial infection. *Nat. Rev. Immunol.* 5, 675–687. doi: 10.1038/nri1684
- Disson, O., Grayo, S., Huillet, E., Nikitas, G., Langa-Vives, F., Dussurget, O., et al. (2008). Conjugated action of two species-specific invasion proteins for fetoplaental listeriosis. *Nature* 455, 1114–1118. doi: 10.1038/nature07303
- Dortet, L., Mostowy, S., and Cossart, P. (2012). *Listeria* and autophagy escape: involvement of InlK, an internalin-like protein. *Autophagy* 8, 132–134. doi: 10.4161/auto.8.1.18218
- Durbin, R. K., Kolenko, S. V., and Durbin, J. E. (2013). Interferon induction and function at the mucosal surface. *Immunol. Rev.* 255, 25–39. doi: 10.1111/immr.12101
- Fox, B. A., Sheppard, P. O., and O'Hara, P. J. (2009). The role of genomic data in the discovery, annotation and evolutionary interpretation of the interferon- λ family. *PLoS ONE* 4:e4933. doi: 10.1371/journal.pone.0004933
- Freitag, N. E., Port, G. C., and Miner, M. D. (2009). *Listeria monocytogenes*—from saprophyte to intracellular pathogen. *Nat. Rev. Microbiol.* 7, 623–628. doi: 10.1038/nrmicro2171
- Gahan, C., and Hill, C. (2005). Gastrointestinal phase of *Listeria monocytogenes* infection. *J. Appl. Microbiol.* 98, 1345–1353. doi: 10.1111/j.1365-2672.2005.02559.x
- Gao, D., Wu, J., Wu, Y. T., Du, F., Aroh, C., Yan, N., et al. (2013). Cyclic GMP-AMP synthase is an innate immune sensor of HIV and other retroviruses. *Science* 341, 903–906. doi: 10.1126/science.1240933
- Garifulin, O., Qi, Z., Shen, H., Patnala, S., Green, M. R., and Boyartchuk, V. (2007). Irf3 polymorphism alters induction of interferon beta in response to *Listeria monocytogenes* infection. *PLoS Genet.* 3:e152. doi: 10.1371/journal.pgen.0030152
- Glimcher, L. H., Townsend, M. J., Sullivan, B. M., and Lord, G. M. (2004). Recent developments in the transcriptional regulation of cytolytic effector cells. *Nat. Rev. Immunol.* 4, 900–911. doi: 10.1038/nri1490
- Godfrey, D. I., and Berzins, S. P. (2007). Control points in NKT-cell development. *Nat. Rev. Immunol.* 7, 505–518. doi: 10.1038/nri2116
- Gouin, E., Adib-Conquy, M., Balestrino, D., Nahori, M.-A., Villiers, V., Colland, F., et al. (2010). The *Listeria monocytogenes* InlC protein interferes with innate immune responses by targeting the I κ B kinase subunit IKK α . *Proc. Nat. Acad. Sci. U.S.A.* 107, 17333–17338. doi: 10.1073/pnas.1007765107
- Goulet, V., Hebert, M., Hedberg, C., Laurent, E., Vaillant, V., De Valk, H., et al. (2012). Incidence of listeriosis and related mortality among groups at risk of acquiring listeriosis. *Clin. Infect. Dis.* 54, 652–660. doi: 10.1093/cid/cir902
- Hagmann, C. A., Herzner, A. M., Abdullah, Z., Zillinger, T., Jakobs, C., Schubert, C., et al. (2013). RIG-I detects triphosphorylated RNA of *Listeria monocytogenes* during infection in non-immune cells. *PLoS ONE* 8:e62872. doi: 10.1371/journal.pone.0062872
- Hamada, S., Umehara, M., Shiono, T., Hara, H., Kishihara, K., Tanaka, K., et al. (2008). Importance of murine Vdelta1 gammadelta T cells expressing interferon- γ and interleukin-17A in innate protection against *Listeria monocytogenes* infection. *Immunology* 125, 170–177. doi: 10.1111/j.1365-2567.2008.02841.x
- Hamming, O. J., Terczynska-Dyla, E. T. N., Vieyres, G., Dijkman, R., Jorgensen, S. E., Akhtar, H., et al. (2013). Interferon lambda 4 signals via the IFN- λ receptor to regulate antiviral activity against HCV and coronaviruses. *EMBO J.* 32, 3055–3065. doi: 10.1038/emboj.2013.232
- Hamon, M. A., Batsche, E., Regnault, B., Tham, T. N., Seveau, S., Muchardt, C., et al. (2007). Histone modifications induced by a family of bacterial toxins. *Proc. Natl. Acad. Sci. U.S.A.* 104, 13467–13472. doi: 10.1073/pnas.0702729104
- Harty, J. T., and Bevan, M. J. (1995). Specific immunity to *Listeria monocytogenes* in the absence of IFN γ . *Immunity* 3, 109–117. doi: 10.1016/1074-7613(95)90163-9
- Heim, M. H. (2013). 25 years of interferon-based treatment of chronic hepatitis C: an epoch coming to an end. *Nat. Rev. Immunol.* 13, 535–542. doi: 10.1038/nri3463
- Henry, T., Kirimanjeswara, G. S., Ruby, T., Jones, J. W., Peng, K., Perret, M., et al. (2010). Type I IFN signaling constrains IL-17A/F secretion by gamma-delta T cells during bacterial infections. *J. Immunol.* 184, 3755–3767. doi: 10.4049/jimmunol.0902065
- Hermant, P., Demarez, C., Mahlaköiv, T., Staeheli, P., Meuleman, P., and Michiels, T. (2014). Human but not mouse hepatocytes respond to interferon- λ *in vivo*. *PLoS ONE* 9:e87906. doi: 10.1371/journal.pone.0087906
- Hoelzer, K., Pouillot, R., and Dennis, S. (2012). Animal models of listeriosis: a comparative review of the current state of the art and lessons learned. *Vet. Res.* 43:18. doi: 10.1186/1297-9716-43-18
- Holland, S. M. (2010). Chronic granulomatous disease. *Clin. Rev. Allerg. Immunol.* 38, 3–10. doi: 10.1007/s12016-009-8136-z
- Honda, K., Takaoka, A., and Taniguchi, T. (2006). Type I interferon gene induction by the interferon regulatory factor family of transcription factors. *Immunity* 25, 349–360. doi: 10.1016/j.immuni.2006.08.009

- Honda, K., Yanai, H., Negishi, H., Asagiri, M., Sato, M., Mizutani, T., et al. (2005). IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* 434, 772–777. doi: 10.1038/nature03464
- Hu, X., and Ivashkin, L. B. (2009). Cross-regulation of signaling pathways by interferon-gamma: implications for immune responses and autoimmune diseases. *Immunity* 31, 539–550. doi: 10.1016/j.immuni.2009.09.002
- Huang, S., Hendriks, W., Althage, A., Hemmi, S., Bluethmann, H., Kamijo, R., et al. (1993). Immune response in mice that lack the interferon-gamma receptor. *Science* 259, 1742–1745. doi: 10.1126/science.8456301
- Isaacs, A., and Lindemann, J. (1957). Virus Interference. I. The Interferon. *Proc. Royal. Soc. B Biol. Sci.* 147, 258–267. doi: 10.1098/rspb.1957.0048
- Iversen, M. B., and Paludan, S. R. (2010). Mechanisms of type III interferon expression. *J. Interferon Cytokine Res.* 30, 573–578. doi: 10.1089/jir.2010.0063
- Jia, T., Leiner, I., Dorothee, G., Brandl, K., and Pamer, E. G. (2009). MyD88 and Type I interferon receptor-mediated chemokine induction and monocyte recruitment during *Listeria monocytogenes* infection. *J. Immunol.* 183, 1271–1278. doi: 10.4049/jimmunol.0900460
- Jouanguy, E., Altare, F., Lamhamedi, S., Revy, P., Emile, J. F., Newport, M., et al. (1996). Interferon-gamma-receptor deficiency in an infant with fatal bacille Calmette-Guérin infection. *N. Engl. J. Med.* 335, 1956–1961. doi: 10.1056/NEJM199612263352604
- Jouanguy, E., Döffinger, R., Dupuis, S., Pallier, A., Altare, F., and Casanova, J.-L. (1999). IL-12 and IFN-gamma in host defense against mycobacteria and salmonella in mice and men. *Curr. Opin. Immunol.* 11, 346–351. doi: 10.1016/S0952-7915(99)80055-7
- Kamijo, R., Le, J., Shapiro, D., Havell, E. A., Huang, S., Aguet, M., et al. (1993). Mice that lack the interferon-gamma receptor have profoundly altered responses to infection with *Bacillus Calmette-Guérin* and subsequent challenge with lipopolysaccharide. *J. Exp. Med.* 178, 1435–1440. doi: 10.1084/jem.178.4.1435
- Kaplan, D. H., Shankaran, V., Dighe, A. S., Stockert, E., Aguet, M., Old, L. J., et al. (1998). Demonstration of an interferon gamma-dependent tumor surveillance system in immunocompetent mice. *Proc. Natl. Acad. Sci. U.S.A.* 95, 7556–7561. doi: 10.1073/pnas.95.13.7556
- Kearney, S. J., Delgado, C., Eshleman, E. M., Hill, K. K., O'Connor, B. P., and Lenz, L. L. (2013). Type I IFNs downregulate myeloid cell IFN-gamma receptor by inducing recruitment of an early growth response 3/NGFI-A binding protein 1 complex that silences ifngr1 transcription. *J. Immunol.* 191, 3384–3392. doi: 10.4049/jimmunol.1203510
- Kernbauer, E., Maier, V., Rauch, I., Müller, M., and Decker, T. (2013). Route of infection determines the impact of type I interferons on innate immunity to *Listeria monocytogenes*. *PLoS ONE* 8:e65007. doi: 10.1371/journal.pone.0065007
- Kernbauer, E., Maier, V., Stoiber, D., Strobl, B., Schneckleithner, C., Sexl, V., et al. (2012). Conditional Stat1 ablation reveals the importance of interferon signaling for immunity to *Listeria monocytogenes* infection. *PLoS Pathog.* 8:e1002763. doi: 10.1371/journal.ppat.1002763
- Kim, B.-H., Shenoy, A. R., Kumar, P., Das, R., Tiwari, S., and MacMicking, J. D. (2011). A family of IFN-gamma-inducible 65-kD GTPases protects against bacterial infection. *Science* 332, 717–721. doi: 10.1126/science.1201711
- Kotenko, S. V. (2011). IFN- λ s. *Curr. Opin. Immunol.* 23, 583–590. doi: 10.1016/j.coi.2011.07.007
- Kotenko, S. V., Gallagher, G., Baurin, V. V., Lewis-Antes, A., Shen, M., Shah, N. K., et al. (2003). IFN- λ s mediate antiviral protection through a distinct class II cytokine receptor complex. *Nat. Immunol.* 4, 69–77. doi: 10.1038/ni875
- Lazarevic, V., Glimcher, L. H., and Lord, G. M. (2013). T-bet: a bridge between innate and adaptive immunity. *Nat. Rev. Immunol.* 13, 777–789. doi: 10.1038/nri3536
- Lebreton, A., Cossart, P., and Bierne, H. (2012). Bacteria tune interferon responses by playing with chromatin. *Virulence* 3, 87–91. doi: 10.4161/viru.3.1.18531
- Lebreton, A., Job, V., Ragon, M., Le Monnier, A., Dessan, A., Cossart, P., et al. (2014). Structural basis for the inhibition of the chromatin repressor BAHD1 by the bacterial nucleomodulin LntA. *MBio*. 5:e00775-13. doi:10.1128/mBio.00775-13
- Lebreton, A., Lakisic, G., Job, V., Fritsch, L., Tham, T. N., Camejo, A., et al. (2011). A bacterial protein targets the BAHD1 chromatin complex to stimulate type III interferon response. *Science* 331, 1319–1321. doi: 10.1126/science.1200120
- Lecuit, M. (2007). Human listeriosis and animal models. *Microbes Infect.* 9, 1216–1225. doi: 10.1016/j.micinf.2007.05.009
- Lecuit, M., Vandormael-Pournin, S., Lefort, J., Huerre, M., Gounon, P., Dupuy, C., et al. (2001). A transgenic model for listeriosis: role of internalin in crossing the intestinal barrier. *Science* 292, 1722–1725. doi: 10.1126/science.1059852
- Levy, D. E., Marié, I. J., and Durbin, J. E. (2011). Induction and function of type I and III interferon in response to viral infection. *Curr. Opin. Virol.* 1, 476–486. doi: 10.1016/j.coviro.2011.11.001
- Lipinski, S., Till, A., Sina, C., Arlt, A., Grasberger, H., Schreiber, S., et al. (2009). DUOX2-derived reactive oxygen species are effectors of NOD2-mediated antibacterial responses. *J. Cell Sci.* 122, 3522–3530. doi: 10.1242/jcs.050690
- Lorber, B. (1997). Listeriosis. *Clin. Infect. Dis.* 24, 1–9. doi: 10.1093/clinids/24.1.1
- Mackness, G. B. (1962). Cellular resistance to infection. *J. Exp. Med.* 116, 381–406. doi: 10.1084/jem.116.3.381
- MacMicking, J. D. (2004). IFN-inducible GTPases and immunity to intracellular pathogens. *Trends Immunol.* 25, 601–609. doi: 10.1016/j.it.2004.08.010
- MacMicking, J. D. (2012). Interferon-inducible effector mechanisms in cell-autonomous immunity. *Nat. Rev. Immunol.* 12, 367–382. doi: 10.1038/nri3210
- Malireddi, R. K. S., and Kanneganti, T.-D. (2013). Role of type I interferons in inflammasome activation, cell death, and disease during microbial infection. *Front. Cell. Infect. Microbiol.* 3:77. doi: 10.3389/fcimb.2013.00077
- Mancuso, G., Midiri, A., Biondo, C., Beninati, C., Zummo, S., Galbo, R., et al. (2007). Type I IFN signaling is crucial for host resistance against different species of pathogenic bacteria. *J. Immunol.* 178, 3126–3133.
- Marçais, A., Viel, S., Grau, M., Henry, T., Marvel, J., and Walzer, T. (2013). Regulation of mouse NK cell development and function by cytokines. *Front. Immunol.* 4:450. doi: 10.3389/fimmu.2013.00450
- Marcello, T., Grakoui, A., Barba Spaeth, G., Machlin, E. S., Kotenko, S. V., Macdonald, M. R., et al. (2006). Interferons α and λ inhibit hepatitis C virus replication with distinct signal transduction and gene regulation kinetics. *Gastroenterology* 131, 1887–1898. doi: 10.1053/j.gastro.2006.09.052
- Martin, F. J., Gomez, M. I., Wetzel, D. M., Memmi, G., O'Seaghdha, M., Soong, G., et al. (2009). *Staphylococcus aureus* activates type I IFN signaling in mice and humans through the Xr repeated sequences of protein A. *J. Clin. Invest.* 119, 1931–1939. doi: 10.1172/JCI35879
- McCaffrey, R. L., Fawcett, P., O'Riordan, M., Lee, K.-D., Havell, E. A., Brown, P. O., et al. (2004). A specific gene expression program triggered by Gram-positive bacteria in the cytosol. *Proc. Natl. Acad. Sci. U.S.A.* 101, 11386–11399. doi: 10.1073/pnas.0403215101
- Monroe, K. M., McWhirter, S. M., and Vance, R. E. (2010). Induction of type I interferons by bacteria. *Cell. Microbiol.* 12, 881–890. doi: 10.1111/j.1462-5822.2010.01478.x
- Mordstein, M., Michiels, T., and Staeheli, P. (2010). What have we learned from the IL28 receptor knockout mouse? *J. Interferon Cytokine Res.* 30, 579–584. doi: 10.1089/jir.2010.0061
- Myers, J. T., Tsang, A. W., and Swanson, J. A. (2003). Localized reactive oxygen and nitrogen intermediates inhibit escape of *Listeria monocytogenes* from vacuoles in activated macrophages. *J. Immunol.* 171, 5447–5453.
- Newman, K. C., and Riley, E. M. (2007). Whatever turns you on: accessory-cell-dependent activation of NK cells by pathogens. *Nat. Rev. Immunol.* 7, 279–291. doi: 10.1038/nri2057
- Newport, M. J., Huxley, C. M., Huston, S., Hawrylowicz, C. M., Oostra, B. A., Williamson, R., et al. (1996). A mutation in the interferon-gamma-receptor gene and susceptibility to mycobacterial infection. *N. Engl. J. Med.* 335, 1941–1949. doi: 10.1056/NEJM199612263352602
- Nikitas, G., Deschamps, C., Disson, O., Niaux, T., Cossart, P., and Lecuit, M. (2011). Transcytosis of *Listeria monocytogenes* across the intestinal barrier upon specific targeting of goblet cell accessible E-cadherin. *J. Exp. Med.* 208, 2263–2277. doi: 10.1084/jem.20110560
- O'Connell, R. M., Saha, S. K., Vaidya, S. A., Bruhn, K. W., Miranda, G. A., Zarnegar, B., et al. (2004). Type I interferon production enhances susceptibility to *Listeria monocytogenes* infection. *J. Exp. Med.* 200, 437–445. doi: 10.1084/jem.20040712
- O'Connell, R. M., Vaidya, S. A., Perry, A. K., Saha, S. K., Dempsey, P. W., and Cheng, G. (2005). Immune activation of type I IFNs by *Listeria monocytogenes* occurs independently of TLR4, TLR2, and receptor interacting protein 2 but involves TNFR-associated NF kappa B kinase-binding kinase 1. *J. Immunol.* 174, 1602–1607.
- Okamoto, M. (1994). Host defense and endogenous interferon-gamma in the intestines during an oral infection with *Listeria monocytogenes*. *Hokkaido Igaku Zasshi* 69, 794–809.

- Ooi, S. T., and Lorber, B. (2005). Gastroenteritis due to *Listeria monocytogenes*. *Clin. Infect. Dis.* 40, 1327–1332. doi: 10.1086/429324
- O’Riordan, M., Yi, C. H., Gonzales, R., Lee, K.-D., and Portnoy, D. A. (2002). Innate recognition of bacteria by a macrophage cytosolic surveillance pathway. *Proc. Natl. Acad. Sci. U.S.A.* 99, 13861–13866. doi: 10.1073/pnas.202476699
- Panne, D. (2008). The enhanceosome. *Curr. Opin. Struct. Biol.* 18, 236–242. doi: 10.1016/j.sbi.2007.12.002
- Panne, D., Maniatis, T., and Harrison, S. C. (2007). An atomic model of the interferon-beta enhanceosome. *Cell* 129, 1111–1123. doi: 10.1016/j.cell.2007.05.019
- Parvatiyar, K., Zhang, Z., Teles, R. M., Ouyang, S., Jiang, Y., Iyer, S. S., et al. (2012). The helicase DDX41 recognizes the bacterial secondary messengers cyclic di-GMP and cyclic di-AMP to activate a type I interferon immune response. *Nat. Immunol.* 13, 1155–1161. doi: 10.1038/ni.2460
- Patel, A. A., Lee-Lewis, H., Hughes-Hanks, J., Lewis, C. A., and Anderson, D. M. (2012). Opposing roles for interferon regulatory factor-3 (IRF-3) and type I interferon signaling during plague. *PLoS Pathog.* 8:e1002817. doi: 10.1371/journal.ppat.1002817
- Pentecost, M., Otto, G., Theriot, J. A., and Amieva, M. R. (2006). *Listeria monocytogenes* invades the epithelial junctions at sites of cell extrusion. *PLoS Pathog.* 2:e3. doi: 10.1371/journal.ppat.0020003
- Personnic, N., Bruck, S., Nahori, M.-A., Toledo-Arana, A., Nikitas, G., Lecuit, M., et al. (2010). The stress-induced virulence protein InlH controls interleukin-6 production during murine listeriosis. *Infect. Immun.* 78, 1979–1989. doi: 10.1128/IAI.01096-09
- Pestka, S. (2007). The interferons: 50 years after their discovery, there is much more to learn. *J. Biol. Chem.* 282, 20047–20051. doi: 10.1074/jbc.R700004200
- Pestka, S., Krause, C. D., and Walter, M. R. (2004). Interferons, interferon-like cytokines, and their receptors. *Immunol. Rev.* 202, 8–32. doi: 10.1111/j.0105-2896.2004.00204.x
- Piehl, J., Thomas, C., Garcia, K. C., and Schreiber, G. (2012). Structural and dynamic determinants of type I interferon receptor assembly and their functional interpretation. *Immunol. Rev.* 250, 317–334. doi: 10.1111/imr.12001
- Pietilä, T. E., Latvala, S., Osterlund, P., and Julkunen, I. (2010). Inhibition of dynamin-dependent endocytosis interferes with type III IFN expression in bacteria-infected human monocyte-derived DCs. *J. Leukoc. Biol.* 88, 665–674. doi: 10.1189/jlb.1009651
- Pontiroli, F., Dussurget, O., Zanoni, I., Urbano, M., Beretta, O., Granucci, F., et al. (2012). The timing of IFN β production affects early innate responses to *Listeria monocytogenes* and determines the overall outcome of lethal infection. *PLoS ONE* 7:e43455. doi: 10.1371/journal.pone.0043455
- Popov, A., Abdullah, Z., Wickenhauser, C., Saric, T., Driesen, J., Hanisch, F.-G., et al. (2006). Indoleamine 2,3-dioxygenase-expressing dendritic cells form suppurative granulomas following *Listeria monocytogenes* infection. *J. Clin. Invest.* 116, 3160–3170. doi: 10.1172/JCI28996
- Prokunina-Olsson, L., Muchmore, B., Tang, W., Pfeiffer, R. M., Park, H., Dickensheets, H., et al. (2013). A variant upstream of *IFNL3* (*IL28B*) creating a new interferon gene *IFNL4* is associated with impaired clearance of hepatitis C virus. *Nat. Genet.* 45, 164–171. doi: 10.1038/ng.2521
- Qiu, H., Fan, Y., Joyee, A. G., Wang, S., Han, X., Bai, H., et al. (2008). Type I IFNs enhance susceptibility to *Chlamydia muridarum* lung infection by enhancing apoptosis of local macrophages. *Immunol. Rev.* 181, 2092–2102.
- Rae, C. S., Geissler, A., Adamson, P. C., and Portnoy, D. A. (2011). Mutations of the *Listeria monocytogenes* peptidoglycan N-deacetylase and O-acetylase result in enhanced lysozyme sensitivity, bacteriolysis, and hyperinduction of innate immune pathways. *Infect. Immun.* 79, 3596–3606. doi: 10.1128/IAI.00077-11
- Rayamajhi, M., Humann, J., Kearney, S., Hill, K. K., and Lenz, L. L. (2010). Antagonistic crosstalk between type I and II interferons and increased host susceptibility to bacterial infections. *Virulence* 1, 418–422. doi: 10.4161/viru.1.5.12787
- Reutterer, B., Stockinger, S., Pilz, A., Soulat, D., Kastner, R., Westermayer, S., et al. (2008). Type I IFN are host modulators of strain-specific *Listeria monocytogenes* virulence. *Cell. Microbiol.* 10, 1116–1129. doi: 10.1111/j.1462-5822.2007.01114.x
- Reynders, A., Yessaad, N., Vu Manh, T.-P., Dalod, M., Fenis, A., Aubry, C., et al. (2011). Identity, regulation and *in vivo* function of gut NKp46(+)ROR γ t(+) and NKp46(+)ROR γ t(-) lymphoid cells. *EMBO J.* 30, 2934–2947. doi: 10.1038/emboj.2011.201
- Robinson, N., McComb, S., Mulligan, R., Dudani, R., Krishnan, L., and Sad, S. (2012). Type I interferon induces necroptosis in macrophages during infection with *Salmonella enterica* serovar Typhimurium. *Nat. Immunol.* 13, 954–962. doi: 10.1038/ni.2397
- Roesler, J., Kofink, B., Wendisch, J., Heyden, S., Paul, D., Friedrich, W., et al. (1999). *Listeria monocytogenes* and recurrent mycobacterial infections in a child with complete interferon-gamma-receptor (IFN γ R1) deficiency: mutational analysis and evaluation of therapeutic options. *Exp. Hematol.* 27, 1368–1374. doi: 10.1016/S0301-472X(99)00077-6
- Rusinova, I., Forster, S., Yu, S., Kannan, A., Masse, M., Cumming, H., et al. (2012). Interferome v2.0: an updated database of annotated interferon-regulated genes. *Nucleic Acids Res.* 41, 1040–1046. doi: 10.1093/nar/gks1215
- Samten, B., Townsend, J. C., Weis, S. E., Bhoumik, A., Klucar, P., Shams, H., et al. (2008). CREB, ATF, and AP-1 transcription factors regulate IFN- γ secretion by human T cells in response to mycobacterial antigen. *J. Immunol.* 181, 2056–2064.
- Sauer, J.-D., Sotelo-Troha, K., von Moltke, J., Monroe, K. M., Rae, C. S., Brubaker, S. W., et al. (2011). The ENU-induced Goldenticket mouse mutant reveals an essential function of Sting in the *in vivo* interferon response to *Listeria monocytogenes* and cyclic-di-nucleotides. *Infect. Immun.* 79, 688–694. doi: 10.1128/IAI.00999-10
- Schindler, C., Levy, D. E., and Decker, T. (2007). JAK-STAT signaling: from interferons to cytokines. *J. Biol. Chem.* 282, 20059–20063. doi: 10.1074/jbc.R700016200
- Schlech, W. F., Lavigne, P. M., Bortolussi, R. A., Allen, A. C., Haldane, E. V., Wort, A. J., et al. (1983). Epidemic listeriosis: evidence for transmission by food. *N. Engl. J. Med.* 308, 203–206. doi: 10.1056/NEJM198301273080407
- Schoenborn, J. R., and Wilson, C. B. (2007). Regulation of interferon- γ during innate and adaptive immune responses. *Adv. Immunol.* 96, 41–101. doi: 10.1016/S0065-2776(07)96002-2
- Schoggins, J. W., MacDuff, D. A., Imanaka, N., Gainey, M. D., Shrestha, B., Eitson, J. L., et al. (2014). Pan-viral specificity of IFN-induced genes reveals new roles for cGAS in innate immunity. *Nature* 505, 691–695. doi: 10.1038/nature12862
- Schoggins, J. W., Wilson, S. J., Panis, M., Murphy, M. Y., Jones, C. T., Bieniasz, P., et al. (2011). A diverse range of gene products are effectors of the type I interferon antiviral response. *Nature* 472, 481–485. doi: 10.1038/nature09907
- Schroder, K. (2003). Interferon-gamma: an overview of signals, mechanisms and functions. *J. Leukoc. Biol.* 75, 163–189. doi: 10.1189/jlb.0603252
- Schwartz, K. T., Carleton, J. D., Quillin, S. D., Portnoy, D. A., and Leber, J. H. (2012). Hyperinduction of host beta interferon by a *Listeria monocytogenes* strain naturally overexpressing the multidrug efflux pump MdrT. *Infect. Immun.* 80, 1537–1545. doi: 10.1128/IAI.06286-11
- Shaughnessy, L. M., and Swanson, J. A. (2007). The role of the activated macrophage in clearing *Listeria monocytogenes* infection. *Front. Biosci.* 12, 2683–2692. doi: 10.2741/2364
- Sheppard, P., Kindsvogel, W., Xu, W., Henderson, K., Schlutsmeyer, S., Whitmore, T. E., et al. (2003). IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat. Immunol.* 4, 63–68. doi: 10.1038/ni873
- Shi, C., and Pamer, E. G. (2011). Monocyte recruitment during infection and inflammation. *Nat. Rev. Immunol.* 11, 762–774. doi: 10.1038/nri3070
- Sonoda, J., Laganier, J., Mehl, I. R., Barish, G. D., Chong, L.-W., Li, X., et al. (2007). Nuclear receptor ERR- α and coactivator PGC-1 β are effectors of IFN- γ -induced host defense. *Genes Dev.* 21, 1909–1920. doi: 10.1101/gad.1553007
- Stavru, F., Archambaud, C., and Cossart, P. (2011). Cell biology and immunology of *Listeria monocytogenes* infections: novel insights. *Immunol. Rev.* 240, 160–184. doi: 10.1111/j.1600-065X.2010.00993.x
- Stetson, D., and Medzhitov, R. (2006). Type I interferons in host defense. *Immunity* 25, 373–381. doi: 10.1016/j.immuni.2006.08.007
- Stockinger, S., Kastner, R., Kernbauer, E., Pilz, A., Westermayer, S., Reutterer, B., et al. (2009). Characterization of the interferon-producing cell in mice infected with *Listeria monocytogenes*. *PLoS Pathog.* 5:e1000355. doi: 10.1371/journal.ppat.1000355
- Stockinger, S., Reutterer, B., Schaljo, B., Schellack, C., Brunner, S., Materna, T., et al. (2004). IFN regulatory factor 3-dependent induction of type I IFNs by intracellular bacteria is mediated by a TLR- and Nod2-independent mechanism. *J. Immunol.* 173, 7416–7425.
- Sun, L., Wu, J., Du, F., Chen, X., and Chen, Z. J. (2013). Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* 339, 786–791. doi: 10.1126/science.1232458

- Swaminathan, B., and Gernersmidt, P. (2007). The epidemiology of human listeriosis. *Microbes Infect.* 9, 1236–1243. doi: 10.1016/j.micinf.2007.05.011
- Taylor, P., Tamura, T., Kong, H. J., Kubota, T., Kubota, M., Borghi, P., et al. (2007). The feedback phase of type I interferon induction in dendritic cells requires interferon regulatory factor 8. *Immunity* 27, 228–239. doi: 10.1016/j.immuni.2007.06.009
- Thale, C., and Kiderlen, A. (2005). Sources of interferon-gamma in early immune response to *Listeria monocytogenes*. *Immunobiol* 210, 673–683. doi: 10.1016/j.imbio.2005.07.003
- Toledo-Arana, A., Dussurget, O., Nikitas, G., Sesto, N., Guet-Revillet, H., Balestrino, D., et al. (2009). The *Listeria* transcriptional landscape from saprophytism to virulence. *Nature* 459, 950–956. doi: 10.1038/nature08080
- Trinchieri, G. (2010). Type I interferon: friend or foe? *J. Exp. Med.* 207, 2053–2063. doi: 10.1084/jem.20101664
- van de Vosse, E., van Dissel, J. T., and Ottenhoff, T. H. (2009). Genetic deficiencies of innate immune signalling in human infectious disease. *Lancet Infect. Dis.* 9, 688–698. doi: 10.1016/S1473-3099(09)70255-5
- van Pesch, V., Lanaya, H., Renaud, J. C., and Michiels, T. (2004). Characterization of the murine alpha interferon gene family. *J. Virol.* 78, 8219–8228. doi: 10.1128/JVI.78.15.8219-8228.2004
- Vázquez-Boland, J. A., Kuhn, M., Berche, P., Chakraborty, T., Domínguez-Bernal, G., Goebel, W., et al. (2001). *Listeria* pathogenesis and molecular virulence determinants. *Clin. Microbiol. Rev.* 14, 584–640. doi: 10.1128/CMR.14.3.584-640.2001
- Walters, K.-A., Joyce, M. A., Thompson, J. C., Smith, M. W., Yeh, M. M., Prohl, S., et al. (2006). Host-specific response to HCV infection in the chimeric SCID-beige/Alb-uPA mouse model: role of the innate antiviral immune response. *PLoS Pathog.* 2:e59. doi: 10.1371/journal.ppat.0020059
- Watanabe, T., Asano, N., Fichtner-Feigl, S., Gorelick, P. L., Tsuji, Y., Matsumoto, Y., et al. (2010). NOD1 contributes to mouse host defense against *Helicobacter pylori* via induction of type I IFN and activation of the ISGF3 signaling pathway. *J. Clin. Invest.* 120, 1645–1662. doi: 10.1172/JCI39481
- Wheelock, E. F., and Sibley, W. A. (1965). Circulating virus, interferon and antibody after vaccination with the 17-D strain of yellow-fever virus. *N. Engl. J. Med.* 273, 194–198. doi: 10.1056/NEJM196507222730404
- Wilson, C. B., and Brooks, D. G. (2013). Decoding the complexity of type I interferon to treat persistent viral infections. *Trends Microbiol.* 21, 634–640. doi: 10.1016/j.tim.2013.10.003
- Wilson, C. B., Rowell, E., and Sekimata, M. (2009). Epigenetic control of T-helper-cell differentiation. *Nat. Rev. Immunol.* 9, 91–105. doi: 10.1038/nri2487
- Witte, C. E., Archer, K. A., Rae, C. S., Sauer, J.-D., Woodward, J. J., and Portnoy, D. A. (2012). Innate immune pathways triggered by *Listeria monocytogenes* and their role in the induction of cell-mediated immunity. *Adv. Immunol.* 113, 135–156. doi: 10.1016/B978-0-12-394590-7.00002-6
- Woodward, J. J., Iavarone, A. T., and Portnoy, D. A. (2010). c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response. *Science* 328, 1703–1705. doi: 10.1126/science.1189801
- Wu, J., Sun, L., Chen, X., Du, F., Shi, H., Chen, C., et al. (2013). Cyclic GMP-AMP is an endogenous second messenger in innate immune signaling by cytosolic DNA. *Science* 339, 826–830. doi: 10.1126/science.1229963
- Yamamoto, T., Hara, H., Tsuchiya, K., Sakai, S., Fang, R., Matsuura, M., et al. (2012). *Listeria monocytogenes* strain-specific impairment of the TetR regulator underlies the drastic increase in cyclic di-AMP secretion and beta interferon-inducing ability. *Infect. Immun.* 80, 2323–2332. doi: 10.1128/IAI.06162-11
- Yang, P., An, H., Liu, X., Wen, M., Zheng, Y., Rui, Y., et al. (2010). The cytosolic nucleic acid sensor LRRFIP1 mediates the production of type I interferon via a β -catenin-dependent pathway. *Nat. Immunol.* 11, 487–494. doi: 10.1038/ni.1876
- Zenewicz, L., and Shen, H. (2007). Innate and adaptive immune responses to *Listeria monocytogenes*: a short overview. *Microbes Infect.* 9, 1208–1215. doi: 10.1016/j.micinf.2007.05.008
- Zheng, Y.-W., Li, H., Yu, J.-P., Zhao, H., Wang, S. E., and Ren, X.-B. (2013). Interferon- λ s: special immunomodulatory agents and potential therapeutic targets. *J. Innate Immun.* 5, 209–218. doi: 10.1159/000345365
- Zwaferink, H., Stockinger, S., Reipert, S., and Decker, T. (2007). Stimulation of inducible nitric oxide synthase expression by beta interferon increases necrotic death of macrophages upon *Listeria monocytogenes* infection. *Infect. Immun.* 76, 1649–1656. doi: 10.1128/IAI.01251-07

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 05 February 2014; accepted: 04 April 2014; published online: 28 April 2014.

Citation: Dussurget O, Bierne H and Cossart P (2014) The bacterial pathogen *Listeria monocytogenes* and the interferon family: type I, type II and type III interferons. *Front. Cell. Infect. Microbiol.* 4:50. doi: 10.3389/fcimb.2014.00050

This article was submitted to the journal *Frontiers in Cellular and Infection Microbiology*.

Copyright © 2014 Dussurget, Bierne and Cossart. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.