



Cpe1786/IscR of *Clostridium perfringens* represses expression of genes involved in Fe-S cluster biogenesis.

Gaelle André, Elise Haudecoeur, Emmanuelle Courtois, Marc Monot, Bruno Dupuy, Dmitry A Rodionov, Isabelle Martin-Verstraete

► To cite this version:

Gaelle André, Elise Haudecoeur, Emmanuelle Courtois, Marc Monot, Bruno Dupuy, et al.. Cpe1786/IscR of *Clostridium perfringens* represses expression of genes involved in Fe-S cluster biogenesis.. Research in Microbiology, 2016, 10.1016/j.resmic.2016.03.002 . pasteur-01370878

HAL Id: pasteur-01370878

<https://pasteur.hal.science/pasteur-01370878>

Submitted on 23 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Manuscript Number:

Title: Cpe1786/IscR of *Clostridium perfringens*, represses expression of genes involved in FeS cluster biogenesis

Article Type: Special Issue Paper

Keywords: Rrf2 repressor; cysteine-dependent regulation; FeS clusters; clostridia; fermentation

Corresponding Author: Prof. Isabelle Martin-Verstraete, PhD

Corresponding Author's Institution: Institut Pasteur

First Author: Isabelle Martin-Verstraete, PhD

Order of Authors: Isabelle Martin-Verstraete, PhD; Gaelle André, PhD; Elise Haudecoeur, PhD; Emmanuelle Courtois, bachelor; Marc Monot, PhD; Bruno Dupuy, PhD; Dmitry Rodionov, PhD

Abstract: Cpe1786 of *Clostridium perfringens* is an Rrf2-type regulator containing the three-cysteine residues coordinating a Fe-S in IscR, the repressor controlling Fe-S homeostasis in enterobacteria. The cpe1786 gene formed an operon with iscSU involved in Fe-S biogenesis and tmrU. This operon was transcribed from a σ A-dependent promoter. We showed that in the heterologous host *B. subtilis*, Cpe1786 renamed IscRCp negatively controlled its own transcription. We constructed an iscR mutant in *C. perfringens*. We then compared the expression profile of the strain 13 and of the iscR mutant. IscRCp controlled the expression of genes involved in Fe-S biogenesis, in amino-acid or sugar metabolisms, in fermentation pathways and in host compounds utilization. We then demonstrated using a ChIP-PCR experiment that IscRCp interacted with its promoter region in vivo in *C. perfringens* and with the promoter of cpe2093 encoding an amino-acid ABC transporter. We utilized a comparative genomic approach to infer a candidate IscR-binding motif in clostridia and reconstruct IscR regulons in clostridia. We showed that point mutations in the conserved motif of 29 bp identified upstream of iscR decreased the cysteine-dependent repression of iscR mediated by IscRCp.

Dear Editor,

Please find enclosed a copy of an article entitled Cpe1786/IscR of *Clostridium perfringens*, represses expression of genes involved in Fe-S cluster biogenesis. This paper described the characterization of IscR in the spore former pathogen, *Clostridium perfringens*. We combined studies in the heterologous host *Bacillus subtilis*, the inactivation of *iscR* in *C. perfringens*, a comparative transcriptome analysis of the wild-type strain and of the *iscR* mutant and a ChIP-PCR experiment to study the role of IscR in *C. perfringens*. This is to our knowledge the first physiological characterization of an IscR regulator in Gram-positive bacteria.

This paper is a contribution to the Special Issue of Research in Microbiology on "Beneficial and detrimental spore-formers" edited by Véronique Broussole, Christina Nielsen-Leroux, Vincent Sanchis, Frédéric Carlin and Didier Lereclus.

We hope that this paper will be of sufficient importance and quality to be considered suitable for publication in Research in Microbiology.

Yours sincerely

Isabelle Martin-Verstraete

Cpe1786/IscR of *Clostridium perfringens*, represses expression of genes involved in Fe-S cluster biogenesis

Gaëlle André^{1,2}, Elise Haudecoeur^{1,2}, Emmanuelle Courtois^{1,2}, Marc Monot¹, Bruno Dupuy¹, Dmitry A Rodionov³ and Isabelle Martin-Verstraete^{1,2*}

1. Institut Pasteur, Laboratoire de Pathogénèse des Bactéries Anaérobies, 28 rue du Docteur Roux, 75015 Paris, France.

2. Univ Paris 7-Denis Diderot, Sorbonne Paris Cité, 75205 Paris, France.

3. Institute for Information Transmission Problems, Russian Academy of Sciences, Moscow 127994, Russia.

Key words: Rrf2 repressor, cysteine-dependent regulation, Fe-S clusters, clostridia, fermentation

* corresponding author. mail: isabelle.martin-verstraete@pasteur.fr. Tel: 33 (1) 40 61 35 61

Abstract

Cpe1786 of *Clostridium perfringens* is an Rrf2-type regulator containing the three-cysteine residues coordinating a Fe-S in IscR, the repressor controlling Fe-S homeostasis in enterobacteria. The *cpe1786* gene formed an operon with *iscSU* involved in Fe-S biogenesis and *tmrU*. This operon was transcribed from a σ^A -dependent promoter. We showed that in the heterologous host *B. subtilis*, Cpe1786 renamed IscR_{Cp} negatively controlled its own transcription. We constructed an *iscR* mutant in *C. perfringens*. We then compared the expression profile of the strain 13 and of the *iscR* mutant. IscR_{Cp} controlled the expression of genes involved in Fe-S biogenesis, in amino-acid or sugar metabolisms, in fermentation pathways and in host compound utilization. We then demonstrated using a ChIP-PCR experiment that IscR_{Cp} interacted with its promoter region *in vivo* in *C. perfringens* and with the promoter of *cpe2093* encoding an amino-acid ABC transporter. We utilized a comparative genomic approach to infer a candidate IscR-binding motif and reconstruct IscR regulons in clostridia. We showed that point mutations in the conserved motif of 29 bp identified upstream of *iscR* decreased the cysteine-dependent repression of *iscR* mediated by IscR_{Cp}.

Introduction

Clostridium perfringens is a Gram-positive, spore forming anaerobic bacterium. Toxinotype A strains are the causative agent of food poisoning and gas gangrene. Numerous toxins and degradative enzymes secreted by this bacterium contribute to the pathogenicity of fatal infection like gas gangrene [30, 34]. The synergistic actions of these toxins and enzymes on the host tissue are needed for the infection and a coordinated regulation of such virulence factors is observed. In *C. perfringens* strain 13, the VirS/VirR two-component system is involved in the global regulation of production of several toxins (α -, θ -, κ -toxin) and of other virulence factors (capsular polysaccharide, sialidase, hyaluronidases or other enzymes able to degrade host compounds) [24, 35]. Genes involved in carbon, energy, base and amino acid metabolisms are also controlled by VirS/VirR [24, 35]. The response regulator VirR directly regulates the expression of *pfoA* encoding the θ -toxin and of three non-coding RNAs, the VR-RNA, VirU and VirT, which in turns control the expression of *plc* and *colA* encoding the α - and κ -toxin, respectively [26, 36]. Less is known about the physiology of *C. perfringens*. The *ubiG* operon involved in methionine to cysteine conversion and in AI-2 production is controlled by VirS/VirR [25]. Several genes involved in sulfur metabolism are regulated in response to cysteine availability in *C. perfringens*.

strain 13 [1]. These genes are controlled by premature termination of transcription [10] through a cysteine specific T-box or a S-box dependent riboswitch. The *ubiG* operon is submitted to a complex regulation [1]: i) an induction during cysteine starvation via a cysteine specific T-box present upstream of *ubiG* that senses the level of charge of tRNA_{cys}; ii) a control by VirR through the VR-RNA and iii) a regulation by VirX, a regulatory RNA, which controls toxin production independently from VirR.

The expression of genes involved in Fe-S cluster biogenesis, in the maintenance of the cell redox and in the fermentation pathways is also induced during cysteine limitation via mechanisms different from the T-box and S-box systems [1]. Among the genes induced during cysteine limitation, there is *cpe1786* encoding a regulator, which is a good candidate to participate in cysteine-dependent regulation in *C. perfringens*. Cpe1786 belongs to the widespread Rrf2 family of transcription factors. Notably, this family of regulators comprises: i) the Fe-S cluster biogenesis regulator, IscR [32]; ii) the NsrR repressor controlling the expression of NO detoxification systems [40]; iii) the global iron-responsive regulator RirA in *Rhizobiaceae* [15]; and iv) the global repressor of cysteine metabolism, CymR [7, 37]. Several mechanisms for the modulation of Rrf2-type regulator activity in response to environmental signals are involved. IscR, NsrR and RirA coordinate a Fe-S cluster via three conserved cysteines [8, 32, 40]. The Fe-S cluster in IscR and NsrR contributes in sensing the pool of Fe-S or the presence of NO. These metal centers are sensitive to oxidation by NO or peroxide but also to iron availability [32]. Interestingly, the cysteine residues conserved other Rrf2-type regulators are absent in CymR, which displays a different mechanism of control of its activity with the formation of a complex with CysK, an *O*-acetyl-serine-thiol-lyase [39]. CysK via its substrate *O*-acetyl-serine, is the sensor of the cysteine pool and transmits this information to CymR.

In this work, we analyzed the regulatory role of the unique Rrf2-type regulator of *C. perfringens*, Cpe1786.

Materials and Methods

Bacterial strains and culture conditions:

Escherichia coli strains were grown in LB medium and *B. subtilis* strains in SP or a sulfur-free minimal medium [7] containing either 1 mM methionine or 1 mM cystine. For the experiments involving expression of genes under the control of the *xylA* promoter, threonine (50 mg L⁻¹) and 0.1% xylose were added. When needed, ampicillin (100 µg/ml) or chloramphenicol (15 µg/ml) was added to *E. coli* cultures while chloramphenicol (5 µg/ml) or spectinomycin (60 µg/ml) was added for *B. subtilis* cultures. *C. perfringens* strains were grown under anaerobic conditions (10% H₂, 10% CO₂, 90% N₂) in TY (30 g L⁻¹ bacto tryptone, 2g L⁻¹ yeast extract, 1 g L⁻¹ thioglycolate, pH7.4), BHI (37 g L⁻¹) or a sulfur-free minimal medium [1] containing either 1 mM cystine or 1 mM homocysteine. When necessary, thiamphenicol (5 µg/ml) or erythromycin (10 µg/ml) was added to *C. perfringens* cultures.

Strain and plasmid constructions

The *cpe1786* promoter regions (-131,+ 61 or -41,+61 from the transcriptional start site) were amplified by PCR with the creation of EcoRI and BamHI sites. These DNA fragments were inserted between the EcoRI and BamHI sites of pAC6 [38] resulting in plasmids pDIA5820 and pDIA5837, respectively. These plasmids linearized by ScaI were used to transform the *B. subtilis* BSIP1798 or BSIP 2018 strains (Table 1).

Plasmid pDIA5820 was used to perform site directed mutagenesis with the Quikchange site-directed mutagenesis Kit (Stratagene). Synthetic oligonucleotides complementary to opposite strands and containing different mutations (a T→G or G→A at position -22 or -19, respectively) were used to amplify pDIA5820 (Table S1). The presence of the mutation was verified by sequencing. A plasmid containing the double mutation (T-22G and G-19A) was obtained by site directed mutagenesis using pDIA5840 (P(T-22G)*cpe1786-lacZ*) as template.

The complete coding sequence of *cpe1786* (+3 to +503 from the transcriptional start site) was amplified by PCR. This fragment was inserted into pXT [21]. The resulting plasmid, pDIA5744, was introduced in strain BSIP1978 ($\Delta cymR$ P(-131 +61)*cpe1786_{CP}-lacZ*) (Table 1).

A DNA fragment corresponding to the *cpe1786* gene (-131 to +497 from the transcriptional start site) fused to a sequence encoding the X-flag motif was amplified by PCR. This PCR product was inserted by TA cloning into the pGEM-Teasy vector (Promega, Madison, USA) to give pDIA5925. A 2.4 kb fragment corresponding to the pCB102 replicon and the *catP* marker of pMTL83151 (PmeI-AscI fragment treated by Klenow) [13] was cloned into the NaeI site of pDIA5925 giving pDIA5928.

The ClosTron gene knockout system [12] was used to inactivate *cpe1786*. Primers IBS, EBS1d, EBS2 to retarget the group II intron on pMTL007 to this gene (Table S1) were designed by the Targetron design software (<http://www.sigmaaldrich.com>). We generated by overlap extension PCR a 353 bp product that would facilitate intron retargeting to *cpe1786*. The PCR product was cloned into pMTL007. The plasmid pMTL007-*cpe1786*-28a obtained was then introduced in *C. perfringens* strain 13 by electroporation. Transformants were selected on BHI agar containing thiamphenicol and then plated on BHI agar containing erythromycin. Chromosomal DNA of clones resistant to erythromycin and sensitive to thiamphenicol was extracted using the kit QIAamp DNA Mini Kit (Qiagen). PCR using the ErmRAM primers confirmed that ErmR phenotype was due to the splicing of the group I intron from the group II intron following integration. PCRs with primers IMV484 and IMV485 flanking the *cpe1786* gene or with IMV484 located in the *cpe1786* gene (Table S1, Fig. S1) and EBSu were then performed to verify the integration of the Ll.LtrB intron in *cpe1786*.

Enzyme assays and volatile fatty acid analysis

β -Galactosidase specific activity was measured as previously described [7]. Concentration of proteins was determined by the method of Bradford. One unit of β -galactosidase is defined as the amount of enzyme that produces 1 nmol min⁻¹ of *O*-nitrophenol (ONP) at 28°C. The mean value of at least three independent experiments was presented.

The end products of fermentation were detected in supernatants of strain 13 and the *cpe1786* mutant after growth in minimal medium with 1 mM cystine for 48h at 37°C by gas-liquid chromatography as previously described [2]. The amount of fatty acids was calculated by comparison with an internal standard [6].

RNA isolation, quantitative RT-PCR and 5'RACE analysis

Total RNA from strain 13 and its derivative *cpe1786* mutant grown in minimal medium with cystine or homocysteine was extracted as previously described [1]. cDNAs were synthesized with Superscript II Reverse Transcriptase (Invitrogen) using 1 μ g of RNA and 1pmol of hexamer oligonucleotide primers (pDN6, Roche). Real-time quantitative PCR was performed with gene-specific primers (Table S1) as previously described [1]. In each sample, the quantity of cDNAs of a gene was normalized to the quantity of cDNAs of the *gyrA* gene. The relative change in gene expression was recorded as the ratio of normalized target concentrations ($\Delta\Delta C_t$) [16].

5'RACE assays were performed on total RNA extracted from strain 13 grown in minimal medium in the presence of homocysteine using a 5'RACE System kit (Invitrogen). After reverse transcription, the cDNAs were treated with Terminal deoxynucleotidyl transferase to add a

polyC tail. PCR was then performed using a primer hybridizing with the polyC tail and gene-specific primers (Table S1). PCR amplification products were sequenced.

DNA-array hybridization and microarray analysis.

The microarray analysis was performed using previously described Agilent microarrays (GEO Database, GPL 9765) [1]. RNA was labeled with either Cy3 or Cy5 fluorescent dye (GE healthcare) using the SuperScript Indirect cDNA labelling system kit (Invitrogen) according to the manufacturer's recommendations. The cDNA synthesis, the hybridization, the washing and the scanning were performed as previously described [1]. All the slides were analyzed using R and limma software (Linear Model for Microarray Data) from Bioconductor project (www.bioconductor.org) and the normalization was performed as previously described [1]. The complete experience dataset was deposited in the GEO database with accession numbers GSE19359.

Chromatine immunoprecipitation of Cpe1786-XFlag bound to DNA

Strains CPIP01 (*cpe1786*) and CPIP11 (*cpe1786* pDIA5927-*cpe1786*-XFlag) grown in minimal medium containing 1 mM cystine were incubated with 1 % formaldehyde 10 minutes at 37°C. Cross-linking was quenched by addition of glycine (125 mM). Cells were then collected by centrifugation, washed twice with 20 mM Tris-HCl pH7.5, 150 mM NaCl and stored at -80°C. Pellets were resuspended in 10 ml of buffer A (50 mM Tris-HCl pH7.5, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% SDS) and sonicated in a water bath sonicator (Bioruptor, Diagenode) to shear DNA to an average size of 300-500 bp. After removal of cells debris, the supernatant was incubated in the presence of magnetic beads (Sigma) on a rotating wheel. The magnetic beads were washed twice with buffer A, once with buffer A plus 500 mM NaCl, once with buffer B (10 mM Tris-HCl pH 7.5, 1 mM EDTA, 0.1 % Nonidet-P40) and once with 10 mM Tris-HCl pH 7.5, 1 mM EDTA. Proteins were then eluted for 24h at 37°C with 100 µl of elution buffer (50 mM Tris-HCl pH 7.5, 10 mM EDTA, 0.5% SDS, 150 mM NaCl, Triton X-100 0.5%). Samples were treated 1h at 56°C in 0.5X elution buffer containing 50 µg of Proteinase K. DNA was purified using the QIAquick PCR purification Kit (QIAGEN). Chromatine immunoprecipitation (ChIP) was analyzed by q-PCRs using primer pairs flanking the *cpe1786*, *cpe0664*, *cpe1031*, *cpe1371*, *cpe2093* and *ldh* promoter regions (Table S1). The reaction mixture contained 1µl of DNA obtained by ChIP from strains CPIP11 or CPIP01, 400 nM primers and 10 µl of SYBR Green Mix (Roche) in a total volume of 20 µl. We used *gyrA* a gene not regulated by Cpe1786 in transcriptome as a control. The ratio of enrichment by ChIP for a target was calculated as follows: $2^{(Ct_{\text{target}} - Ct_{\text{gyrA}})_{\text{CPIP11}} / (Ct_{\text{target}} - Ct_{\text{gyrA}})_{\text{CPIP01}}}$. The means of two

independent experiments was obtained.

Results

The Cpe1786 regulator of *C. perfringens*

Cpe1786 of *C. perfringens*, a Rrf2-type regulator, shares 50 % identity with the cysteine metabolism repressor CymR from *B. subtilis* [7] and 37 % and 49 % identity with the Fe-S cluster biogenesis regulator IscR from *E. coli* [33] and *Thermincola potens* [31], respectively. An alignment of CymR-type regulators, IscR-type regulators and Cpe1786 (Fig. S2) indicates that the three-cysteine residues coordinating the Fe-S cluster in IscR are conserved in Cpe1786 [8]. Three genes are located downstream of *cpe1786*. The *cpe1785/iscS* and *cpe1784/iscU* genes encode a cysteine desulfurase and a scaffold protein for Fe-S assembly, respectively [3] while *trmU* (*cpe1783*) encodes an enzyme involved in thio-uridylation of tRNAs (Fig. S3A). The expression of these genes increased during cysteine starvation [1] suggesting that they can form an operon. RT-PCR experiments using primers hybridizing with *cpe1786* and *iscS*, *iscS* and *iscU* or *iscU* and *trmU* were performed and confirmed that these genes were cotranscribed (Fig. S1B). So, *cpe1786* of *C. perfringens* forms an operon with an *iscS*-type gene and with *trmU* as found in *B. subtilis* [7] and with *iscS* and *iscU* as observed in *E. coli* and in *T. potens* [31, 32] (Fig. S3A). We then characterized the *cpe1786* promoter by determining its transcriptional start site by 5'RACE analysis. Transcription is initiated 39 bp upstream of the translational start site. -10 (TATAAT) and -35 (TTGACA) boxes that perfectly match with the consensus for σ^A -dependent promoters are found upstream of this transcriptional start site (Fig. 1).

Cpe1786 controls *cpe1786* transcription in the heterologous host *B. subtilis*

To determine whether Cpe1786 controls its own transcription, we constructed a transcriptional fusion between the *cpe1786* promoter region (-131,+61) and *lacZ* and tested the effect of the addition of Cpe1786 on the expression of this fusion in the heterologous host *B. subtilis*. The fusion was integrated at the *amyE* locus of a *B. subtilis* $\Delta cymR$ mutant or a $\Delta cymR$ mutant containing at the *thrC* locus a copy of P_{xylA} -*cpe1786*. The resulting strains were grown in minimal medium in the presence of cystine or methionine corresponding to conditions of cysteine limitation [39]. The P(-131,+61)*cpe1786-lacZ* fusion was constitutively expressed in a *B. subtilis* *cymR* mutant. The introduction of Cpe1786 in this mutant led to a cysteine-dependent repression of this fusion (Table 2) as observed for *cpe1786* expression in *C. perfringens* [1]. This result indicates that Cpe1786 represses its own transcription as observed for IscR in *E. coli* [33]

and we renamed this regulator IscR_{Cp}. The expression of a second P(-41, +61)*cpe1786-lacZ* fusion was also repressed in the presence of cysteine indicating that the DNA binding motif of IscR_{Cp} is located between position -41 and +61 from the transcriptional start site as classically observed for repressors.

Global regulation by the Cpe1786/IscR regulator in *C. perfringens*

To study the role of IscR_{Cp} in *C. perfringens*, we constructed a *cpe1786/iscR* mutant using the Clostron system [12]. The group II intron was inserted into the *iscR* gene in antisense orientation immediately after the 28th nucleotide in its coding sequence. To verify the insertion of the group II intron into *iscR*, PCRs were carried out using a primer internal of the target gene and the intron-specific primer EBSu or primers flanking the insertion site of the intron (Fig. S1C). To identify genes regulated by IscR_{Cp} in *C. perfringens*, we further compared the expression profiles of strain 13 and the *iscR* mutant after growth in a minimal medium in the presence of cystine. 250 genes were differentially expressed in these two strains. We confirmed these transcriptome data by qRT-PCR analysis for several genes (Table 3). Some of the controlled genes including those with expression ratio above three-fold between these two strains, those associated with iron-sulfur clusters biogenesis, carbon or sulfur metabolism and the degradation of host compounds are presented in Table 3. Most of these genes (62 %) were derepressed in an *iscR* mutant as expected for the inactivation of a repressor. A large set of genes regulated by IscR_{Cp} (50 %) was also regulated in response to cysteine availability [1].

Regulation of genes involved in iron-sulfur clusters biogenesis

The expression of genes involved in Fe-S cluster biogenesis was up-regulated in the *iscR* mutant (Table 3). IscR_{Cp} regulates its own transcription in response to cysteine availability in *B. subtilis* (Table 2). Our transcriptome data also showed a 2-fold increase of *iscS*, *iscU* and *trmU* expression in the *iscR* mutant. This up-expression of genes located downstream from *cpe1786/iscR* might be due to the presence of a promoter in the group II intron or in the *erm* cassette and indicated the absence of a major polar effect of the *iscR* gene disruption on the expression of the downstream genes. The expression of *cpe0664*, encoding a 114 amino-acid protein belonging to the HesB family that probably corresponds to an A-type carrier required for Fe-S biogenesis [41], increased in the *iscR* mutant (12.5-fold in transcriptome/27-fold in qRT-PCR). This gene is also induced during cysteine starvation [1]. IscR_{Cp} controls *iscS*, *iscU* and *cpe0664* involved in Fe-S clusters biogenesis likely in response to cysteine availability.

The expression of *cpe2092-cpe2093* and *cpe1371* increased in the mutant in our transcriptome (Table 3). Cpe2092 and Cpe2093, which correspond to the ATP binding cassette and a solute binding protein fused to a permease of an ABC transporter, respectively, share 61 %, 47 % and 44% identity with YxeO, YxeN and YxeM from *B. subtilis*. The *yxe* operon, which is induced under conditions of cysteine limitation and belongs to the CymR regulon, is probably involved in the uptake and degradation of sulfur compounds including S-methyl-cysteine [5]. Cpe1371 shares similarity with symporters. The expression *cpe1371* is also 5-fold derepressed during cysteine starvation [1] suggesting that Cpe1371 could also play a role in the uptake of sulfur compounds. So, IscR_{Cp} might regulate the uptake systems for sulfur-containing metabolites that might be required for the maintenance of pools of cysteine, the sulfur donor for Fe-S cluster biogenesis.

Regulation of fermentation pathways

The expression of *ldh* encoding the lactate dehydrogenase increased 38-fold in transcriptome and 130-fold in qRT-PCR in the *iscR* mutant compared to the strain 13 (Table 3 and Fig. 2A). By contrast, the expression of the *cpe2297-cpe2301* operon, encoding the enzymes responsible for acetyl-CoA to butyryl-CoA conversion decreased in the mutant compared to strain 13 (0.1 to 0.3-fold). These genes are also regulated in response to cysteine availability [1]. To confirm the role of IscR_{Cp} in the control of fermentation pathways, we analyzed the end products of fermentation of strain 13 and the *iscR* mutant grown 48h in minimal medium containing cystine. The butyrate production decreased 2.5-fold in the *iscR* mutant (Fig. 2B). By contrast, lactate production drastically increased in the *iscR* mutant in agreement with the huge derepression of *ldh* transcription in this mutant. So, the modulation of expression of key genes of fermentation pathways correlates with changes in the amount of lactate and butyrate produced.

Regulation of genes involved in host compounds utilization

NagL is a hyaluronidase (μ -toxins), which is probably involved in hyaluronic acid degradation, a host component. The *nagL* expression was induced in the *iscR* mutant (3.75 in transcriptome/ 8.8 in qRT-PCR) and during cysteine limitation [1] suggesting the involvement of IscR_{Cp} in its regulation in response to cysteine availability. The *cpe0818* and *cpe0866* genes were also more expressed in the *iscR* mutant (9.9- and 5.8-fold in transcriptome/16- and 15-fold in qRT-PCR) (Table 3). Cpe0866, which shares similarity with α -N-acetyl-glucosaminidase, could also degrade host compounds. Finally, Cpe0818 encodes a protein similar to endo- β -N-acetyl-

glucosaminidases. These enzymes could be involved either in peptidoglycan hydrolysis such as *B. subtilis* LytD protein or in the hydrolysis between two N-acetyl-glucosamine residues of glycoproteins. Two endo- β -N-acetyl-glucosaminidases that are active on (Man)₆(GlcNAc)₂Asn and/or (Man)₅(GlcNAc)₂Asn substrates exist in *C. perfringens* [14]. Cpe0818 probably degrades glycoproteins to provide mannose and N-Acetyl-glucosamine to *C. perfringens*. Interestingly, an operon encoding a PTS system (Cpe1463 to Cpe1466) belonging to the Mannose/Fructose family of PTS and a gene encoding a phosphomannomutase (Cpe1873) were also up-regulated in the *iscR* mutant. It is tempting to speculate that Cpe0818, the PTS system and the phosphomannomutase are involved in host glycoprotein degradation and in the uptake and utilization of the released sugars and that IscR_{Cp} coordinately regulates the corresponding genes.

Identification of IscR_{Cp} direct targets *in vivo* by chromatine immunoprecipitation

Among the large set of genes negatively controlled by IscR_{Cp} in transcriptome, we would like to identify some direct targets. For this purpose, we tested by ChIP the binding *in vivo* of IscR_{Cp} to a selection of controlled promoters identified in transcriptome. The IscR_{Cp} protein was modified by addition of a C-terminal 3XFlag-tag. The modified gene was expressed under the control of its own promoter. Strain CPIP11 (*iscR* pDIA5928-*iscR*-XFlag) and CPIP01 (*iscR*) used as a control were grown in minimal medium in the presence of cystine that corresponds to conditions of repression by IscR_{Cp}. After *in vivo* cross-linking, DNA fragments bound to IscR_{Cp}-XFlag were enriched by immunoprecipitation using monoclonal antibodies raised against the XFlag. Immunoprecipitated DNAs were used as templates to amplify the promoter regions of 6 genes derepressed in the *iscR* mutant in transcriptome. We then performed q-PCR experiments and compared the relative quantity of DNA retained in strain CPIP11 compared to strain CPIP01 (*iscR*). The data were standardized using *gyrA*, a gene not regulated by IscR_{Cp}, as a control. After ChIP, we detected a 25 +/-1 and 4.5 +/-1 fold enrichment for the *iscR* and the *cpe2093* promoter regions, respectively with strain CPIP11 compared to strain CPIP01. This clearly indicated that IscR_{Cp} controls its own transcription by binding to its promoter region and is a direct regulator of *cpe2093-cpe2092* expression. By contrast, we observed an enrichment factor between 1 and 1.5 in ChIP experiments for the promoter regions of *ldh*, *cpe0664*, *cpe1031* and *cpe1371*. For these genes, it is therefore difficult to discriminate between an absence of binding or a weak interaction of IscR_{Cp} to some of these promoters in our conditions.

Identification of an IscR_{Cp} binding motif

To identify a putative binding motif for IscR_{Cp}, we first compared the promoter regions of *iscR* and *cpe2093*. We identified a conserved sequence upstream of these two promoters (Fig. S3B). In the *iscR* promoter region, the location of this motif (Fig. 1) is in agreement with the requirement for the presence of a DNA sequence between position -41 and +61 to observe a negative control by IscR_{Cp} of its own transcription in *B. subtilis* (Table 2). Interestingly, this motif is very similar to the IscR binding sites of the *iscR* promoter regions of *E. coli* and *T. potens* [9, 31] and is conserved in the *iscR* promoter regions of several clostridia (see discussion). The alignment of the *iscR* promoters of *E. coli*, *T. potens* and *C. perfringens* and the *cpe2093* promoter allowed proposing a conserved motif, AWWGTTGACMAWW-TRMTSGGNWWT (Fig. 3SB). In all cases, this motif overlaps the -35 boxes of the promoters. To confirm the involvement of this motif in *iscR*_{Cp} regulation, two point mutations were introduced in conserved nucleotides in this sequence. The *PiscR-lacZ* fusions containing mutations were introduced at the *amyE* locus of a *B. subtilis* $\Delta cymR$ *thrC*::P_{xyIA}-*iscR*_{Cp} strain. The level of β -galactosidase activity was determined after growth in the presence of methionine or cystine (Table 2). The replacement of the T at position -22 by a G or the G at position -19 by a A (Fig. 1) led to a partial derepression of *iscR* expression in the presence of cystine while in a double mutant (T-22G/G-19A), the expression of *iscR* was only two-fold repressed in the presence of cystine instead of 9-fold for the fusion containing the wild-type promoter region. These results are in agreement with a role of the conserved motif for the cystine-dependent repression of *iscR*_{Cp}, a repression mediated by IscR_{Cp}.

The RegPredict web-server [23] was subsequently used to search for similar DNA motifs and reconstruct candidate IscR regulons in the genomes of 16 *Clostridium* spp. (Table S2). A constructed positional-weight-matrix for the identified IscR-binding motif was applied to upstream gene regions in the genomic sequences of clostridia to identify additional candidate sites (Fig. 3). One or two copies of IscR motif were found upstream of the *iscRSU* gene cluster in each analyzed genome. We also identified a second potential IscR binding site located downstream of the transcriptional start site of *iscR*_{Cp} (Fig. 1 and 3). Additional candidate sites were found upstream of the *suf* operons in *C. acetobutylicum* and *C. kluyveri*, and of the *cysK* gene in *C. cellulolyticum* and *C. beijerinckii*. Finally, we found additional IscR sites upstream of genes controlled by IscR_{Cp} in the obtained *C. perfringens* transcriptome (Table 3). These include potential IscR binding sites in the promoter regions of *cpe0664* encoding an A type carrier for Fe-S clusters biogenesis and the lactate dehydrogenase gene *ldh* (Fig. 3 and Table S2). We mapped the promoters of *cpe0664* and *ldh* by RACE (Fig. S4). In both cases, the potential IscR

binding site is located downstream from the transcriptional start site for *ldh* or overlaps the -35 box for *cpe0664*. This is in agreement with the repression of their expression by IscR_{Cp}.

Discussion

In *E. coli* and several other bacteria, genes involved in Fe-S cluster biogenesis are regulated in response to Fe-S availability through the Fe-S regulatory protein IscR, and are induced during iron starvation and oxidative stress [32]. By contrast, only few data are available concerning the control of Fe-S cluster synthesis in Gram-positive bacteria, in anaerobic bacteria [31] or in response to sulfur availability [11]. In *C. perfringens*, we have previously shown a coordinated regulation in response to cysteine availability of genes involved in Fe-S production (*iscS*, *iscU*, *cpe0664*) [1]. We demonstrate in this work that the IscR_{Cp} repressor mediates this control. Other genes encoding transporters (Cpe2093-Cpe2092 and Cpe1371) that might be involved in supplying for sulfur required for Fe-S biogenesis are also under the coordinated control of IscR_{Cp}. These regulations may allow *C. perfringens* maintaining its pools of Fe-S clusters, which play a crucial role in the physiology of clostridia lacking the heme synthesis machinery [17]. Interestingly, IscR_{Cp} also controls the expression of genes involved in fermentation pathways (Fig. 2) and we observe accordingly a decrease of butyrate production and a drastic increase of lactate production in the *iscR* mutant. The pyruvate to acetyl-CoA conversion is catalyzed by the pyruvate ferredoxin-oxido-reductase (PFOR). This Fe-S enzyme forms CO₂ and acetyl-CoA by oxidizing pyruvate and reducing a 2Fe-4S ferredoxin. This step implies the utilization of several Fe-S clusters while lactate production by the lactate dehydrogenase does not. So, we propose that during cysteine limitation [1], IscR_{Cp} could reroute the fermentation metabolism increasing *ldh* expression to reoxidize the NADH produced during glycolysis and limiting the utilization of Fe-S clusters. It is intriguing to note that in *C. acetobutylicum*, lactate production increases under conditions of iron limitation [4] while in *C. perfringens* *ldh* expression increases under conditions of sulfur limitation [1]. In both cases, IscR that is probably able to sense iron and sulfur availability might be involved in this control.

We then establish using ChIP experiments that IscR_{Cp} directly interacts with the promoter region of *iscR_{Cp}* and of *cpe2093* *in vivo* in *C. perfringens*. We further identify a conserved motif overlapping the -35 boxes of these two promoters that share similarities with the IscR binding motifs of *E. coli*, *Erwinia chrysanthemi*, *Pseudomonas aeruginosa* and *T. potens* [9, 28, 29, 31]. We also propose that a second site is present in the *iscR* promoter region of *C. perfringens* as observed in *E. coli* and *T. potens*. We further show that this motif is conserved upstream of *iscR*

in clostridia (Fig. 3 and Table S2). For *cpe0664* encoding an A-type scaffold protein involved in Fe-S biogenesis and *ldh*, we identify DNA sequences located at position allowing repression (Fig. 3 and S4) that share similarities to the IscR_{Cp} binding motif. However, we fail to detect a clear binding of IscR_{Cp} to the *cpe0664* or the *ldh* promoter region *in vivo* in our conditions. It is known that the binding of several proteins to a promoter region may lead to false negatives in ChIP experiments [20]. So, it is interesting to note that a Rex binding site is identified in the *ldh* promoter region [27] and we cannot exclude that interference might exist between IscR and Rex. In conclusion, *in vitro* experiments will be required to determine if IscR_{Cp} binds to the candidate IscR binding motifs in the *cpe0664* and *ldh* promoter regions. This will deserve further investigations.

Other genes involved in stress response, in metabolism or encoding proteins of unknown functions are regulated both by IscR_{Cp} and in response to cysteine availability [1]. A second class of genes including *cpe1031* encoding a regulator, genes involved in host compound utilization or encoding proteins of unknown function seems to be regulated only by IscR_{Cp}. In *E. coli* and probably also in *T. potens*, two classes of IscR-controlled genes with distinct binding motifs exist. Type I sites are only bound by IscR associated to a Fe-S cluster while type II sites interact both with Apo-IscR and IscR containing a Fe-S cluster [9, 22, 31, 32]. The possible existence of type II promoters in *C. perfringens* and the determination of direct or indirect regulation by IscR_{Cp} for most of the IscR-controlled genes remain to be established and will deserve further investigations.

The IscR regulator is absent in aerobic firmicutes but is present in several clostridia. With a few exceptions, the *iscR* gene is in operon with *iscS* and *iscU* (Fig. 3). The presence of an IscR binding motif upstream of these *iscR* operons strongly suggests a conserved mechanism of control of Fe-S cluster homeostasis in these clostridia. Interestingly, a putative *iscR* binding site is also present upstream of the *suf* operons in *C. acetobutylicum* and *C. kluyveri*. In *C. acetobutylicum*, the Isc system is probably absent while in *C. kluyveri* like in *E. coli* and *T. potens*, both the Isc and Suf machineries are present. In *E. coli*, the *suf* genes are positively controlled by IscR in aerobic conditions and IscR is able to bind to the *suf* promoter region in *E. coli* and *T. potens* through a type II binding site [9, 31]. Moreover, we also identify a potential IscR binding motif upstream of *cysK* of *C. cellulolyticum* and *C. beijerinckii* (Fig. 3 and Table S2). While the *cysK* gene encoding the O-acetylserine-thiol-lyase in *C. perfringens* is induced under conditions of cysteine starvation through a cysteine specific T-box [1], the *cysK* gene in *C. cellulolyticum* and *C. beijerinckii* might be under the control of IscR. Cysteine is the sulfur donor

for Fe-S biogenesis and the possible existence of a coordinated regulation of the *isc* genes and the *cysK* gene might allow supplying for sulfur required for Fe-S production.

Finally, the control by IscR_{Cp} of genes encoding proteins involved in host compound degradation suggests that IscR_{Cp} might play a role during the host colonization or the infection by *C. perfringens*. This is in agreement with recent works showing that in several pathogenic Gram-negative bacteria, IscR controls expression of factors involved in virulence or allowing adaptation to hostile conditions encounters in the host [18, 28].

Acknowledgments. We are thankful to Olga Soutourina for helpful discussion. This research was supported by a grant from the Institut Pasteur (PTR N°256). G. André was the recipient of a grant from the Ministère de l'enseignement supérieur et de la recherche and from the Pasteur-Weizmann foundation. D.Rodionov was supported by the Russian Foundation for Basic Research (14-04-91154).

Figure legends

Figure 1. *C. perfringens* *cpe1786/iscR_{Cp}* promoter region.

The initiation of transcription start site « +1 » identified by 5'RACE is indicated by an arrow. The -10 and -35 boxes of the σ^A -dependent promoter are underlined. The ribosome-binding site is indicated in bold while the translational start site of *cpe1786* is indicated in bold and italic. In the genome of *C. perfringens* 13, the translational start codon of *cpe1786* is a GTG corresponding to a valine leading to a truncation of 25 amino acids and the absence of the $\alpha 1$ helix of the WHTH motif (Fig. S2). We rather proposed the ATG preceded by a ribosome binding site (GAGG). This start codon corresponds to the start of CPF_2040, the homologs of Cpe1786 in the genome of *C. perfringens* ATCC 13124 [19]. Broken arrows indicate the end points of the *iscR-lacZ* fusions. The conserved motif present upstream of the *iscR* genes in clostridia is boxed. Point mutations obtained in this motif are indicated. A second potential IscR binding motif (see Table S2) is over-lined.

Figure 2. Effect of *iscR_{Cp}* gene inactivation on fermentation pathways. (A) Genes involved in fermentation pathways differentially expressed in an *iscR* mutant compared to strain 13. Arrows ↗ and ↘ indicated genes whose expression increased or decreased in an *iscR* mutant compared to strain 13. (B) Changes in metabolic end products in the *C. perfringens* strain 13 and the *iscR* mutant. Chromatography analysis from these two strains after 48 h of growth in minimal medium containing cystine was performed. The mean and standard error of two experiments are shown.

Figure 3. Candidate IscR regulons reconstructed by comparative genomics approach in *Clostridia* spp. (A) Chromosomal clusters of candidate IscR-regulon genes in *Clostridia* and *Thermincola potens*. Homologous genes are marked by the same color. Potential IscR-binding sites are shown by red arrows. (B) Consensus sequence logo of the predicted IscR-binding sites in *Clostridia*. The logo was generated from the sequence alignments with WebLogo (<http://weblogo.berkeley.edu>).

Table 1 Strains used in this study

Strain	Genotype	Origin
<i>E. coli</i>		
DH5	F- ϕ 80 <i>lacZ</i> Δ M15 Δ (<i>lacZYA-argF</i>)U169 <i>deoR recA1 endA1 hsdR17</i> (r_k^- , m_k^-) <i>phoA supE44 thi-1 gyrA96 relA1 λ^-</i>	Invitrogen
Top10	F ⁻ <i>mcrA</i> D(<i>mrr-hsdRMS-mcrBC</i>) f80 <i>lacZ</i> Δ M15 D <i>lacX74 deoR, recA1 araD139 D(ara-leu)</i> 7697 <i>galK rpsL</i> (StrR) <i>endA1 nupG</i>	Invitrogen
<i>B. subtilis</i>		
168	<i>trpC2</i>	Laboratory stock
BSIP1798	<i>trpC2 ΔcymR_{BS} amyE::aphA3 lacZ</i>	[7]
BSIP1978	<i>trpC2 ΔcymR_{BS} amyE::P(-131 +61) <i>iscR</i>_{Cp}-<i>lacZ cat</i></i>	pDIA5820→BSIP1798
BSIP1981	<i>trpC2 ΔcymR_{BS} amyE::P(-131 +61)<i>iscR</i>_{Cp}-<i>lacZ cat thrC::PxylA-iscR</i>_{Cp} <i>spc</i></i>	pDIA5744→BSIP1978
BSIP2018	<i>trpC2 ΔcymR_{BS} amyE::aphA3 lacZ thrC::PxylA-iscR</i> _{Cp} <i>spc</i>	PAC7→BSIP1981
BSIP2020	<i>trpC2 ΔcymR_{BS} amyE ::P(-41,+61)<i>iscR</i>_{Cp}-<i>lacZ thrC::PxylA-iscR</i>_{Cp} <i>spc</i></i>	PDIA5837→BSIP2018
BSIP2022	<i>trpC2 ΔcymR_{BS} amyE ::PiscR_{Cp}T-22G-<i>lacZ cat thrC::PxylA-iscR</i>_{Cp} <i>spc</i></i>	pDIA5840 →BSIP2018
BSIP2024	<i>trpC2 ΔcymR_{BS} amyE ::PiscR_{Cp}G-19A-<i>lacZ cat thrC ::PxylA-iscR</i>_{Cp} <i>spc</i></i>	pDIA5842 →BSIP2018
BSIP2029	<i>trpC2 ΔcymR_{BS} amyE ::PiscR_{Cp}T-22G/G-19A-<i>lacZ cat thrC ::PxylA-iscR</i>_{Cp} <i>spc</i></i>	pDIA5846 →BSIP2018
<i>C. perfringens</i>		
13	Wild-type	[34]
CPIP01	<i>cpe1786/iscR::erm</i>	pMTL007-Cpe1786-28a → strain 13
CPIP11	<i>cpe1786/iscR::erm</i> pDIA5928- <i>iscR</i> -Xflag	pDIA5928→ CPIP01
Plasmids		
pMTL007	group II intron, ErmBtdRAM2 and ltrA ORF from pMTL20lacZTTErmBtdRAM2, Cm ^R /Tm ^R	[12]
pMTL83151	PCB102 replicon, Cm ^R	[13]
pAC6		[38]
pXT	Ap ^R , Sp ^R	[21]
pMTL007- <i>cpe1786</i> -28a	Cm ^R /Tm ^R	This work
pDIA5744	pXT- <i>iscR</i> _{Cp}	This work
pDIA5820	pAC6 P(-131, +61) <i>iscR</i> _{Cp} - <i>lacZ</i>	This work
pDIA5837	pAC6 P(-41, +61) <i>iscR</i> _{Cp} - <i>lacZ</i>	This work
pDIA5840	pAC6 P(-131 +61) T-22G <i>iscR</i> _{Cp} - <i>lacZ</i>	This work
pDIA5842	pAC6 P(-131 +61) G-19A <i>iscR</i> _{Cp} - <i>lacZ</i>	This work
pDIA5846	pAC6 P(-131 +61) T-22G / G-19A <i>iscR</i> _{Cp} - <i>lacZ</i>	This work

pDIA5852	pAC6 P(-131 +61) T-15C <i>iscR_{Cp}-lacZ</i>	This work
pDIA5925	pGEMTeasy- <i>iscR</i> -XFlag	This work
pDIA5928	pGEMTeasy/pCB3 replicon/cat- <i>iscR</i> -XFlag	This work

(1) Ec, Bs and Cp correspond to *E. coli*, *B. subtilis* and *C. perfringens*, respectively

(2) *cat*, *erm* and *spc* encode proteins leading to chloramphenicol (Cm), erythromycin (Em) or spectinomycin (Sp) resistance. Tm corresponds to thiamphenicol.

(3) the arrow indicates a construction by transformation

Table 2. Regulation of *iscR_{Cp}* expression by *IscR_{Cp}* in *B. subtilis*.

Strain	Relevant genotype	β-galactosidase activity (nmol ONP/min/mg of protein)	
		Methionine	Cystine
BSIP1978	$\Delta cymR_{Bs}$ <i>amyE</i> ::P(-131,+61) <i>iscR_{Cp}-lacZ</i>	1070+/-90	1025+/-60
BSIP1981	$\Delta cymR_{Bs}$ <i>amyE</i> ::P(-131,+61) <i>iscR_{Cp}-lacZ thrC</i> ::Pxyl- <i>iscR_{Cp}</i>	1085+/-2	115+/-20
BSIP2020	$\Delta cymR_{Bs}$ <i>amyE</i> ::P(-41,+61) <i>iscR_{Cp}-lacZ thrC</i> ::Pxyl- <i>iscR_{Cp}</i>	730+/-17	58.5+/-1.5
BSIP2022	$\Delta cymR_{Bs}$ <i>amyE</i> ::P(-131,+61) <i>iscR_{Cp}-T-22G-lacZ thrC</i> ::Pxyl- <i>iscR_{Cp}</i>	1025+/-30	370+/-20
BSIP2024	$\Delta cymR_{Bs}$ <i>amyE</i> ::P(-131,+61) <i>iscR_{Cp}-G-19A-lacZ thrC</i> ::Pxyl- <i>iscR_{Cp}</i>	1070+/-30	380+/-5
BSIP2029	$\Delta cymR_{Bs}$ <i>amyE</i> ::P(-131,+61) <i>iscR_{Cp}-T-22G/G-19A-lacZ thrC</i> ::Pxyl- <i>iscR_{Cp}</i>	1050+/-24	515+/-10

β-galactosidase activity was measured on crude extracts from strains grown in minimal medium in the presence of 1 mM methionine or 1 mM cystine. *iscR_{Cp}*= *cpe1786*. For strains BSIP1981, BSIP2020, BSIP2022, BSIP2024 and BSIP2029 0.1% xylose and 50 mg/L of threonine were added.

Table 3. Genes differentially expressed between strain 13 of *C. perfringens* and a *iscR_{Cp}* mutant.

gene (Synonym)	Functions/similarities	Transcriptome	quantitative RT-PCR
		<i>iscR_{Cp}</i> /13	<i>iscR_{Cp}</i> /13
Fe-S clusters and redox functions			
<i>cpe0664</i>	HesB-like protein (114 aa)	12.5	27
<i>cpe1785 (iscS)</i>	Cysteine desulfurase, Fe-S clusters biosynthesis	2.1	
<i>cpe1784 (iscU)</i>	Fe-S clusters assembly	2	
<i>cpe1783 (trmU)</i>	Methylaminomethyl-2-Thiouridylate-Methyltransferase	2	
<i>cpe2511 (fer)</i>	Ferredoxin [3Fe-4S]	0.29	0.2
<i>cpe0855 (rubY)</i>	rubrerythrin	0.17	
Transporters, membrane or exported proteins			
<i>cpe2092</i>	Amino acid ABC transporter, ATP binding cassette	10.2	26
<i>cpe2093</i>	Amino acid ABC transporter, permease	5.5	
<i>cpe2295 (lepW)</i>	Type I Signal peptidase	6.6	
<i>cpe1371</i>	Na ⁺ -dependent symporter	3.6	7.5
<i>cpe1621</i>	putative cation efflux protein	3.4	
<i>cpe1343</i>	probable galactoside ABC transporter	0.1	
<i>cpe1341</i>	probable galactoside ABC transporter	0.21	
Regulators			
<i>cpe1031</i>	ArsR-SmtB family regulator	7.8	20

<i>cpe2304</i>	ArsR-SmtB family regulator	1.99	
Carbon metabolism			
<i>cpe0103 (ldh)</i>	L-lactate dehydrogenase	38.6	130
<i>cpe2531(adhE)</i>	Aldehyde alcohol dehydrogenase	3.03	
<i>cpe2195 (atoB)</i>	Acetyl-CoA acetyltransferase	0.13	
<i>cpe2297</i>	3-hydroxybutyryl-CoA dehydrogenase	0.163	
<i>cpe2298 (fixB)</i>	α subunit electron transfer flavoprotein	0.22	
<i>cpe2299 (etfB)</i>	β subunit electron transfer flavoprotein	0.29	
<i>cpe2300 (bcd)</i>	Acyl-CoA dehydrogenase	0.21	
<i>cpe2301 (crt)</i>	3-hydroxybutyryl-CoA dehydratase	0.33	0.5
<i>cpe0892</i>	butanol dehydrogenase NADPH dependent	0.618	
<i>cpe2347 (buk)</i>	butyrate kinase	0.63	
<i>cpe2348 (ptb)</i>	phosphotransbutyrylase	0.625	
<i>cpe1185 (pfk)</i>	6-phosphofructokinase	0.5	
<i>cpe1299 (eno)</i>	enolase	0.54	
<i>cpe2149 (pykA)</i>	Pyruvate kinase	0.59	
<i>cpe2267</i>	glucose-6-phosphate isomerase	1.65	
<i>cpe1463</i>	PTS system IID component (IIDMan)	2.2	
<i>cpe1464</i>	PTS system IIC component (IICMan)	1.45	
<i>cpe1465</i>	PTS system IIB component (IIDMan)	1.53	
<i>cpe1466</i>	PTS system IIA component (IIAMan)	2.65	
<i>cpe1873(manB)</i>	phosphomannomutase	1.587	
Other metabolisms			
<i>cpe1050</i>	5'-MTA/SAH nucleosidase	3.8	
<i>cpe0056</i>	5'-MTA/SAH nucleosidase	1.58	
<i>cpe1573</i>	Probable glutamate cysteine ligase	2.9	
<i>cpe0447 (lguL)</i>	Lactoylglutathione lyase	2	
Host compounds degradation			
<i>cpe0818</i>	endo- β -N-acetyl-glucosaminidase	9.9	16
<i>cpe0866</i>	α -N-acetyl-glucosaminidase	5.8	15
<i>cpe1523 (nagL)</i>	hyaluronidase	3.8	8.8
Unknown function			
<i>cpe0554</i>	Unknown	6.7	
<i>cpe2261</i>	Unknown	5.3	
<i>cpe2262</i>	Unknown	4.35	
<i>cpe2063</i>	Unknown	5.2	
<i>cpe1079</i>	Unknown	5.2	
<i>cpe1257</i>	Unknown	4.6	
<i>cpe1875</i>	Unknown	4.3	
<i>cpe0106</i>	Unknown	3.5	
<i>cpe0105</i>	Unknown	3.15	
<i>cpe1169</i>	Unknown	3.42	
<i>cpe1735</i>	Unknown	3.66	
<i>cpe1173</i>	Unknown	0.15	
<i>cpe2111</i>	Unknown	0.26	
<i>cpe1078</i>	Unknown	0.27	
<i>cpe0098</i>	Unknown	0.29	
<i>cpe1472</i>	Unknown	0.29	

466 Gene names and functions correspond to those indicated in the GenoList data-base
467 (<http://genodb.pasteur.fr/cgi-bin/WebObjects/GenoList>). A gene was considered as differentially
468 expressed when the p-value is < 0.05 using the statistical analysis described in Materials and

Methods. Controlled genes with expression ratio above three-fold between strains 13 and *iscR_{CP}* and those associated with iron-sulfur clusters biogenesis, redox functions, carbon and sulfur metabolisms and the degradation of host compounds are presented in this Table. We confirmed the transcriptome data by qRT-PCR analysis for several genes.

References

- [1] Andre G, Haudecoeur E, Monot M, Ohtani K, Shimizu T, Dupuy B, et al. Global regulation of gene expression in response to cysteine availability in *Clostridium perfringens*. BMC Microbiol 2010;10:234.
- [2] Antunes A, Camiade E, Monot M, Courtois E, Barbut F, Sernova NV, et al. Global transcriptional control by glucose and carbon regulator CcpA in *Clostridium difficile*. Nucleic acids research 2012;40:10701-18.
- [3] Ayala-Castro C, Saini A, Outten FW. Fe-S cluster assembly pathways in bacteria. Microbiol Mol Biol Rev 2008;72:110-25, table of contents.
- [4] Bahl H, Gottwald M, Kuhn A, Rale V, Andersch W, Gottschalk G. Nutritional Factors Affecting the Ratio of Solvents Produced by *Clostridium acetobutylicum*. Appl Environ Microbiol 1986;52:169-72.
- [5] Burguiere P, Auger S, Hullo MF, Danchin A, Martin-Verstraete I. Three different systems participate in L-cystine uptake in *Bacillus subtilis*. J Bacteriol 2004;186:4875-84.
- [6] Carlier JP, Sellier N. Gas chromatographic-mass spectral studies after methylation of metabolites produced by some anaerobic bacteria in spent media. J Chromatogr 1989;493:257-73.
- [7] Even S, Burguiere P, Auger S, Soutourina O, Danchin A, Martin-Verstraete I. Global control of cysteine metabolism by CymR in *Bacillus subtilis*. J Bacteriol 2006;188:2184-97.
- [8] Fleischhacker AS, Stubna A, Hsueh KL, Guo Y, Teter SJ, Rose JC, et al. Characterization of the [2Fe-2S] cluster of *Escherichia coli* transcription factor IscR. Biochemistry 2012;51:4453-62.
- [9] Giel JL, Rodionov D, Liu M, Blattner FR, Kiley PJ. IscR-dependent gene expression links iron-sulphur cluster assembly to the control of O₂-regulated genes in *Escherichia coli*. Mol Microbiol 2006;60:1058-75.
- [10] Grundy FJ, Henkin TM. The T box and S box transcription termination control systems. Front Biosci 2003;8:20-31.
- [11] Gyaneshwar P, Paliy O, McAuliffe J, Popham DL, Jordan MI, Kustu S. Sulfur and nitrogen limitation in *Escherichia coli* K-12: specific homeostatic responses. J Bacteriol 2005;187:1074-90.
- [12] Heap JT, Pennington OJ, Cartman ST, Carter GP, Minton NP. The ClosTron: a universal gene knock-out system for the genus *Clostridium*. J Microbiol Methods 2007;70:452-64.
- [13] Heap JT, Pennington OJ, Cartman ST, Minton NP. A modular system for *Clostridium* shuttle plasmids. J Microbiol Methods 2009;78:79-85.
- [14] Ito S, Muramatsu T, Kobata A. Endo-beta-N-acetylglucosaminidases acting on carbohydrate moieties of glycoproteins: purification and properties of the two enzymes with different specificities from *Clostridium perfringens*. Arch Biochem Biophys 1975;171:78-86.
- [15] Johnston AW, Todd JD, Curson AR, Lei S, Nikolaidou-Katsaridou N, Gelfand MS, et al. Living without Fur: the subtlety and complexity of iron-responsive gene regulation in the symbiotic bacterium *Rhizobium* and other alpha-proteobacteria. Biometals 2007;20:501-11.
- [16] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 2001;25:402-8.
- [17] Meyer J. Clostridial iron-sulphur proteins. J Mol Microbiol Biotechnol 2000;2:9-14.

- 517 [18] Miller HK, Auerbuch V. Bacterial iron-sulfur cluster sensors in mammalian pathogens.
518 Metallomics 2015;7:943-56.
- 519 [19] Myers GS, Rasko DA, Cheung JK, Ravel J, Seshadri R, DeBoy RT, et al. Skewed
520 genomic variability in strains of the toxigenic bacterial pathogen, *Clostridium perfringens*.
521 Genome Res 2006;16:1031-40.
- 522 [20] Myers KS, Park DM, Beauchene NA, Kiley PJ. Defining bacterial regulons using ChIP-
523 seq. Methods 2015;86:80-8.
- 524 [21] Nair S, Derre I, Msadek T, Gaillot O, Berche P. CtsR controls class III heat shock gene
525 expression in the human pathogen *Listeria monocytogenes*. Mol Microbiol 2000;35:800-11.
- 526 [22] Nesbit AD, Giel JL, Rose JC, Kiley PJ. Sequence-specific binding to a subset of IscR-
527 regulated promoters does not require IscR Fe-S cluster ligation. J Mol Biol 2009;387:28-41.
- 528 [23] Novichkov PS, Kazakov AE, Ravcheev DA, Leyn SA, Kovaleva GY, Sutormin RA, et al.
529 RegPrecise 3.0--a resource for genome-scale exploration of transcriptional regulation in
530 bacteria. BMC Genomics 2013;14:745.
- 531 [24] Ohtani K, Hirakawa H, Tashiro K, Yoshizawa S, Kuhara S, Shimizu T. Identification of a
532 two-component VirR/VirS regulon in *Clostridium perfringens*. Anaerobe 2010;16:258-64.
- 533 [25] Ohtani K, Takamura H, Yaguchi H, Hayashi H, Shimizu T. Genetic analysis of the *ycgJ*-
534 *metB-cysK-ygaG* operon negatively regulated by the VirR/VirS system in *Clostridium*
535 *perfringens*. Microbiol Immunol 2000;44:525-8.
- 536 [26] Okumura K, Ohtani K, Hayashi H, Shimizu T. Characterization of genes regulated directly
537 by the VirR/VirS system in *Clostridium perfringens*. J Bacteriol 2008;190:7719-27.
- 538 [27] Ravcheev DA, Li X, Latif H, Zengler K, Leyn SA, Korostelev YD, et al. Transcriptional
539 regulation of central carbon and energy metabolism in bacteria by redox-responsive repressor
540 Rex. J Bacteriol 2012;194:1145-57.
- 541 [28] Rincon-Enriquez G, Crete P, Barras F, Py B. Biogenesis of Fe/S proteins and
542 pathogenicity: IscR plays a key role in allowing *Erwinia chrysanthemi* to adapt to hostile
543 conditions. Mol Microbiol 2008;67:1257-73.
- 544 [29] Romsang A, Duang-Nkern J, Leesukon P, Saninjuk K, Vattanaviboon P, Mongkolsuk S.
545 The iron-sulphur cluster biosynthesis regulator IscR contributes to iron homeostasis and
546 resistance to oxidants in *Pseudomonas aeruginosa*. PLoS One 2014;9:e86763.
- 547 [30] Rood JJ. Virulence genes of *Clostridium perfringens*. Ann Rev Microbiol 1998;52:333-60.
- 548 [31] Santos JA, Alonso-Garcia N, Macedo-Ribeiro S, Pereira PJ. The unique regulation of iron-
549 sulfur cluster biogenesis in a Gram-positive bacterium. Proc Natl Acad Sci U S A
550 2014;111:E2251-60.
- 551 [32] Santos JA, Pereira PJ, Macedo-Ribeiro S. What a difference a cluster makes: The
552 multifaceted roles of IscR in gene regulation and DNA recognition. Biochim biophys acta
553 2015;1854:1101-12.
- 554 [33] Schwartz CJ, Giel JL, Patschkowski T, Luther C, Ruzicka FJ, Beinert H, et al. IscR, an Fe-
555 S cluster-containing transcription factor, represses expression of *Escherichia coli* genes
556 encoding Fe-S cluster assembly proteins. Proc Natl Acad Sci U S A 2001;98:14895-900.
- 557 [34] Shimizu T, Ohtani K, Hirakawa H, Ohshima K, Yamashita A, Shiba T, et al. Complete
558 genome sequence of *Clostridium perfringens*, an anaerobic flesh-eater. Proc Natl Acad Sci
559 USA 2002;99:996-1001.
- 560 [35] Shimizu T, Shima K, Yoshino K, Yonezawa K, Hayashi H. Proteome and transcriptome
561 analysis of the virulence genes regulated by the VirR/VirS system in *Clostridium perfringens*. J
562 Bacteriol 2002;184:2587-94.
- 563 [36] Shimizu T, Yaguchi H, Ohtani K, Banu S, Hayashi H. Clostridial VirR/VirS regulon
564 involves a regulatory RNA molecule for expression of toxins. Mol Microbiol 2002;43:257-65.

- [37] Soutourina O, Poupel O, Coppee JY, Danchin A, Msadek T, Martin-Verstraete I. CymR, the master regulator of cysteine metabolism in *Staphylococcus aureus*, controls host sulfur source utilization and plays a role in biofilm formation. *Mol Microbiol* 2009;73:194-211.
- [38] Stülke J, Martin-Verstraete I, Zagorec M, Rose M, Klier A, Rapoport G. Induction of the *Bacillus subtilis ptsGHI* operon by glucose is controlled by a novel antiterminator, GlcT. *Mol Microbiol* 1997;25:65-78.
- [39] Tanous C, Soutourina O, Raynal B, Hullo MF, Mervelet P, Gilles AM, et al. The CymR regulator in complex with the enzyme CysK controls cysteine metabolism in *Bacillus subtilis*. *J Biol Chem* 2008;283:35551-60.
- [40] Tucker NP, Le Brun NE, Dixon R, Hutchings MI. There's NO stopping NsrR, a global regulator of the bacterial NO stress response. *Trends Microbiol* 2010;18:149-56.
- [41] Vinella D, Brochier-Armanet C, Loiseau L, Talla E, Barras F. Iron-sulfur (Fe/S) protein biogenesis: phylogenomic and genetic studies of A-type carriers. *PLoS Genet* 2009;5:e1000497.

Supplementary data

Figure S1. Genetic organization of the *cpe1786/iscR* locus and inactivation of the *cpe1786* gene.

The ClosTron delivery system (A) encoded on plasmid pMTL007 consists of a group II intron (black arrow) with an internal Retrotransposition-Activated Marker (RAM) conferring erythromycin resistance (white arrow) that is itself interrupted by the *td* group I intron (hatched box) [12]. The group II intron is retargeted to the *cpe1786* gene (grey arrow) by altering the sequence of the IBS/EBS region using overlap PCR. Splicing of the group II intron into *cpe1786* disrupts it and the splicing out of the *td* group I intron from the *erm* RAM restores a functional *ermB* gene allowing positive selection of *cpe1786* mutants. B) Analysis of genetic organization of the *cpe1786* locus by RT-PCR experiments. cDNA were synthesized using RNA extracted from cells grown in the presence of homocysteine. Couples of primers used for PCRs are indicated in panel A. Agarose gel corresponds to the migration of the different PCR fragments obtained from RNA without (lane 1,3,5) or with (lane 2,4,6) a reverse transcriptase treatment. Lane 1 and 2: IMV489-IMV490, lane 3 and 4: GA54-IMV491, lane 5 and 6: IMV492-493.

C) Confirmation of *cpe1786* knockout by PCR was performed using chromosomal DNA of *C. perfringens* CP13 (lane 1, 3 and 5) or an erythromycin resistant clones containing an intron in the *cpe1786* gene (lane 2, 4 and 6) and the primer pairs RAM-F/RAM-R (lane 1 and 2), IMV485/EBSu (lane 3 and 4) and IMV485-IMV484. PCR products were visualized on 1% w/v agarose gel.

Figure S2. Alignment of CymR and IscR-type regulators. The alignments were performed using the CLUSTALW algorithm. SA: *Staphylococcus aureus*; BS: *B. subtilis*; CA: *C. acetobutylicum*; CP: *C. perfringens*; CD: *C. difficile*; TP: *Thermincola potens*; EC: *E. coli*; ERWCT: *Erwinia carotovora*. The winged helix-turn-helix (HTH) motif is indicated by a blue

arrow and the amino acids shown to interact with DNA of type 2 sites in *E. coli* are underlined. The Glutamate at position 43 proposed to be involved in discrimination between type 2 and type 1 site is indicated in blue. The region containing the Fe-S binding motif corresponds to the red arrow. The 3 conserved cysteine residues coordinating the Fe-S cluster are indicated in red while the histidine residue involved in Fe-S coordination in *E. coli* is indicated in green [8].


Figure S3. The *iscR* and *cymR* operons. **A.** Comparison of the genetic organization of the *cpe1786* operon of *C. perfringens* with that of the *cymR* operon of *B. subtilis* and the *iscR* operons of *E. coli* and *T. potens*. Genes encoding regulators are indicated by blue arrows. The % of identity between each regulator and Cpe1786 is indicated. **B.** IscR binding motif. Alignment of IscR sites experimentally validated in *E. coli* [9], *T. potens* [31] and *C. perfringens* (this work). The -35 boxes are underlined. The promoters have been mapped for *iscR* and *yadR* of *E. coli* [9] and *iscR* of *C. perfringens* (Fig 1).

Figure S4. Mapping of *ldh* and *cpe0664* promoters by 5'RACE. The promoters region of *ldh* (*cpe0103*) and *cpe0664* encoding an A-type scaffold protein for Fe-S assembly are presented. The ATG of *ldh* and *cpe0664* are indicated in green. The ribosome bonding sites (RBS) are underlined and the -10 and -35 corresponding to SigA consensus are boxed. The IscR binding motifs identified using Reg-Predict [23] are indicated by blue arrows. In the case of *cpe0664*, a second motif with a score <5 is also indicated.

Table S1. List of oligonucleotides

Table S2. Genomic identification of candidate IscR-binding sites in *Clostridia* spp.

Figure
[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)





Figure S1
[Click here to download high resolution image](#)






Figure S2
[Click here to download high resolution image](#)





Figure S3

[Click here to download high resolution image](#)



B

iscR E.C ATAGTTGACCAATTACTCGGGAAT
iscR T.P ATAGTTGACCAATTACTCGGGAAT
iscR C.P ATTGTTGACAAATTACTCGGTTTT
cpe2093 AAAGTTGACAATTTTGATGGAATT
 AWWGTTGACMAWWTRMTSGGNWWT

Figure S4
[Click here to download high resolution image](#)




Table S1. Oligonucleotides used in this study.

Name	sequence	function
Quantitative RT-PCR		
GA99	TGCAGGATTTGTTGGTTCAA	5' <i>ldh</i>
GA100	GCACCTTGTGCTAGGTCCAT	3' <i>ldh</i>
GA103	GGTAGCAATGAGCGATGATG	5' <i>cpe0664</i>
GA104	TCCGTTCCCAACAAAGTTTA	3' <i>cpe0664</i>
GA105	CCTGGCTCTTTTGAAGGTGT	5' <i>cpe1371</i>
GA106	ACAACCATACCGCATCCATT	3' <i>cpe1371</i>
GA108	TCCCAATGTTACCCTTTTGA	5' <i>cpe2092</i>
GA109	TTGCAAAGCTCATTTCATGG	3' <i>cpe2092</i>
GA110	CTCCTTACAAGGCACCAGGA	5' <i>cpe0818</i>
GA111	CCCTTTGGCTACTGGAACAC	3' <i>cpe0818</i>
GA112	ATGCCTGAAACAATGCCTTC	5' <i>cpe0866</i>
GA113	TTGATCCCAATCCCAGAAAG	3' <i>cpe0866</i>
GA114	TGGTACTGCTCCAAC TTTCG	5' <i>cpe1031</i>
GA115	TTCCACCAACAAC TACAGCATC	3' <i>cpe1031</i>
GA124	GAGGGAAACATTGGGGTTTT	5' <i>crt</i>
GA125	ATCAGCACCTGCAACAAATG	3' <i>crt</i>
GA126	TCAAAAGCAAGCTGAGTGGA	5' <i>nagL</i>
GA127	CATTGTTGCATGTCCTGTCC	3' <i>nagL</i>
GA30	TGCCAGAATAGTTGGGGAAG	5' <i>gyrA</i>
GA31	TACCATGTCCGTCAACAAGC	3' <i>gyrA</i>
5'RACE		
GA130	TTCATAGTCGCTCATTAATGGAT	5'RACE <i>cpe0664</i>
GA131	AGAAGAATCAATAACAAAAGTTAA	5'RACE <i>cpe0664</i> int
GA132	TACTAAAAGTATTGAGTTTGGGC	5'RACE <i>ldh</i>
GA133	GTTATTATAACTATGTCAGAG	5'RACE <i>ldh</i> int
GA51	TCCTCCTTGAGCACCTCTAA	5'RACE <i>cpe1786</i>
GA62	GTCTAGTCGCACAACAA	5'RACE <i>cpe1786</i> int
Plasmid constructions		
pMTL007-F	TTAAGGAGGTGTATTTTCATATGACCATGATTACG	intron variable region
pMTL007-R	AGGGTATCCCCAGTTAGTGTTAAGTCTTGG	intron variable region
RAM-F	ACGCGTTATATTGATAAAAAATAAATAGTGGG	Erm cassette
RAM-R	ACGCGTGCGACTCATAGAATTATTTCTCCCG	Erm cassette
EBSu	CGAAATTAGAACTTGCGTTCAGTAAAC	intron internal
TBD043	AAAAAAGCTTATAATTATCCTTAGGATACGATACTGTGCGC CCAGATAGGGTG	IBS <i>cpe1786</i>
TBD044	CAGATTGTACAAATGTGGTGATAACAGATAAGTCGATACT GGTAACTTACCTTTCTTTGT	EBS1d <i>cpe1786</i>
TBD045	TGAACGCAAGTTTCTAATTTTCGATTATCCTCGATAGAGGA AAGTGTCT	EBS2 <i>cpe1786</i>
IMV484	GACAAATTTACTCGGTTTTGAT	5' <i>cpe1786</i>
IMV485	CTCTATTATTCATTTTCTTCACTC	3' <i>cpe1786</i>
IMV390	GGCGAATTCGTAGTATAAAAGCCTACTGG	<i>cpe1786</i> -P-131 EcoRI
IMV391	GGCGAATTCCTTATTGTTGACAAATTTACT	<i>cpe1786</i> P-41 EcoRI
IMV394	GGCGGATCCCTTCCTTTGGTAGATAA	<i>cpe1786</i> P+61 BamHI
IMV439	TTGTTGACAAATTTACGCGGTTTGTGATATTATA	T-22G P- <i>cpe1786</i>
IMV440	TATAATATCAAAACCGCGTAAATTTGTCAACAA	T-22G P- <i>cpe1786</i>
IMV441	TTGACAAATTTACTCGATTTTGTGATATTATAATC	G-19A P- <i>cpe1786</i>
IMV442	GATTATAATATCAAAATCGAGTAAATTTGTCAA	G-19A P- <i>cpe1786</i>
IMV455	TTGTTGACAAATTTACGCGATTTTGTGATATTATAATC	T-22G/G-19A P- <i>cpe1786</i>
IMV456	GATTATAATATCAAAATCGCGTAAATTTGTCAACAA	T-22G G-19A P- <i>cpe1786</i>
EH18	AAATCTAGATCACTACTTG7CA7CGTCATCCTTGTAAGTCGATGTCATGATCT TTATAATCACCGTCATGGTCTTTGTAGTCTTCATTTCTTCACTCCTACT	Fusion of a XFlag at the end of <i>cpe1786</i>
ChIP-PCR		
IMV524	ACGAAATAAAAAATCAAAGTTGACAAT	<i>cpe2093</i> ChIP-PCR

IMV525	TTGCTACAACCAACTAATACCATCA	<i>cpe2093</i> ChIP-PCR
IMV520	GAAATTTCCATTTTAAATTCCTACTAAAA	<i>cpe0664</i> ChIP-PCR
IMV521	AGCATCATCGCTCATTGCTA	<i>cpe0664</i> ChIP-PCR
EH40	TTGACAAATTTACTCGGTTTTGA	<i>cpe1786</i> ChIP-PCR
EH41	ATCAACCATGGCTTTTACAC	<i>cpe1786</i> ChIP-PCR
EH44	GAGGTGGCTTATGGAAGATAAAA	<i>cpe1031</i> ChIP-PCR
EH45	CTAGCTATTCCCTTTGCACACA	<i>cpe1031</i> ChIP-PCR
EH50	GTGTTGGTGCCGCTTTAGG	<i>cpe1371</i> ChIP-PCR
EH51	GACCTGTAGTTCCAAGTATAAAAACAA	<i>cpe1371</i> ChIP-PCR

Table S2. Genomic identification of candidate IscR-binding sites in Clostridia spp.

Genome	First gene locus tag	regulated operon	Site Position	Site Score	Candidate IscR-binding site
Clostridium perfringens str. 13	CPE1786	iscRSU-trmU	-80	5.8	TTATTGTTGACAAATTTACTCGGTTTTGA
			-39	5	ATAAAGTACATCGAATTAGTCAGATTTAG
	CPE0103	ldh	-46	5.5	ATAATTCAGAGTAATTAAGTAGGAATTAG
	CPE0664	hesB	-101	5.3	TAAATTCCTACTAAAATAGTCTTGATTTA
			-76	4.8	TTATTTTGTTTAAAAATTATAAATTTAA
	CPE2093	yxeONM	-69	5.3	TCAAAGTTGACAATTTTGATGGGAATTAA
Clostridium acetobutylicum ATCC 824	CAC1675	iscR	-41	5.8	ACAAAGTAGAGTATTATGTGCGGAATTAA
	CAC3288	sufCBDSE	-58	5.4	TGAATTCGACTTAAATGGTCAAGATTAA
Clostridium novyi NT	NT01CX_228	iscRSU-trmU	-82	5.9	TTAATGTTGACAAAAACACTCGGATTGA
			-41	5.6	AGAATTCATAGTATAAACTCGGATTAA
Clostridium kluyveri DSM 555	CKL_1317	iscRSU	-71	5.8	TTATTGTTGACAAGTTTACTCTGATTGA
	CKL_0828	sufCBDSE	-41	5.1	TTAAATATGAGTCAAATACTCGGATTAA
Clostridium butyricum 5521	CBY_2622	iscRSU-trmU	-61	5.3	CAATAGCCGAGTATTTTGATAAGATATAA
Clostridium beijerincki NCIMB 8052	Cbei_1097	iscRSU-trmU	-105	5.7	CAAAAGCCGAGTATTTTGGTAAGATATAA
	Cbei_4356	cysK	-55	5.2	TTATAACATAGTAAACTGATAAGAATTAT
Clostridium tetani E88	CTC01049	iscRSU-trmU	-92	5.2	TTATAGTTGACAGGAATACTCGGATTGT
			-52	5.5	TGAAAGTTTAGTAAAGCAGTAGACTTTCA
Clostridium botulinum A str. ATCC 35 CBO2569		iscRSU	-82	4.8	ATATTGTTGACAGGTTTAGTCTGATTGTC
			-43	6	ATAAATCATAGTAAAGTGGTCAACATTAA
Clostridium bartlettii DSM 16795	CLOBAR_021	iscRSU-trmU	-113	6.6	TTAATTCTGACTAAAATAGTCAAAATTAA
			-88	5.7	TTAATGTTGACAAAAACACTCAAGTATAA
			-41	4.7	AGAAATCCTACTAAAAAGTAGGAATAAG
Clostridium hiranonis DSM 13275	CLOHIR_011	iscRSU-trmU	-103	5.9	ACAAATCATACTAAAATAGTAGGAATATAA
			-77	5.7	TAATACTTGACAAAAATAGTCGGATATAA
			-43	6	TTAAACCATAGTAAACAGTAGGAATTAA
Clostridium difficile 630	CD1278	iscRSU-trmU	-98	6	TTAATCTTGACAAAATTAGTAGGGTATAA
			-53	5.4	GTAATTCCTACTAAACAGTATGAATTAG
Clostridium sp. OhILAs	Clos_1671	iscRSU	1	5.2	TGAATGTTGACAAAAATAGTAGGGTATAC
	Clos_0333	cysK	-43	5	ATAAAACCTATCGGTTTACTATGAATTAA
Clostridium cellulolyticum H10	Ccel_1933	iscRSU-trmU	-93	6.1	TTAAATCATAGTAATTTAATTGGAATTAA
	Ccel_3186	cysK	-106	5.2	TTAATTCATACTAATTAGATAGGAAATAT
Clostridium nexile DSM 1787	CLONEX_030	iscRSU-trmU	-51	5.9	ATAAACCTAGTAATTCAGTAGGAATTAA
Clostridium scindens ATCC 35704	CLOSCI_0139	iscRSU-trmU	18	5.7	AGAAATCCTAGCAAATTGCTAGGAATTAA
Clostridium sp. SS2/1	CLOSS21_00	iscRSU-trmU	-38	5.4	TGAAATCCGAGTTATATAGTCGGGATTGA
Thermincola potens JR	TherJR_1914	iscRSU-trmU	-120	5.2	AAAAATCTTAGTATTTTAGTTGGAATTTT
			-50	4.8	CCATAACCGAGTGCAACACTAGGAATTAA