



**HAL**  
open science

## A Eukaryotic-like Serine/Threonine Kinase Protects Staphylococci against Phages

Florence Depardieu, Jean-Philippe Didier, Aude Bernheim, Andrew Sherlock, Henrik Molina, Bertrand Duclos, David Bikard

► **To cite this version:**

Florence Depardieu, Jean-Philippe Didier, Aude Bernheim, Andrew Sherlock, Henrik Molina, et al.. A Eukaryotic-like Serine/Threonine Kinase Protects Staphylococci against Phages. *Cell Host and Microbe*, 2016, 20 (4), pp.471 - 481. 10.1016/j.chom.2016.08.010 . pasteur-01402430

**HAL Id: pasteur-01402430**

**<https://pasteur.hal.science/pasteur-01402430>**

Submitted on 24 Nov 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# **A eukaryotic-like Serine/Threonine kinase protects Staphylococci against viruses**

Florence Depardieu<sup>1</sup>, Jean-Philippe Didier<sup>2\*</sup>, Aude Bernheim<sup>3, 1</sup>, Andrew Sherlock<sup>4</sup>, Henrik Molina<sup>5</sup>, Bertrand Duclos<sup>2</sup> and David Bikard<sup>1</sup>

## **Affiliations**

<sup>1</sup>Synthetic Biology Group, Microbiology Department, Institut Pasteur, Paris, 75015, France.

<sup>2</sup>Institute of Molecular and Supramolecular Chemistry and Biochemistry, University of Lyon-CNRS, Villeurbanne, 69100

<sup>3</sup>AgroParisTech, Paris, 75005, France.

<sup>4</sup>Laboratory of Bacteriology, The Rockefeller University, New York, NY 10065, USA.

<sup>5</sup>Proteomics Resource Center, The Rockefeller University, New York, NY 10065, USA.

**Co-corresponding authors:** David Bikard, david.bikard@pasteur.fr, Bertrand Duclos, bertrand.duclos@univ-lyon1.fr

**Lead Contact:** David Bikard

**\*Present address:** Unilabs SA, – 1296 Coppet, Switzerland

## **Summary**

Organisms from all domains of life are infected by viruses, the most abundant biological entities on earth. In eukaryotes, Serine/Threonine kinases play a central role in antiviral response. Bacteria, however, are not commonly known to use protein phosphorylation as part of their defense against phages. Here we identify Stk2, a Serine/Threonine kinase that provides efficient immunity against bacteriophages in staphylococci through abortive infection. A phage protein of unknown function activates the kinase activity of Stk2 when expressed in the cell. This leads to the phosphorylation of several proteins involved in translation, global transcription control, cell cycle control, stress response, DNA topology, DNA repair and central metabolism. Cells die as a consequence of Stk2 activation, thereby preventing phage propagation to the rest of the bacterial population. Our work shows that mechanisms of viral defense that rely on protein phosphorylation constitute an antiviral strategy conserved across domains of life.

## 1 Introduction

2 The arms race between bacteria and phages has led to the evolution of many bacterial defense  
3 systems that can act at every stage of the phage life cycle, blocking phage adsorption, DNA  
4 injection, degrading phage DNA, and interfering with phage replication or the production of  
5 phage proteins (Labrie et al., 2010). These defense systems are mechanistically diverse and  
6 can vary considerably among bacterial species or even among different isolates of a particular  
7 species. At a glance, bacterial defense against phages has little in common with eukaryotic  
8 antiviral systems. In plants, defense is primarily conducted via RNA-mediated interference  
9 (RNAi), while in vertebrates, pattern-recognition receptors (PRR) recognize nucleic acids and  
10 proteins from pathogens and activate the interferon, proinflammatory and adaptive immune  
11 responses (Kanneganti, 2010; Palm and Medzhitov, 2009; Pumplin and Voinnet, 2013; Sadler  
12 and Williams, 2008). Serine/Threonine kinases (STKs) play a critical role at different stages  
13 of the antiviral response in both plants and vertebrates. They behave as switches that are  
14 activated by phosphorylation of one or several residues in an activation loop (Huse and  
15 Kuriyan, 2002). Some STKs, such as the interferon-induced, double-stranded RNA-activated  
16 protein kinase (PKR) in mammals, can also directly sense and interfere with viruses (Yan and  
17 Chen, 2012). Upon activation by dsRNA, PKR phosphorylates the translation initiation factor  
18 eIF2 $\alpha$ , blocking translation and viral protein synthesis. A similar mechanism was also recently  
19 described in plants. The NIK1 STK of *Arabidopsis* was shown to phosphorylate the ribosomal  
20 protein L10 and globally suppress translation as an antiviral immunity strategy (Zorzatto et  
21 al., 2015).

22  
23 STKs were assumed for a long time to exist only in eukaryotes, but Eukaryotic-like STKs  
24 (eSTKs) have now been found in most bacterial clades where they have been implicated in a  
25 variety of functions including cell cycle control, exit of dormancy, cell wall synthesis, cell  
26 division, control of the central metabolism and virulence (Pereira et al., 2011). Unlike  
27 eukaryotes, bacteria are generally not known to use STKs in viral defense. An exception to  
28 this is the *pgl* phage defense system from *Streptomyces coelicolor* and the related BREX  
29 (Bacteriophage Exclusion) systems, which are thought to exist in many unrelated bacteria  
30 (Goldfarb et al., 2015; Hoskisson et al., 2015; Sumbly and Smith, 2002). These systems  
31 include a STK known as PglW for which kinase activity was confirmed *in vitro*; however, its  
32 exact role in the defense pathway is not yet understood (Hoskisson et al., 2015). Also of note  
33 is the discovery of a prophage-encoded tyrosine kinase which excludes superinfection by  
34 phage HK97 in *Escherichia coli* (Friedman et al., 2011).

35  
36 Here we report the discovery of an eSTK involved in phage defense in *Staphylococci*. The  
37 Stk2 protein is activated when a specific phage protein, PacK, is present in the cell. The  
38 activation of Stk2 results in cell death through phosphorylation of proteins involved in essential  
39 cellular processes, including translation, transcription, control of cell cycle and others.  
40 Infected cells die before releasing infectious phages thereby protecting neighboring bacteria.  
41 This altruistic defense strategy is known as abortive infection (Abi) and can be performed by  
42 mechanically diverse systems (Chopin et al., 2005). *Staphylococci* carry another STK known as  
43 Stk1 or PknB, which is important for cell-wall structure, antimicrobial resistance, and  
44 virulence (Beltramini et al., 2009; Debarbouille et al., 2009; Donat et al., 2009; Truong-  
45 Bolduc et al., 2008). The existence of a second STK, known as Stk2, present in only some  
46 isolates of *S. aureus* was noted in a few studies but its function remained  
47 mysterious (Debarbouille et al., 2009; Didier et al., 2010). We now demonstrate that Stk2  
48 provides immunity against bacteriophages through an Abi mechanism. Interestingly, the Stk1  
49 kinase is also involved in this defense pathway, suggesting a phosphorylation cascade  
50 reminiscent of Eukaryotic viral defense pathways.

51

## 52 **Results**

53

54 **Discovery of *stk2*, a bacteriophage defense kinase.** We isolated a novel temperate phage of  
55 *Staphylococcus epidermidis*, CNP<sub>x</sub> a 43kb Siphoviridae with 90.2% overall nucleotide  
56 identity to phage CNPH82 (Daniel et al., 2007). CNP<sub>x</sub> was isolated on strain LM1680, a  
57 derivative of *S. epidermidis* RP62A carrying a large deletion that includes a type III-A  
58 CRISPR system and a type I restriction modification (RM) system (Hatoum-Aslan et al.,  
59 2014). Interestingly, CNP<sub>x</sub> does not infect strain RP62A, suggesting that the region deleted in  
60 LM1680 contains a defense system, possibly the CRISPR or the type I RM, that blocks  
61 infection by this phage. To narrow down the position of the defense system, we tested the  
62 ability of CNP<sub>x</sub> to infect a collection of RP62A derivatives, obtained by Marraffini and  
63 colleagues, that carry various deletions of this region (Jiang et al., 2013). This analysis  
64 allowed us to identify a ~16kb candidate region that carries the phage defense system (Figure  
65 1A). To our surprise, this region did not include the type I RM system or the CRISPR system;  
66 instead, it contained a number of hypothetical proteins as well as an operon involved in  
67 potassium transport and associated regulatory genes (Table S1). Genes coding for the  
68 hypothetical proteins were cloned either alone or two at a time on plasmid pC194 which is  
69 present in ~15 copies in the cell (Novick, 1989). The resulting plasmids (pDB31, pDB32,  
70 pDB33 and pDB34) were introduced by electrotransformation in strain LM1680, and the  
71 bacteria were tested for sensitivity to phage CNP<sub>x</sub>. The pDB31 plasmid carrying gene  
72 SERP2479 provided strong resistance (EOP < 10<sup>-5</sup>) while the other genes carried by plasmids  
73 pDB32, pDB33 and pDB34 did not have any effect on the susceptibility of the strain to the  
74 phage (Figure 1B).

75

76 SERP2479 contains a serine/threonine kinase domain that is easily identified by prediction  
77 algorithm such as hhmer or CD-search (Figure 2A) (Finn et al., 2011; Marchler-Bauer and  
78 Bryant, 2004). Proteins identical to SERP2479 are found in several strains of *Staphylococcus*  
79 *aureus* (Figure 1C). In particular, protein SA0077 whose sequence is 100% identical to  
80 SERP2479 was previously described in *S. aureus* strain N315. It was named Stk2 and shown  
81 to phosphorylate the virulence regulator SarA *in vitro*, but could not be assigned a  
82 physiological role (Didier et al., 2010). Closely related proteins are also present in more  
83 distantly related Firmicutes, including some *Bacilli* and *Streptococci* (Table S2). In addition  
84 to the kinase domain, a distinctive feature of these proteins is the presence of large N-term  
85 and C-term domains of unidentified fold or function.

86

87 **Stk2 provides phage resistance in staphylococci.** To investigate the scope and significance  
88 of our finding, we decided to test whether *stk2* could also block phage infection in *S. aureus*.  
89 The pDB31 plasmid, carrying *stk2* under the control of its wild-type promoter, was introduced  
90 by electrotransformation in several *S. aureus* strains (TB4, NCTC8325-4 and RN4220) which  
91 do not carry *stk2* in their chromosome (Bae et al., 2006; Nair et al., 2011). Stk2  
92 provided strong resistance to phage phiNM1 in all backgrounds (Table S3). Strain RN4220 is  
93 a derivative of *S. aureus* strain NCTC8325 that is easy to manipulate but is generally not  
94 considered to be a good model strain for the study of bacterial virulence (Nair et al., 2011).  
95 Nonetheless, since the phage resistance phenotype provided by *stk2* was identical in all tested  
96 strains, we decided to use RN4220 for the remaining of this study. We tested the ability of  
97 plasmid pDB31 to confer resistance to infection by different phages in the RN4220  
98 background, including five Siphoviridae (phage 80alpha, phage 85, phiNM1, phiNM2 and  
99 phiNM4) and one Twort-like Myoviridae (phage Staph1N) (Bae et al., 2006; Lobočka et al.,

100 2012). Stk2 provided resistance against all tested Siphoviridae but not against the Twort-like  
101 phage (Table S3).

102

103 **Characterization of the Stk2 kinase activity.** The *stk2* gene was cloned with a 6x His N-  
104 terminal tag in plasmid pET15b and introduced in *E. coli* BL21 (DE3). Upon induction with  
105 IPTG, efficient overproduction of His6-Stk2 fusion protein was obtained, though in the form  
106 of inclusion bodies. The His6-Stk2 product was then purified by denaturation/renaturation  
107 method using guanidinium chloride, followed by a step of purification on an affinity column.  
108 Finally, the linked His6 was removed through proteolysis by thrombin (Figure  
109 2B). Autophosphorylation activity was tested in the presence of various divalent cations:  
110 Mn<sup>2+</sup>, Mg<sup>2+</sup>, Ca<sup>2+</sup>, Zn<sup>2+</sup> and Co<sup>2+</sup> (Figure 2C). It was observed that purified Stk2 was  
111 significantly labelled *in vitro* in the presence of [ $\gamma$ -<sup>32</sup>P] ATP and Mn<sup>2+</sup> (Figure 2C, lane 4).  
112 The ability of Stk2 to autophosphorylate in these conditions indicates that it displays intrinsic  
113 kinase activity. To exclude the possibility of contamination by an exogenous kinase that  
114 might interfere with the phosphorylation of Stk2, the invariant lysine 152 involved in the  
115 binding of the ATP phosphoryl donor, was mutated to isoleucine. As expected Stk2-  
116 K152I could no longer autophosphorylate (Figure 2B, lane 5).

117

118 The phosphoamino acid content of the labelled protein was determined after acid hydrolysis  
119 and two-dimensional analysis (Duclos et al., 1991). Both phosphoserine and  
120 phosphothreonine were revealed on the corresponding autoradiogram (Figure 2D), indicating  
121 that Stk2 was modified exclusively on these two types of residues. NanoLC/nano-  
122 spray/tandem mass spectrometry was then used for the identification of phosphorylated  
123 peptides and for the localization of the phosphorylation sites in Stk2 (Molle et al., 2006). Nine  
124 phosphorylation sites could be identified including three sites in the activation loop of Stk2  
125 (S272, T275 and T278) (Figure 2A). Various mutated proteins were produced and purified,  
126 and the effect of substitution of the different residues to alanine on kinase activity was  
127 analyzed by measuring autophosphorylation activity (Figure 2E). Kinase activity was  
128 completely abolished when the substitution was on T147, S272, T275, and T278. In contrast,  
129 substitution of S246 had no effect on the activity of Stk2, and replacement of S163, S337 and  
130 S360 unexpectedly increased Stk2 activity. Altogether these results show that Stk2 is indeed  
131 an active STK.

132

133 To confirm *in vivo* that the kinase activity of Stk2 is required for the phage defense phenotype,  
134 we mutated conserved residues: K152 in the ATP-binding loop, the predicted catalytic  
135 aspartate D243, and the T275 autophosphorylation residue in the activation loop (Figure 2A).  
136 All alleles were introduced in *S. aureus* strain RN4220 and tested for sensitivity to phage  
137 phiNM1 (Figure 3A). As expected, all mutants showed sensitivity to the phage.

138

139 **Stk2 triggers cell death.** Several assays were performed to understand the mechanism of  
140 protection provided by Stk2. We first tested whether it could affect phage adsorption. Cells  
141 carrying plasmid pDB31 or the control pC194 were both able to adsorb 99% of the phiNM1  
142 phage particles. However, infection of growing cells carrying *stk2*(pDB31) by phage phiNM1  
143 led to an interruption in the growth of the culture (Figure 3B). This result stands in clear  
144 contrast to the lysis observed for cells that lack *stk2* and are sensitive to the phage. Consistent  
145 with this observation, ~3000x fewer phage particles are recovered after infection of cells  
146 carrying *stk2* relative to cells without *stk2* (Figure 3C). Also, measurement of the efficiency of  
147 center of infection (ECOI) in the presence of *stk2* indicates that only 0.4±0.08% of cells  
148 receiving the phiNM1 phage are able to release functional phiNM1 particles. Plating a culture  
149 of cells carrying *stk2* after infection reveals that most cells are dead as only ~5% of cells form

150 colonies (Figure 3C). This shows that *stk2* mediates cell death and acts as an abortive infection  
151 (Abi) system, killing bacteria upon infection to prevent phage propagation (Abedon, 2012).

152

153 It is worth noting that the number of bacteria that survive phiNM1 infection is similar in the  
154 presence and absence of *stk2* (Figure 3C). Wild-type RN4220 cells can survive phiNM1  
155 infection when the phage enters lysogeny and integrates in the genome. Interestingly, analysis  
156 of cells that survive phiNM1 infection in the presence of *stk2* revealed that some (3/8) had  
157 lysogenized phage phiNM1, while the remainder (5/8) most likely did not receive a phiNM1  
158 phage particle, eliminated the phage without dying or mutated its receptor (Figure S1A).

159 These results suggest that *stk2* kills staphylococci only if the phage enters its lytic cycle. To  
160 confirm this, we sought to induce the lytic cycle of phage phiNM1 lysogenized in cells  
161 carrying *stk2* or a control plasmid (Figure 3D). Prophages were induced with mitomycin C. In  
162 the presence of *stk2*, the culture stopped growing but did not lyse. In agreement with this  
163 observation,  $3 \times 10^4$ -fold fewer phage particles were recovered after induction of cells carrying  
164 *stk2* compared to cells carrying the control plasmid (Figure 3D). Note that in the absence of  
165 mitomycin C, phage phiNM1 is spontaneously induced at a lower rate. Under these  
166 conditions, the presence of *stk2* also limits the number of phages released. Altogether, these  
167 experiments demonstrate that presence of phage DNA is not recognized by Stk2; instead, the  
168 Abi phenotype of Stk2 is only triggered during the lytic cycle of the phage.

169

170 **Identification of the Stk2 activation factor.** In the pDB31 plasmid used here, *stk2* is  
171 expressed under the control of its WT promoter. To understand whether the transcriptional  
172 control of *stk2* is important for the Abi phenotype, we cloned *stk2* under the control of a  
173 Ptet promoter (Table S5). Resistance to phage was only observed upon induction of Stk2, and  
174 overexpression of Stk2 on its own did not lead to any growth defect (Figure S1B). These  
175 results clearly show that the Abi phenotype is only induced in the presence of the phage, and  
176 that the natural transcriptional control of *stk2* is not required. Thus, as expected, Stk2 likely  
177 behaves as a protein switch that is activated upon phage infection. While Stk2 is able to  
178 rapidly auto-phosphorylate *in vitro*, we believe that it is not active in the absence of phage  
179 infection *in vivo*.

180

181 To understand what might activate Stk2, we isolated phage mutants capable of  
182 infecting *S. aureus* strain RN4220 carrying pDB31. The efficiency of plaquing (EOP) of phage  
183 phiNM1 on cells carrying *stk2* is only  $5 \times 10^{-7}$  (Table S3). Nonetheless, some plaques can be  
184 recovered and propagated on cells expressing *stk2* (Figure S2A). Phages isolated in this way  
185 retain the ability to infect cells carrying *stk2* even after being passaged on cells lacking *stk2*,  
186 suggesting that the new phenotype is the result of mutation and not epigenetic variation.  
187 These phiNM1 mutants form small plaques and are harder to propagate than the wild-type  
188 phage (data not shown). Two mutant phages were sequenced (phiNM1k1 and phiNM1k2), and  
189 both carried independent miss-sense mutations (V200M and K30E) in the same gene,  
190 SAPPV1\_GP14. These results indicate that this protein likely activates Stk2. SAPPV1\_GP14  
191 contains a P-loop NTPase domain frequently found in proteins involved in molecular motion.  
192 An interesting observation enabled us to link the function of this gene to either DNA  
193 replication or packaging. When sequencing phage phiNM1k1 and phiNM1k2 we observed  
194 that coverage was highest shortly after the packaging site of the phage and then slowly  
195 dropped over the rest of the sequence (Figure 4A). This contrasts sharply with the wild-type  
196 phage which shows almost uniform coverage throughout the sequence. A possible explanation  
197 for this skewed coverage is that the capsids from which DNA was purified do not all contain  
198 the full phage genome; instead, most capsids only contain the part of the phage genome that is  
199 packaged first. Phage particles which contain only part of the genome would likely not be

200 functional and might even lack the tail. As we did not purify intact phage particles before  
201 DNA extraction, we recovered DNA from both functional phage particles and any  
202 incompletely assembled particles present in our samples. Thus, the skewed coverage could be  
203 explained by random premature termination of phage DNA packaging. Such premature  
204 termination could either be due to a defect of the packaging machinery itself, or a defect of  
205 the phage DNA replication or concatemer formation. The position of SAPPV1\_GP14 in the  
206 replication cluster of the phage supports the later hypothesis.

207  
208 Because of its phenotype in DNA packaging, we decided to call the SAPPV1\_GP14 gene  
209 *pacK*. To confirm that this phage protein is sufficient to trigger Stk2, we cloned *pacK* under  
210 the control of an inducible Ptet promoter on plasmid pE194, giving plasmid pAS10 (Figure  
211 4B). Upon induction with anhydrotetracycline, cell death was observed only when *stk2* was  
212 present in the cells (Figure 4C). The V200M and K30E mutations identified in the mutant  
213 phages were also tested in this assay and abolished the Abi phenotype (Figure S2B). These  
214 experiments confirm that PacK is sufficient to trigger Stk2-mediated cell death.

215  
216 It is interesting to note that Stk2 can provide resistance to phages which do not carry PacK  
217 (see Tables S3 and S4), suggesting that it can be activated by other phage proteins. In  
218 particular, this is true for *S. epidermidis* phage CNP $\alpha$  which was used in this study to first  
219 identify Stk2 (Figure 1), as well as *S. aureus* phage 80 $\alpha$ , phage 85 and phiNM2 (Table  
220 S3). In phage phiNM2, gene AVT76\_GP14 encodes a protein with 43% identity to PacK, but  
221 no homologous proteins exist in phage 80 $\alpha$ , phage 85 or CNP $\alpha$ . To identify how these  
222 phages activate Stk2, we isolated mutants of phiNM2 and phage 85 capable of infecting *S.*  
223 *aureus* cells expressing *stk2*. Sequencing of these mutants revealed a H230T mutation in gene  
224 AVT76\_GP14 of phage phiNM2 and a K97G mutation in gene ST85ORF023 of phage 85. To  
225 confirm that these phage genes encode activators of Stk2, we cloned them under the control of a  
226 Ptet promoter on plasmid pE194, producing plasmids pFD16 and pFD20, respectively (Table  
227 S5). After induction with anhydrotetracycline, cells were killed in the presence of *stk2* but not  
228 in its absence (Figure 4C). The mutations identified in these genes were also confirmed to  
229 abolish Stk2-mediated cell death (data not shown). These results show that, in addition to  
230 PacK, two other phage proteins can activate Stk2.

231  
232 **Identification of Stk2 phosphorylation targets.** Our results suggest that Stk2 triggers an Abi  
233 phenotype through phosphorylation of one or several host proteins. To identify the  
234 phosphorylation target(s) of Stk2, we first characterized *S. aureus* colonies that survive the  
235 induction of *pacK* in the presence of Stk2 with the goal of identifying mutants of the target  
236 proteins. Unfortunately, all of the 36 analyzed colonies carried mutations either in the *stk2* or  
237 *pacK* genes, but no other mutant could be identified (data not shown). This result suggests  
238 that several mutations might be required to survive Stk2 activation; these would occur at a  
239 lower frequency than point mutations in *stk2* or *pacK*.

240  
241 We then performed a phosphoproteome analysis of cells expressing *pacK* in either the  
242 presence or absence of *stk2*. Expression of PacK was induced from plasmid pAS10 (Ptet-  
243 *pacK*) in exponentially growing cells. After 30 min of induction, proteins were precipitated  
244 and digested followed by titanium dioxide-based phosphopeptide enrichment (Larsen et al.,  
245 2005). To confidently differentiate basal and Stk2-induced phosphorylation events, we  
246 labeled the different proteomes with mass spectrometry-differentiable stable isotopes of  
247 dimethyl (Boersema et al., 2009). We identified 32 phosphopeptides that could only be found  
248 in the presence of Stk2 (Table 1). These include several proteins related to translation,  
249 including elongation factors Tu and P, 50S ribosomal proteins L6, L5, L31 and the MetG

250 methionine-tRNA ligase. This extensive phosphorylation of the translation machinery likely  
251 indicates that translation is modified after Stk2 activation. In addition, we identified proteins  
252 involved in global transcription control, cell cycle control, stress response, DNA topology,  
253 DNA repair and central metabolism. This suggests a coordinated response influencing many  
254 aspects of the cellular machinery, and a general shift towards stress response and growth  
255 arrest. Phosphopeptides corresponding to Stk2 itself could also be identified. Three residues  
256 are phosphorylated in the activation loop (S272, T275, T278), as well as a serine S163  
257 between the P-loop and the catalytic site. These residues were also identified in the *in*  
258 *vitro* autophosphorylation assay (Figure 2). Phosphorylation of the trigger PacK protein could  
259 also be identified at residue S176, suggesting that Stk2 interacts with PacK directly. Mutation  
260 of the S176 residue to alanine did not have any impact on the Abi phenotype (Figure S3).

261  
262 **Role of Stk1 in the Stk2-mediated immunity.** The question of whether the phosphopeptides  
263 identified are directly phosphorylated by Stk2 remains to be investigated. Indeed, it is possible  
264 that the activation of Stk2 results in the activation of Stk1 which would lead to secondary  
265 phosphorylation events. For instance, the elongation factor P identified in our analysis was  
266 previously reported as a target of Stk1 (Lomas-Lopez et al., 2007). In further support of this  
267 possible role of Stk1 in the phage defense phenotype, we detected the phosphorylation of Stk1  
268 at two threonines in the activation loop (T164 and T166). The phosphopeptide carrying these  
269 residues was 1.5-fold more abundant in the presence of Stk2 than in its absence (Table S7).  
270 To investigate whether Stk1 could play a role in Stk2-mediated phage defense, plasmid  
271 pDB31 carrying *stk2* was introduced by electrotransformation in *S. aureus* strain NCTC8325-  
272 4 and in the corresponding *stk1* deletion mutant (Debarbouille et al., 2009). The EOP of phage  
273 phiNM1 on cells carrying both *stk1* and *stk2* is  $4.4 \times 10^{-6}$ , but when only *stk2* is present the  
274 EOP jumps to  $2 \times 10^{-2}$  (Figure 5A). Thus, in the absence of *stk1*, we can still observe some  
275 protective effect of *stk2*, but ultimately *stk1* is required for efficient immunity. We also  
276 investigated whether *stk2* could trigger cell death in the absence of *stk1*. *S. aureus* NCTC8325-  
277 4 and the *stk1* mutant were electrotransformed with both pDB31 (*stk2*) and pAS10 (Ptet-  
278 *pacK*) plasmids. Upon induction of PacK expression, cells were killed with identical  
279 efficiencies regardless of the presence of *stk1* (Figure 5B). This demonstrates that while *stk1*  
280 is necessary for efficient antiviral immunity, it is not required for Stk2-mediated cell death.

## 281 282 **Discussion**

283  
284 Recent bioinformatics analyses have led to the discovery that bacteriophage defense systems  
285 frequently cluster together in bacterial genomes (Makarova et al., 2011). Here we report the  
286 discovery of a novel defense system in close proximity to the type III CRISPR and type I RM  
287 system of *S. epidermidis* RP62A. SERP2479, or Stk2, is responsible for abortive infection and  
288 cell death upon phage infection. Stk2 belongs to the family of eukaryotic-like serine/threonine  
289 kinases but differs from previously described eSTKs in its lack of transmembrane or PASTA  
290 domains. We were able to confirm the kinase activity of Stk2 *in vitro* and identified nine auto-  
291 phosphorylated residues. Four of these residues were corroborated *in vivo*, including three in  
292 the activation loop (S272, T275, T278), as well as a serine (S163) close to the ATP-binding  
293 region (Figure 2A). All three residues of the activation loop are essential for *in*  
294 *vitro* autophosphorylation, while an S163A mutation actually increased the kinase activity,  
295 suggesting a regulatory role. The phosphorylation of several residues in the activation loop of  
296 eSTKs has been reported before and seems to be a common feature of these kinases (Young et  
297 al., 2003).

298

299 Our data suggest that while Stk2 is able to autophosphorylate *in vitro*, it is only activated in the  
300 presence of a phage protein *in vivo*. We identified three such phage proteins by analyzing the  
301 genomes of mutant phages able to propagate on cells carrying *stk2*: Gene SAPPV1\_GP14  
302 (*pacK*) from phage phiNM1, geneAVT76\_GP14 from phage phiNM2, a distant homologue of  
303 *pacK* with 43% protein identity, and gene ST85ORF023 from phage 85, which shows no  
304 identity to PacK. The PacK protein carries a P-loop NTPase domain and leads to a defect in  
305 phage DNA packaging when mutated. This defect could either be due to problems in DNA  
306 replication leading to DNA molecules in a bad conformation for packaging, or to a defect in  
307 packaging itself. It is currently identified in databases as the chromosomal replication initiator  
308 DnaA. However, we believe this to be a simple case of incorrect annotation, as no significant  
309 homologies can be found between PacK and DnaA proteins. The function of ST85ORF023 is  
310 not known, and no protein domain of known function can be identified. These activator  
311 genes are located within the phage lytic operon, which likely explains why Stk2-induced cell  
312 death is not triggered when the phage enters lysogeny (Figure 3D). However, the induction of  
313 a lysogenic phage in cells carrying *stk2* also leads to cell death. The ability of Stk2 to  
314 tolerate prophages while maintaining an active defense against the phage lytic cycle is  
315 reminiscent of the similar capacity of type III CRISPR systems to tolerate lysogenic phages  
316 (Goldberg et al., 2014).

317  
318 An analysis of Staphylococcal phages in the RefSeq database shows that 23% carry  
319 homologues of PacK, while 13% carry homologues of ST85ORF023 (Table S4). Phage CNP<sub>x</sub>,  
320 which was used in this study to first identify Stk2 in *S. epidermidis* carries a protein with 85%  
321 identity to ST85ORF023. Interestingly, a blast analysis only identified activators of Stk2 in  
322 Siphoviridae; none were found in other phage families. The ability of Stk2 to sense multiple  
323 phage proteins is fascinating, and leads to deeper questions about the regulation of the kinase  
324 activity.

325  
326 We hypothesize that the activation of Stk2 by PacK is the result of a direct interaction  
327 between the two proteins, as PacK itself phosphorylated during the response. After sensing  
328 the presence of the infecting phage, the activation of Stk2 results in the phosphorylation of a  
329 large number of proteins involved in several core functions of the cell including translation,  
330 transcription and cell division (Table 1). The modulation of the activity of these proteins  
331 through phosphorylation is likely responsible for cell death, preventing phage propagation  
332 concurrently. It is worthwhile to note that while the SarA protein was previously reported as a  
333 phosphorylation target of Stk2, we did not identify it in our analysis (Didier et al., 2010). This  
334 could be explained by the fact that a serine-rich segment of SarA was not covered by our  
335 mass-spectrometry analysis (data not shown).

336  
337 Another STK known as Stk1 (sometimes also named PknB or PrkC), present in all  
338 Staphylococci, is also involved in this antiviral defense pathway. Deletion of *stk1* strongly  
339 impairs the ability of Stk2 to protect *S. aureus* against phages (Figure 5A). Nonetheless, the  
340 activation of Stk2 leads to cell death even in the absence of Stk1 (Figure 5B). Thus, the role  
341 of Stk1 in this defense pathway is likely to ensure that phage particles are not produced before  
342 cells are killed by Stk2. It remains to be investigated whether this occurs by accelerating cell  
343 death, slowing down the phage, or some other mechanism. The phosphoproteome analysis  
344 performed here does not allow for differentiation of direct targets of Stk2 from targets  
345 phosphorylated by Stk1 as a result of Stk2 activation. Future work will focus on elucidating  
346 the molecular interaction between Stk2 and PacK, for which we only provide circumstantial  
347 evidence, as well as deciphering the exact phosphorylation cascade occurring during the  
348 response.

349  
350 In sum, we provide strong evidence for a bacterial antiviral defense pathway involving a  
351 complex phosphorylation cascade and resulting in cell death through the modification of  
352 several essential cellular pathways. Other bacterial eSTKs have been shown to target different  
353 components of translation (EF-Tu, EF-P), transcription (various sigma and anti-sigma  
354 factors), cell division machinery (FtsZ) and central metabolism (Pereira et al., 2011), but none  
355 so far have been linked to phage defense. On the contrary, some phages have been described  
356 as using STKs in order to manipulate the host translation machinery for their own benefit  
357 (Robertson and Nicholson, 1992). It is also interesting to note that STKs play critical roles in  
358 the antiviral defense of eukaryotes. In particular, there are striking similarities between Stk2,  
359 the mammalian PKR and the plant NIK1. All three STKs are activated by viral infection and  
360 target the translation machinery. Moreover, PKR not only inhibits the initiation of translation  
361 through phosphorylation of eIF-2 $\alpha$  (Meurs et al., 1990), can also trigger cell death through  
362 apoptosis (Dai et al., 2012; Hsu et al., 2004; Stark et al., 1998). Viral defense strategies which  
363 involve the Serine/Threonine phosphorylation of essential cellular pathways thus exist in both  
364 eukaryotes and bacteria. eSTKs have also recently been identified in archaea (Kennelly,  
365 2014). In particular the Ph0512p kinase from *Pyrococcus horikoshii* OT3 was shown to  
366 phosphorylate the archaeal homolog of eIF2 $\alpha$  (aIF2 $\alpha$ ) *in vitro* (Tahara et al., 2004). It is  
367 tempting to hypothesize that Ph0512p and other archaeal kinases could also be involved in  
368 viral defense, making this a universal strategy conserved across all domains of life.

369

## 370 **Experimental Procedures**

### 371 **Bacterial strains and growth conditions**

372 *S. epidermidis* LM1680 (Hatoum-Aslan et al., 2014), *S. aureus* RN4220 (Nair et al., 2011) and  
373 derivative strains were grown in TSB media at 37°C with shaking at 200 rpm. *S. epidermidis*  
374 LM1680 and *S. aureus* RN4220 were used as hosts for recombinant plasmids.  
375 Chloramphenicol (10  $\mu$ g/mL), erythromycin (10  $\mu$ g/ml), and ampicillin (100  $\mu$ g/ml) were  
376 added to the medium to prevent loss of plasmids derived from pC194, pE194 (Horinouchi and  
377 Weisblum, 1982a, b), and pET15b (Novagen), respectively. *E. coli* BL21 (DE3) AD494  
378 (Novagen) was used for expression of recombinant proteins, and grown in LB medium  
379 supplemented with 100  $\mu$ g/mL ampicillin at 37°C.

380

### 381 **Isolation of phage CNP<sub>x</sub>**

382 Phage CNP<sub>x</sub> (GenBank accession KU598975) was isolated as a plaque on a soft-agar lawn of  
383 *S. epidermidis* LM1680 that was infected with phage CNPH82 (Daniel et al., 2007). LM1680  
384 is resistant to phage CNPH82 and the isolation of the CNP<sub>x</sub> was a single fortuitous event that  
385 might have occurred via contamination with an environmental phage and recombination with  
386 CNPH82. Indeed CNP<sub>x</sub> shares close to 100% homology with CNPH82 over 65% of its  
387 genome length, but carries a divergent segment of ~13 kb in the region of the genome  
388 containing the lysogenic operon and the early lytic genes.

389

### 390 **Introduction of plasmids in Staphylococci**

391 Plasmid constructions are detailed in the supplemental experimental procedures. Lists of  
392 plasmids and oligonucleotides used in this study are provided in supplementary Tables S5 and  
393 S6. After DNA assembly all plasmids were first electroporated in *S. aureus* strain RN4220.  
394 Briefly, cells were grown to an optical density (600nm) of 0.8 and washed three times in cold  
395 water and concentrated 100x in 10% glycerol. Electroporation of dialyzed DNA was  
396 performed in 2mm cuvettes using the following settings: 100 $\Omega$ , 2.5kV, 25 $\mu$ F. In order to  
397 introduce plasmids in other Staphylococci strains, plasmids were extracted from RN4220

398 using the NucleoSpin Plasmid kit (Macherey Nagel) with the following modification: 4ug of  
399 lysostaphin (Ambi) was added to the A1 buffer, and cells were incubated 1H at 37°C in this  
400 buffer before resuming the protocol as described. Plasmids extracted from RN4220 can then  
401 be introduced in other Staphylococci through electroporation following the same protocol.

402

### 403 **Overproduction and purification of Stk2 and derivatives**

404 Plasmids pET15Ω*stk2(sa0077)* and derivative mutants were introduced into *E. coli*  
405 BL21λDE3/AD494. The transformants were grown in 1 liter of LB medium with shaking at  
406 25°C until OD<sub>600</sub> = 0.5, IPTG (0.5 mM) was added to induce protein production, and  
407 incubation was pursued for 6 h at 25°C. Cells were then harvested by centrifugation at 3000g  
408 for 10 min. Since Stk2 and its mutants were not soluble and retained in inclusion bodies, a  
409 step of denaturation/renaturation using guanidine chloride according to London(London et al.,  
410 1974)and Goldberg (Goldberg et al., 1996)was performed before the purification on a Ni-  
411 NTA column.

412

### 413 **In vitro phosphorylation assay**

414 Phosphotransfer to Stk2 and its derivatives purified was performed in a buffer containing 25  
415 mM Tris-HCl, pH7.5, 1 mM DTT, 2.5 mM MnCl<sub>2</sub>, 10 mM ATP and 5 μCi of [γ-<sup>32</sup>P]-ATP  
416 and incubated at 37°C 10 min to 30 min following the substrate. The reaction was stopped by  
417 the addition of 20% Laemmli 5X (Sigma), followed by electrophoresis on SDS-PAGE and  
418 autoradiography.

419

### 420 **Phosphoamino acid analysis**

421 The method used to detect acid-stable phosphoamino acids was described previously (Duclos et  
422 al., 1991).

423

### 424 **Phage production**

425 Phages were mixed with *S. aureus* RN4220 in soft TSA agar supplemented with CaCl<sub>2</sub>  
426 (5 mM) and then poured on top of TSA agar plate supplemented with CaCl<sub>2</sub> (5 mM). The  
427 plates were incubated overnight at 37°C. Soft TSA agar lawns were then resuspended in PBS  
428 solution (1X), centrifuged and the lysate containing the phage was filtered on a 0.22 μm filter.  
429 To measure phage titers, serial dilutions were spotted on a soft agar lawn of RN4220 in TSA  
430 agar supplemented with CaCl<sub>2</sub> (5 mM) and PFUs were quantified after incubation overnight  
431 at 37°C.

432

### 433 **Efficiency of plaquing assays**

434 Phage lysates containing ~10<sup>7</sup> PFU/μl of CNPX, phage 80alpha, phage 85, phiNM1, phiNM2,  
435 phiNM4 or Staph1N were serially diluted and spotted on soft TSA agar lawns supplemented  
436 with 5 mM of CaCl<sub>2</sub> and containing either *S. epidermidis* LM1680 or *S. aureus* RN4220, TB4,  
437 8325-4, or 8325-4(Δ*stk1*) cells containing the indicated plasmids. PFUs were quantified after  
438 incubation overnight at 37°C.

439

### 440 **Efficiency of center of infection (ECOI)**

441 RN4220 cells carrying plasmid pC194 or pDB31 were grown to an optical density of 0.6 and  
442 incubated 10 min at 37°C with phage phiNM1 at an MOI of 5. Cells were then washed twice  
443 in fresh TSB to remove unbound phages and plated on a lawn of RN4220 cells. ECOI was  
444 obtained by dividing the number of plaques (or center of infections) obtained after infecting  
445 cells carrying pDB31 by the number of plaques obtained with cells carrying pC194.

446

### 447 **Phage DNA isolation and sequencing**

448 Samples of phage lysates were treated with DNase and RNase to a final volume of 200  $\mu$ l for  
449 30 min at 37°C followed by treatment with EDTA (pH 8.0, 5 mM) and proteinase K (0,5  
450 mg/ml) for 30 min at 37°C. Phage DNA was then purified using a PCR purification kit  
451 (Macherey Nagel). Phage DNA was sequenced using the Nextera library preparation kit from  
452 Illumina and sequenced on a MiSeq device.

453

#### 454 **Adsorption assay**

455 Recipient RN4220 cells were grown to an OD of 0.6 and incubated with phage phiNM1 at an  
456 MOI of 1 for 10min. Cells were then centrifuged and the number of phages remaining in the  
457 lysates was quantified ( $n_{ad}$ ). Adsorption efficiency is computed as  $(1 - n_{ad}/n_{tot})$  where  $n_{tot}$  is the  
458 total number of phages added to the sample and reported as percentages.

459

#### 460 **Growth curves**

461 *S. aureus* strains (RN4220, RN4220/pC194, RN4220/pDB31, RN4220/pDB275,  
462 RN4220/pFD6) were grown in triplicate overnight at 37°C and diluted 1:100 in 200  $\mu$ l of TSB  
463 broth in a 96-well microplate that was incubated at 37°C with shaking in an Infinite M200  
464 PRO reader (TECAN). Absorbance was measured at 600 nm every 10 min. For RN4220,  
465 RN4220/pC194, and RN4220/pDB31, after 1h of growth ( $OD_{600} \approx 0.2$ ), 10  $\mu$ l of phiNM1  
466 ( $4 \cdot 10^7$  PFU/ $\mu$ l) phage was added. For RN4220, RN4220/pDB275, and RN4220/pFD6, when  
467  $OD_{600}$  reached 0.2, the cultures were induced by anhydrotetracycline (aTc) during 1h and  
468 then phage 80alpha ( $5 \cdot 10^7$  PFU/ $\mu$ l) was added.

469

#### 470 **Prophage induction**

471 *S. aureus* strains (RN4220, RN4220::phiNM1, RN4220::phiNM1/pC194,  
472 RN4220::phiNM1/pDB31) were grown in triplicate overnight at 37°C, diluted 1:100 in TSB  
473 broth and incubated at 37°C with shaking. When cultures reached  $OD_{600} \approx 0.4$ , mitomycin  
474 C was added at a final concentration of 2  $\mu$ g/ml. After 3h of incubation in the presence or in  
475 the absence of mitomycin C, the samples were serially diluted and plated to quantify the  
476 number of surviving bacteria. Samples were also centrifuged to recover the supernatant and  
477 measure the phage titer.

478

#### 479 **Induction of candidate activators of Stk2**

480 *S. aureus* strains were grown in triplicate overnight at 37°C, diluted 1:100 in TSB broth and  
481 incubated at 37°C with shaking. When cultures reached  $OD_{600} \approx 0.2$ , aTc was added at a final  
482 concentration of 0.5  $\mu$ g/ml. All the strains were grown in parallel without aTc as a control.  
483 After 1h30 of incubation in the presence or in the absence of aTc, the samples were serially  
484 diluted and 5  $\mu$ l was spotted in TSA with appropriate antibiotics to count viable bacteria.

485

#### 486 **Mass Spectrometry**

487 Mass spectrometry methods are detailed in the supplemental experimental procedures.

488

489

490

491 **Author Contributions**

492 F.D., J.-P.D, A.S. and D.B. performed the experiments. B.D., F.D. and D.B. wrote the  
493 manuscript. H.M. performed the mass spectrometry experiments and analyzed the data. A.B.  
494 performed the phylogenetic analysis.

495

496 **Acknowledgments**

497 We are indebted to Dr. Luciano Marraffini for reagents and support, and to the group of Dr.  
498 RomainKoszul for their help with phage sequencing. This study has received funding from  
499 the French Government's Investissementd'Avenir program, Laboratoire d'Excellence  
500 "Integrative Biology of Emerging Infectious Diseases" (grant n°ANR-10-LABX-62-IBEID).

## References

- 501 Abedon, S.T. (2012). Bacterial 'immunity' against bacteriophages. *Bacteriophage* 2, 50-54.
- 502 Bae, T., Baba, T., Hiramatsu, K., and Schneewind, O. (2006). Prophages of *Staphylococcus aureus*  
503 Newman and their contribution to virulence. *Molecular microbiology* 62, 1035-1047.
- 504 Beltramini, A.M., Mukhopadhyay, C.D., and Pancholi, V. (2009). Modulation of Cell Wall Structure  
505 and Antimicrobial Susceptibility by a *Staphylococcus aureus* Eukaryote-Like Serine/Threonine Kinase  
506 and Phosphatase. *Infection and immunity* 77, 1406-1416.
- 507 Boersema, P.J., Raijmakers, R., Lemeer, S., Mohammed, S., and Heck, A.J.R. (2009). Multiplex peptide  
508 stable isotope dimethyl labeling for quantitative proteomics. *Nature protocols* 4, 484-494.
- 509 Chopin, M.C., Chopin, A., and Bidnenko, E. (2005). Phage abortive infection in lactococci: variations  
510 on a theme. *Current opinion in microbiology* 8, 473-479.
- 511 Dai, R.Y., Yan, D.M., Li, J., Chen, S.K., Liu, Y.P., Chen, R., Duan, C.Y., Wei, M., Li, H., and He, T. (2012).  
512 Activation of PKR/eIF2a signaling cascade is associated with dihydrotestosterone-induced cell cycle  
513 arrest and apoptosis in human liver cells. *J Cell Biochem* 113, 1800-1808.
- 514 Daniel, A., Bonnen, P.E., and Fischetti, V.A. (2007). First complete genome sequence of two  
515 *Staphylococcus epidermidis* bacteriophages. *Journal of bacteriology* 189, 2086-2100.
- 516 Debarbouille, M., Dramsi, S., Dussurget, O., Nahori, M.A., Vaganay, E., Jouvion, G., Cozzone, A.,  
517 Msadek, T., and Duclos, B. (2009). Characterization of a Serine/Threonine Kinase Involved in  
518 Virulence of *Staphylococcus aureus*. *Journal of bacteriology* 191, 4070-4081.
- 519 Didier, J.P., Cozzone, A.J., and Duclos, B. (2010). Phosphorylation of the virulence regulator SarA  
520 modulates its ability to bind DNA in *Staphylococcus aureus*. *FEMS microbiology letters* 306, 30-36.
- 521 Donat, S., Streker, K., Schirmeister, T., Rakette, S., Stehle, T., Liebeke, M., Lalk, M., and Ohlsen, K.  
522 (2009). Transcriptome and Functional Analysis of the Eukaryotic-Type Serine/Threonine Kinase PknB  
523 in *Staphylococcus aureus*. *Journal of bacteriology* 191, 4056-4069.
- 524 Duclos, B., Marcandier, S., and Cozzone, A.J. (1991). Chemical-Properties and Separation of  
525 Phosphoamino Acids by Thin-Layer Chromatography and or Electrophoresis. *Methods in enzymology*  
526 201, 10-21.
- 527 Finn, R.D., Clements, J., and Eddy, S.R. (2011). HMMER web server: interactive sequence similarity  
528 searching. *Nucleic acids research* 39, W29-W37.
- 529 Friedman, D.I., Mozola, C.C., Beeri, K., Ko, C.C., and Reynolds, J.L. (2011). Activation of a prophage-  
530 encoded tyrosine kinase by a heterologous infecting phage results in a self-inflicted abortive  
531 infection. *Molecular microbiology* 82, 567-577.
- 532 Goldberg, G.W., Jiang, W., Bikard, D., and Marraffini, L.A. (2014). Conditional tolerance of temperate  
533 phages via transcription-dependent CRISPR-Cas targeting. *Nature* 514, 633-637.
- 534 Goldberg, M.E., ExpertBezancon, N., Vuillard, L., and Rabilloud, T. (1996). Non-detergent  
535 sulphobetaines: A new class of molecules that facilitate in vitro protein renaturation. *Fold Des* 1, 21-  
536 27.
- 537 Goldfarb, T., Sberro, H., Weinstock, E., Cohen, O., Doron, S., Charpak-Amikam, Y., Afik, S., Ofir, G.,  
538 and Sorek, R. (2015). BREX is a novel phage resistance system widespread in microbial genomes.  
539 *Embo Journal* 34, 169-183.
- 540 Hatoum-Aslan, A., Maniv, I., Samai, P., and Marraffini, L.A. (2014). Genetic characterization of  
541 antiplasmid immunity through a type III-A CRISPR-Cas system. *Journal of bacteriology* 196, 310-317.
- 542 Horinouchi, S., and Weisblum, B. (1982a). Nucleotide-Sequence and Functional Map of Pc194, a  
543 Plasmid That Specifies Inducible Chloramphenicol Resistance. *Journal of bacteriology* 150, 815-825.
- 544 Horinouchi, S., and Weisblum, B. (1982b). Nucleotide-Sequence and Functional Map of Pe194, a  
545 Plasmid That Specifies Inducible Resistance to Macrolide, Lincosamide, and Streptogramin Type-B  
546 Antibiotics. *Journal of bacteriology* 150, 804-814.
- 547 Hoskisson, P.A., Sumbly, P., and Smith, M.C.M. (2015). The phage growth limitation system in  
548 *Streptomyces coelicolor* A(3)2 is a toxin/antitoxin system, comprising enzymes with DNA  
549 methyltransferase, protein kinase and ATPase activity. *Virology* 477, 100-109.

550 Hsu, L.C., Park, J.M., Zhang, K.Z., Luo, J.L., Maeda, S., Kaufman, R.J., Eckmann, L., Guiney, D.G., and  
551 Karin, M. (2004). The protein kinase PKR is required for macrophage apoptosis after activation of  
552 Toll-like receptor 4. *Nature* 428, 341-345.

553 Huse, M., and Kuriyan, J. (2002). The conformational plasticity of protein kinases. *Cell* 109, 275-282.

554 Jiang, W., Maniv, I., Arain, F., Wang, Y., Levin, B.R., and Marraffini, L.A. (2013). Dealing with the  
555 evolutionary downside of CRISPR immunity: bacteria and beneficial plasmids. *PLoS genetics* 9,  
556 e1003844.

557 Kanneganti, T.D. (2010). Central roles of NLRs and inflammasomes in viral infection. *Nature Reviews*  
558 *Immunology* 10, 688-698.

559 Kennelly, P.J. (2014). Protein Ser/Thr/Tyr Phosphorylation in the Archaea\*. *Journal of Biological*  
560 *Chemistry* 289, 9480-9487.

561 Labrie, S.J., Samson, J.E., and Moineau, S. (2010). Bacteriophage resistance mechanisms. *Nature*  
562 *reviews Microbiology* 8, 317-327.

563 Larsen, M.R., Thingholm, T.E., Jensen, O.N., Roepstorff, P., and Jorgensen, T.J.D. (2005). Highly  
564 selective enrichment of phosphorylated peptides from peptide mixtures using titanium dioxide  
565 microcolumns. *Molecular & Cellular Proteomics* 4, 873-886.

566 Lobočka, M., Hejnowicz, M.S., Dabrowski, K., Gozdek, A., Kosakowski, J., Witkowska, M., Ulatowska,  
567 M.I., Weber-Dabrowska, B., Kwiatek, M., Parasion, S., *et al.* (2012). Genomics of Staphylococcal  
568 Twort-like Phages - Potential Therapeutics of the Post-Antibiotic Era. *Adv Virus Res* 83, 143-216.

569 Lomas-Lopez, R., Paracuellos, P., Riberty, M., Cozzone, A.J., and Duclos, B. (2007). Several enzymes of  
570 the central metabolism are phosphorylated in *Staphylococcus aureus*. *FEMS microbiology letters* 272,  
571 35-42.

572 London, J., Skrzynia, C., and Goldberg, M.E. (1974). Renaturation of *Escherichia-Coli* Tryptophanase  
573 after Exposure to 8 M Urea - Evidence for Existence of Nucleation Centers. *European Journal of*  
574 *Biochemistry* 47, 409-415.

575 Makarova, K.S., Wolf, Y.I., Snir, S., and Koonin, E.V. (2011). Defense Islands in Bacterial and Archaeal  
576 Genomes and Prediction of Novel Defense Systems. *Journal of bacteriology* 193, 6039-6056.

577 Marchler-Bauer, A., and Bryant, S.H. (2004). CD-Search: protein domain annotations on the fly.  
578 *Nucleic acids research* 32, W327-W331.

579 Meurs, E., Chong, K., Galabru, J., Thomas, N.S.B., Kerr, I.M., Williams, B.R.G., and Hovanessian, A.G.  
580 (1990). Molecular-Cloning and Characterization of the Human Double-Stranded-Rna Activated  
581 Protein-Kinase Induced by Interferon. *Cell* 62, 379-390.

582 Molle, V., Zanella-Cleon, I., Robin, J.P., Mallejac, S., Cozzone, A.J., and Becchi, M. (2006).  
583 Characterization of the phosphorylation sites of *Mycobacterium tuberculosis* serine/threonine  
584 protein kinases, PknA, PknD, PknE, and PknH by mass spectrometry. *Proteomics* 6, 3754-3766.

585 Nair, D., Memmi, G., Hernandez, D., Bard, J., Beaume, M., Gill, S., Francois, P., and Cheung, A.L.  
586 (2011). Whole-Genome Sequencing of *Staphylococcus aureus* Strain RN4220, a Key Laboratory Strain  
587 Used in Virulence Research, Identifies Mutations That Affect Not Only Virulence Factors but Also the  
588 Fitness of the Strain. *Journal of bacteriology* 193, 2332-2335.

589 Novick, R.P. (1989). Staphylococcal Plasmids and Their Replication. *Annu Rev Microbiol* 43, 537-565.

590 Palm, N.W., and Medzhitov, R. (2009). Pattern recognition receptors and control of adaptive  
591 immunity. *Immunol Rev* 227, 221-233.

592 Pereira, S.F.F., Goss, L., and Dworkin, J. (2011). Eukaryote-Like Serine/Threonine Kinases and  
593 Phosphatases in Bacteria. *Microbiol Mol Biol R* 75, 192-+.

594 Pumplín, N., and Voinnet, O. (2013). RNA silencing suppression by plant pathogens: defence,  
595 counter-defence and counter-counter-defence. *Nature Reviews Microbiology* 11, 745-760.

596 Robertson, E.S., and Nicholson, A.W. (1992). Phosphorylation of *Escherichia-Coli* Translation  
597 Initiation-Factors by the Bacteriophage-T7 Protein-Kinase. *Biochemistry-Us* 31, 4822-4827.

598 Sadler, A.J., and Williams, B.R.G. (2008). Interferon-inducible antiviral effectors. *Nature Reviews*  
599 *Immunology* 8, 559-568.

600 Stark, G.R., Kerr, I.M., Williams, B.R.G., Silverman, R.H., and Schreiber, R.D. (1998). How cells respond  
601 to interferons. *Annual review of biochemistry* 67, 227-264.

602 Sumbly, P., and Smith, M.C.M. (2002). Genetics of the phage growth limitation (Pgl) system of  
603 *Streptomyces coelicolor* A3(2). *Molecular microbiology* *44*, 489-500.

604 Tahara, M., Ohsawa, A., Saito, S., and Kimura, M. (2004). In vitro phosphorylation of initiation factor  
605 2 alpha (aIF2 alpha) from hyperthermophilic Archaeon *Pyrococcus horikoshii* OT3. *J Biochem* *135*,  
606 479-485.

607 Truong-Bolduc, Q.C., Ding, Y.P., and Hooper, D.C. (2008). Posttranslational Modification Influences  
608 the Effects of MgrA on norA Expression in *Staphylococcus aureus*. *Journal of bacteriology* *190*, 7375-  
609 7381.

610 Yan, N., and Chen, Z.J.J. (2012). Intrinsic antiviral immunity. *Nat Immunol* *13*, 214-222.

611 Young, T.A., Delagoutte, B., Endrizzi, J.A., Falick, A.M., and Alber, T. (2003). Structure of  
612 *Mycobacterium tuberculosis* PknB supports a universal activation mechanism for Ser/Thr protein  
613 kinases. *Nat Struct Biol* *10*, 168-174.

614 Zorzatto, C., Machado, J.P.B., Lopes, K.V.G., Nascimento, K.J.T., Pereira, W.A., Brustolini, O.J.B., Reis,  
615 P.A.B., Calil, I.P., Deguchi, M., Sachetto-Martins, G., *et al.* (2015). NIK1-mediated translation  
616 suppression functions as a plant antiviral immunity mechanism. *Nature* *520*, 679-U227.

## Figure Legends

617

618 **Figure 1. Discovery of a novel phage defense system in *S. epidermidis* strain RP62A.** A) Strains with various deletions in the region of the CRISPR locus (numbered as in Jiang et al.,  
619 2013) were screened for sensitivity to bacteriophage CNP<sub>x</sub>. ORFs in the genomic region are  
620 represented in green. Each horizontal line represents a strain and the line is discontinued in  
621 the deleted region. Blue lines indicate strains sensitive to phage CNP<sub>x</sub> while red lines indicate  
622 resistance. The region that is sufficient to provide resistance is highlighted in purple. The  
623 exact position of the candidate region is 2535598-2551561 and the ORFs it contains are  
624 described in Table S1. B) ORFs contained in the candidate region were cloned either alone or  
625 two at a time on plasmid pC194 to give plasmids pDB31 (*serp2479, stk2*), pDB32 (*serp2480*),  
626 pDB33 (*serp2481+serp2482*) and pDB34 (*serp2483+serp2484*)(see Table S1). Efficiency of  
627 plaquing (EOP) of phage CNP<sub>x</sub> is reported on *S. epidermidis* strain LM1680 containing these  
628 different plasmids (mean+sd, n=3). No plaques were recovered in bacteria carrying gene  
629 *Serp2479 (stk2)* (EOP detection limit of 10<sup>-5</sup>). C) A tree of Staphylococci was constructed  
630 based on the assembled complete genomes available in GenBank. Red indicates the presence  
631 of proteins with a minimum of 90% identity to Stk2 (SERP2479) over 100% of the sequence  
632 length. More distantly related proteins were also identified in Staphylococci that are not  
633 present in this tree and are reported in Table S2.

635 **Figure 2. Stk2 is an active Serine/Threonine kinase.** A) Schematic presentation of the  
636 position of phosphoresidues in protein Stk2 and location of different domains. The kinase  
637 domain, the activation loop, the cystein-rich region, and the position of the residues K152 and  
638 D243 are shown. The position of phosphoresidues is indicated by dotted bars. Residues in red  
639 were also shown to be phosphorylated *in vivo* (see Table S7). B) SDS-PAGE analysis of  
640 purified Stk2 (lane 2) and Stk2 mutant K152I (lane 3) after staining with Coomassie blue.  
641 Molecular mass standards are shown on the left (lane 1). Autophosphorylation of Stk2 (lane  
642 4) and mutant K152I (lane 5) in the presence of radioactive [ $\gamma$ -32P] ATP is shown.  
643 Radioactive molecules were detected by autoradiography. C) Effect of cations on  
644 Stk2 autophosphorylation activity *in vitro*. D) 2D-analysis of phosphorylated amino acids in  
645 Stk2. The acid-stable phosphoamino acids from [ $\gamma$ -32P]-labelled Stk2 were separated by  
646 electrophoresis in the first dimension (1D) followed by ascending chromatography in the  
647 second dimension (2D). (P-Tyr) phosphotyrosine, (P-Ser) phosphoserine, and (P-Thr)  
648 phosphothreonine were located by ninhydrin staining. Phosphorylated molecules were  
649 revealed by autoradiography. E) Effect mutations on the kinase activity of Stk2. Purified wild-  
650 type and mutants of Stk2 were incubated in the presence of radioactive [ $\gamma$ -32P] ATP, protein  
651 were separated by SDS-PAGE, and radioactive molecules were detected by autoradiography.

652 **Figure 3. The Stk2 kinase triggers cells death during the phage lytic cycle.** A) EOP of  
653 phage phiNM1 on RN4220 *S. aureus* cells in the presence of the WT Stk2 protein (pDB31) or  
654 various mutants: K152Q (pDB81), D243A (pDB82) and T275A (pDB83). B) Growth curve  
655 of *S. aureus* RN4220 carrying control plasmid pC194 or pDB31 (pC194 $\Delta$ *stk2*). Phage  
656 phiNM1 was added after 1H of growth at a multiplicity of infection(MOI) of 10. C) PFU and  
657 CFU recovered after infection of RN4220 cells carrying *stk2* or not. Cells were grown to OD

658 = 0.2 and incubated with phiNM1 for 2H. Cells were then plated on TSA to measure CFUs,  
659 and the filtered supernatant was spotted on a top-agar lawn of RN4220 cells to measure PFUs.  
660 Upon infection with phiNM1 and in the presence of *stk2*, cells are killed but the phage is not  
661 amplified. D) PFU and CFU recovered after induction with mitomycin C (MMC) of growing  
662 RN4220 carrying a phiNM1 lysogenic phage or not, in the presence or absence of *stk2*. Upon  
663 induction, cells carrying a lysogenic phage are killed regardless of the presence of *stk2*, but  
664 the production of phage is inhibited in the presence of *stk2*. Note that PFU are recovered even  
665 in the absence of MMC due to the spontaneous induction of the phage. See also Figure S1.

666 **Figure 4. A phage protein triggers Stk2-mediated cell death.** A) Two mutants of phage  
667 phiNM1 able to infect RN4220 cells carrying *stk2* were isolated and sequenced. Coverage  
668 along the phage genome is plotted and mutations are highlighted. Both phages carry a  
669 mutation in gene *pacK*(SAPPV1\_GP14), V200M and K30E for phages phiNM1K1 and  
670 phiNM1K2 respectively. B) To test whether PacK is sufficient to activate Stk2, the *pacK* gene  
671 was cloned under the control of a Ptet promoter on a pE194 vector giving pAS10, and  
672 introduced in cells carrying plasmid pDB31. C) RN4220 cells carrying genes  
673 *pacK*(SAPPV1\_GP14) from phage phiNM1 (pAS10), AVT76\_gp14 from phage phiNM2  
674 (pFD16) or ST85ORF023 from phage 85 (pFD20) under the control of a Ptet promoter,  
675 together with plasmids pC194 or pDB31 were grown to OD~0.2 and induced with  
676 anhydrotetracycline. After 1H of induction cells were plated and colonies quantified. See also  
677 Figure S2.

678 **Figure 5. Stk1 is required for efficient Stk2-mediated immunity.** A) EOP of phage  
679 phiNM1 is reported against *S. aureus* strain NCTC8325-4 harboring *stk1* or not ( $\Delta$ *stk1*) in the  
680 presence of *stk2* (pDB31) or a control plasmid (pC194). B) NCTC8325-4 cells harboring *stk1*  
681 or not ( $\Delta$ *stk1*) in the presence of *stk2* (pDB31) or a control plasmid (pC194) were  
682 transformed with plasmid pAS10 expressing *pacK* under the control of Ptet promoter. Cells  
683 were grown to OD  $\approx$  0.2 and induced with anhydrotetracycline. After 2H of induction, cells  
684 were plated and colonies quantified. Induction of *pacK* expression triggers cell death in the  
685 presence of *stk2* regardless of the presence or absence of *stk1*.

**Table 1.** List of proteins phosphorylated upon expression of PacK and in the presence of Stk2 and which are never found phosphorylated in the absence of Stk2. A detailed list of all phosphopeptides identified with match metrics and measured ratios are provided in Table S7.

	Gene Name	Protein Descriptions	Accession #
Transcription	greA	Transcription elongation factor GreA	A6QHF1
	sigA	RNA polymerase sigma factor SigA	P0A0J0
	nusA	Transcription termination/antitermination protein NusA	Q2G2D2
Translation	metG	Methionine--tRNA ligase	A6QEE3
	tuf	Elongation factor Tu	A6QEK0
	efp	Elongation factor P	A6QH73
	rpmE2	50S ribosomal protein L31 type B	A6QJW4
	rplF	50S ribosomal protein L6	A6QJ77
	rplE	50S ribosomal protein L5	A6QJ80
Cell cycle	ftsZ	Cell division protein FtsZ	A6QG86
	sepF	Cell division protein SepF	Q2FZ86
	gpsB	Cell cycle protein GpsB	Q2FYI5
Stress response	clpX	ATP-dependent Clp protease ATP-binding subunit ClpX	Q2FXQ7
	clpB	Chaperone protein ClpB	Q2FZS8
	ydaG / yzzA	General stress protein 26	Q2FVN7
	AQ00_RS06590	alkaline shock protein (Asp23) / Stress response regulator gls24 homolog	Q2FZ59
	tela	Tellurite/ Toxin anion resistance protein	Q2FYM7
DNA topology and repair	ssb	Single-stranded DNA-binding protein	A6QE48
	parE	DNA topoisomerase 4 subunit B	A6QGQ7
	mutS	DNA mismatch repair protein MutS	Q2FYZ9
Central metabolism and Biosynthesis	hemL1	Glutamate-1-semialdehyde 2,1-aminomutase 1 (protoporphyrin-IX biosynthesis)	A6QHK1
	pgk	Phosphoglycerate kinase (Glycolysis)	A6QF82
	pgi	Glucose-6-phosphate isomerase (Glycolysis pathway)	Q2FZU0
	dltA	D-alanine--poly(phosphoribitol) ligase subunit 1 (LTA biosynthesis)	Q2FZW6
Others	pacK / SAPPV1_GP14	pacK (Stk2 activation protein)	A6QDW1
	stk2	SERP2479	Q5HK71
	ylaL	Uncharacterized protein	A6QFW6
	esxA	Virulence factor EsxA (ESAT-6-like protein)	Q2G189
	obg	GTPase	Q2FXT1
	phoP	Alkaline phosphatase synthesis two-component response regulator	Q2FXN6
	AQ00_RS00105	short-chain dehydrogenase	Q2FV41