



**HAL**  
open science

## 4'-Methoxyphenacyl-Assisted Synthesis of $\beta$ -Kdo Glycosides

Marcelina Mazur, Barbara Barycza, Hanitra Andriamboavonjy, Serge Lavoie, Marielle Tamigney Kenfack, Anaïs Laroussarie, Yves Blériot, Charles Gauthier

### ► To cite this version:

Marcelina Mazur, Barbara Barycza, Hanitra Andriamboavonjy, Serge Lavoie, Marielle Tamigney Kenfack, et al.. 4'-Methoxyphenacyl-Assisted Synthesis of  $\beta$ -Kdo Glycosides. *Journal of Organic Chemistry*, 2016, 81 (22), pp.10585-10599. <10.1021/acs.joc.6b01431>. <pasteur-01421190>

**HAL Id: pasteur-01421190**

**<https://pasteur.hal.science/pasteur-01421190v1>**

Submitted on 12 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

## 4'-Methoxyphenacyl-Assisted Synthesis of $\beta$ -Kdo Glycosides

Marcelina Mazur,<sup>†,‡</sup> Barbara Barycza,<sup>†,‡</sup> Hanitra Andriamboavonjy,<sup>†</sup> Serge Lavoie,<sup>§</sup> Marielle

Tamigney Kenfack,<sup>†</sup> Anaïs Laroussarie,<sup>†</sup> Yves Blériot,<sup>†</sup> and Charles Gauthier<sup>\*,†,§,||</sup>


<sup>†</sup>Institut de Chimie IC2MP, CNRS-UMR 7285, Équipe Synthèse Organique, Université de Poitiers, 4 rue Michel Brunet, 86073 Poitiers Cedex 9, France

<sup>‡</sup>Department of Chemistry, Wrocław University of Environmental and Life Sciences, Norwida 25, 50-375 Wrocław, Poland

<sup>§</sup>Laboratoire LASEVE, Département des Sciences Fondamentales, Université du Québec à Chicoutimi, 555 boul. de l'Université, Chicoutimi (Québec), Canada G7H 2B1

<sup>||</sup>INRS-Institut Armand-Frappier, Université du Québec, 531 boul. des Prairies, Laval (Québec), Canada H7V 1B7

### TABLE OF CONTENTS/ABSTRACT GRAPHIC




1  
2  
3 **ABSTRACT:** 3-Deoxy- $\beta$ -D-*manno*-oct-2-ulosonic acid ( $\beta$ -Kdo) glycosides are mainly found in  
4 capsular polysaccharides (CPS) and extracellular exopolysaccharides (EPS) from Gram-negative  
5 bacteria. These compounds have profound biological implications in immune response and act as  
6 virulence factors. We have developed a novel methodology for the stereoselective synthesis of  $\beta$ -  
7 Kdo glycosides via the use of a 4'-methoxyphenacyl (Phen) auxiliary group at the C1 position of  
8 a peracetylated  $\beta$ -Kdo thioglycoside. Under the promotion of NIS/AgOTf in acetonitrile, a series  
9 of Kdo glycosides was synthesized in good yield and  $\beta$ -selectivity while minimizing the  
10 formation of undesirable glycals. Stereoselectivity of the glycosylation was shown to be  
11 modulated by various factors such as promotor, solvent, anomeric ratio of donor, nature of  
12 acceptor, and Phen substitution. Chemoselective cleavage of the Phen group was performed  
13 under the action of Zn/HOAc. DFT calculations together with experimental results suggested that  
14  $\alpha$ -triflate and a six-membered  $\alpha$ -spiroPhen are plausible intermediates of the reaction, accounting  
15 for the enhanced formation of  $\beta$ -Kdo glycosides. The developed methodology could be applied to  
16 the synthesis of  $\beta$ -Kdo-containing glycans from pathogenic bacteria.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## INTRODUCTION

3-Deoxy-D-manno-oct-2-ulosonic acid (Kdo) glycosides are mainly found in the surface polysaccharides of bacteria.<sup>1</sup> Kdo glycosides with the  $\alpha$ -configuration are present in virtually all of the lipopolysaccharide (LPS) core regions of Gram-negative bacteria, playing a crucial role in the structural integrity of bacterial membranes.<sup>2</sup> In contrast,  $\beta$ -Kdo glycosides occur far less frequently within LPS. Rare occurrences include: 1) LPS core regions from *Alteromonas macleodii*;<sup>3</sup> 2) LPS O-antigen (OAg) from *Providencia alcalifaciens*;<sup>4</sup> and 3) non-reducing end of LPS OAg from *Klebsiella pneumoniae* serotype O12.<sup>5</sup> The most frequent occurrence of  $\beta$ -Kdo glycosides is within the repeating unit of capsular polysaccharides (CPS),<sup>1</sup> which are known as virulence factors and are involved in protection from host immune mechanisms.<sup>6</sup> For instance, *Kingella kingae*, a Gram-negative bacteria causing septic arthritis, osteomyelitis, and bacteremia in young children, produces a CPS featuring a repeating disaccharide comprised of a Kdo residue in the  $\beta$ -configuration (Figure 1).<sup>7</sup>  $\beta$ -Kdo glycosides have also been found in extracellular exopolysaccharides (EPS), such as the one expressed by the 'Tier 1 Select Agent' *Burkholderia pseudomallei*, the causative agent of melioidosis.<sup>8-10</sup> Cytidine monophosphate (CMP)-Kdo,<sup>11,12</sup> the activated sugar nucleotide processed by Kdo glycosyltransferases, is another important example of a naturally occurring compound bearing a  $\beta$ -Kdo unit. Recently, Whitfield and co-workers<sup>13</sup> have highlighted the presence of poly-Kdo linkers, containing alternating  $\beta$ -(2 $\rightarrow$ 4) and  $\beta$ -(2 $\rightarrow$ 7) linkages, at the reducing end of CPS from various Gram-negative pathogens including *Escherichia coli*, *Campylobacter jejuni*, *Haemophilus influenzae*, *Neisseria meningitidis*, and *Pasteurella multocida*. Enzymes involved in the biosynthesis of these poly-Kdo linkers have been characterized as novel retaining Kdo transferases (KpsC and KpsS).<sup>14,15</sup> Owing to the structural significance and biological importance of  $\beta$ -Kdo residues in bacterial polysaccharides,

straightforward synthetic routes towards  $\beta$ -Kdo glycosides are needed.<sup>16,17</sup> Access to these compounds in pure and homogeneous forms would further the development of vaccines, diagnostics and therapeutics against some clinically relevant bacterial pathogens.<sup>1,2,18</sup>



**Figure 1.** Naturally occurring  $\beta$ -Kdo-containing glycans from bacteria.

The synthesis of Kdo glycosides is not a trivial task, and shares similarities with the glycosylation of *N*-acetylneuraminic acid (Neu5Ac).<sup>19</sup> The lack of a hydroxyl group at the C3 position, which hampers the use of the neighboring group participation effect, the presence of an electron withdrawing carboxylic acid at C1, which deactivates and hinders the anomeric position, and the undesirable formation of 2,3-glycals are the main issues regarding the synthesis of Kdo glycosides.<sup>16,17</sup> In the last few years, novel methodologies have been implemented allowing access to  $\alpha$ -Kdo glycosides in excellent yields and stereoselectivity. In this respect, it is worth mentioning the use of 5,7-*O*-di-*tert*-butylsilylene protected thioglycoside<sup>20</sup> and 3-iodo fluoride<sup>21-23</sup> donors. Yet, the synthetic chemistry of Kdo glycosides having the opposite thermodynamically less stable  $\beta$ -configuration (OR group in equatorial rather than axial position) still requires improvements.<sup>16</sup>

1  
2  
3  
4  
5 Towards this aim, Ling and co-workers,<sup>24</sup> relying on the pioneering work of Takahashi,<sup>25</sup>  
6 developed a novel class of 4,5;7,8-di-*O*-isopropylidene protected 1-C-arylglycal donors, which,  
7 upon treatment with *N*-iodosuccinimide (NIS), led to the stereoselective formation of  $\beta$ -Kdo  
8 glycosides. Yet, their approach required a supplemental reductive deiodination step followed by  
9 an oxidative transformation in order to provide the carboxylic acid at C1. Recently, Mong and co-  
10 workers<sup>26</sup> partially resolved the latter issue by preparing Kdo glycal donors bearing a preinstalled  
11 carboxylate at C1. NIS-mediated glycosylation of these glycals in a DCM/CH<sub>3</sub>CN mixture led to  
12  $\beta$ -Kdo glycosides in a  $\beta/\alpha$  ratio of up to 20:1 following radical deiodination. In both previous  
13 cases, the presence of a 4,5-*O*-isopropylidene group locking the pyranose ring into a skew-boat  
14 conformation was found to be essential for providing high  $\beta$ -stereoselectivity. Using unlocked  
15 perbenzylated or peracetylated Kdo glycals led to the opposite *trans*-diaxial selectivity with  
16 regard to the C3 iodine atom and C2 OR group.<sup>25,27</sup>

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36 There have been few reports regarding the synthesis of  $\beta$ -Kdo glycosides with ‘non-glycal’  
37 donors. van Boom and co-workers<sup>28,29</sup> were the first to show that reacting peracetylated  $\beta$ -Kdo  
38 thioglycosides with NIS/TfOH could provide Kdo glycosides in the major  $\beta$ -configuration when  
39 3-amino-*N*-benzyloxycarbonyl-1-propanol was used as an acceptor. More recently, an interesting  
40 study by the group of Oscarson<sup>27</sup> revealed that peracylated  $\beta$ -Kdo thioglycosides were suitable  
41 donors for the formation of  $\beta$ -Kdo glycosides bearing 2-(4-trifluoroacetamidophenyl)ethyl as a  
42 spacer when DMTST or IBr/AgOTf were used as promoters. In all these studies, however, no  
43 systematic evaluations of the glycosylation conditions were performed and no mechanistic details  
44 were provided. Herein, we report a novel approach for the stereoselective synthesis of  $\beta$ -Kdo  
45 glycosides involving the use of a long-range participating 4'-methoxyphenacyl (Phen) auxiliary

1  
2  
3 group at the C1 position. Glycosylation conditions were thoroughly investigated both in the  
4 presence and in the absence of the Phen group, e.g. promoters, leaving groups, anomeric  
5 configuration of donors, solvents, nature of acceptors, addition order of reagents, and counter-  
6 anions. On the basis of DFT calculations and experimental details, we also propose plausible  
7 intermediates accounting for the formation of both  $\alpha$ - and  $\beta$ -Kdo glycosides under the optimized  
8 reaction conditions.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

## 20 RESULTS AND DISCUSSION


21  
22  
23  
24 **Synthetic Approach.** Long-range participating effects through the use of an auxiliary group have  
25 been previously described for the synthesis of  $\alpha$ -Neu5Ac glycosides, which display structural  
26 similarities with  $\beta$ -Kdo glycosides. The presence of ester chains at the C1 position of Neu5Ac,  
27 such as 2-methylthioethyl, 2-phenylthioethyl,<sup>30</sup> and *N,N*-dimethylglycolamide,<sup>31,32</sup> was shown to  
28 enhance the  $\alpha$ -selectivity of the glycosylation reaction via the stabilization of the oxocarbenium  
29 ion from the  $\beta$ -axial orientation, thereby favoring the attack of the nucleophile from the  $\alpha$ -face.  
30 Enhanced  $\alpha$ -selectivities were also observed for Neu5Ac thioesters<sup>33</sup> and 2-cyanoethyl esters<sup>34</sup> in  
31 conjunction with CH<sub>3</sub>CN, presumably through a mechanism involving stabilization of the  $\beta$ -  
32 oriented nitrilium ion.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 Inspired by these previous studies, we have devised an analogous approach to tackle the problem  
49 of  $\beta$ -Kdo glycosides synthesis via the use of a 4'-methoxyphenacyl (Phen) auxiliary group at the  
50 C1 position of peracetylated Kdo donors. Our choice was driven by two important factors: 1) the  
51 enhanced electronic density of the ketone functionality that would be likely to participate  
52 favorably in the course of the glycosylation reaction, and 2) the orthogonality of phenacyl  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 groups<sup>35</sup> with several base- and acid-sensitive protecting groups that would be an asset over  
4  
5 previously reported auxiliary groups.  
6  
7

8  
9  
10 Our working hypothesis is depicted in Figure 2. Once the Kdo donor is activated by a suitable  
11  
12 electrophilic promoter, the oxocarbenium ion (glycosyl cation)<sup>36</sup> will be formed. According to  
13  
14 Woerpel,<sup>37-40</sup> the attack of the electron-rich ketone from the  $\alpha$ -face of the  ${}^5H_4$  half-chair  
15  
16 conformer would be favored in order to minimize the destabilizing 1,3-diaxial interactions. The  
17  
18 resulting six-membered  $\alpha$ -spiro compound, which could be found either as a covalent or contact  
19  
20 ion pair intermediate, would then be attacked by the acceptor preferentially from the opposite  $\beta$ -  
21  
22 face leading to enhanced  $\beta$ -selectivity for the formation of Kdo glycosides. DFT calculations at  
23  
24 the B3LYP/6-311++G(2d,2p) level of theory tend to support this hypothesis since the  $\alpha$ -spiro  
25  
26 intermediate was found to be energetically favored compared to the  $\beta$ -spiro intermediate by 13.0  
27  
28 kJ·mol<sup>-1</sup> (see Figure S1 and Table S1). Moreover, these two intermediates were at least 36.8  
29  
30 kJ·mol<sup>-1</sup> more stable than the free oxocarbenium ion. Nevertheless, it is important to point out  
31  
32 that, in the case of a Curtin-Hammett scenario in which there is a rapid exchange between both  
33  
34 intermediates, the major product could also arise from the higher energy ground state  
35  
36 intermediate.<sup>41,42</sup> Furthermore, it has to be stressed out that a conformational change from the  
37  
38 more stable *Z*-ester to the less stable *E*-ester must occur in order to allow the phenacyl ketone to  
39  
40 approach the glycosyl cation. According to the literature, the energy difference for the *Z/E* ester  
41  
42 isomerization is about 12.5 kJ·mol<sup>-1</sup>.<sup>43</sup> This energy barrier can be lowered by using polar  
43  
44 solvents, such as acetonitrile, or when an electron-withdrawing group is attached to the R'  
45  
46 position of a RCO<sub>2</sub>R' ester, such as a phenacyl group.<sup>44</sup> In the case of our work, the *Z/E* ester  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

isomerization would be beneficial in terms of energy because it opens the way for the formation of the spiro intermediates, which are more stable than the free oxocarbenium ion.




**Figure 2.** Proposed approach for the synthesis of  $\beta$ -Kdo glycosides through the use of a 4'-methoxyphenacyl (Phen) C1-auxiliary group. A = activating group; E = electrophile. The 3D structures were obtained by DFT geometrical optimization at the B3LYP/6-311++G(2d,2p) level of theory (hydrogen atoms have been omitted for the sake of clarity).

**Synthesis of Kdo Donors.** The synthesis of peracetylated Kdo thioglycoside and fluoride donors **2-5** bearing participating (Phen) or non-participating (Bn) ester groups was investigated first (Scheme 1). Crystalline ammonium Kdo was obtained through the modified Cornforth procedure<sup>45-47</sup> using an optimized methodology recently reported by Kosma.<sup>48</sup> This allowed us to prepare gram quantities of pure Kdo in a reliable manner. Ammonium Kdo was subjected to acetylation under standard conditions (Ac<sub>2</sub>O, py, DMAP, rt) leading to peracetylated **1**<sup>49</sup> with nearly quantitative yield. Performing the reaction at more elevated temperatures (>30 °C) generated substantial amounts of Kdo lactone.<sup>50</sup> Then, two different routes were studied for the synthesis of thioglycosides **2** and **3**. Esterification with 4'-methoxyphenacyl bromide (PhenBr) or BnBr in the presence of Cs<sub>2</sub>CO<sub>3</sub> followed by glycosylation with EtSH under the action of BF<sub>3</sub>·OEt<sub>2</sub> produced donors **2** and **3** with 15 and 67% yield, respectively, both predominantly featuring the  $\beta$ -configuration as expected<sup>27</sup> ( $\beta/\alpha$  9:1 for **2**, and 7:1 for **3**). The low yield obtained for the Phen derivative **2** was due to the formation of an undesirable by-product, presumably a dithioketal coming from the addition of two EtSH molecules on the activated ketone (LRMS:  $m/z$  [M + Na] calcd for C<sub>31</sub>H<sub>44</sub>O<sub>12</sub>S<sub>3</sub> 727.2; found 727.7). Another route was then investigated in which free acid **1** was first refluxed in DCE with EtSH and BF<sub>3</sub>·OEt<sub>2</sub> followed by esterification of the resulting thioglycoside. Using this route, donors **2** and **3** were obtained in convenient yields (45 and 50%, respectively), but with different anomeric ratios than route A ( $\beta/\alpha$  ~1:1). Fluoride donors **4** and **5** were synthesized via a similar approach. Regioselective fluorination at the anomeric position was performed by treatment of peracetylated **1** with HF·py 7:3 followed by standard esterification to provide donors **4** and **5** with 49 and 45% yield, respectively.<sup>51</sup> The exclusive  $\alpha$ -configuration of these fluoride donors was confirmed by <sup>19</sup>F NMR analysis (<sup>3</sup>J<sub>F,H3ax</sub> = 34.6 Hz, <sup>3</sup>J<sub>F,H3eq</sub> = 6.0 Hz).


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Scheme 1. Synthesis of Kdo Thioglycoside and Fluoride Donors




28 In order to provide donor **2** with the same anomeric ratio as **3** ( $\beta/\alpha$  7:1, route A) and to improve  
29 the yield, an additional methodology was investigated for comparison purposes (Scheme 2).  
30 Therefore, hydrogenolysis of thioglycoside **3** followed by esterification under the above-  
31 mentioned conditions provided donor **2** with a good yield (76%) without erosion of  
32 diastereoselectivity ( $\beta/\alpha$  7:1). Furthermore, this approach allowed us to synthesize Kdo  
33 thioglycoside donors bearing 3'-methoxy (**6**), 2'-methoxy (**7**) as well as unsubstituted (**8**)  
34 phenacyl groups all having the same  $\beta/\alpha$  ratio (7:1).  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

47 **Scheme 2. Synthesis of  $\beta$ -Kdo Thioglycoside Bearing Diversely Substituted Phenacyl**  
48 **Groups**  
49  
50




16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

**Synthesis of  $\beta$ -Kdo Glycosides.** With Kdo donors (**2-8**) in hand, study of their glycosylation behavior for the selective formation of  $\beta$ -Kdo glycosides was investigated next. Using Phen thioglycoside **2** in a  $\sim$ 1:1  $\beta/\alpha$  ratio together with 5-amino-*N*-benzyloxycarbonyl-1-pentanol (**9**,<sup>52</sup> Figure 3) as a model acceptor, we first screened different thiophilic promoters (Table 1). Glycosylation reactions in entries 1 to 8 were conducted in the non-participating solvent DCE at  $-10$  °C in the presence of water scavenging 4 Å molecular sieves, and the  $\beta/\alpha$  selectivity ratio was evaluated by <sup>1</sup>H NMR analysis. At this temperature, all promoters were shown to give excellent conversions (>95%), with the exception of MeOTf (60%), while  $\beta$ -anomeric selectivity varied significantly.



49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure 3.** Glycosyl acceptors (**9-14**) used in this study.

Table 1. Synthesis of  $\beta$ -Kdo Glycosides: Promoter and Solvent Screening

entry	promoter	solvent	conv. (%) <sup>a</sup>	selectivity ratio <sup>a</sup>		
				15 $\beta$	15 $\alpha$	16
1	IBr/AgOTf	DCE	>95	1.5	1.0	nd <sup>b</sup>
2	MeOTf	DCE	60	2.4	1.0	0.8
3	Me <sub>2</sub> S <sub>2</sub> /MeOTf <sup>c</sup>	DCE	>95	3.7	1.0	2.0
4	Me <sub>2</sub> S <sub>2</sub> /Tf <sub>2</sub> O	DCE	>95	3.0	1.0	1.0
5	NCS/TfOH	DCE	>95	5.0	1.0	5.7
6	NIS/TfOH	DCE	>95	5.9	1.0	1.5
7	NIS/TfOH <sup>d</sup>	DCE	41	6.0	1.0	nd
8	NIS/AgOTf	DCE	>95	4.7	1.0	0.6
9	NIS/AgOTf	Et <sub>2</sub> O	>95	5.8	1.0	nd
10	<b>NIS/AgOTf</b>	<b>CH<sub>3</sub>CN</b>	<b>&gt;95 (84)<sup>e</sup></b>	<b>7.9</b>	<b>1.0</b>	<b>nd</b>
11	NIS/AgOTf <sup>f</sup>	CH <sub>3</sub> CN	>95	nd	nd	1.0
12	NIS/AgOTf <sup>f</sup>	DCE	>95	nd	nd	1.0
13	NIS/AgClO <sub>4</sub>	CH <sub>3</sub> CN	>95	3.2	1.0	nd
14	NIS/AgBF <sub>4</sub>	CH <sub>3</sub> CN	>95	3.2	1.0	nd

<sup>a</sup>Determined by <sup>1</sup>H NMR analysis of the crude reaction mixture. <sup>b</sup>Not detected. <sup>c</sup>*In situ* formed DMTST. <sup>d</sup>Performed at -40 °C. <sup>e</sup>Performed at 1.25 mmol scale. Value in parentheses corresponds to the isolated yield of **15 $\beta$**  and **15 $\alpha$** . <sup>f</sup>Donor and reagents were premixed before the addition of acceptor **9** (preactivation conditions).

In contrast with the results of Oscarson,<sup>27</sup> the use of IBr/AgOTf<sup>53</sup> gave the lowest  $\beta/\alpha$  ratio (1.5:1.0) without formation of glycal **16** (entry 1). We interpreted this result as a competitive attack of the nucleophilic bromide anion at the anomeric center, forming a reactive  $\beta$ -bromide species<sup>54</sup> that can be displaced from the  $\alpha$ -face by the acceptor. Using Me<sub>2</sub>S<sub>2</sub>/MeOTf (DMTST)<sup>55</sup> or Me<sub>2</sub>S<sub>2</sub>/Tf<sub>2</sub>O<sup>56</sup> provided better  $\beta$ -selectivity (up to 3.7:1.0) but glycal formation was observed


1  
2  
3 (entries 3 and 4). Enhanced formation of Kdo glycoside **15 $\beta$**  was found with *N*-chlorosuccinimide  
4 (NCS)/TfOH as the promoter ( $\beta/\alpha$  5.0:1.0) but, again, substantial amounts of glycal **16** were  
5 formed (entry 5). Switching to NIS<sup>57-59</sup> significantly decreased the formation of **16** but kept good  
6  $\beta$ -selectivity (entry 6). Conducting the reaction at  $-40$  °C with NIS/TfOH prevented glycal  
7 formation (entry 7); however, conversion of the donor was not complete (~41%).  
8  
9

10  
11  
12  
13  
14  
15  
16  
17 We next examined the use of NIS/AgOTf<sup>59</sup> as the promoter, which gave the best results in terms  
18 of  $\beta$ -selectivity while minimizing glycal formation (entry 8). This result was somewhat  
19 unexpected since Oscarson<sup>27</sup> showed that a similar peracetylated methyl ester Kdo thioglycoside  
20 only furnished elimination product following treatment by NIS/AgOTf in DCM. We then  
21 explored the use of well-known participating solvents such as Et<sub>2</sub>O and CH<sub>3</sub>CN (entries 9 and  
22 10).<sup>60</sup> The outcome of these reactions was found to be promising: complete conversion of donor  
23 **2**, enhanced  $\beta$ -selectivity, and no formation of glycal were observed. The reaction in CH<sub>3</sub>CN was  
24 performed at 1.25 mmol scale providing Kdo glycoside **15** in 84% yield with a  $\beta/\alpha$  ratio of  
25 7.9:1.0. We hypothesized that preactivation conditions<sup>61</sup> would be valuable in order to favor the  
26 formation of the  $\alpha$ -spiro intermediate and would thereby potentially enhance the  $\beta$ -selectivity.  
27 Unfortunately, premixing donor **2** with NIS/AgOTf before adding acceptor **9** led exclusively to  
28 glycal **16**, either in participating (CH<sub>3</sub>CN) or non-participating (DCE) solvents (entries 11 and  
29 12). The effect of counter-anions<sup>62,63</sup> was also studied. Therefore, promoters containing anions  
30 less-coordinating than OTf<sup>-</sup> such as NIS/AgClO<sub>4</sub> and NIS/AgBF<sub>4</sub> were evaluated (entries 13 and  
31 14). Using these promoters,  $\beta$ -selectivity decreased by more than two-fold, implying that the  
32 reaction intermediates were sensitive to the strength of the coordinating anion. On the basis of  
33 this result, we can hypothesize that a covalently-bound (or contact-ion pair) triflate could be one  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 of the intermediates involved in the glycosylation reaction although this has not been  
4  
5 experimentally proven.  
6  
7

8  
9  
10 These results were then compared to those obtained with donor **3** ( $\beta/\alpha$  ratio of  $\sim 1:1$ ) bearing a  
11 non-participating benzyl ester at C1 (Table 2). The advantage of using the Phen auxiliary group  
12 was clearly demonstrated here. Indeed, under the promotion of NIS/AgOTf, enhanced  $\beta$ -  
13 selectivity was obtained with donor **2** compared to donor **3** (entries 1 to 4). Notably, the  $\beta/\alpha$  ratio  
14 increased by more than two-fold when CH<sub>3</sub>CN was used as the solvent (7.9:1.0). Impact of the  
15 starting anomeric ratio of donors **2** and **3** was investigated next (entries 5 to 10). As shown in  
16 previous studies,<sup>28,29</sup> performing the reaction with Kdo thioglycosides as major  $\beta$ -anomer  
17 (7.0:1.0) significantly improved  $\beta$ -selectivity (up to  $\beta/\alpha$  11.0:1.0 in CH<sub>3</sub>CN, entry 8). For all of  
18 these reactions, donor **2** bearing a Phen group provided better  $\beta$ -selectivity than benzyl ester **3**  
19 and no glycal (**16** or **18**) was detected. As previously mentioned, the use of IBr instead of NIS  
20 decreased selectivity. Next, reactions were performed with  $\alpha$ -fluoride donors **4** and **5** in order to  
21 probe the impact of the leaving group (entries 11 and 12). Six equiv. of BF<sub>3</sub>·OEt<sub>2</sub> were needed to  
22 ensure full conversion of these fluorides.<sup>64</sup> Similarly to thioglycosides,  $\beta$ -selectivity was  
23 enhanced when the Phen group-containing donor **4** was used in comparison with donor **5**  
24 although the formation of glycals **16** and **18** was predominant.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 2. Synthesis of  $\beta$ -Kdo Glycosides: Influence of Phenacyl Group and Anomeric Ratio of Donors**




entry	donor ( $\beta/\alpha$ ratio)	solvent	conv. (%) <sup>a</sup>	selectivity ratio <sup>a</sup>		
				$\beta$	$\alpha$	glycal
1	<b>3</b> (1:1)	DCE	90	3.8	1.0	nd <sup>b</sup>
2	<b>2</b> (1:1)	DCE	>95	4.7	1.0	0.6
3	<b>3</b> (1:1)	CH <sub>3</sub> CN	93	3.4	1.0	nd
4	<b>2</b> (1:1)	CH <sub>3</sub> CN	>95	7.9	1.0	nd
5	<b>3</b> (7:1)	DCE	>95	4.8	1.0	nd
6	<b>2</b> (7:1)	DCE	>95	10.8	1.0	nd
7	<b>3</b> (7:1)	CH <sub>3</sub> CN	>95 (94) <sup>c</sup>	6.3	1.0	nd
8	<b>2</b> (7:1)	<b>CH<sub>3</sub>CN</b>	<b>&gt;95</b>	<b>11.0</b>	<b>1.0</b>	<b>nd</b>
9	<b>3</b> (7:1)	CH <sub>3</sub> CN <sup>d</sup>	>95	3.0	1.0	nd
10	<b>2</b> (7:1)	CH <sub>3</sub> CN <sup>d</sup>	>95	4.0	1.0	nd
11	<b>5</b> (0:1)	DCM <sup>e</sup>	>95	1.0	4.0	1.1
12	<b>4</b> (0:1)	DCM <sup>e</sup>	>95	1.0	1.0	1.4

<sup>a</sup>Determined by <sup>1</sup>H NMR analysis of the crude reaction mixture. <sup>b</sup>Not detected. <sup>c</sup>Performed at the gram scale. Value in parentheses corresponds to the isolated yield of **17 $\beta$**  and **17 $\alpha$** . <sup>d</sup>I<sub>2</sub> was used instead of NIS. <sup>e</sup>Fluoride donors **4** and **5** were activated with BF<sub>3</sub>·OEt<sub>2</sub> (6.0 equiv) and the reaction performed at 0 °C for 2 h.

The impact of phenacyl substitution on  $\beta$ -selectivity was also examined. As depicted in Scheme 3, thioglycoside donors **6** and **7**, bearing the methoxy group at positions 3' and 2' on the aromatic ring, respectively, as well as previously described **2** and unsubstituted **8** were coupled with acceptor **9** through the optimized glycosylation conditions, thereby generating Kdo glycosides **15**, and **19-21** with yields ranging from 73 to 80%. Owing to the electron-donating properties of the

methoxy group, it was anticipated that 4' and 2'-substituted Phen derivatives **2** and **7** would give the highest  $\beta/\alpha$  ratios on an electronic effect basis. Although this was true for donor **2**, only moderate  $\beta$ -selectivity (4.6:1.0) was obtained with donor **7**. Moreover, similar  $\beta/\alpha$  ratios were obtained for donors **6** and **8** (~6.4:1.0), as expected. These results mean that both electronic and steric effects could be responsible for the formation of the plausible  $\alpha$ -spiro intermediate, which accounts for the enhanced  $\beta$ -selectivity.


### Scheme 3. Impact of Phenacyl Substitution on $\beta/\alpha$ Ratio



Having studied various parameters modulating  $\beta$ -selectivity, we next investigated the general scope of the glycosylation reaction. In order to do so, a series of acceptors (Figure 3) featuring primary, secondary, or tertiary alcohols, including 1-nonanol (**10**), cyclohexanol (**11**), 2-adamantanol (**12**), 1-adamantanol (**13**), and methyl 2,3-di-*O*-benzyl- $\alpha$ -D-glucopyranoside (**14**) was reacted with thioglycoside donors **2** and **3** under the optimized conditions, *i.e.* NIS/AgOTf, CH<sub>3</sub>CN, -10 °C (Scheme 4). Using 1.4 equiv. of acceptors **10-14**, the reactions provided Kdo glycosides **22-31** with fair to very good yields (40-87%) while minimizing the formation of

glycols **16** and **18**. In all cases,  $\beta$ -Kdo glycosides were formed predominantly with the exception of the glycosylation of donor **3** with 1-adamantanol (**13**) that was moderately  $\alpha$ -selective ( $\beta/\alpha$  1.0:1.6). Once again, the effect of the Phen auxiliary group at C1 significantly enhanced  $\beta$ -selectivity in all cases (up to two-fold) compared to the non-participating benzyl ester. It is worth mentioning that using Phen donor **2** with acceptor **13** led to an inversion in selectivity giving a slight excess of  $\beta$ -Kdo glycoside **28** ( $\beta/\alpha$  1.6:1.0). Moreover, glycosylation with 4,6-diol **14** was fully regioselective at the C6 primary position.

#### Scheme 4. Scope of the Synthesis of $\beta$ -Kdo Glycosides using Different Acceptors




**Determination of Anomeric Configuration of Kdo Glycosides.** Since Kdo glycosides lack an anomeric proton at the C2 position, determination of anomeric configuration is not as

1  
2  
3 straightforward as it is for other glycosides and cannot rely only on  $^1\text{H}$  NMR analysis. One of the  
4  
5 most accurate method is the determination of the coupling constant between carbonyl carbon at  
6  
7 C1 and axial proton at C3.<sup>49</sup> For Kdo glycosides adopting a  $^5C_2$  conformation, which is the case  
8  
9 for glycosides **22-31**, a  $^3J_{\text{C1,H3ax}}$  value of 5.0-7.0 Hz is indicative of a  $\beta$ -configuration while a  
10  
11 value  $\leq 1.0$  Hz denotes an  $\alpha$ -configuration. Thus, the  $\alpha$ - or  $\beta$ -anomeric configuration of Kdo  
12  
13 glycosides **22-31** was determined via examination of this coupling constant obtained from an  
14  
15 undecoupled 150 MHz  $^{13}\text{C}$  NMR experiment. As expected,  $^3J_{\text{C1,H3ax}}$  values were found to be  
16  
17 between 5.0-7.0 Hz for  $\beta$ -glycosides and  $\leq 1.0$  Hz for  $\alpha$ -glycosides. Furthermore, an interesting  
18  
19 empirical observation was made by comparing the  $^1\text{H}$  NMR data of  $\alpha$ - and  $\beta$ -Kdo glycosides. We  
20  
21 found that the two geminal protons at C8 were closer (or superimposed) for  $\beta$ -Kdo glycosides  
22  
23 while the difference of the chemical shifts ( $\Delta\delta$ ) between H-8a and H-8b were more pronounced  
24  
25 for  $\alpha$ -Kdo glycosides ( $\Delta\delta$  from 0.46 to 0.54 ppm). This statement was true for all of the  
26  
27 synthesized Kdo glycosides. However,  $\Delta\delta$  values between H-3ax and H-3eq were not always  
28  
29 smaller for  $\alpha$ -glycosides, especially for 1- and 2-adamantyl glycosides **26-29**, and thus, as  
30  
31 recently emphasized by Mong,<sup>26</sup> this empirical method could not be reliably used for determining  
32  
33 the anomeric configuration of Kdo glycosides.  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 **Deprotection of Phenacyl Group.** As previously mentioned, one of the main advantage of using  
44  
45 a Phen auxiliary group lies in its possible chemoselective cleavage in the presence of other  
46  
47 protecting groups. As examples, Kdo glycosides **22** and **15** were reacted with activated Zn  
48  
49 powder in the presence of AcOH at 35 °C for 2 h (Scheme 5) to produce free carboxylic acids **32**  
50  
51 and **33** with very good yields (85 and 89%, respectively). The Phen derivatives were thus  
52  
53 deprotected chemoselectively in the presence of acetyl and NHCbz groups showing the  
54  
55 orthogonality of this auxiliary functionality. Importantly, other reaction conditions that have not  
56  
57  
58  
59  
60


been tested in the course of this study could also be suitable for the selective cleavage of phenacyl groups including  $(\text{Bu}_3\text{Sn})_2\text{O}$  in refluxing DCE; TBAF in THF;  $\text{H}_2$ , Pd/C; and photodeprotection.<sup>65</sup>

### Scheme 5. Zn-Mediated Cleavage of Phenacyl Group



**Proposed Mechanism.** On the basis of the above experimental results, DFT calculations, and literature precedent,<sup>66</sup> reaction mechanism and plausible intermediates were proposed, accounting for the formation of both  $\alpha$ - and  $\beta$ -Kdo glycosides. As shown in Figure 4, NIS would react with AgOTf to form an electrophilic species that would activate Kdo thioglycoside donors **2** or **3**. Following activation, oxocarbenium ion **I** would be formed together with EtSI and a proton acceptor imine. DFT calculations showed that, owing to the highly unstable nature of intermediate **I**, it is likely that acetyl group at C4 would stabilize ion **I** from the  $\beta$ -face, thereby forming dioxalenium ion intermediate **II** in the  $B_{3,6}$  conformation (Figure S1).  $\alpha$ -Triflate **III** would also be a plausible intermediate, which would be stabilized by the electron-withdrawing nature of acetyl groups and carboxylate at C1. Calculations revealed that triflate **III** would exist in the  ${}^5C_2$  conformation and be a stable intermediate. As previously discussed,  $\alpha$ -spiroPhen **IV** would be a plausible intermediate, which would be formed by the attack of the Phen activated ketone on the  $\alpha$ -face of oxocarbenium ion **I**. For Kdo donors bearing a Phen group, all these intermediates (**I** to **IV**) would exist in equilibrium reacting in different ways with the acceptor ( $\text{R}^1\text{OH}$ ). Glycosylation of oxocarbenium ion **I**, according to Woerpel model, as well as  $\beta$ -

dioxalenium ion **II** would produce  $\alpha$ -Kdo glycoside as the major anomer. On the other hand, glycosylation of  $\alpha$ -triflate **III** and  $\alpha$ -spiroPhen **IV** would occur from the  $\beta$ -face leading to the preferential formation of  $\beta$ -Kdo glycoside.  $\alpha$ -Triflate **III** would thus represent a plausible intermediate, accounting for the good  $\beta$ -selectivity obtained with Kdo donor **3** bearing a non-participating benzyl ester. Limitations of this mechanistic pathway include difficulties in explaining the impact of the starting anomeric ratio of donors giving enhanced  $\beta$ -selectivity for  $\beta$ -Kdo thioglycosides in comparison with their  $\alpha$ -counterparts (retention of configuration). Moreover, we cannot rule out the possibility that the generated *N*-succinimide (NHS) would trap the oxocarbenium ion forming a transient *N*-glycoside that could be displaced by the glycosyl acceptor.<sup>67-70</sup>



1  
2  
3 **Figure 4.** Proposed mechanism and plausible intermediates for the synthesis of  $\beta$ -Kdo  
4 glycosides. Dashed lines mean that species **II** to **IV** can be found either as covalent or contact-ion  
5  
6 pair intermediates.  
7  
8  
9

## 10 11 12 13 **CONCLUSIONS**

14  
15  
16  
17 In summary, a novel methodology was developed for the stereoselective formation of  $\beta$ -Kdo  
18 glycosides by taking advantage of the long-range participating effect of a 4'-methoxyphenacyl  
19 auxiliary group at the C1 position of a peracetylated Kdo thioglycoside donor. In addition to the  
20 positive effect of the Phen group, various parameters were shown to be crucial for enhancing the  
21  $\beta$ -selectivity of the reaction, including NIS/AgOTf as promoter, CH<sub>3</sub>CN as solvent,  $\beta$ -anomeric  
22 configuration of the donor, as well as *para*-substitution of the Phen aromatic ring. The optimized  
23 glycosylation conditions were applied to the synthesis of a series of Kdo glycosides, providing  
24 good yields and  $\beta$ -selectivity while minimizing the formation of undesirable glycals.  
25 Interestingly, chemoselective deprotection of the Phen group was achieved using activated  
26 Zn/HOAc, which represents an advantage over previously reported C1 auxiliary groups.  $\alpha$ -  
27 Triflate and six-membered  $\alpha$ -spiroPhen were postulated as plausible intermediates, accounting  
28 for the enhanced  $\beta$ -stereoselectivity obtained with Kdo thioglycoside donors bearing non-  
29 participating (Bn) or participating (Phen) groups at C1. The developed methodology could find  
30 application for the synthesis of  $\beta$ -Kdo-containing oligosaccharides from pathogenic bacteria.  
31  
32 Work towards this aim is currently in progress in our laboratory.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

## 55 **EXPERIMENTAL SECTION**

56  
57  
58  
59  
60

**General Methods.** All starting materials and reagents were purchased from commercial sources, and used as received without further purification. Air and water sensitive reactions were performed in heat gun-dried glassware under Ar atmosphere. Moisture sensitive reagents were introduced via a dry syringe. Anhydrous solvents were supplied over molecular sieves, and used as received. Petroleum ether (PE) refers to the 40-60 °C boiling fraction. Powdered 4 Å molecular sieves (MS) were activated before use by heating with a heat gun for  $\geq 5$  min under high vacuum. Reactions were monitored by thin-layer chromatography (TLC) with silica gel 60 F<sub>254</sub> 0.25 mm pre-coated aluminium foil plates. Compounds were visualized by using UV<sub>254</sub> and/or orcinol (1 mg·mL<sup>-1</sup>) in a 10% H<sub>2</sub>SO<sub>4</sub>(aq) solution and/or Hanessian's stain [2.5 g (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>·4H<sub>2</sub>O, 1.0 g Ce(NH<sub>4</sub>)<sub>4</sub>(SO<sub>4</sub>)<sub>4</sub>·2H<sub>2</sub>O, 90 mL H<sub>2</sub>O, 10 mL H<sub>2</sub>SO<sub>4</sub>] with heating. Normal-phase flash column chromatography was performed on silica gel 60 Å (15-40 μm). NMR spectra were recorded at 297 K in the indicated solvent (CDCl<sub>3</sub> or MeOD) with 400 or 600 MHz instruments, employing standard software provided by the manufacturer. <sup>1</sup>H and <sup>13</sup>C NMR spectra were referenced to tetramethylsilane (TMS,  $\delta_{\text{H}} = \delta_{\text{C}} = 0.00$  ppm) as internal reference for spectra in CDCl<sub>3</sub> and MeOD. Assignments were based on <sup>1</sup>H, <sup>13</sup>C, uncoupled <sup>13</sup>C, DEPT-135, COSY, HSQC, and HMBC experiments. High-resolution mass spectra (HRMS) were recorded on an ESI-Q-TOF mass spectrometer.

**Computational Method.** Reaction intermediates were first modeled using Sparta'10 V1.1.0 software package (Wavefunction Inc.). For each intermediate, a conformational distribution was generated through stochastic Monte-Carlo guided searches at the molecular mechanics (MMFF) level of theory.<sup>71</sup> Conformers within 25 kJ·mol<sup>-1</sup> of the most stable conformer were subjected to geometry optimizations with Gaussian 09.E01 software.<sup>72</sup> Calculations using DFT with the

1  
2  
3 hybrid B3LYP functional<sup>73,74</sup> and 6-31G(d,p) basis set<sup>75</sup> were performed with IEF-PCM model  
4  
5 solvent (CH<sub>3</sub>CN).<sup>76</sup> The most stable conformer of each intermediate was then further optimized  
6  
7 at the B3LYP/6-311++G(2d,2p) level of theory.<sup>77,78</sup> In these cases, the Grimme's empirical  
8  
9 dispersion correction was applied (D3 version).<sup>79</sup> Energy minima were confirmed at 263 K by  
10  
11 vibrational analysis at the same level of theory, which also allowed for calculation of the Gibbs  
12  
13 free energies. The 3D structures were rendered using PyMOL.  
14  
15  
16  
17  
18  
19

20 *Ammonium 2,4,5,7,8-Penta-O-acetyl-3-deoxy- $\alpha$ -D-manno-oct-2-ulopyranosylonate (1).*

21  
22 Crystalline ammonium Kdo<sup>48</sup> (1.39 g, 5.45 mmol, 1.0 equiv) was suspended in anhydrous py (55  
23  
24 mL), and then Ac<sub>2</sub>O (55 mL) followed by DMAP (6.6 mg, 54  $\mu$ mol, 0.01 equiv) were added. The  
25  
26 suspension was stirred for 16 h at rt under Ar after which time the solution was found to be  
27  
28 homogeneous. The mixture was concentrated under reduced pressure, keeping the temperature  
29  
30 below 50 °C, and coevaporated with toluene (3 $\times$ ). The residue was purified by silica gel flash  
31  
32 chromatography (DCM/MeOH 1:0 to 6:4) to give peracetylated Kdo (**1**, 2.40 g, 95%, ratio  $\alpha/\beta$  >  
33  
34 95:5) as a yellow oil. The physical and analytical data of **1**<sup>49</sup> were in agreement with those  
35  
36 published in the literature.  
37  
38  
39  
40  
41  
42

43 *4'-Methoxyphenacyl (Ethyl 4,5,7,8-Tetra-O-acetyl-3-deoxy-2-thio- $\alpha,\beta$ -D-manno-oct-2-*

44  
45 *ulopyranosid)onate (2). Route A:* 2-Bromo-4'-methoxyacetophenone (733 mg, 3.20 mmol, 1.6  
46  
47 equiv), TBAI (111 mg, 301  $\mu$ mol, 0.15 equiv) and Cs<sub>2</sub>CO<sub>3</sub> (1.33 g, 4.08 mmol, 2.0 equiv) were  
48  
49 successively added to a solution of peracetylated **1** (900 mg, 2.01 mmol, 1.0 equiv) in anhydrous  
50  
51 DMF (20 mL). The mixture was stirred for 16 h at rt under Ar. The suspension was filtered over  
52  
53 Celite, rinsed, and diluted with EtOAc (50 mL). The organic phase was washed with a saturated  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 NH<sub>4</sub>Cl(aq) solution (25 mL), and H<sub>2</sub>O (25 mL). The aqueous phase was back extracted with  
4  
5 EtOAc (25 mL). The combined organic layers were dried over MgSO<sub>4</sub>, and the solvents were  
6  
7 concentrated under reduced pressure. The residue was purified by silica gel flash chromatography  
8  
9 (PE/EtOAc 8:2 to 6:4) to give 4'-methoxyphenacyl (2,4,5,7,8-penta-*O*-acetyl-3-deoxy- $\alpha$ -D-  
10  
11 *manno*-oct-2-ulopyranosyl)onate (452 mg, 59%) as a white amorphous powder:  $[\alpha]_D^{20} = +84$  (*c*  
12  
13 0.54, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.91–7.86 (m, 2H, *CH*-Ar), 6.99–6.95 (m, 2H, *CH*-  
14  
15 Ar), 5.46 (d, *J* = 16.0 Hz, 1H, *CHH*Phen), 5.43–5.38 (m, 2H, H-4, H-5), 5.35 (d, *J* = 16.0 Hz, 1H,  
16  
17 *CHH*Phen), 5.25 (ddd, *J*<sub>6,7</sub> = 9.8 Hz, *J*<sub>7,8b</sub> = 3.6 Hz, *J*<sub>7,8a</sub> = 2.3 Hz, 1H, H-7), 4.48 (dd, *J*<sub>8a,8b</sub> = 12.4  
18  
19 Hz, *J*<sub>7,8a</sub> = 2.2 Hz, 1H, H-8a), 4.18 (dd, *J*<sub>6,7</sub> = 9.9 Hz, *J*<sub>5,6</sub> = 1.0 Hz, 1H, H-6), 4.13 (dd, *J*<sub>8a,8b</sub> =  
20  
21 12.4 Hz, *J*<sub>7,8b</sub> = 3.6 Hz, 1H, H-8b), 2.49 (dd, *J*<sub>3ax,3eq</sub> = 13.1 Hz, *J*<sub>3ax,4</sub> = 11.8 Hz, 1H, H-3ax), 2.39  
22  
23 (dd, *J*<sub>3ax,3eq</sub> = 13.2 Hz, *J*<sub>3eq,4</sub> = 5.1 Hz, 1H, H-3eq), 2.17, 2.12, 2.05, 2.02, 2.00 (all s, 15H, 5  $\times$   
24  
25 COCH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  189.7 (CH<sub>2</sub>CO), 170.5, 170.2, 170.1, 169.5, 167.9 (5  $\times$   
26  
27 COCH<sub>3</sub>), 165.8 (C-1), 164.3 (C-Ar), 130.1, 130.0 (CH-Ar), 127.0 (C-Ar), 114.2, 114.2 (CH-Ar),  
28  
29 97.3 (C-2), 69.5 (C-6), 67.3 (C-7), 66.6 (CH<sub>2</sub>CO), 65.9 (C-4), 63.9 (C-5), 62.1 (C-8), 55.6  
30  
31 (OCH<sub>3</sub>), 31.6 (C-3), 20.7, 20.7, 20.6, 20.6, 20.6 (5  $\times$  COCH<sub>3</sub>); HRMS (ESI-TOF) *m/z* [M +  
32  
33 NH<sub>4</sub>]<sup>+</sup> calcd for C<sub>27</sub>H<sub>36</sub>NO<sub>15</sub> 614.2082; found 614.2079; *m/z* [M + Na]<sup>+</sup> calcd for  
34  
35 C<sub>27</sub>H<sub>32</sub>NaO<sub>15</sub> 619.1633; found 619.1635. Ethanethiol (12  $\mu$ L, 170  $\mu$ mol, 2.0 equiv) was added to  
36  
37 a solution of the latter phenacyl (50 mg, 84  $\mu$ mol, 1.0 equiv) in anhydrous DCM (1.7 mL). The  
38  
39 solution was cooled to 0 °C; then, BF<sub>3</sub>·OEt<sub>2</sub> (16  $\mu$ L, 130  $\mu$ mol, 1.5 equiv) was added. The  
40  
41 mixture was stirred for 24 h under Ar, while gradually being warmed to rt. The solution was  
42  
43 diluted with DCM, and a saturated NaHCO<sub>3</sub>(aq) solution was added for neutralization. Then,  
44  
45 iodine was added until a dark color persisted. The excess of iodine was reduced by washing the  
46  
47 organic phase with a freshly prepared 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>(aq) solution until the red color disappeared.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The solution was poured into a separatory funnel, and the aqueous layer was extracted with DCM. The pooled organic phases were dried over  $\text{MgSO}_4$  and filtered, and the solvents were concentrated under reduced pressure. The residue was purified by silica gel flash chromatography (PE/EtOAc 1:0 to 5:55) to give thioglycoside **2** (13 mg, 25%) in a 9.0:1.0  $\beta/\alpha$  anomeric mixture.  $[\alpha]_D^{20} = +50$  ( $c$  0.83,  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  7.93–7.87 (m, 2H,  $\text{CH-Ar}$ ), 6.95 (m, 2H,  $\text{CH-Ar}$ ), 5.54–5.38 (m, 3H,  $\text{CHHPhen}$ , H-4, H-5), 5.37–5.28 (m, 1H,  $\text{CHHPhen}$ ), 5.28–5.20 (m, 1H, H-7), 4.36–4.33 (m, 2H, H-8a, H-8b), 4.24 (dd,  $J_{6,7} = 9.6$  Hz,  $J_{5,6} = 1.1$  Hz, 1H, H-6), 3.89 (s, 3H,  $\text{OCH}_3$ ), 2.91–2.77 (m, 1H,  $\text{SCHH}$ ), 2.74–2.66 (m, 1H,  $\text{SCHH}$ ), 2.63 (dd,  $J_{3\text{eq},3\text{ax}} = 12.6$  Hz,  $J_{3\text{eq},4} = 4.2$  Hz, 1H, H-3eq), 2.23 (t,  $J_{3\text{ax},3\text{eq}} \approx J_{3\text{ax},4} = 12.9$  Hz, 1H, H-3ax), 2.13, 2.03, 1.99, 1.97 (s, 12H,  $4 \times \text{COCH}_3$ ), 1.26 (t,  $J = 7.5$  Hz, 3H,  $\text{SCH}_2\text{CH}_3$ );  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ,  $\alpha$ -anomer)  $\delta$  7.93–7.87 (m, 2H,  $\text{CH-Ar}$ ), 6.95 (m, 2H,  $\text{CH-Ar}$ ), 5.54–5.38 (m, 3H,  $\text{CHHPhen}$ , H-4, H-5), 5.37–5.28 (m, 1H,  $\text{CHHPhen}$ ), 5.28–5.20 (m, 1H, H-7), 4.63 (dd,  $J_{8\text{a},8\text{b}} = 12.3$  Hz,  $J_{7,8\text{a}} = 2.4$  Hz, 1H, H-8a), 4.53 (dd,  $J_{6,7} = 9.7$  Hz,  $J_{5,6} = 1.3$  Hz, 1H, H-6), 4.12 (dd,  $J_{8\text{a},8\text{b}} = 12.3$  Hz,  $J_{7,8\text{b}} = 3.6$  Hz, 1H, H-8b), 3.89 (s, 3H,  $\text{OCH}_3$ ), 2.91–2.77 (m, 1H,  $\text{SCHH}$ ), 2.59–2.53 (m, 1H,  $\text{SCHH}$ ), 2.50 (dd,  $J_{3\text{ax},3\text{eq}} = 11.9$  Hz,  $J_{3\text{ax},4} = 10.6$  Hz, 1H, H-3ax), 2.33 (dd,  $J_{3\text{ax},3\text{eq}} = 13.7$  Hz,  $J_{3\text{eq},4} = 4.8$  Hz, 1H, H-3eq), 2.10, 2.08, 2.03, 2.01, (s, 12H,  $4 \times \text{COCH}_3$ ), 1.26 (t,  $J = 7.5$  Hz, 3H,  $\text{SCH}_2\text{CH}_3$ );  $^{13}\text{C NMR}$  (100 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  189.3 (CO), 170.8, 170.6, 169.9, 169.7 ( $4 \times \text{COCH}_3$ ), 167.8 (C-1), 164.3 (C-Ar), 130.1 (CH-Ar), 126.7 (C-Ar), 114.2 (CH-Ar), 83.9 (C-2), 72.1 (C-6), 67.9 (C-7), 67.2 (C-4), 66.9 ( $\text{CH}_2$ ), 64.4 (C-5), 62.5 (C-8), 55.6 ( $\text{OCH}_3$ ), 32.8 (C-3), 22.6 ( $\text{SCH}_2\beta$ ), 20.8–20.7 ( $4 \times \text{COCH}_3$ ), 14.1 ( $\text{SCH}_2\text{CH}_3$ );  $^{13}\text{C NMR}$  (100 MHz,  $\text{CDCl}_3$ ,  $\alpha$ -anomer)  $\delta$  189.6 (CO), 170.5, 170.4, 169.9, 169.7 ( $4 \times \text{COCH}_3$ ), 167.7 (C-1), 164.2 (C-Ar), 130.1 (CH-Ar), 127.0 (C-Ar), 114.2 (CH-Ar), 85.1 (C-2), 68.3 (C-6), 67.6 (C-7), 67.0 (C-4), 66.6 ( $\text{CH}_2$ ), 64.4 (C-5), 61.9 (C-8), 55.6 ( $\text{OCH}_3$ ), 31.9 (C-3), 23.5 ( $\text{SCH}_2$ ), 20.8–20.7

1  
2  
3 (4 × COCH<sub>3</sub>), 13.6 (SCH<sub>2</sub>CH<sub>3</sub>); HRMS (ESI-TOF)  $m/z$  [M + H]<sup>+</sup> calcd for C<sub>27</sub>H<sub>35</sub>O<sub>13</sub>S 599.1793;  
4  
5 found 599.1789;  $m/z$  [M + NH<sub>4</sub>]<sup>+</sup> calcd for C<sub>27</sub>H<sub>38</sub>NO<sub>13</sub>S 616.2058; found 616.2059;  $m/z$  [M +  
6  
7 Na]<sup>+</sup> calcd for C<sub>27</sub>H<sub>34</sub>NaO<sub>13</sub>S 621.1612; found 621.1611. *Route B*: Ethanethiol (261 μL, 3.53  
8  
9 mmol, 5.0 equiv) was added to a solution of peracetylated **1**<sup>49</sup> (328 mg, 705 μmol, 1.0 equiv) in  
10  
11 anhydrous DCE (7.1 mL) at rt under Ar. The solution was cooled to 0 °C and BF<sub>3</sub>·OEt<sub>2</sub> (183 μL,  
12  
13 1.48 mmol, 2.1 equiv) was slowly added. The mixture was refluxed for 40 min and then cooled at  
14  
15 rt prior adding Et<sub>3</sub>N (206 μL, 1.48 mmol, 2.1 equiv). The solvents were concentrated under  
16  
17 reduced pressure to give a residue, which was used in the next step without further purification.  
18  
19 2-Bromo-4'-methoxyacetophenone (322 mg, 1.41 mmol, 2.0 equiv) was added to a solution of  
20  
21 the crude thioglycoside in anhydrous DMF (5.6 mL) followed by Cs<sub>2</sub>CO<sub>3</sub> (69 mg, 212 μmol, 0.3  
22  
23 equiv) and the mixture was stirred for 2 h at rt under Ar. The suspension was filtered over Celite,  
24  
25 rinsed, and diluted with EtOAc (50 mL). The organic phase was washed with a saturated  
26  
27 NH<sub>4</sub>Cl(aq) solution (25 mL) and H<sub>2</sub>O (25 mL). The combined organic layers were dried over  
28  
29 MgSO<sub>4</sub>, and the solvents were concentrated under reduced pressure. The residue was purified by  
30  
31 silica gel flash chromatography (PE/EtOAc 9:1 to 6:4) to give thioglycoside **6** (190 mg, 45%, two  
32  
33 steps) as a yellow oil in a ~1:1 β/α anomeric mixture. *Route C: Representative Procedure for the*  
34  
35 *Synthesis of Phenacyl Derivatives starting from Benzyl Ester 3*. Thioglycoside **3** (965 mg, 1.79  
36  
37 mmol, 1.0 equiv, ratio β/α 7:1) was dissolved in anhydrous DCE/MeOH (35 mL, 1:4 v/v). The  
38  
39 solution was degassed with Ar, and 10% Pd/C (965 mg) was added. The suspension was stirred  
40  
41 under an atmosphere of H<sub>2</sub> at rt for 2 h. The mixture was filtered over Celite to remove the  
42  
43 catalyst, and the cake was rinsed with MeOH and DCM. The solvents were concentrated under  
44  
45 reduced pressure to give a residue (804 mg, quant.) as a yellow oil, which was used in the next  
46  
47 step without further purification. 2-Bromo-4'-methoxyacetophenone (61 mg, 268 μmol, 1.5  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 equiv) was added to a solution of crude carboxylic acid (80 mg, 180  $\mu\text{mol}$ ) in anhydrous DMF  
4 (0.9 mL) followed by  $\text{Cs}_2\text{CO}_3$  (64 mg, 196  $\mu\text{mol}$ , 1.1 equiv). The mixture was stirred for 2 h at rt  
5  
6 under Ar. The suspension was filtered over Celite, rinsed, and diluted with EtOAc (100 mL). The  
7  
8 organic phase was washed with a saturated  $\text{NH}_4\text{Cl}(\text{aq})$  solution (50 mL) and  $\text{H}_2\text{O}$  (50 mL). The  
9  
10 combined organic layers were dried over  $\text{MgSO}_4$ , and the solvents were concentrated under  
11  
12 reduced pressure. The residue was purified by silica gel flash chromatography (PE/EtOAc 1:0 to  
13  
14 6:4) to give thioglycoside **2** (81 mg, 76%) as a colorless oil in a 7:1  $\beta/\alpha$  anomeric mixture.  
15  
16  
17  
18  
19

20  
21  
22 *Benzyl (Ethyl 4,5,7,8-Tetra-O-acetyl-3-deoxy-2-thio- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid)onate (3).*  
23

24  
25 *Route A:* BnBr (848  $\mu\text{L}$ , 7.09 mmol, 2.2 equiv) followed by  $\text{Cs}_2\text{CO}_3$  (420 mg, 1.29 mmol, 0.4  
26  
27 equiv) were added to a solution of peracetylated **1**<sup>49</sup> (1.5 g, 3.22 mmol, 1.0 equiv) in anhydrous  
28  
29 DMF (16 mL). The mixture was stirred for 2 h at rt under Ar. The suspension was filtered over  
30  
31 Celite, rinsed, and diluted with EtOAc (50 mL). The organic phase was washed with a saturated  
32  
33  $\text{NH}_4\text{Cl}(\text{aq})$  solution (25 mL) and  $\text{H}_2\text{O}$  (25 mL). The combined organic layers were dried over  
34  
35  $\text{MgSO}_4$ , and the solvents were concentrated under reduced pressure. The residue was purified by  
36  
37 silica gel flash chromatography (PE/EtOAc 8:2 to 7:3) to give benzyl (2,4,5,7,8-penta-O-acetyl-  
38  
39 3-deoxy- $\alpha$ -D-manno-oct-2-ulopyranosyl)onate (1.26 g, 73%) as a white foam. The physical and  
40  
41 analytical data of **3**<sup>80</sup> were in agreement with those published in the literature. Ethanethiol (286  
42  
43  $\mu\text{L}$ , 3.86 mmol, 2.0 equiv) was added to a solution of the latter benzyl ester (1.04 g, 1.93 mmol,  
44  
45 1.0 equiv) in anhydrous DCE (10 mL). The solution was cooled to 0  $^\circ\text{C}$ ; then,  $\text{BF}_3\cdot\text{OEt}_2$  (357  $\mu\text{L}$ ,  
46  
47 2.90 mmol, 1.5 equiv) was slowly added. The mixture was stirred for 2 h under Ar, while  
48  
49 gradually being warmed to rt. The solution was diluted with DCM, and a saturated  $\text{NaHCO}_3(\text{aq})$   
50  
51 solution was added for neutralization. Then, iodine was added until a dark color persisted. The  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 excess of iodine was reduced by washing the organic phase with a freshly prepared 10%  
4  
5 Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>(aq) solution until the red color disappeared. The solution was poured into a separatory  
6  
7 funnel, and the aqueous layer was extracted with DCM. The pooled organic phases were dried  
8  
9 over MgSO<sub>4</sub> and filtered, and the solvents were concentrated under reduced pressure. The residue  
10  
11 was purified by silica gel flash chromatography (PE/EtOAc 85:15 to 75:25) to give thioglycoside  
12  
13 **3** (956 mg, 92%) as a 7.0:1.0 β/α anomeric mixture. The physical and analytical data of **3**<sup>10</sup> were  
14  
15 in agreement with those published in the literature. *Route B:* Ethanethiol (307 μL, 4.14 mmol, 5.0  
16  
17 equiv) was added to a solution a peracetylated **1**<sup>49</sup> (386 mg, 829 μmol, 1.0 equiv) in anhydrous  
18  
19 DCE (8.0 mL) at rt under Ar. The solution was cooled to 0 °C and BF<sub>3</sub>·OEt<sub>2</sub> (95 μL, 770 μmol,  
20  
21 2.0 equiv) was slowly added. The mixture was refluxed for 40 min and then cooled at rt prior  
22  
23 adding Et<sub>3</sub>N (230 μL, 4.15 mmol, 5.0 equiv). The solvents were concentrated under reduced  
24  
25 pressure to give a residue, which was used in the next step without further purification. Benzyl  
26  
27 bromide (218 μL, 1.82 mmol, 2.2 equiv) was added to a solution of the crude thioglycoside in  
28  
29 anhydrous DMF (4.1 mL) followed by Cs<sub>2</sub>CO<sub>3</sub> (108 mg, 332 μmol, 0.4 equiv) and the mixture  
30  
31 was stirred for 2 h at rt under Ar. The suspension was filtered over Celite, rinsed, and diluted with  
32  
33 EtOAc (50 mL). The organic phase was washed with a saturated NH<sub>4</sub>Cl(aq) solution (25 mL) and  
34  
35 H<sub>2</sub>O (25 mL). The combined organic layers were dried over MgSO<sub>4</sub>, and the solvents were  
36  
37 concentrated under reduced pressure. The residue was purified by silica gel flash chromatography  
38  
39 (PE/EtOAc 9:1 to 6:4) to give thioglycoside **3** (224 mg, 50%, two steps) as a yellow oil in a ~1:1  
40  
41 β/α anomeric mixture.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 *4'-Methoxyphenacyl* (Fluoride *4,5,7,8-Tetra-O-acetyl-3-deoxy-α-D-manno-oct-2-*  
54  
55 *ulopyranosyl)onate (4)*. HF.py (5.0 mL, ca. 70% HF, ca. 30% py) was added to a solution of  
56  
57  
58  
59  
60

peracetylated **1**<sup>49</sup> (675 mg, 1.59 mmol, 1.0 equiv) in anhydrous DCM (16 mL) at 0 °C under Ar. The mixture was stirred for 3 h, while gradually being warmed to rt. Ice-cold water (10 mL) was added and the phases were mixed and left separated. The aqueous phase was acidified with 10% HCl(aq) until pH ~3 and extracted with DCM (3 × 10 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and the solvents were concentrated under reduced pressure to give a crude fluoride (391 mg, 63%) as a colorless oil after being placed under high vacuum for 3 h. A portion of the latter compound (191 mg, 450 μmol, 1.0 equiv) was dissolved in anhydrous DMF (2.3 mL). 2-Bromo-4'-methoxyacetophenone (226 mg, 989 μmol, 2.2 equiv) followed by Cs<sub>2</sub>CO<sub>3</sub> (59 mg, 180 μmol, 0.4 equiv) were added and the mixture was stirred for 2 h at rt under Ar. The suspension was filtered over Celite, rinsed, and diluted with EtOAc (50 mL). The organic phase was washed with a saturated NH<sub>4</sub>Cl(aq) solution (25 mL) and H<sub>2</sub>O (25 mL). The combined organic layers were dried over MgSO<sub>4</sub>, and the solvents were concentrated under reduced pressure. The residue was purified by silica gel flash chromatography (PE/EtOAc 1:0 to 6:4) to give α-fluoride **4** (192 mg, 77%) as a yellow oil.  $[\alpha]_D^{20} = +35$  (*c* 0.75, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.91–7.87 (m, 2H, CH-Ar), 7.00–6.96 (m, 2H, CH-Ar), 5.51–5.44 (m, 3H, H-5, CH<sub>2</sub>Phen), 5.39 (ddd,  $J_{3ax,4} = 12.1$  Hz,  $J_{4,5} = 5.4$  Hz,  $J_{3eq,4} = 3.0$  Hz, 1H, H-4), 5.26 (dddd,  $J_{6,7} = 9.6$  Hz,  $J_{7,8b} = 4.4$  Hz,  $J_{7,8a} = 2.3$  Hz,  $J_{5,7} = 0.8$  Hz, 1H, H-7), 4.51 (dd,  $J_{8a,8b} = 12.4$  Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-8a), 4.42 (dd,  $J_{6,7} = 9.7$  Hz,  $J_{5,6} = 1.3$  Hz, 1H, H-6), 4.17 (dd,  $J_{8a,8b} = 12.4$  Hz,  $J_{7,8b} = 4.4$  Hz, 1H, H-8b), 3.89 (s, 3H, OCH<sub>3</sub>), 2.63–2.41 (m, 2H, H-3ax, H-3eq), 2.12, 2.09, 2.02, 2.02 (all s, 12H, 4 × COCH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 188.9 (COPhen), 170.7, 170.5, 170.0, 169.8 (4 × COCH<sub>3</sub>), 164.3 (C-1,  $^2J_{C1,F} = 41.8$  Hz), 130.2 (2 × CH-Ar), 126.9 (C-Ar), 114.4 (2 × CH-Ar), 108.3 (C-2,  $^1J_{C2,F} = 232$  Hz), 70.7 (C-6,  $^3J_{C6,F} = 1.5$  Hz), 67.5 (C-7), 67.1 (CH<sub>2</sub>Phen), 65.6 (C-4), 64.1 (C-5), 62.2 (C-8), 55.7 (OCH<sub>3</sub>), 30.7 (C-3,  $^2J_{C3,F} = 27.3$  Hz), 20.84 (2C), 20.78,

20.77 (4 × COCH<sub>3</sub>); <sup>19</sup>F (376 MHz, CDCl<sub>3</sub>) δ 376.5 (dd, <sup>3</sup>J<sub>F,H3ax</sub> = 34.6 Hz, <sup>3</sup>J<sub>F,3eq</sub> = 6.0 Hz); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup> calcd for C<sub>25</sub>H<sub>29</sub>FNaO<sub>13</sub> 579.1484; found 579.1494.

*Benzyl (Fluoride 4,5,7,8-Tetra-O-acetyl-3-deoxy-α-D-manno-oct-2-ulopyranosyl)onate (5).*

HF.py (5.0 mL, ca. 70% HF, ca. 30% py) was added to a solution of peracetylated **1**<sup>49</sup> (675 mg, 1.59 mmol, 1.0 equiv) in anhydrous DCM (16 mL) at 0 °C under Ar. The mixture was stirred for 3 h, while gradually being warmed to rt. Ice-cold water (10 mL) was added and the phases were mixed and left separated. The aqueous phase was acidified with 10% HCl(aq) until pH ~3 and extracted with DCM (3 × 10 mL). The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and the solvents were concentrated under reduced pressure to give a crude fluoride (391 mg, 63%) as a colorless oil after being placed under high vacuum for 3 h. A portion of the latter compound (191 mg, 450 μmol, 1.0 equiv) was dissolved in anhydrous DMF (2.3 mL). Benzyl bromide (118 μL, 989 μmol, 2.2 equiv) followed by Cs<sub>2</sub>CO<sub>3</sub> (59 mg, 180 μmol, 0.4 equiv) were added and the mixture was stirred for 2 h at rt under Ar. The suspension was filtered over Celite, rinsed, and diluted with EtOAc (50 mL). The organic phase was washed with a saturated NH<sub>4</sub>Cl(aq) solution (25 mL) and H<sub>2</sub>O (25 mL). The combined organic layers were dried over MgSO<sub>4</sub>, and the solvents were concentrated under reduced pressure. The residue was purified by silica gel flash chromatography (PE/EtOAc 1:0 to 55:45) to give α-fluoride **5** (159 mg, 71%) as a colorless oil. [ $\alpha$ ]<sub>D</sub><sup>20</sup> = +50 (*c* 1.9, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.44–7.33 (m, 5H, CH-Ar), 5.44–5.42 (m, 1H, H-5), 5.33 (ddd, *J*<sub>3ax,4</sub> = 11.9 Hz, *J*<sub>4,5</sub> = 5.4 Hz, *J*<sub>3eq,4</sub> = 3.0 Hz, 1H, H-4), 5.29 (d, *J* = 10.1 Hz, 1H, CHHPh), 5.28 (d, *J* = 9.4 Hz, 1H, CHHPh), 5.21 (dddd, *J*<sub>6,7</sub> = 9.7 Hz, *J*<sub>7,8b</sub> = 4.4 Hz, *J*<sub>7,8a</sub> = 2.3 Hz, *J*<sub>5,7</sub> = 1.0 Hz, 1H, H-7), 4.49 (dd, *J*<sub>8a,8b</sub> = 12.3 Hz, *J*<sub>7,8a</sub> = 2.3 Hz, 1H, H-8a), 4.38 (dd, *J*<sub>6,7</sub> = 9.7 Hz, *J*<sub>5,6</sub> = 1.3 Hz, 1H, H-6), 4.13 (dd, *J*<sub>8a,8b</sub> = 12.3 Hz, *J*<sub>7,8b</sub> = 4.4 Hz,

1  
2  
3 1H, H-8b), 2.44–2.23 (m, 2H, H-3ax, H-3eq), 2.10, 2.07, 2.01, 2.00 (all s, 12H, 4 × COCH<sub>3</sub>); <sup>13</sup>C  
4 NMR (100 MHz, CDCl<sub>3</sub>) δ 170.7, 170.4, 169.9, 169.8 (4 × COCH<sub>3</sub>), 164.4 (C-1, <sup>2</sup>J<sub>C1,F</sub> = 29.7  
5 Hz), 134.7 (C-Ar), 128.9, 128.8, 128.3 (5 × CH-Ar), 108.1 (C-2, <sup>1</sup>J<sub>C2,F</sub> = 232 Hz), 70.7 (C-6,  
6 <sup>3</sup>J<sub>C6,F</sub> = 2.2 Hz), 68.2 (CH<sub>2</sub>Ph), 67.4 (C-7), 65.7 (C-4), 64.1 (C-5), 62.2 (C-8), 30.3 (C-3, <sup>2</sup>J<sub>C3,F</sub> =  
7 27.6 Hz), 20.82 (2C), 20.77, 20.7 (4 × COCH<sub>3</sub>); <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ 376.5 (dd, <sup>3</sup>J<sub>F,H3ax</sub>  
8 = 34.1 Hz, <sup>3</sup>J<sub>F,H3eq</sub> = 5.4 Hz); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup> calcd for  
9 C<sub>23</sub>H<sub>27</sub>FNaO<sub>11</sub> 521.1430; found 521.1421; *m/z* [2M + Na]<sup>+</sup> calcd for C<sub>46</sub>H<sub>54</sub>F<sub>2</sub>NaO<sub>22</sub> 1019.2967;  
10 found 1019.2944.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24  
25 *3'-Methoxyphenacyl (Ethyl 4,5,7,8-Tetra-O-acetyl-3-deoxy-2-thio- $\alpha,\beta$ -D-manno-oct-2-*  
26 *ulopyranosid)onate (6)*. Thioglycoside **3** ( $\beta/\alpha$  7:1, 163 mg, 303  $\mu$ mol, 1.0 equiv) was reacted  
27 according to the representative procedure for the synthesis of phenacyl derivatives starting from  
28 benzyl ester **3** and gave thioglycoside **6** (147 mg, 81%, two steps,  $\beta/\alpha$  7:1) as a white foam.  
29  
30  
31  
32  
33  
34  $[\alpha]_D^{20} = +74$  (*c* 0.23, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>,  $\beta$ -anomer) δ 7.49–7.45 (m, 2H, CH-  
35 Ar), 7.44–7.39 (m, 1H, CH-Ar), 7.20–7.16 (m, 1H, CH-Ar), 5.54 (d, *J* = 16.2 Hz, 1H,  
36 CHHPhen), 5.48–5.41 (m, 2H, H-5, H-4), 5.37 (d, *J* = 16.2 Hz, 1H, CHHPhen), 5.23 (ddd, *J*<sub>6,7</sub> =  
37 9.6 Hz, *J*<sub>7,8b</sub> = 5.2 Hz, *J*<sub>7,8a</sub> = 2.9 Hz, 1H, H-7), 4.36–4.33 (m, 2H, H-8a, H-8b), 4.22 (d, *J*<sub>6,7</sub> = 9.6  
38 Hz, *J*<sub>5,6</sub> = 1.0 Hz, 1H, H-6), 3.87 (s, 3H, OCH<sub>3</sub>), 2.91–2.81 (m, 1H, SCHH), 2.73–2.66 (m, 1H,  
39 SCHH), 2.63 (ddd, *J*<sub>3ax,3eq</sub> = 12.5 Hz, *J*<sub>3eq,4</sub> = 4.2 Hz, *J*<sub>3eq,5</sub> = 0.9 Hz, 1H, H-3eq), 2.24 (t, *J*<sub>3ax,3eq</sub> ≈  
40 *J*<sub>3ax,4</sub> ≈ 12.5 Hz, 1H, H-3ax), 2.13, 2.03, 2.00, 1.98 (all s, 12H, 4 × COCH<sub>3</sub>), 1.27 (t, *J* = 7.6 Hz,  
41 3H, SCH<sub>2</sub>CH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>,  $\beta$ -anomer) δ 190.9 (COPhen), 170.9, 169.9, 169.8,  
42 167.9 (4 × COCH<sub>3</sub>), 167.9 (C-1), 160.3, 135.2 (2 × C-Ar), 130.1, 121.0, 120.3, 112.1 (4 × CH-  
43 Ar), 84.1 (C-2), 72.3 (C-6), 68.1 (C-7), 67.4 (C-4), 67.3 (CH<sub>2</sub>Phen), 64.5 (C-5), 62.7 (C-8), 55.7  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

(OCH<sub>3</sub>), 33.0 (C-3), 23.6 (SCH<sub>2</sub>), 20.9–20.8 (4 × COCH<sub>3</sub>), 14.2 (SCH<sub>2</sub>CH<sub>3</sub>); HRMS (ESI-TOF)  $m/z$  [M + Na]<sup>+</sup> calcd for C<sub>27</sub>H<sub>34</sub>NaO<sub>13</sub>S 621.1612; found 621.1618;  $m/z$  [2M + Na]<sup>+</sup> calcd for C<sub>54</sub>H<sub>68</sub>NaO<sub>26</sub>S<sub>2</sub> 1219.3332; found 129.3336.

*2'-Methoxyphenacyl* (Ethyl 4,5,7,8-Tetra-O-acetyl-3-deoxy-2-thio- $\alpha,\beta$ -D-manno-oct-2-*ulopyranosid)onate* (7). Thioglycoside **3** ( $\beta/\alpha$  7:1, 150 mg, 279  $\mu$ mol, 1.0 equiv) was reacted according to the representative procedure for the synthesis of phenacyl derivatives starting from benzyl ester **3** and gave thioglycoside **7** (134 mg, 80%, two steps,  $\beta/\alpha$  7:1) as a white foam.  $[\alpha]_D^{20} = +52$  (*c* 0.7, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>,  $\beta$ -anomer)  $\delta$  8.03–8.00 (m, 1H, CH-Ar), 7.58–7.53 (m, 1H, CH-Ar), 7.09–6.99 (m, 2H, CH-Ar), 5.48–5.43 (m, 2H, H-5, H-4), 5.41 (d, *J* = 17.1 Hz, 1H, CHHPhen), 5.32 (d, *J* = 17.1 Hz, 1H, CHHPhen), 5.26–5.20 (m, 1H, H-7), 4.36–4.34 (m, 2H, H-8a, H-8b), 4.26 (dd, *J*<sub>6,7</sub> = 9.6 Hz, *J*<sub>5,6</sub> = 1.1 Hz, 1H, H-6), 3.98 (s, 3H, OCH<sub>3</sub>), 2.93–2.83 (m, 1H, SCHH), 2.77–2.67 (m, 1H, SCHH), 2.63 (ddd, *J*<sub>3ax,3eq</sub> = 12.5 Hz, *J*<sub>3eq,4</sub> = 4.6 Hz, *J*<sub>3eq,5</sub> = 1.0 Hz, 1H, H-3eq), 2.22 (t, *J*<sub>3ax,3eq</sub>  $\approx$  *J*<sub>3ax,4</sub>  $\approx$  12.5 Hz, 1H, H-3ax), 2.13, 2.03, 1.99 (all s, 12H, 4 × COCH<sub>3</sub>), 1.27 (t, *J* = 7.5 Hz, 3H, SCH<sub>2</sub>CH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>,  $\beta$ -anomer)  $\delta$  191.6 (COPhen), 170.9, 170.7, 170.0, 169.8 (4 × COCH<sub>3</sub>), 168.0 (C-1), 159.8 (C-Ar), 135.4, 131.5 (2 × CH-Ar), 123.9 (C-Ar), 121.3, 111.7 (2 × CH-Ar), 84.2 (C-2), 72.2 (C-6), 71.2 (CH<sub>2</sub>Phen), 68.1 (C-7), 67.4 (C-4), 64.6 (C-5), 62.8 (C-8), 55.8 (OCH<sub>3</sub>), 33.0 (C-3), 23.6 (SCH<sub>2</sub>CH<sub>3</sub>), 20.9–20.8 (4 × COCH<sub>3</sub>), 14.2 (SCH<sub>2</sub>CH<sub>3</sub>); HRMS (ESI-TOF)  $m/z$  [M + Na]<sup>+</sup> calcd for C<sub>27</sub>H<sub>34</sub>NaO<sub>13</sub>S 621.1612; found 621.1622;  $m/z$  [2M + Na]<sup>+</sup> calcd for C<sub>54</sub>H<sub>68</sub>NaO<sub>26</sub>S<sub>2</sub> 1219.3332; found 1219.3340.

1  
2  
3  
4 *Phenacyl (Ethyl 4,5,7,8-Tetra-O-acetyl-3-deoxy-2-thio- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid)onate*  
5  
6 (8). Thioglycoside **3** ( $\beta/\alpha$  7:1, 174 mg, 322  $\mu$ mol, 1.0 equiv) was reacted according to the  
7  
8 representative procedure for the synthesis of phenacyl derivatives starting from benzyl ester **3** and  
9  
10 gave thioglycoside **8** (129 mg, 70%, two steps,  $\beta/\alpha$  7:1) as a white foam.  $[\alpha]_D^{20} = +58$  (*c* 0.35,  
11  
12 CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>,  $\beta$ -anomer)  $\delta$  7.95–7.91 (m, 2H, CH-Ar), 7.67–7.62 (m, 1H,  
13  
14 CH-Ar), 7.55–7.49 (m, 2H, CH-Ar), 5.55 (d, *J* = 16.2 Hz, 1H, CHHPhen), 5.48–5.43 (m, 2H, H-  
15  
16 5, H-4), 5.39 (d, *J* = 16.2 Hz, 1H, CHHPhen), 5.23 (ddd, *J*<sub>6,7</sub> = 9.5 Hz, *J*<sub>7,8b</sub> = 3.9 Hz, *J*<sub>7,8a</sub> = 2.8  
17  
18 Hz, 1H, H-7), 4.36–4.33 (m, 2H, H-8a, H-8b), 4.22 (dd, *J*<sub>6,7</sub> = 9.6 Hz, *J*<sub>5,6</sub> = 1.0 Hz, 1H, H-6),  
19  
20 2.91–2.81 (m, 1H, SCHH), 2.73–2.65 (m, 1H, SCHH), 2.64 (ddd, *J*<sub>3ax,3eq</sub> = 12.5 Hz, *J*<sub>3eq,4</sub> = 4.2  
21  
22 Hz, *J*<sub>3eq,5</sub> = 1.0 Hz, 1H, H-3eq), 2.24 (t, *J*<sub>3ax,3eq</sub>  $\approx$  *J*<sub>3ax,4</sub>  $\approx$  12.5 Hz, 1H, H-3ax), 2.13, 2.03, 2.00,  
23  
24 1.98 (all s, 12H, 4  $\times$  COCH<sub>3</sub>), 1.27 (t, *J* = 7.6 Hz, 3H, SCH<sub>2</sub>CH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>,  
25  
26  $\beta$ -anomer)  $\delta$  191.1 (COPhen), 170.9, 170.7, 170.0, 169.8 (4  $\times$  COCH<sub>3</sub>), 167.9 (C-1), 134.4 (CH-  
27  
28 Ar), 133.9 (C-Ar), 129.1 (2C), 127.9 (2C, 4  $\times$  CH-Ar), 84.1 (C-2), 72.3 (C-6), 68.1 (C-7), 67.4  
29  
30 (C-4), 67.2 (CH<sub>2</sub>Phen), 64.5 (C-5), 62.7 (C-8), 33.0 (C-3), 23.6 (SCH<sub>2</sub>), 20.90, 20.89, 20.85,  
31  
32 20.80 (4  $\times$  COCH<sub>3</sub>), 14.2 (SCH<sub>2</sub>CH<sub>3</sub>); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup> calcd for C<sub>26</sub>H<sub>32</sub>NaO<sub>12</sub>S  
33  
34 591.1507; found 591.1508; *m/z* [2M + Na]<sup>+</sup> calcd for C<sub>52</sub>H<sub>64</sub>NaO<sub>24</sub>S<sub>2</sub> 1159.3121; found  
35  
36 1159.3117.  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 **General Procedure for Glycosylation with Thioglycoside Donors.** Freshly activated powdered  
47  
48 4 Å molecular sieves (4 mg·mg<sup>-1</sup> of acceptor) was added to a solution of thioglycoside **2** or **3** (1  
49  
50 equiv), glycosyl acceptor **9–14** (1.4–2.0 equiv), and NIS (2.0 equiv) in anhydrous CH<sub>3</sub>CN (20  
51  
52 mL·mmol<sup>-1</sup>). The mixture was stirred for 1 h at rt under Ar. Then, the suspension was cooled to –  
53  
54 10 °C; the flask was protected from light, and AgOTf (1.0 equiv) was added in one portion. The  
55  
56  
57  
58  
59  
60

1  
2  
3 mixture was stirred for 30 min at  $-10\text{ }^{\circ}\text{C}$  under Ar.  $\text{Et}_3\text{N}$  (2.0 equiv) was added to quench the  
4  
5 reaction. The suspension was filtered over Celite, rinsed with DCM, and the solvents were  
6  
7 concentrated under reduced pressure. The residue was purified by silica gel flash chromatography  
8  
9 to give a mixture of Kdo glycoside anomers.  
10  
11

12  
13  
14 **General Procedure for Glycosylation with Fluoride Donors.** Freshly activated powdered 4 Å  
15  
16 molecular sieves (4 mg·mg<sup>-1</sup> of acceptor) was added to a solution of fluoride **4** or **5** (1 equiv) and  
17  
18 acceptor **9** (2.0 equiv) in anhydrous DCM (25 mL·mmol<sup>-1</sup>). The mixture was stirred for 1 h at rt  
19  
20 under Ar. Then, the suspension was cooled to  $0\text{ }^{\circ}\text{C}$  and  $\text{BF}_3\cdot\text{OEt}_2$  (6.0 equiv) was slowly added.  
21  
22 The mixture was stirred from  $0\text{ }^{\circ}\text{C}$  to rt for 2 h or until TLC had showed complete conversion of  
23  
24 the donor. Then, the suspension was filtered over Celite, rinsed and diluted with DCM. The  
25  
26 organic phase was washed with a saturated  $\text{NaHCO}_3(\text{aq})$  solution and brine. The organic phase  
27  
28 was dried over  $\text{MgSO}_4$ , filtered, and the solvents were concentrated under reduced pressure. The  
29  
30 residue was purified by silica gel flash chromatography to give a mixture of Kdo glycoside  
31  
32 anomers together with glycal.  
33  
34  
35  
36  
37  
38  
39  
40

41 *4'-Methoxyphenacyl* [2-(5-Amino-N-benzyloxycarbonyl-1-pentyl) 4,5,7,8-Tetra-O-acetyl-3-  
42  
43 deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid]onate (**15 $\beta$**  and **15 $\alpha$** ) and *4'-Methoxyphenacyl* 4,5,7,8-  
44  
45 Tetra-O-acetyl-2,6-anhydro-3-deoxy-D-manno-oct-2-enosonate (**16**). Thioglycoside **2** (750 mg,  
46  
47 1.25 mmol, 1.0 equiv,  $\beta/\alpha \sim 1:1$ ) and acceptor **9** (595 mg, 2.51 mmol, 2.0 equiv) were reacted  
48  
49 according to the general procedure for glycosylation with thioglycoside donors and gave **15** (827  
50  
51 mg, 84%,  $\beta/\alpha$  7.9:1.0) as a colorless oil.  $[\alpha]_{\text{D}}^{20} = -11$  ( $c$  1.4,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ,  
52  
53  $\beta$ -anomer)  $\delta$  7.92–7.86 (m, 2H, CH-Ar), 7.38–7.28 (m, 5H, CH-Ar), 6.98–6.92 (m, 2H, CH-Ar),  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 5.44 (d,  $J = 16.0$  Hz, 1H, *CHHPhen*), 5.40–5.38 (m, 1H, H-5), 5.38 (d,  $J = 16.0$  Hz, 1H,  
4 *CHHPhen*), 5.32 (ddd,  $J_{3ax,4} = 13.0$  Hz,  $J_{3eq,4} = 4.6$  Hz,  $J_{4,5} = 3.0$  Hz, 1H, H-4), 5.21 (ddd,  $J_{6,7} =$   
5 13.0 Hz,  $J_{7,8a} = 4.6$  Hz,  $J_{7,8b} = 3.0$  Hz, 1H, H-7), 5.11–5.07 (m, 2H,  $COCH_2Ph$ ), 5.02–4.93 (m,  
6 1H, *NHCbz*), 4.42–4.29 (m, 3H, H-8a, H-8b, H-6), 3.82 (dt,  $J = 9.2, 6.3$  Hz, 1H, H-1a'), 3.52 (dt,  
7  $J = 9.2, 6.3$  Hz, 1H, H-1b'), 3.24–3.18 (m, 2H, H-5ab'), 2.52 (dd,  $J_{3eq,3ax} = 12.5$  Hz,  $J_{3eq,4} = 4.7$   
8 Hz, 1H, H-3eq), 2.16 (t,  $J_{3eq,3ax} \approx J_{3ax,4} \approx 12.8$  Hz, 1H, H-3ax), 2.11, 2.02, 2.00, 1.99 (all s, 12H, 4  
9  $\times COCH_3$ ), 1.68–1.51 (m, 4H, H-2ab', H-4ab'), 1.47–1.37 (m, 2H, H-3ab');  $^{13}C$  NMR (100 MHz,  
10  $CDCl_3$ ,  $\beta$ -anomer)  $\delta$  189.1 (*COPhen*), 170.8, 170.5, 169.9, 169.7 ( $4 \times COCH_3$ ), 167.5 (C-1,  
11  $^3J_{Cl,H3ax} = 5.2$  Hz), 164.3 ( $COCH_2Ph$ ), 156.5 (C-Ar), 136.7 (C-Ar), 130.1, 128.5, 128.0 (CH-Ar),  
12 126.7 (C-Ar), 114.2 (CH-Ar), 99.5 (C-2), 70.8 (C-6), 68.1 (C-7), 67.2 (C-4), 66.6, 66.5 ( $2 \times$   
13  $CH_2$ ), 64.4 (C-1'), 64.3 (C-5), 62.7 (C-8), 55.6 ( $OCH_3$ ), 40.9 (C-5'), 32.6 (C-3), 29.7, 29.1 (C-2',  
14 C-4'), 23.1 (C-3'), 20.80, 20.75 (2C), 20.73 ( $4 \times COCH_3$ ); HRMS (ESI-TOF)  $m/z$   $[M + H]^+$  calcd  
15 for  $C_{38}H_{48}NO_{16}$  774.2968; found 774.2969;  $m/z$   $[M + NH_4]^+$  calcd for  $C_{38}H_{51}N_2O_{16}$  791.3233;  
16 found 791.3237;  $m/z$   $[M + Na]^+$  calcd for  $C_{38}H_{47}NNaO_{16}$  796.2787; found 796.2788. Analytical  
17 data for glycal **16**:  $[\alpha]_D^{20} = +33$  ( $c$  2.7,  $CHCl_3$ );  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.93–7.89 (m, 2H,  
18 *CH-Ar*), 7.00–6.95 (m, 2H, *CH-Ar*), 6.05 (t,  $J_{3,4} \approx J_{3,5} \approx 2.0$  Hz, 1H, H-3), 5.76 (ddd,  $J_{4,5} = 4.5$   
19 Hz,  $J_{3,4} = 2.2$  Hz,  $J_{4,6} = 1.3$  Hz, 1H, H-4), 5.51 (ddd,  $J_{4,5} = 4.5$  Hz,  $J_{3,5} = 1.7$  Hz,  $J_{5,6} = 1.1$  Hz, 1H,  
20 H-5), 5.45 (d,  $J = 16.0$  Hz, 1H, *CHHPhen*), 5.39 (d,  $J = 16.0$  Hz, 1H, *CHHPhen*), 5.29 (ddd,  $J_{6,7} =$   
21 9.7 Hz,  $J_{7,8b} = 3.9$  Hz,  $J_{7,8a} = 2.5$  Hz, 1H, H-7), 4.63 (dd,  $J_{8a,8b} = 12.4$  Hz,  $J_{7,8a} = 2.5$  Hz, 1H, H-  
22 8a), 4.41 (dt,  $J_{6,7} = 9.7$  Hz,  $J_{5,6} \approx J_{4,6} \approx 1.1$  Hz, 1H, H-6), 4.24 (dd,  $J_{8a,8b} = 12.4$  Hz,  $J_{7,8b} = 3.9$  Hz,  
23 1H, H-8b), 2.11, 2.09, 2.05, 2.04 (all s, 12H,  $4 \times COCH_3$ );  $^{13}C$  NMR (100 MHz,  $CDCl_3$ )  $\delta$  189.8  
24 (*COPhen*), 170.7, 170.5, 170.2, 169.6 ( $4 \times COCH_3$ ), 164.4 (C-Ar), 160.6 (C-1), 144.3 (C-2),  
25 130.3 ( $2 \times CH-Ar$ ), 128.7 (C-Ar), 114.3 ( $2 \times CH-Ar$ ), 108.8 (C-3), 73.6 (C-6), 67.5 (C-7), 66.6  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

(CH<sub>2</sub>Ph), 64.9 (C-4), 62.0 (C-8), 60.9 (C-5), 55.7 (OCH<sub>3</sub>), 20.9, 20.8, 20.74, 20.70 (4 × COCH<sub>3</sub>); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup> calcd for C<sub>25</sub>H<sub>28</sub>NaO<sub>13</sub> 559.1422; found 559.1445.

*Benzyl* [2-(5-Amino-N-benzyloxycarbonyl-1-pentyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid]onate (**17 $\beta$**  and **17 $\alpha$** ) and *Benzyl* 4,5,7,8-Tetra-O-acetyl-2,6-anhydro-3-deoxy-D-manno-oct-2-enosonate (**18**). Thioglycoside **3** (1.14 g, 2.10 mmol, 1.0 equiv,  $\beta/\alpha$  7:1) and acceptor **9** (998 mg, 4.21 mmol, 2.0 equiv) were reacted according to the general procedure for glycosylation with thioglycoside donors and gave **17 $\beta$**  (1.21 g, 81%) and **17 $\alpha$**  (192 mg, 13%) both as yellow oils ( $\beta/\alpha$  6.3:1.0). The physical and analytical data of **17 $\beta$**  and **17 $\alpha$**  were in agreement with those published in the literature.<sup>10</sup> Analytical data for glycal **18**: [ $\alpha$ ]<sub>D</sub><sup>20</sup> = +1.3 (*c* 1.9, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.41–7.34 (m, 5H, CH-Ar), 5.92 (t,  $J_{3,4} \approx J_{3,5} \approx 2.0$  Hz, 1H, H-3), 5.73–5.70 (m, 1H, H-4), 5.49–5.46 (m, 1H, H-5), 5.30–5.24 (m, 3H, H-7, CH<sub>2</sub>Ph), 4.63 (dd,  $J_{8a,8b} = 12.3$  Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-8a), 4.36 (br d,  $J_{6,7} = 9.7$  Hz, 1H, H-6), 4.24 (dd,  $J_{8a,8b} = 12.3$  Hz,  $J_{7,8b} = 4.0$  Hz, 1H, H-8b), 2.09 (s, 6H, 2 × COCH<sub>3</sub>), 2.04, 2.03 (all s, 6H, 2 × COCH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  170.6, 170.5, 170.2, 169.6 (4 × COCH<sub>3</sub>), 160.9 (C-1), 144.7 (C-2), 135.3 (C-Ar), 128.8, 128.7, 128.5 (5 × CH-Ar), 107.9 (C-3), 73.4 (C-6), 67.39 (C-7), 67.37 (CH<sub>2</sub>Ph), 64.8 (C-4), 62.0 (C-8), 60.8 (C-5), 20.9, 20.8, 20.75, 20.68 (4 × COCH<sub>3</sub>); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup> calcd for C<sub>23</sub>H<sub>26</sub>NaO<sub>11</sub> 501.1367; found 501.1380.

*3'-Methoxyphenacyl* [2-(5-Amino-N-benzyloxycarbonyl-1-pentyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid]onate (**19**). Thioglycoside **6** (45 mg, 75  $\mu$ mol, 1.0 equiv,  $\beta/\alpha$  7:1) and acceptor **9** (31 mg, 110  $\mu$ mol, 1.5 equiv) were reacted according to the general procedure for glycosylation with thioglycoside donors and gave **19** (42 mg, 73%,  $\beta/\alpha$  6.6:1.0) as

1  
2  
3 a yellow oil.  $[\alpha]_D^{20} = +31$  ( $c$  5.7,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  7.49–7.29  
4 (m, 8H,  $\text{CH-Ar}$ ), 7.19–7.14 (m, 1H,  $\text{CH-Ar}$ ), 5.49 (d,  $J = 16.2$  Hz, 1H,  $\text{CHHPhen}$ ), 5.42 (d,  $J =$   
5 16.2 Hz, 1H,  $\text{CHHPhen}$ ), 5.41–5.39 (m, 1H, H-5), 5.31 (ddd,  $J_{3\text{ax},4} = 13.0$  Hz,  $J_{3\text{eq},4} = 4.6$  Hz,  $J_{4,5}$   
6 = 3.1 Hz, 1H, H-4), 5.22 (ddd,  $J_{6,7} = 9.5$  Hz,  $J_{7,8\text{b}} = 4.8$  Hz,  $J_{7,8\text{a}} = 2.4$  Hz, 1H, H-7), 5.15–5.05  
7 (m, 3H,  $\text{CH}_2\text{Ph}$ ,  $\text{NHCbz}$ ), 4.42–4.30 (m, 3H, H-8a, H-8b, H-6), 3.88–3.79 (m, 1H, H-1a'), 3.86 (s,  
8 3H,  $\text{OCH}_3$ ), 3.56–3.48 (m, 1H, H-1b'), 3.24–3.17 (m, 2H, H-5ab'), 2.53 (dd,  $J_{3\text{ax},3\text{eq}} = 12.5$  Hz,  
9  $J_{3\text{eq},4} = 4.6$  Hz, 1H, H-3eq), 2.17 (t,  $J_{3\text{ax},3\text{eq}} \approx J_{3\text{ax},4} \approx 12.5$  Hz, 1H, H-3ax), 2.12, 2.03, 2.00 (all s,  
10 12H,  $4 \times \text{COCH}_3$ ), 1.67–1.50, 1.47–1.38 (all m, 6H, H-2', H-3', H-4');  $^{13}\text{C}$  NMR (100 MHz,  
11  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  190.7 (COPhen), 170.9, 170.6, 170.0, 169.9 ( $4 \times \text{COCH}_3$ ), 167.5 (C-1),  
12 160.1, 136.8, 135.1 ( $3 \times \text{C-Ar}$ ), 130.1, 128.6, 128.1, 120.9, 120.3, 112.1 ( $\text{CH-Ar}$ ), 99.5 (C-2),  
13 70.9 (C-6), 68.2 (C-7), 67.3 (C-4), 67.1 ( $\text{CH}_2\text{Phen}$ ), 66.6 ( $\text{CH}_2\text{Ph}$ ), 64.5 (C-1'), 64.3 (C-5), 62.8  
14 (C-8), 55.6 ( $\text{OCH}_3$ ), 41.0 (C-5'), 32.7 (C-3), 29.8, 29.2, 23.1 (C-2', C-3', C-4'), 20.9–20.8 ( $4 \times$   
15  $\text{COCH}_3$ ); HRMS (ESI-TOF)  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{38}\text{H}_{48}\text{NO}_{16}$  774.2968; found 774.2973;  $m/z$   
16  $[\text{M} + \text{Na}]^+$  calcd for  $\text{C}_{38}\text{H}_{47}\text{NNaO}_{16}$  796.2787; found 796.2792;  $m/z$   $[2\text{M} + \text{Na}]^+$  calcd for  
17  $\text{C}_{76}\text{H}_{94}\text{N}_2\text{NaO}_{32}$  1569.5682; found 1569.5688.

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42 *2'-Methoxyphenacyl* [2-(5-Amino-N-benzoyloxycarbonyl-1-pentyl) 4,5,7,8-Tetra-O-acetyl-3-  
43 deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid]onate (**20**). Thioglycoside **7** (45 mg, 75  $\mu\text{mol}$ , 1.0 equiv,  
44  $\beta/\alpha$  7:1) and acceptor **9** (31 mg, 110  $\mu\text{mol}$ , 1.5 equiv) were reacted according to the general  
45 procedure for glycosylation with thioglycoside donors and gave **20** (47 mg, 80%,  $\beta/\alpha$  4.6:1.0) as  
46 a colorless oil.  $[\alpha]_D^{20} = +31$  ( $c$  4.6,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  8.00–7.97  
47 (m, 1H,  $\text{CH-Ar}$ ), 7.58–7.52 (m, 1H,  $\text{CH-Ar}$ ), 7.37–7.29 (m, 5H,  $\text{CH-Ar}$ ), 7.06–6.99 (m, 2H,  $\text{CH-}$   
48 Ar), 5.40–5.38 (m, 1H, H-5), 5.36–5.33 (m, 2H,  $\text{CH}_2\text{Phen}$ ), 5.32 (ddd,  $J_{3\text{ax},4} = 12.5$  Hz,  $J_{3\text{eq},4} = 5.3$   
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Hz,  $J_{4,5} = 3.1$  Hz, 1H, H-4), 5.22 (ddd,  $J_{6,7} = 9.5$  Hz,  $J_{7,8b} = 5.0$  Hz,  $J_{7,8a} = 2.2$  Hz, 1H, H-7), 5.10–  
4 5.07 (m, 2H,  $CH_2Ph$ ), 4.41 (dd,  $J_{8a,8b} = 12.3$  Hz,  $J_{7,8a} = 2.1$  Hz, 1H, H-8a), 4.37 (dd,  $J_{6,7} = 9.6$  Hz,  
5 6  $J_{5,6} = 1.1$  Hz, 1H, H-6), 4.32 (dd,  $J_{8a,8b} = 12.3$  Hz,  $J_{7,8b} = 5.1$  Hz, 1H, H-8b), 3.97 (s, 3H,  $OCH_3$ ),  
7 8 3.86–3.80 (m, 1H, H-1a'), 3.59–3.52 (m, 1H, H-1b'), 3.25–3.17 (m, 2H, H-5ab'), 2.52 (dd,  $J_{3ax,3eq}$   
9 = 12.5 Hz,  $J_{3eq,4} = 4.8$  Hz, 1H, H-3eq), 2.15 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx 12.5$  Hz, 1H, H-3ax), 2.11, 2.02,  
10 2.00 (all s, 12H, 4 ×  $COCH_3$ ), 1.68–1.50, 1.47–1.37 (all m, 6H, H-2', H-3', H-4');  $^{13}C$  NMR (100  
11 MHz,  $CDCl_3$ ,  $\beta$ -anomer)  $\delta$  191.3 ( $COPhen$ ), 171.0, 170.7, 170.1, 169.9 (4 ×  $COCH_3$ ), 167.7 (C-  
12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36  
1), 159.7, 135.4, 123.8 (3 × C-Ar), 135.4, 131.4, 128.6, 128.2, 128.1, 121.3, 111.7 (CH-Ar), 99.6  
2), 71.0 ( $CH_2Phen$ ), 70.8 (C-6), 68.3 (C-7), 67.3 (C-4), 66.6 ( $CH_2Ph$ ), 64.6 (C-1'), 64.4 (C-5),  
62.9 (C-8), 55.7 ( $OCH_3$ ), 41.1 (C-5'), 32.8 (C-3), 29.8, 29.2, 23.2 (C-2', C-3', C-4'), 20.9–20.8 (4  
×  $COCH_3$ ); HRMS (ESI-TOF)  $m/z$   $[M + H]^+$  calcd for  $C_{38}H_{48}NO_{16}$  774.2968; found 774.2985;  
 $m/z$   $[M + Na]^+$  calcd for  $C_{38}H_{47}NNaO_{16}$  796.2787; found 796.2804;  $m/z$   $[2M + Na]^+$  calcd for  
 $C_{76}H_{94}N_2NaO_{32}$  1569.5682; found 1569.5711.

37 *Phenacyl* [2-(5-Amino-N-benzyloxycarbonyl-1-pentyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-  
38 *manno-oct-2-ulopyranosid]onate* (**21**). Thioglycoside **8** (45 mg, 75  $\mu$ mol, 1.0 equiv,  $\beta/\alpha$  7:1) and  
39 acceptor **9** (31 mg, 110  $\mu$ mol, 1.5 equiv) were reacted according to the general procedure for  
40 glycosylation with thioglycoside donors and gave **21** (45 mg, 77%,  $\beta/\alpha$  6.2:1.0) as a colorless oil.  
41  $[\alpha]_D^{20} = +36$  ( $c$  4.4,  $CHCl_3$ );  $^1H$  NMR (400 MHz,  $CDCl_3$ ,  $\beta$ -anomer)  $\delta$  7.94–7.89 (m, 2H,  $CH$ -  
42 Ar), 7.66–7.60 (m, 1H,  $CH$ -Ar), 7.53–7.47 (m, 2H,  $CH$ -Ar), 7.38–7.29 (m, 5H,  $CH$ -Ar), 5.50 (d,  
43  $J = 16.2$  Hz, 1H,  $CHHPhen$ ), 5.43 (d,  $J = 16.2$  Hz, 1H,  $CHHPhen$ ), 5.41–5.38 (m, 1H, H-5), 5.32  
44 (ddd,  $J_{3ax,4} = 13.0$  Hz,  $J_{3eq,4} = 4.6$  Hz,  $J_{4,5} = 3.0$  Hz, 1H, H-4), 5.22 (ddd,  $J_{6,7} = 9.6$  Hz,  $J_{7,8b} = 4.9$   
45 Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-7), 5.11–5.06 (m, 2H,  $CH_2Ph$ ), 5.03 (t,  $J = 5.8$  Hz, 1H,  $NHCbz$ ), 4.42–  
46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

1  
2  
3 4.31 (m, 3H, H-8a, H-8b, H-6), 3.86–3.79 (m, 1H, H-1a'), 3.55–3.49 (m, 1H, H-1b'), 3.25–3.17  
4  
5 (m, 2H, H-5ab'), 2.53 (dd,  $J_{3ax,3eq} = 12.5$  Hz,  $J_{3eq,4} = 4.7$  Hz, 1H, H-3eq), 2.17 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx$   
6  
7 12.5 Hz, 1H, H-3ax), 2.11, 2.03, 2.00, 1.99 (all s, 12H, 4 × COCH<sub>3</sub>), 1.67–1.51, 1.47–1.38 (all m,  
8  
9 6H, H-2', H-3', H-4'); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>, β-anomer) δ 190.9 (COPhen), 171.0, 170.6,  
10  
11 170.0, 169.9 (4 × COCH<sub>3</sub>), 167.6 (C-1), 136.8, 133.8 (2 × C-Ar), 134.3, 129.1, 128.6, 128.2,  
12  
13 128.0 (CH-Ar), 99.6 (C-2), 70.9 (C-6), 68.2 (C-7), 67.3 (C-4), 67.0 (CH<sub>2</sub>Phen), 66.7 (CH<sub>2</sub>Ph),  
14  
15 64.6 (C-1'), 64.4 (C-5), 62.8 (C-8), 41.1 (C-5'), 32.8 (C-3), 29.7, 29.2, 23.2 (C-2', C-3', C-4'),  
16  
17 20.92–20.85 (4 × COCH<sub>3</sub>); HRMS (ESI-TOF)  $m/z$  [M + H]<sup>+</sup> calcd for C<sub>37</sub>H<sub>46</sub>NO<sub>15</sub> 744.2862;  
18  
19 found 744.2878;  $m/z$  [M + Na]<sup>+</sup> calcd for C<sub>37</sub>H<sub>45</sub>NNaO<sub>15</sub> 766.2681; found 766.2699;  $m/z$  [2M +  
20  
21 Na]<sup>+</sup> calcd for C<sub>74</sub>H<sub>90</sub>N<sub>2</sub>NaO<sub>30</sub> 1509.5471; found 1509.5505.  
22  
23  
24  
25  
26  
27  
28  
29

30 *4'-Methoxyphenacyl* [2-(1-Nonyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy-α,β-D-manno-oct-2-  
31  
32 *ulopyranosid]onate (22)*. Thioglycoside **2** (50 mg, 84 μmol, 1.0 equiv, β/α 7:1) and 1-nonanol  
33  
34 (**10**, 22 μL, 130 μmol, 1.5 equiv) were reacted according to the general procedure for  
35  
36 glycosylation with thioglycoside donors and gave **22** (37 mg, 65%, β/α 5.3:1.0) as a yellow oil.  
37  
38  $[\alpha]_D^{20} = +48$  (c 0.4, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, β-anomer) δ 7.92–7.88 (m, 2H, CH-  
39  
40 Ar), 6.99–6.95 (m, 2H, CH-Ar), 5.47 (d,  $J = 15.8$  Hz, 1H, CHHPhen), 5.42–5.39 (m, 1H, H-5),  
41  
42 5.36 (d,  $J = 15.8$  Hz, 1H, CHHPhen), 5.35 (ddd,  $J_{3ax,4} = 13.0$  Hz,  $J_{3eq,4} = 4.6$  Hz,  $J_{4,5} = 3.1$  Hz,  
43  
44 1H, H-4), 5.22 (ddd,  $J_{7,8a} = 9.6$  Hz,  $J_{7,8b} = 4.3$  Hz,  $J_{6,7} = 3.1$  Hz, 1H, H-7), 4.41–4.35 (m, 3H, H-  
45  
46 8a, H-8b, H-6), 3.89 (s, 3H, OCH<sub>3</sub>), 3.81 (dt,  $J = 9.0, 6.5$  Hz, 1H, H-1a'), 3.48 (dt,  $J = 9.0, 6.8$   
47  
48 Hz, 1H, H-1b'), 2.54 (ddd,  $J_{3eq,3ax} = 12.5$  Hz,  $J_{3eq,4} = 4.8$  Hz,  $J_{3,5} = 0.8$  Hz, 1H, H-3eq), 2.17 (t,  
49  
50  $J_{3eq,3ax} \approx J_{3ax,4} \approx 12.7$  Hz, 1H, H-3ax), 2.12, 2.02, 2.00 (all s, 12H, 4 × COCH<sub>3</sub>), 1.64–1.56 (m,  
51  
52 2H, H-2'), 1.36–1.24 (m, 12H, H-3', H-4', H-5', H-6', H-7', H-8'), 0.88 (t,  $J = 6.8$  Hz, 3H, H-9');  
53  
54  
55  
56  
57  
58  
59  
60

<sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>, β-anomer) δ 189.1 (COPhen), 170.9, 170.6, 170.0, 169.9 (4 × COCH<sub>3</sub>), 167.7 (C-1, <sup>3</sup>J<sub>Cl,H3ax</sub> = 5.2 Hz), 164.4 (C-Ar), 130.2 (2 × CH-Ar), 126.9 (C-Ar), 114.3 (2 × CH-Ar), 99.6 (C-2), 70.9 (C-6), 68.3 (C-7), 67.4 (C-4), 66.7 (CH<sub>2</sub>Phen), 65.0 (C-1'), 64.5 (C-5), 62.8 (C-8), 55.7 (OCH<sub>3</sub>), 32.8 (C-3), 32.0, 29.8, 29.6, 29.5, 29.4, 26.0, 22.8 (C-2', C-3', C-4', C-5', C-6', C-7', C-8'), 20.91, 20.86 (2C), 20.83 (4 × COCH<sub>3</sub>), 14.2 (C-9'); HRMS (ESI-TOF) *m/z* [M + H]<sup>+</sup> calcd for C<sub>34</sub>H<sub>49</sub>O<sub>14</sub> 681.3117; found 681.3091; *m/z* [M + Na]<sup>+</sup> calcd for C<sub>34</sub>H<sub>48</sub>NaO<sub>14</sub> 703.2936; found 703.2913; *m/z* [2M + Na]<sup>+</sup> calcd for C<sub>68</sub>H<sub>96</sub>NaO<sub>28</sub> 1385.5980; found 1386.5926.

*Benzyl* [2-(1-Nonyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy-α,β-D-manno-oct-2-ulopyranosid]onate (**23**). Thioglycoside **3** (50 mg, 93 μmol, 1.0 equiv, β/α 7:1) and 1-nonanol (**10**, 24 μL, 140 μmol, 1.5 equiv) were reacted according to the general procedure for glycosylation with thioglycoside donors and gave **23** (37 mg, 87%, β/α 3.5:1.0) as a yellow oil. [α]<sub>D</sub><sup>20</sup> = +42 (c 5.2, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, β-anomer) δ 7.39–7.34 (m, 5H, CH-Ar), 5.28–5.16 (m, 4H, H-5, H-7, CH<sub>2</sub>Ph), 4.87 (ddd, *J*<sub>3ax,4</sub> = 13.2 Hz, *J*<sub>3eq,4</sub> = 4.6 Hz, *J*<sub>4,5</sub> = 3.0 Hz, 1H, H-4), 4.35–4.33 (m, 2H, H-8a, H-8b), 4.20 (dd, *J*<sub>6,7</sub> = 9.6 Hz, *J*<sub>5,6</sub> = 1.4 Hz, 1H, H-6), 3.70 (dt, *J* = 9.1, 6.6 Hz, 1H, CHH), 3.17 (dt, *J* = 9.1, 6.8 Hz, 1H, CHH), 2.40 (dd, *J*<sub>3eq,3ax</sub> = 12.5 Hz, *J*<sub>3eq,4</sub> = 4.5 Hz, 1H, H-3eq), 2.10 (t, *J*<sub>3eq,3ax</sub> ≈ *J*<sub>3ax,4</sub> ≈ 12.5 Hz, 1H, H-3ax), 2.10, 2.09, 2.00, 1.98 (all s, 12H, 4 × COCH<sub>3</sub>), 1.51–1.44 (m, 2H, CH<sub>2</sub>), 1.33–1.19 (m, 12H, 6 × CH<sub>2</sub>), 0.88 (t, *J* = 6.8 Hz, 3H, CH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>, β-anomer) δ 170.9, 170.7, 170.0, 169.9 (4 × COCH<sub>3</sub>), 168.0 (C-1, <sup>3</sup>J<sub>Cl,H3ax</sub> = 6.4 Hz), 135.1 (C-Ar), 128.9–128.5 (CH-Ar), 99.6 (C-2), 70.8 (C-6), 68.2 (C-7), 67.8 (CH<sub>2</sub>Ph), 67.5 (C-4), 65.0 (CH<sub>2</sub>), 64.3 (C-5), 62.7 (C-8), 32.7 (C-3), 32.0, 29.68, 29.65, 29.5, 29.4, 26.0, 22.8 (7 × CH<sub>2</sub>), 20.92, 20.86, 20.84, 20.82 (4 × COCH<sub>3</sub>), 14.3 (CH<sub>3</sub>); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup>

1  
2  
3 calcd for  $C_{32}H_{46}NaO_{12}$  645.2881; found 645.2887;  $m/z$   $[2M + Na]^+$  calcd for  $C_{64}H_{92}NaO_{24}$   
4  
5 1267.5871; found 1267.5890.  
6  
7  
8

9  
10 *4'-Methoxyphenacyl (2-Cyclohexyl 4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-manno-oct-2-*  
11 *ulopyranosid)onate (24)*. Thioglycoside **2** (50 mg, 84  $\mu$ mol, 1.0 equiv,  $\beta/\alpha$  7:1) and cyclohexanol  
12  
13 (**11**, 13  $\mu$ L, 130  $\mu$ mol, 1.5 equiv) were reacted according to the general procedure for  
14  
15 glycosylation with thioglycoside donors and gave **24** (40 mg, 76%,  $\beta/\alpha$  4.0:1.0) as a yellow oil.  
16  
17  
18  
19  
20  $[\alpha]_D^{20} = +43$  ( $c$  4.3,  $CHCl_3$ );  $^1H$  NMR (400 MHz,  $CDCl_3$ ,  $\beta$ -anomer)  $\delta$  7.92–7.88 (m, 2H, *CH-*  
21 *Ar*), 6.99–6.94 (m, 2H, *CH-Ar*), 5.47 (d,  $J = 15.9$  Hz, 1H, *CHHPhen*), 5.41–5.40 (m, 1H, H-5),  
22  
23 5.36 (ddd,  $J_{3ax,4} = 12.9$  Hz,  $J_{3eq,4} = 4.7$  Hz,  $J_{4,5} = 3.2$  Hz, 1H, H-4), 5.33 (d,  $J = 15.8$  Hz, 1H,  
24  
25 *CHHPhen*), 5.19 (ddd,  $J_{6,7} = 9.6$  Hz,  $J_{7,8b} = 4.9$  Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-7), 4.40 (dd,  $J_{8a,8b} = 12.1$   
26  
27 Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-8a), 4.38 (dd,  $J_{6,7} = 9.6$  Hz,  $J_{5,6} = 1.4$  Hz, 1H, H-6), 4.33 (dd,  $J_{8a,8b} =$   
28  
29 12.3 Hz,  $J_{7,8b} = 4.9$  Hz, 1H, H-8b), 3.89–3.83 (m, 1H, *CHCy*), 3.88 (s, 3H, *OCH*<sub>3</sub>), 2.53 (ddd,  
30  
31  $J_{3ax,3eq} = 12.4$  Hz,  $J_{3eq,4} = 4.7$  Hz,  $J_{3eq,5} = 0.7$  Hz, 1H, H-3eq), 2.16 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx 12.4$  Hz,  
32  
33 1H, H-3ax), 2.11, 2.02, 1.999, 1.994 (all s, 12H, 4  $\times$  *COCH*<sub>3</sub>), 1.95–1.14 (m, 10H, 5  $\times$  *CH*<sub>2</sub>*Cy*);  
34  
35  
36  
37  
38  
39  $^{13}C$  NMR (100 MHz,  $CDCl_3$ ,  $\beta$ -anomer)  $\delta$  189.2 (*COPhen*), 170.8, 170.6, 170.1, 169.8 (4  $\times$   
40  
41 *COCH*<sub>3</sub>), 168.2 (C-1,  $^3J_{C1,H3ax} = 5.2$  Hz), 164.4 (C-*Ar*), 130.2 (2  $\times$  *CH-Ar*), 126.9 (C-*Ar*), 114.3  
42  
43 (2  $\times$  *CH-Ar*), 99.8 (C-2), 74.5 (*CHCy*), 70.8 (C-6), 68.4 (C-7), 67.4 (C-4), 66.7 (*CH*<sub>2</sub>*Phen*), 64.5  
44  
45 (C-5), 62.8 (C-8), 55.6 (*OCH*<sub>3</sub>), 34.8, 33.6 (2  $\times$  *CH*<sub>2</sub>*Cy*), 33.1 (C-3), 25.5, 24.5, 24.4 (3  $\times$  *CH*<sub>2</sub>*Cy*),  
46  
47 20.90, 20.85 (2C), 20.82 (4  $\times$  *COCH*<sub>3</sub>); HRMS (ESI-TOF)  $m/z$   $[M + Na]^+$  calcd for  
48  
49  $C_{31}H_{40}NaO_{14}$  659.2310; found 659.2315;  $m/z$   $[2M + Na]^+$  calcd for  $C_{62}H_{80}NaO_{28}$  1295.4728;  
50  
51  
52 found 1295.4732.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 *Benzyl (2-Cyclohexyl 4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid)onate*  
4  
5  
6 (25). Thioglycoside **3** (50 mg, 93  $\mu$ mol, 1.0 equiv,  $\beta/\alpha$  7:1) and cyclohexanol (**11**, 15  $\mu$ L, 140  
7  
8  $\mu$ mol, 1.5 equiv) were reacted according to the general procedure for glycosylation with  
9  
10 thioglycoside donors and gave **25** (43 mg, 80%,  $\beta/\alpha$  2.2:1.0) as a yellow oil.  $[\alpha]_D^{20} = +69$  (*c* 0.69,  
11  
12  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  7.41–7.33 (m, 5H, *CH-Ar*), 5.27–5.25 (m, 1H,  
13  
14 H-5), 5.26 (d,  $J = 12.0$  Hz, 1H, *CHHPh*), 5.18 (d,  $J = 12.0$  Hz, 1H, *CHHPh*), 5.16 (ddd,  $J_{6,7} = 9.5$   
15  
16 Hz,  $J_{7,8b} = 4.6$  Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-7), 4.85 (ddd,  $J_{3ax,4} = 13.2$  Hz,  $J_{3eq,4} = 4.6$  Hz,  $J_{4,5} = 3.0$   
17  
18 Hz, 1H, H-4), 4.37 (dd,  $J_{8a,8b} = 12.3$  Hz,  $J_{7,8a} = 2.4$  Hz, 1H, H-8a), 4.31 (dd,  $J_{8a,8b} = 12.3$  Hz,  $J_{7,8b}$   
19  
20 = 4.7 Hz, 1H, H-8b), 4.15 (dd,  $J_{6,7} = 9.5$  Hz,  $J_{5,6} = 1.4$  Hz, 1H, H-6), 3.66–3.58 (m, 1H, *CHCy*),  
21  
22 2.40 (ddd,  $J_{3ax,3eq} = 12.5$  Hz,  $J_{3eq,4} = 4.6$  Hz,  $J_{3eq,5} = 0.8$  Hz, 1H, H-3eq), 2.09 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx$   
23  
24 12.3 Hz, 1H, H-3ax), 2.09, 2.06, 2.00, 1.97 (all s, 12H, 4  $\times$   $\text{COCH}_3$ ), 1.94–1.00 (m, 10H, 5  $\times$   
25  
26  $\text{CH}_2\text{Cy}$ );  $^{13}\text{C NMR}$  (100 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  170.7, 170.6, 169.94, 169.91 (4  $\times$   $\text{COCH}_3$ ),  
27  
28 168.3 (C-1,  $^3J_{\text{C1,H3ax}} = 6.2$  Hz), 134.9, (C-Ar), 128.7–128.3 (*CH-Ar*), 99.8 (C-2), 74.6 (*CHCy*),  
29  
30 70.7 (C-6), 68.3 (C-7), 67.7 ( $\text{CH}_2\text{Ph}$ ), 67.2 (C-4), 64.2 (C-5), 62.6 (C-8), 34.8 ( $\text{CH}_2\text{Cy}$ ), 33.3  
31  
32 ( $\text{CH}_2\text{Cy}$ ), 33.0 (C-3), 25.4, 24.4, 24.3 (3  $\times$   $\text{CH}_2\text{Cy}$ ), 20.88, 20.82, 20.79, 20.77 (4  $\times$   $\text{COCH}_3$ );  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*4'-Methoxyphenacyl [2-(2'-Adamantyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-manno-oct-2-*  
*ulopyranosid]onate (26)*. Thioglycoside **2** (50 mg, 84  $\mu$ mol, 1.0 equiv,  $\beta/\alpha$  7:1) and 2-  
adamantanol (**12**, 19 mg, 130  $\mu$ mol, 1.5 equiv) were reacted according to the general procedure  
for glycosylation with thioglycoside donors and gave **26** (34 mg, 58%,  $\beta/\alpha$  4.3:1.0) as a colorless  
oil.  $[\alpha]_D^{20} = +7$  (*c* 2.7,  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  7.93–7.88 (m, 2H, *CH-*

Ar), 6.99–6.94 (m, 2H, CH-Ar), 5.46 (d,  $J = 15.9$  Hz, 1H, CHHPhen), 5.43–5.35 (m, 2H, H-4, H-5), 5.33 (d,  $J = 15.9$  Hz, 1H, CHHPhen), 5.17 (ddd,  $J_{6,7} = 9.5$  Hz,  $J_{7,8b} = 4.8$  Hz,  $J_{7,8a} = 2.2$  Hz, 1H, H-7), 4.39–4.34 (m, 2H, H-8a, H-6), 4.29 (dd,  $J_{8a,8b} = 12.2$  Hz,  $J_{7,8b} = 5.0$  Hz, 1H, H-8b), 4.08–4.04 (m, 1H, CH-Ad), 3.89 (s, 3H, OCH<sub>3</sub>), 2.55 (ddd,  $J_{3ax,3eq} = 12.0$  Hz,  $J_{3eq,4} = 4.7$  Hz,  $J_{3eq,5} = 0.9$  Hz, 1H, H-3eq), 2.20 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx 12.0$  Hz, 1H, H-3ax), 2.16–2.06 (m, 2H, CH<sub>2</sub>-Ad), 2.12, 2.02, 2.00 (all s, 12H, 4 × COCH<sub>3</sub>), 2.06–2.01 (m, 1H, CH-Ad), 1.85–1.68 (m, 9H, 3 × CH<sub>2</sub>-Ad, 3 × CH-Ad), 1.55–1.45 (m, 2H, CH<sub>2</sub>-Ad); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>, β-anomer)  $\delta$  189.3 (COPhen), 170.8, 170.6, 170.1, 169.8 (4 × COCH<sub>3</sub>), 168.2 (C-1,  $^3J_{C1,H3ax} = 5.2$  Hz), 164.3 (C-Ar), 130.2 (2 × CH-Ar), 127.0 (C-Ar), 114.3 (2 × CH-Ar), 99.8 (C-2), 78.2 (CH-Ad), 70.7 (C-6), 68.4 (C-7), 67.5 (C-4), 66.6 (CH<sub>2</sub>Phen), 64.5 (C-5), 62.8 (C-8), 55.7 (OCH<sub>3</sub>), 37.7, 37.0, 36.8 (3 × CH<sub>2</sub>-Ad), 34.4, 33.5 (2 × CH-Ad), 33.2 (C-3), 31.69, 31.67 (2 × CH<sub>2</sub>-Ad), 27.4, 27.1 (2 × CH-Ad), 20.92, 20.88 (2C), 20.85 (4 × COCH<sub>3</sub>); HRMS (ESI-TOF)  $m/z$  [M + Na]<sup>+</sup> calcd for C<sub>35</sub>H<sub>44</sub>NaO<sub>14</sub> 711.2623; found 711.2604;  $m/z$  [2M + Na]<sup>+</sup> calcd for C<sub>70</sub>H<sub>88</sub>NaO<sub>28</sub> 1399.5354; found 1399.5309.

*Benzyl* [2-(2'-Adamantyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid]onate (**27**). Thioglycoside **3** (40 mg, 74  $\mu$ mol, 1.0 equiv,  $\beta/\alpha$  7:1) and 2-adamantanol (**12**, 16 mg, 110  $\mu$ mol, 1.5 equiv) were reacted according to the general procedure for glycosylation with thioglycoside donors and gave **27** (35 mg, 74%,  $\beta/\alpha$  2.4:1.0) as a yellow oil.  $[\alpha]_D^{20} = +54$  ( $c$  3.3, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, β-anomer)  $\delta$  7.40–7.33 (m, 5H, CH-Ar), 5.27–5.25 (m, 1H, H-5), 5.24 (d,  $J = 12.1$  Hz, 1H, CHHPh), 5.16 (d,  $J = 12.2$  Hz, 1H, CHHPh), 5.14 (ddd,  $J_{6,7} = 9.5$  Hz,  $J_{7,8b} = 4.7$  Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-7), 4.87 (ddd,  $J_{3ax,4} = 13.2$  Hz,  $J_{3eq,4} = 4.6$  Hz,  $J_{4,5} = 3.0$  Hz, 1H, H-4), 4.34 (dd,  $J_{8a,8b} = 12.2$  Hz,  $J_{7,8a} = 2.2$  Hz, 1H, H-8a),

1  
2  
3 4.27 (dd,  $J_{8a,8b} = 12.2$  Hz,  $J_{7,8b} = 4.8$  Hz, 1H, H-8b), 4.13 (dd,  $J_{6,7} = 9.5$  Hz,  $J_{5,6} = 1.4$  Hz, 1H, H-  
4 6), 3.86–3.82 (m, 1H, CH-Ad), 2.43 (dd,  $J_{3ax,3eq} = 12.5$  Hz,  $J_{3eq,4} = 4.6$  Hz, 1H, H-3eq), 2.13 (t,  
5  $J_{3ax,3eq} \approx J_{3ax,4} \approx 12.9$  Hz, 1H, H-3ax), 2.10, 2.08, 2.00, 1.98 (all s, 12H, 4 × COCH<sub>3</sub>), 2.11–1.96  
6 (m, 2H, CH<sub>2</sub>-Ad), 1.98–1.94 (m, 1H, CH-Ad), 1.79–1.68 (m, 3H, 2 × CH-Ad, CHHAd), 1.67–  
7 1.58 (m, 4H, 2 × CH<sub>2</sub>-Ad), 1.51–1.33 (m, 4H, CH<sub>2</sub>-Ad, CH-Ad, CHH-Ad); <sup>13</sup>C NMR (100 MHz,  
8 CDCl<sub>3</sub>, β-anomer) δ 170.8, 170.7, 170.05, 169.98 (4 × COCH<sub>3</sub>), 168.4 (C-1,  $^3J_{C1,H3ax} = 6.3$  Hz),  
9 135.1 (C-Ar), 128.8–128.7 (CH-Ar), 99.8 (C-2), 78.2 (CH-Ad), 70.7 (C-6), 68.4 (C-7), 67.6  
10 (CH<sub>2</sub>Ph), 67.4 (C-4), 64.3 (C-5), 62.7 (C-8), 37.6, 36.9, 36.7 (3 × CH<sub>2</sub>-Ad), 34.3, 33.3 (2 × CH-  
11 Ad), 33.1 (C-3), 31.7, 31.6 (2 × CH<sub>2</sub>-Ad), 27.4, 27.0 (2 × CH-Ad), 20.93, 20.89, 20.86, 20.85 (4  
12 × COCH<sub>3</sub>); HRMS (ESI-TOF)  $m/z$  [M + Na]<sup>+</sup> calcd for C<sub>33</sub>H<sub>42</sub>NaO<sub>12</sub> 653.2568; found 653.2557;  
13  $m/z$  [M + K]<sup>+</sup> calcd for C<sub>33</sub>H<sub>42</sub>KO<sub>12</sub> 669.2308; found 669.2295;  $m/z$  [2M + Na]<sup>+</sup> calcd for  
14 C<sub>66</sub>H<sub>84</sub>NaO<sub>24</sub> 1283.5245; found 1283.5208.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 *4'-Methoxyphenacyl* [2-(1'-Adamantyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy-α,β-D-manno-oct-2-  
35 ulopyranosid]onate (**28**). Thioglycoside **2** (50 mg, 84 μmol, 1.0 equiv, β/α 7:1) and 1-  
36 adamantanol (**13**, 19 mg, 130 μmol, 1.5 equiv) were reacted according to the general procedure  
37 for glycosylation with thioglycoside donors and gave **28** (23 mg, 40%, β/α 1.6:1.0) as a colorless  
38 oil.  $[\alpha]_D^{20} = +34$  (*c* 1.9, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, β-anomer) δ 7.94–7.89 (m, 2H,  
39 CH-Ar), 7.00–6.94 (m, 2H, CH-Ar), 5.49 (d,  $J = 15.8$  Hz, 1H, CHHPhen), 5.40–5.37 (m, 1H, H-  
40 5), 5.33–5.18 (m, 3H, CHHPhen, H-4, H-7), 4.57 (dd,  $J_{6,7} = 9.6$  Hz,  $J_{5,6} = 1.4$  Hz, 1H, H-6), 4.48  
41 (dd,  $J_{8a,8b} = 12.2$  Hz,  $J_{7,8a} = 2.2$  Hz, 1H, H-8a), 4.32 (dd,  $J_{8a,8b} = 12.2$  Hz,  $J_{7,8b} = 5.2$  Hz, 1H, H-  
42 8b), 3.89 (s, 3H, OCH<sub>3</sub>), 2.51 (ddd,  $J_{3ax,3eq} = 12.5$  Hz,  $J_{3eq,4} = 4.7$  Hz,  $J_{3eq,5} = 0.8$  Hz, 1H, H-3eq),  
43 2.21 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx 12.5$  Hz, 1H, H-3ax), 2.15–2.10 (m, 3H, 3 × CH-Ad), 2.11, 2.026, 2.015,  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 1.99 (all s, 12H, 4 × COCH<sub>3</sub>), 1.96–1.90 (m, 6H, 3 × CH<sub>2</sub>-Ad), 1.66–1.58 (m, 6H, 3 × CH<sub>2</sub>-Ad);  
4  
5 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, α-anomer) δ 7.94–7.89 (m, 2H, CH-Ar), 7.00–6.94 (m, 2H, CH-Ar),  
6  
7 5.53 (d, *J* = 15.8 Hz, 1H, CHHPhen), 5.42–5.36 (m, 2H, H-5, H-4), 5.33–5.18 (m, 2H, CHHPhen,  
8  
9 H-7), 4.69 (dd, *J*<sub>8a,8b</sub> = 12.3 Hz, *J*<sub>7,8a</sub> = 2.7 Hz, 1H, H-8a), 4.39 (dd, *J*<sub>6,7</sub> = 9.4 Hz, *J*<sub>5,6</sub> = 1.5 Hz,  
10  
11 1H, H-6), 4.15 (dd, *J*<sub>8a,8b</sub> = 12.3 Hz, *J*<sub>7,8b</sub> = 3.8 Hz, 1H, H-8b), 3.89 (s, 3H, OCH<sub>3</sub>), 2.32–2.23 (m,  
12  
13 2H, H-3eq, H-3ax), 2.15–2.10 (m, 3H, 3 × CH-Ad), 2.08, 2.07, 1.99, 1.98 (all s, 12H, 4 ×  
14  
15 COCH<sub>3</sub>), 1.96–1.90 (m, 6H, 3 × CH<sub>2</sub>-Ad), 1.66–1.58 (m, 6H, 3 × CH<sub>2</sub>-Ad); <sup>13</sup>C NMR (100 MHz,  
16  
17 CDCl<sub>3</sub>, β-anomer) δ 189.5 (COPhen), 170.8, 170.7, 170.3, 170.1 (4 × COCH<sub>3</sub>), 169.9 (C-1,  
18  
19 <sup>3</sup>*J*<sub>C1,H3ax</sub> = overlapping), 164.3 (C-Ar), 130.3 (2 × CH-Ar), 127.0 (C-Ar), 114.2 (2 × CH-Ar), 98.9  
20  
21 (C-2), 79.2 (C-Ad), 71.3 (C-6), 68.5 (C-7), 67.2 (C-4), 66.6 (CH<sub>2</sub>Phen), 64.6 (C-5), 63.2 (C-8),  
22  
23 55.7 (OCH<sub>3</sub>), 43.6 (2C), 42.9 (3 × CH<sub>2</sub>-Ad), 36.3 (2C), 36.1 (3 × CH<sub>2</sub>-Ad), 35.5 (C-3), 31.22  
24  
25 (2C), 31.17 (3 × CH-Ad), 21.0–20.8 (4 × COCH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>, α-anomer) δ  
26  
27 189.8 (COPhen), 170.7, 170.6, 170.2, 169.8 (4 × COCH<sub>3</sub>), 169.5 (C-1, <sup>3</sup>*J*<sub>C1,H3ax</sub> < 1.0 Hz), 164.3  
28  
29 (C-Ar), 130.3 (2 × CH-Ar), 127.3 (C-Ar), 114.3 (2 × CH-Ar), 97.9 (C-2); 78.2 (C-Ad), 68.9 (C-  
30  
31 6), 68.6 (C-7), 66.8 (C-4), 66.4 (CH<sub>2</sub>Phen), 65.0 (C-5), 61.9 (C-8), 55.7 (OCH<sub>3</sub>), 43.6 (2C), 42.9  
32  
33 (3 × CH<sub>2</sub>-Ad), 36.3 (2C), 36.1 (3 × CH<sub>2</sub>Phen), 35.2 (C-3), 31.23 (2C), 31.17 (3 × CH-Ad), 21.0–  
34  
35 20.8 (4 × COCH<sub>3</sub>); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup> calcd for C<sub>35</sub>H<sub>44</sub>NaO<sub>14</sub> 711.2623; found  
36  
37 711.2604; *m/z* [2M + Na]<sup>+</sup> calcd for C<sub>70</sub>H<sub>88</sub>NaO<sub>28</sub> 1399.5354; found 1399.5310.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Benzyl* [2-(1'-Adamantyl) 4,5,7,8-Tetra-O-acetyl-3-deoxy-α,β-D-manno-oct-2-  
ulopyranosid]onate (**29**). Thioglycoside **3** (50 mg, 93 μmol, 1.0 equiv, β/α 7:1) and 1-  
adamantanol (**13**, 21 mg, 140 μmol, 1.5 equiv) were reacted according to the general procedure  
for glycosylation with thioglycoside donors and gave **29** (30 mg, 51%, β/α 1.0:1.6) as a yellow

oil.  $[\alpha]_D^{20} = +43$  (*c* 3.4,  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  7.44–7.32 (m, 5H, *CH-Ar*), 5.28–5.15 (m, 4H, H-5, H-7,  $\text{CH}_2\text{Ph}$ ), 4.77 (ddd,  $J_{3\text{ax},4} = 13.4$  Hz,  $J_{3\text{eq},4} = 4.5$  Hz,  $J_{4,5} = 2.9$  Hz, 1H, H-4), 4.50 (dd,  $J_{6,7} = 9.6$  Hz,  $J_{5,6} = 1.5$  Hz, 1H, H-6), 4.46 (dd,  $J_{8\text{a},8\text{b}} = 12.3$  Hz,  $J_{7,8\text{a}} = 2.1$  Hz, 1H, H-8a), 4.30 (dd,  $J_{8\text{a},8\text{b}} = 12.3$  Hz,  $J_{7,8\text{b}} = 5.2$  Hz, 1H, H-8b), 2.31 (ddd,  $J_{3\text{ax},3\text{eq}} = 12.6$  Hz,  $J_{3\text{eq},4} = 4.5$  Hz,  $J_{3\text{eq},5} = 0.8$  Hz, 1H, H-3eq), 2.12 (t,  $J_{3\text{ax},3\text{eq}} \approx J_{3\text{ax},4} \approx 12.6$  Hz, 1H, H-3ax), 2.10, 2.09, 2.01, 1.96 (all s, 12H,  $4 \times \text{COCH}_3$ ), 2.05–1.97 (m, 3H,  $3 \times \text{CH-Ad}$ ), 1.81–1.75 (m, 6H,  $3 \times \text{CH}_2\text{-Ad}$ ), 1.57–1.42 (m, 6H,  $3 \times \text{CH}_2\text{-Ad}$ );  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ,  $\alpha$ -anomer)  $\delta$  7.44–7.32 (m, 5H, *CH-Ar*), 5.39–5.31 (m, 2H, H-5, H-4), 5.25–5.15 (m, 3H,  $\text{CH}_2\text{Ph}$ , H-7), 4.68 (dd,  $J_{8\text{a},8\text{b}} = 12.3$  Hz,  $J_{7,8\text{a}} = 2.7$  Hz, 1H, H-8a), 4.34 (dd,  $J_{6,7} = 9.5$  Hz,  $J_{5,6} = 1.5$  Hz, 1H, H-6), 4.14 (dd,  $J_{8\text{a},8\text{b}} = 12.3$  Hz,  $J_{7,8\text{b}} = 3.8$  Hz, 1H, H-8b), 2.19 (ddd,  $J_{3\text{ax},3\text{eq}} = 12.4$  Hz,  $J_{3\text{eq},4} = 4.7$  Hz,  $J_{3\text{eq},5} = 1.0$  Hz, 1H, H-3eq), 2.06, 2.05, 1.98, 1.96 (all s, 12H,  $4 \times \text{COCH}_3$ ), 2.05–1.97 (m, 3H,  $3 \times \text{CH-Ad}$ ), 1.92 (t,  $J_{3\text{ax},3\text{eq}} \approx J_{3\text{ax},4} \approx 12.4$  Hz, 1H, H-3ax), 1.81–1.75 (m, 6H,  $3 \times \text{CH}_2\text{-Ad}$ ), 1.57–1.42 (m, 6H,  $3 \times \text{CH}_2\text{-Ad}$ );  $^{13}\text{C NMR}$  (100 MHz,  $\text{CDCl}_3$ ,  $\beta$ -anomer)  $\delta$  170.8, 170.7, 170.1, 169.9 ( $4 \times \text{COCH}_3$ ), 169.8 (C-1,  $^3J_{\text{C}1,\text{H}3\text{ax}} = \text{overlapping}$ ), 134.9 (C-Ar), 129.0–128.8 ( $5 \times \text{CH-Ar}$ ), 98.7 (C-2), 79.0 (C-Ad), 71.3 (C-6), 68.4 (C-7), 67.7 ( $\text{CH}_2\text{Ph}$ ), 66.9 (C-4), 64.4 (C-5), 63.2 (C-8), 43.5, 42.8 (2C,  $3 \times \text{CH}_2\text{-Ad}$ ), 36.2, 36.0 (2C,  $3 \times \text{CH}_2\text{-Ad}$ ), 35.3 (C-3), 31.1, 31.0 (2C,  $3 \times \text{CH-Ad}$ ), 20.9–20.8 ( $4 \times \text{COCH}_3$ );  $^{13}\text{C NMR}$  (100 MHz,  $\text{CDCl}_3$ ,  $\alpha$ -anomer)  $\delta$  170.59, 170.55, 170.51, 170.1 ( $4 \times \text{COCH}_3$ ), 169.4 (C-1,  $^3J_{\text{C}1,\text{H}3\text{ax}} < 1.0$  Hz), 134.8 (C-Ar), 129.0–128.8 ( $5 \times \text{CH-Ar}$ ), 97.6 (C-2), 78.1 (C-Ad), 68.8 (C-6), 68.6 (C-7), 67.6 ( $\text{CH}_2\text{Ph}$ ), 66.8 (C-4), 65.0 (C-5), 61.9 (C-8), 43.5, 42.8 (2C,  $3 \times \text{CH}_2\text{-Ad}$ ), 36.2, 36.0 (2C,  $3 \times \text{CH}_2\text{-Ad}$ ), 35.0 (C-3), 31.1, 31.0 (2C,  $3 \times \text{CH-Ad}$ ), 20.9–20.8 ( $4 \times \text{COCH}_3$ ); HRMS (ESI-TOF)  $m/z$   $[\text{M} + \text{Na}]^+$  calcd for  $\text{C}_{33}\text{H}_{42}\text{NaO}_{12}$  653.2568; found 653.2561;  $m/z$   $[2\text{M} + \text{Na}]^+$  calcd for  $\text{C}_{66}\text{H}_{84}\text{NaO}_{24}$  1283.5245; found 1283.5226.

1  
2  
3  
4  
5  
6 *4'*-Methoxyphenacyl [(4,5,7,8-Tetra-*O*-acetyl-3-deoxy- $\alpha,\beta$ -*D*-manno-oct-2-ulopyranosyl)onate]-  
7  
8 (2 $\rightarrow$ 6)-(Methyl 2,3-di-*O*-benzyl- $\alpha$ -*D*-glucopyranoside) (**30**). Thioglycoside **2** (35 mg, 59  $\mu$ mol,  
9  
10 1.0 equiv,  $\beta/\alpha$  7:1) and methyl 2,3-di-*O*-benzyl- $\alpha$ -*D*-glucopyranoside (**14**, 30 mg, 80  $\mu$ mol, 1.4  
11  
12 equiv) were reacted according to the general procedure for glycosylation with thioglycoside  
13  
14 donors and gave **30** (36 mg, 61%,  $\beta/\alpha$  5.0:1.0) as a yellow oil.  $[\alpha]_D^{20} = +26$  (*c* 1.7, CHCl<sub>3</sub>); <sup>1</sup>H  
15  
16 NMR (400 MHz, CDCl<sub>3</sub>,  $\beta$ -anomer)  $\delta$  7.92–7.85 (m, 2H, *CH*-Ar), 7.43–7.26 (m, 10H, *CH*-Ar),  
17  
18 7.00–6.95 (m, 2H, *CH*-Ar), 5.44 (d, *J* = 15.8 Hz, 1H, *CHH*Phen), 5.37–5.35 (m, 1H, H-5), 5.36  
19  
20 (d, *J* = 15.8 Hz, 1H, *CHH*Phen), 5.25 (ddd, *J*<sub>3ax,4</sub> = 13.0 Hz, *J*<sub>3eq,4</sub> = 4.6 Hz, *J*<sub>4,5</sub> = 2.9 Hz, 1H, H-  
21  
22 4), 5.20 (ddd, *J*<sub>6,7</sub> = 9.4 Hz, *J*<sub>7,8b</sub> = 5.1 Hz, *J*<sub>7,8a</sub> = 2.1 Hz, 1H, H-7), 4.97 (d, *J* = 11.1 Hz, 1H,  
23  
24 *CHH*Ph), 4.87 (d, *J* = 11.1 Hz, 1H, *CHH*Ph), 4.79 (d, *J* = 12.1 Hz, 1H, *CHH*Ph), 4.67 (d, *J*<sub>1,2</sub> =  
25  
26 3.7 Hz, 1H, H-1Glc), 4.66 (d, *J* = 12.1 Hz, 1H, *CHH*Ph), 4.41 (dd, *J*<sub>8a,8b</sub> = 12.5 Hz, *J*<sub>7,8a</sub> = 2.3 Hz,  
27  
28 1H, H-8a), 4.37–4.29 (m, 2H, H-8b, H-6), 4.06 (dd, *J*<sub>6a,6b</sub> = 11.0 Hz, *J*<sub>5,6a</sub> = 2.1 Hz, 1H, H-6aGlc),  
29  
30 3.89 (s, 3H, OCH<sub>3</sub>Phen), 3.87–3.80 (m, 2H, H-6bGlc, H-3Glc), 3.79–3.64 (m, 2H, H-5Glc, H-  
31  
32 4Glc), 3.53 (dd, *J*<sub>2,3</sub> = 9.6 Hz, *J*<sub>1,2</sub> = 3.5 Hz, 1H, H-2Glc), 3.42 (s, 3H, OCH<sub>3</sub>Glc), 2.50 (dd,  
33  
34 *J*<sub>3ax,3eq</sub> = 12.6 Hz, *J*<sub>3eq,4</sub> = 4.8 Hz, 1H, H-3eq), 2.22 (t, *J*<sub>3ax,3eq</sub>  $\approx$  *J*<sub>3ax,4</sub>  $\approx$  12.6 Hz, 1H, H-3ax), 2.08,  
35  
36 2.02, 2.00, 1.99 (all s, 12H, 4  $\times$  COCH<sub>3</sub>); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>,  $\beta$ -anomer)  $\delta$  189.6  
37  
38 (COPhen), 171.1, 170.6, 170.0, 169.9 (4  $\times$  COCH<sub>3</sub>), 167.4 (C-1, <sup>3</sup>*J*<sub>C1,H3ax</sub> = 5.6 Hz), 164.5 (C-Ar),  
39  
40 139.1, 138.4 (2  $\times$  C-Ar), 130.3–127.7 (CH-Ar), 126.7 (C-Ar), 114.3 (CH-Ar), 99.8 (C-2), 98.4  
41  
42 (C-1Glc), 81.9 (C-3Glc), 79.6 (C-2Glc), 75.8, 73.3 (2  $\times$  CH<sub>2</sub>Ph), 71.1 (C-6), 70.4 (C-5Glc), 70.2  
43  
44 (C-4Glc), 68.2 (C-7), 67.3 (C-4), 66.8 (CH<sub>2</sub>Phen), 64.3 (C-5), 63.7 (C-6Glc), 62.9 (C-8), 55.7  
45  
46 (OCH<sub>3</sub>Phen), 55.3 (OCH<sub>3</sub>Glc), 32.4 (C-3), 20.9 (2C), 20.81, 20.76 (4  $\times$  COCH<sub>3</sub>); HRMS (ESI-  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

TOF)  $m/z$   $[M + Na]^+$  calcd for  $C_{46}H_{54}NaO_{19}$  933.3152; found 933.3122;  $m/z$   $[2M + Na]^+$  calcd for  $C_{92}H_{108}NaO_{38}$  1843.6411; found 1843.6353.

*Benzyl [(4,5,7,8-Tetra-O-acetyl-3-deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosyl)onate]-(2 $\rightarrow$ 6)-(Methyl 2,3-di-O-benzyl- $\alpha$ -D-glucopyranoside) (31)*. Thioglycoside **3** (35 mg, 65  $\mu$ mol, 1.0 equiv,  $\beta/\alpha$  7:1) and methyl 2,3-di-O-benzyl- $\alpha$ -D-glucopyranoside (**14**, 33 mg, 88  $\mu$ mol, 1.4 equiv) were reacted according to the general procedure for glycosylation with thioglycoside donors and gave **31** (48 mg, 78%,  $\beta/\alpha$  3.7:1.0) as a yellow oil.  $[\alpha]_D^{20} = +33$  ( $c$  3.6,  $CHCl_3$ );  $^1H$  NMR (400 MHz,  $CDCl_3$ ,  $\beta$ -anomer)  $\delta$  7.40–7.28 (m, 15H,  $CH$ -Ar), 5.26–5.24 (m, 1H, H-5), 5.21–5.19 (m, 2H,  $CH_2Ph$ ), 5.15 (ddd,  $J_{6,7} = 9.3$  Hz,  $J_{7,8b} = 4.7$  Hz,  $J_{7,8a} = 2.3$  Hz, 1H, H-7), 4.97 (d,  $J = 11.2$  Hz, 1H,  $CHHPh$ ), 4.88 (ddd,  $J_{3ax,4} = 13.1$  Hz,  $J_{3eq,4} = 4.6$  Hz,  $J_{4,5} = 2.9$  Hz, 1H, H-4), 4.78 (d,  $J = 12.0$  Hz, 1H,  $CHHPh$ ), 4.77 (d,  $J = 11.2$  Hz, 1H,  $CHHPh$ ), 4.65 (d,  $J = 12.2$  Hz, 1H,  $CHHPh$ ), 4.61 (d,  $J_{1,2} = 3.6$  Hz, 1H, H-1Glc), 4.34 (dd,  $J_{8a,8b} = 12.5$  Hz,  $J_{7,8a} = 2.4$  Hz, 1H, H-8a), 4.29 (dd,  $J_{8a,8b} = 12.4$  Hz,  $J_{7,8b} = 4.8$  Hz, 1H, H-8b), 4.13 (dd,  $J_{6,7} = 9.5$  Hz,  $J_{5,6} = 1.3$  Hz, 1H, H-6), 3.99 (dd,  $J_{6a,6b} = 11.0$  Hz,  $J_{5,6a} = 2.0$  Hz, 1H, H-6aGlc), 3.78 (t,  $J_{2,3} \approx J_{3,4} \approx 9.2$  Hz, 1H, H-3Glc), 3.72–3.65 (m, 1H, H-5Glc), 3.59 (dd,  $J_{6a,6b} = 11.0$  Hz,  $J_{5,6b} = 5.2$  Hz, 1H, H-6bGlc), 3.53–3.44 (m, 2H, H-4Glc, H-2Glc), 3.38 (s, 3H,  $OCH_3$ ), 2.40 (dd,  $J_{3ax,3eq} = 12.5$  Hz,  $J_{3eq,4} = 4.6$  Hz, 1H, H-3eq), 2.15 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx 12.5$  Hz, 1H, H-3ax), 2.064, 2.062, 2.00, 1.98 (all s, 12H,  $4 \times COCH_3$ );  $^{13}C$  NMR (100 MHz,  $CDCl_3$ ,  $\beta$ -anomer)  $\delta$  171.1, 170.5, 169.90, 169.88 ( $4 \times COCH_3$ ), 167.6 (C-1,  $^3J_{C1,H3ax} = 6.2$  Hz), 139.0, 138.2, 134.9 ( $3 \times C$ -Ar), 128.9–127.8 ( $15 \times CH$ -Ar), 99.8 (C-2), 98.2 (C-1Glc), 81.7 (C-3Glc), 79.7 (C-2Glc), 75.6, 73.3 ( $2 \times CH_2Ph$ ), 71.0 (C-6), 70.3, 70.2 (C-5Glc, C-4Glc), 68.1 (C-7), 67.9 ( $CH_2Ph$ ), 67.1 (C-4), 64.08 (C-6Glc), 64.06 (C-5), 62.7 (C-8), 55.2 ( $OCH_3$ ), 32.2 (C-3), 20.9, 20.82, 20.79, 20.7 ( $4 \times COCH_3$ ); HRMS (ESI-TOF)  $m/z$   $[M + Na]^+$  calcd for

1  
2  
3  $C_{44}H_{52}NaO_{17}$  875.3097; found 875.3117;  $m/z$   $[2M + Na]^+$  calcd for  $C_{88}H_{104}NaO_{34}$  1727.6301;  
4  
5 found 1727.6336.  
6  
7  
8  
9

10 *2-(1-Nonyl) (4,5,7,8-Tetra-O-acetyl-3-deoxy-β-D-manno-oct-2-ulopyranosid)onic Acid (32)*. Kdo  
11 glycoside **22** (25 mg, 37 μmol, 1.0 equiv) was dissolved in 90% AcOH(aq) (1.1 mL) and the  
12 solution was heated to 35 °C. Freshly activated zinc powder (170 mg) was added in portions  
13 during 2 h. The mixture was filtered over Celite, rinsed with a 90% AcOH(aq) solution (5 mL)  
14 and a solution of EtOH/EtOAc (4 × 10 mL, 1:1 v/v). The solvents were concentrated under  
15 reduced pressure to afford a residue, which was purified by silica gel flash chromatography  
16 (DCM/MeOH 1:0 to 8:2) to give carboxylic acid **32** (17 mg, 89%) as a white amorphous powder.  
17  
18  $[\alpha]_D^{20} = +48$  (*c* 1.8, MeOH);  $^1H$  NMR (400 MHz, MeOD)  $\delta$  5.22 (br s, 1H, H-5), 5.17 (br t,  $J =$   
19 7.7 Hz, 1H, H-7), 5.00 (dt,  $J_{3ax,4} = 12.6$  Hz,  $J_{3eq,4} = 3.6$  Hz, 1H, H-4), 4.51 (d,  $J_{8a,8b} = 12.0$  Hz,  
20 1H, H-8a), 4.39 (d,  $J = 9.5$  Hz, 1H, H-6), 4.27 (dd,  $J_{8a,8b} = 12.0$  Hz,  $J_{7,8b} = 6.7$  Hz, 1H, H-8b),  
21 3.75 (dd,  $J = 15.6, 6.7$  Hz, 1H, H-1a'), 3.49 (dd,  $J = 15.2, 6.7$  Hz, 1H, H-1b'), 2.39 (dd,  $J_{3eq,3ax} =$   
22 11.8 Hz,  $J_{3eq,4} = 4.0$  Hz, 1H, H-3eq), 1.98 (t,  $J_{3eq,3ax} \approx J_{3ax,4} \approx 11.8$  Hz, 1H, H-3ax), 2.07, 2.03,  
23 1.98, 1.93 (all s, 12H, 4 × COCH<sub>3</sub>), 1.60–1.52 (m, 2H, H-2ab'), 1.39–1.23 (m, 12H, H-3ab', H-  
24 4ab', H-5ab', H-6ab', H-7ab', H-8ab'), 0.90 (t,  $J = 6.8$  Hz, 3H, H-9');  $^{13}C$  NMR (100 MHz,  
25 MeOD)  $\delta$  172.54, 172.45, 171.7, 171.6 (4 × COCH<sub>3</sub>), 102.3 (C-2), 72.0 (C-6), 70.0 (C-7), 69.7  
26 (C-4), 66.1 (C-5), 65.4 (C-1'), 64.7 (C-8), 33.9 (C-3), 33.0, 31.0, 30.7, 30.5, 30.4, 27.2, 23.7 (C-  
27 2', C-3', C-4', C-5', C-6', C-7', C-8'), 20.7 (4 × COCH<sub>3</sub>), 14.4 (C-9'); HRMS (ESI-TOF)  $m/z$   $[M$   
28 + Na]<sup>+</sup> calcd for  $C_{25}H_{40}NaO_{12}$  555.2412; found 555.2423;  $m/z$   $[2M + Na]^+$  calcd for  $C_{50}H_{80}NaO_{24}$   
29 1087.4932; found 1087.4953.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 *4'-Methoxyphenacyl* [2-(5-Amino-N-benzyloxycarbonyl-1-pentyl) 4,5,7,8-Tetra-O-acetyl-3-  
4  
5  
6 *deoxy- $\alpha,\beta$ -D-manno-oct-2-ulopyranosid]onic Acid (**33**). Kdo glycoside **15** (25 mg, 32  $\mu$ mol, 1.0  
7  
8 equiv) was dissolved in 90% AcOH(aq) (1.0 mL) and the solution was heated to 35 °C. Freshly  
9  
10 activated zinc powder (170 mg) was added in portions over 2 h. The mixture was filtered over  
11  
12 Celite, rinsed with a 90% AcOH(aq) solution (5 mL) and a solution of EtOH/EtOAc (4  $\times$  10 mL,  
13  
14 1:1 v/v). The solvents were concentrated under reduced pressure to afford a residue, which was  
15  
16 purified by silica gel flash chromatography (DCM/MeOH 1:0 to 8:2) to give carboxylic acid **33**  
17  
18 (17 mg, 85%) as a white amorphous powder.  $[\alpha]_D^{20} = +35$  (*c* 1.1, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz,  
19  
20 CDCl<sub>3</sub>)  $\delta$  7.38–7.30 (m, 5H, CH-Ar), 5.29 (br s, 1H, H-5), 5.19 (dt,  $J_{6,7} = 9.6$  Hz,  $J_{7,8a} \approx J_{7,8b} \approx$   
21  
22 3.1 Hz, 1H, H-7), 5.16–5.00 (m, 4H, CH<sub>2</sub>Ph, H-4, NHCbz), 4.40–4.32 (m, 1H, H-8a), 4.26 (d,  $J_{6,7}$   
23  
24 = 9.6 Hz, 1H, H-6), 3.80–3.73 (m, 1H, H-1a'), 3.65 (t,  $J_{7,8a} \approx J_{7,8b} \approx 6.6$  Hz, 1H, H-8b), 3.59–3.49  
25  
26 (m, 1H, H-1b'), 3.24–3.07 (m, 2H, H-5ab'), 2.41 (dd,  $J_{3ax,3eq} = 12.3$  Hz,  $J_{3eq,4} = 4.3$  Hz, 1H, H-  
27  
28 3eq), 2.06 (t,  $J_{3ax,3eq} \approx J_{3ax,4} \approx 12.3$  Hz, 1H, H-3ax), 2.10, 2.07, 2.01, 1.98 (all s, 12H, 4  $\times$   
29  
30 COCH<sub>3</sub>), 1.67–1.35 (m, 6H, H-2ab', H-3ab', H-4ab'); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  171.2,  
31  
32 170.7, 170.2, 170.0 (4  $\times$  COCH<sub>3</sub>), 136.5 (C-Ar), 128.6–128.3 (CH-Ar), 99.7 (C-2), 70.8 (C-6),  
33  
34 66.2 (C-7), 67.6 (C-4), 67.1 (CH<sub>2</sub>Ph), 64.3 (C-5), 63.7 (C-1'), 62.9 (C-8), 41.2 (C-5'), 32.3 (C-3),  
35  
36 29.9, 29.3, 23.0 (C-2', C-3', C-4'), 20.93–20.88 (4  $\times$  COCH<sub>3</sub>); HRMS (ESI-TOF) *m/z* [M + Na]<sup>+</sup>  
37  
38 calcd for C<sub>29</sub>H<sub>39</sub>NNaO<sub>14</sub> 648.2263; found 648.2280.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48*

## 49 ASSOCIATED CONTENT

### 50 Supporting Information

51  
52  
53  
54 The Supporting Information is available free of charge on the ACS Publications website at DOI:  
55  
56  
57  
58  
59  
60

1  
2  
3 NMR spectra for new compounds and computation results of reaction intermediates  
4  
5 (PDF)  
6  
7  
8  
9

## 10 AUTHOR INFORMATION

### 11 Corresponding Author

12 \*E-mail: charles.gauthier@iaf.inrs.ca  
13  
14

### 15 Notes

16  
17 The authors declare no competing financial interest.  
18  
19  
20  
21  
22  
23

## 24 ACKNOWLEDGMENTS

25  
26 This work was supported by grants from the Agence Nationale de la Recherche Program Jeunes  
27 Chercheuses Jeunes Chercheurs (ANR-JCJC-12-JS07-0003-01), the Direction Générale de  
28 l'Armement (DGA) French Minister of Defence, and the Natural Sciences and Engineering  
29 Research Council of Canada (NSERC). M. Z. thanks the Wroclaw Center of Biotechnology for  
30 PDoc funding within the framework of the Leading National Research Center (KNOW). This  
31 research has been enabled by the use of computing resources provided by WestGrid and Compute  
32 Canada. We thank Jean-François Gallard, from Institut de Chimie des Substances Naturelles  
33 CNRS-ICSN UPR 2301, for performing undecoupled  $^{13}\text{C}$  NMR spectra, and Emily R. Schwartz,  
34 from Georgia Institute of Technology, for proofreading this manuscript.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

## 50 REFERENCES

- 51  
52 (1) Knirel, Y. A.; Shevelev, S. D.; Perepelov, A. V. *Mendeleev Commun.* **2011**, *21*, 173.  
53  
54 (2) Lodowska, J.; Wolny, D.; Weglarz, L. *Can. J. Microbiol.* **2013**, *59*, 645.  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (3) Liparoti, V.; Molinaro, A.; Sturiale, L.; Garozzo, D.; Nazarenko, E. L.; Gorshkova, R. P.;  
4  
5 Ivanova, E. P.; Shevchenko, L. S.; Lanzetta, R.; Parrilli, M. *Eur. J. Org. Chem.* **2006**, 4710.  
6  
7 (4) Ovchinnikova, O. G.; Liu, B.; Guo, D.; Kocharova, N. A.; Shashkov, A. S.; Chem, M.;  
8  
9 Feng, L.; Rozalski, A.; Knirel, Y. A.; Wang, L. *Microbiology* **2012**, *158 (Pt 4)*, 1024.  
10  
11 (5) Vinogradov, E.; Frirdich, E.; MacLean, L. L.; Perry, M. B.; Petersen, B. O.; Duus, J. Ø.;  
12  
13 Whitfield, C. *J. Biol. Chem.* **2002**, *277*, 25070.  
14  
15 (6) Willis, L. M.; Whitfield, C. *Carbohydr. Res.* **2013**, *378*, 35.  
16  
17 (7) Starr, K. F.; Porsch, E. A.; Heiss, C.; Black, I.; Azadi, P.; St. Geme III, J. W. *PLoS One*  
18  
19 **2013**, *8*, e75409.  
20  
21 (8) Masoud, H.; Ho, M.; Schollaardt, T.; Perry, M. B. *J. Bacteriol.* **1997**, *179*, 5663.  
22  
23 (9) Nimtz, M.; Wray, V.; Domke, T.; Brenneke, B.; Häussler, S.; Steinmetz, I. *Eur. J.*  
24  
25 *Biochem.* **1997**, *250*, 608.  
26  
27 (10) Laroussarie, A.; Barycza, B.; Andriamboavonjy, H.; Tamigney Kenfack, M.; Bleriot, Y.;  
28  
29 Gauthier, C. *J. Org. Chem.* **2015**, *80*, 10386.  
30  
31 (11) Unger, F. M. *Adv. Carbohydr. Chem. Biochem* **1981**, *38*, 323.  
32  
33 (12) Lin, C.-H.; Murray, B. W.; Ollmann, I. R.; Wong, C.-H. *Biochemistry* **1997**, *36*, 780.  
34  
35 (13) Willis, L. M.; Stupak, J.; Richards, M. R.; Lowary, T. L.; Li, J.; Whitfield, C. *Proc. Natl.*  
36  
37 *Acad. Sci. USA* **2013**, *110*, 7868.  
38  
39 (14) Willis, L. M.; Whitfield, C. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 20753.  
40  
41 (15) Ovchinnikova, O. G.; Mallette, E.; Koizumi, H.; Lowary, T. L.; Kimber, M. S.; Whitfield,  
42  
43 C. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, E3120.  
44  
45 (16) Kosma, P. *Tetrahedron Lett.* **2016**, *57*, 2133.  
46  
47 (17) Pradhan, T. K.; Mong, K. K. *Isr. J. Chem.* **2015**, *55*.  
48  
49 (18) Tytgat, H. L. P.; Lebeer, S. *Microbiol. Mol. Biol. Rev.* **2014**, *78*, 372.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- (19) Ress, D. K.; Linhardt, R. J. *Curr. Org. Synth.* **2004**, *1*, 31.
- (20) Huang, J.-S.; Huang, W.; Meng, X.; Wang, X.; Gao, P.-C.; Yang, J. S. *Angew. Chem. Int. Ed.* **2015**, *54*, 10894.
- (21) Pokorny, B.; Kosma, P. *Org. Lett.* **2015**, *17*, 110.
- (22) Pokorny, B.; Kosma, P. *Chem. – Eur. J.* **2015**, *21*, 305.
- (23) Pokorny, B.; Kosma, P. *ChemistryOpen* **2015**, *4*, 722.
- (24) Qian, Y.; Feng, J.; Parvez, M.; Ling, C.-C. *J. Org. Chem.* **2012**, *77*, 96.
- (25) Tanaka, H.; Takahashi, D.; Takahashi, T. *Angew. Chem. Int. Ed.* **2006**, *45*, 770.
- (26) Pradhan, T. K.; Lin, C. C.; Mong, K. K. *Org. Lett.* **2014**, *16*, 1474.
- (27) Mannerstedt, K.; Ekelöf, K.; Oscarson, S. *Carbohydr. Res.* **2007**, *342*, 631.
- (28) Boons, G. J. P. H.; van Delft, F. L.; van der Klein, P. A. M.; van der Marel, G. A.; van Boom, J. H. *Tetrahedron* **1992**, *48*, 885.
- (29) van der Klein, P. A. M.; Filemon, W.; Boons, G. J. P. H.; Veeneman, G. H.; van der Marel, G. A.; van Boom, J. H. *Tetrahedron* **1992**, *48*, 4649.
- (30) Takahashi, T.; Tsukamoto, H.; Yamada, H. *Tetrahedron Lett.* **1997**, *38*, 8223.
- (31) Haberman, J. M.; Gin, D. Y. *Org. Lett.* **2001**, *3*, 1665.
- (32) Haberman, J. M.; Gin, D. Y. *Org. Lett.* **2003**, *5*, 2539.
- (33) Hanashima, S.; Akai, S.; Sato, K.-I. *Tetrahedron Lett.* **2008**, *49*, 5111.
- (34) Ishiwata, A.; Ito, Y. *Synlett* **2003**, *9*, 1339.
- (35) Hendrickson, J. B.; Kandall, C. *Tetrahedron Lett.* **1970**, *5*, 343.
- (36) Martin, A.; Arda, A.; Désiré, J.; Martin-Mingot, A.; Probst, N.; Sinaÿ, P.; Jiménez-Barbero, J.; Thibaudeau, S.; Blériot, Y. *Nat. Chem.* **2016**, *8*, 186.
- (37) Romero, J. A. C.; Tabacco, S. A.; Woerpel, K. A. *J. Am. Chem. Soc.* **2000**, *122*, 168.

- 1  
2  
3 (38) Ayala, L.; Lucero, C. G.; Romero, J. A. C.; Tabacco, S. A.; Woerpel, K. A. *J. Am. Chem.*  
4  
5 *Soc.* **2003**, *125*, 15521.  
6  
7  
8 (39) Chamberland, S.; Ziller, J. W.; Woerpel, K. A. *J. Am. Chem. Soc.* **2005**, *127*, 5322.  
9  
10 (40) Lucero, C. G.; Woerpel, K. A. *J. Org. Chem.* **2006**, *71*, 2641.  
11  
12 (41) Codée, J. D. C.; van den Bos, L. J.; de Jong, A.-R.; Dinkelaar, J.; Lodder, G.; Overkleef,  
13  
14 H. S.; van der Marel, G. A. *J. Org. Chem.* **2009**, *74*, 30.  
15  
16 (42) Cumpstey, I. *Org. Biomol. Chem.* **2012**, *10*, 2503.  
17  
18 (43) Dugave, C.; Demange, L. *Chem. Rev.* **2003**, *103*, 2475.  
19  
20 (44) Pawar, D. M.; Khalil, A. A.; Hooks, D. R.; Collins, K.; Elliott, T.; Stafford, J.; Smith, L.;  
21  
22 Noe, E. A. *J. Am. Chem. Soc.* **1998**, *120*, 2108.  
23  
24 (45) Ghalambor, M. A.; Heath, E. C. *Biochem. Biophys. Res. Commun.* **1963**, *11*, 288.  
25  
26 (46) Shirai, R.; Ogura, H. *Tetrahedron Lett.* **1989**, *30*, 2263.  
27  
28 (47) Cornforth, J. W.; Firth, M. E.; Gottschalk, A. *Biochem. J.* **1958**, *68*, 57.  
29  
30 (48) Mikula, H.; Blaukopf, M.; Sixta, G.; Stanetty, C.; Kosma, P. In *Carbohydrate Chemistry:*  
31  
32 *Proven Synthetic Methods*; van der Marel, G., Codee, J., Eds.; CRC Press: Boca Raton, 2014;  
33  
34 Vol. 2, 207 pp.  
35  
36 (49) Unger, F. M.; Stix, D.; Schulz, G. *Carbohydr. Res.* **1980**, *80*, 191.  
37  
38 (50) Guo, X., *PhD Thesis*, ETH Zurich, 2011.  
39  
40 (51) Hashimoto, S.; Hayashi, M.; Naylor, R. *Tetrahedron Lett.* **1984**, *25*, 1379.  
41  
42 (52) Patel, M. K.; Vijayakrishnan, B.; Koeppe, J. R.; Chalker, J. M.; Doores, K. J.; Davis, B.  
43  
44 *G. Chem. Commun.* **2010**, *46*, 9119.  
45  
46 (53) Meijer, A.; Ellervik, U. *J. Org. Chem.* **2002**, *67*, 7407.  
47  
48 (54) Lemieux, R. U.; Hendriks, K. B.; Stick, R. V.; James, K. *J. Am. Chem. Soc.* **1975**, *97*,  
49  
50 4056.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (55) Fügedi, P.; Garegg, P. J. *Carbohydr. Res.* **1986**, *149*, C9.  
4  
5 (56) Tatai, J.; Fügedi, P. *Org. Lett.* **2007**, *9*, 4647.  
6  
7 (57) Veeneman, G. H.; van Leeuwen, S. H.; van Boom, J. H. *Tetrahedron Lett.* **1990**, *31*, 1331.  
8  
9 (58) Konradsson, P.; Mootoo, D. R.; McDevitt, R. E.; Fraser-Reid, B. *J. Chem. Soc. Chem. Commun.* **1990**, 270.  
10  
11 (59) Konradsson, P.; Udodong, U. E.; Fraser-Reid, B. *Tetrahedron Lett.* **1990**, *31*, 4313.  
12  
13 (60) Demchenko, A. V. *Synlett* **2003**, *9*, 1225.  
14  
15 (61) Yang, L.; Qin, Q.; Ye, X. S. *Asian J. Org. Chem.* **2013**, *2*, 30.  
16  
17 (62) Baek, J. Y.; Lee, B.-Y.; Jo, M. G.; Kim, K. S. *J. Am. Chem. Soc.* **2009**, *131*, 17705.  
18  
19 (63) Hasty, S. J.; Ranade, S. C.; Demchenko, A. V. *Rep. Org. Chem.* **2014**, *4*, 1.  
20  
21 (64) Imoto, M.; Kusunose, N.; Matsuura, Y.; Kusumoto, S.; Shiba, T. *Tetrahedron Lett.* **1987**,  
22  
23 *28*, 6277.  
24  
25 (65) Kocieński, P. J. *Protecting groups*, 3rd Edition; Georg Thieme Verlag: Stuttgart, 2005.  
26  
27 (66) Zhang, Y.; Knapp, S. *J. Org. Chem.* **2016**, *81*, 2228.  
28  
29 (67) Krog-Jensen, C.; Oscarson, S. *J. Org. Chem.* **1996**, *61*, 1234.  
30  
31 (68) Grayson, E. J.; Ward, S. J.; Hall, A. L.; Rendle, P. M.; Gamblin, D. P.; Batsanov, A. S.;  
32  
33 Davis, B. G. *J. Org. Chem.* **2005**, *70*, 9740.  
34  
35 (69) Bai, Y.; Lowary, T. L. *J. Org. Chem.* **2006**, *71*, 9672.  
36  
37 (70) Crich, D.; Hu, T.; Cai, F. *J. Org. Chem.* **2008**, *73*, 8942.  
38  
39 (71) Willoughby, P. H.; Jansma, M. J.; Hoye, T. R. *Nat. Protoc.* **2014**, *9*, 643.  
40  
41 (72) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J.  
42  
43 R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.;  
44  
45 Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.;  
46  
47 Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.;  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Vreven, T.; Montgomery, J. A. J.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers,  
4 E.; Kudin, K. N.; Staroverov, V. N.; Keith, T.; Kobayashi, R.; Normand, J.; Raghavachari, K.;  
5 Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene,  
6 M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R.  
7 E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.;  
8 Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.;  
9 Daniels, A. D.; Farkas, O.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J.; Revision D.01  
10 ed. 2013.

11  
12  
13 (73) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 765.

14  
15 (74) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648.

16  
17 (75) Hariharan, P. C.; Pople, J. A. *Theoret. Chim. Acta* **1973**, *28*, 213.

18  
19 (76) Tomasi, J.; Mennucci, B.; Cancès, E. *J. Mol. Struct. THEOCHEM* **1999**, *464*, 211.

20  
21 (77) Krishnan, R.; Binkley, J. S.; Seeger, R.; Pople, J. A. *J. Chem. Phys.* **1980**, *72*, 650.

22  
23 (78) Clark, T.; Chandrasekhar, J.; Spitznagel, G. W.; Schleyer, P. V. R. *J. Comput. Chem.*  
24  
25 **1983**, *4*, 294.

26  
27 (79) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. *J. Chem. Phys.* **2010**, *132*, 154104.

28  
29 (80) Nakamoto, S.-I.; Achiwa, K. *Chem. Pharm. Bull.* **1987**, *35*, 4537.