



HAL
open science

Lack of evidence for Zika virus transmission by Culex mosquitoes

Christopher M Roundy, Sasha R Azar, Aaron C Brault, Gregory D Ebel, Anna-Bella Failloux, Ildefonso Fernandez-Salas, Uriel Kitron, Laura D Kramer, Ricardo Lourenço-De-Oliveira, Jorge E Osorio, et al.

► **To cite this version:**

Christopher M Roundy, Sasha R Azar, Aaron C Brault, Gregory D Ebel, Anna-Bella Failloux, et al.. Lack of evidence for Zika virus transmission by Culex mosquitoes. *Emerging microbes & infections*, 2017, 6, pp.e90. 10.1038/emi.2017.85 . pasteur-01677831

HAL Id: pasteur-01677831

<https://pasteur.hal.science/pasteur-01677831>

Submitted on 8 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

LETTER TO THE EDITOR

Lack of evidence for Zika virus transmission by *Culex* mosquitoes

Christopher M Roundy¹, Sasha R Azar¹, Aaron C Brault², Gregory D Ebel³, Anna-Bella Failloux⁴, Ildfonso Fernandez-Salas⁵, Uriel Kitron⁶, Laura D Kramer⁷, Ricardo Lourenço-de-Oliveira⁸, Jorge E Osorio⁹, Igor D Paploski¹⁰, Gonzalo M Vazquez-Prokopec⁶, Guilherme S Ribeiro¹¹, Scott A Ritchie¹², Laura B Tauro¹⁰, Nikos Vasilakis¹³ and Scott C Weaver¹

Emerging Microbes & Infections (2017) 6, e90; doi:10.1038/emi.2017.85; published online 18 October 2017

Dear Editor,

Since Zika virus (ZIKV) emerged in the Americas, major research efforts have been focused on identifying the mosquito species responsible for transmission. While almost all published results support *Aedes aegypti* and potentially *Ae. albopictus* as urban vectors, a recent article¹ suggests that *Culex quinquefasciatus* may serve as a ZIKV vector in Recife, Brazil, a region that has experienced a high incidence of infection. Accurately identifying the vector of a pathogen enables public health agencies to implement appropriate control strategies and inform citizens of proper prevention measures. Additionally, establishing the vector for an emerging pathogen paves the way for researchers to advance our understanding of virus–vector interactions and pursue novel methods of control. In contrast, erroneously incriminating a vector could lead to misdirected use of limited government funds, diversion of research efforts and misinforming the public through misdirected media and educational programs.

Traditional criteria for arthropod vector incrimination include: (i) demonstration of feeding or other effective contact with pathogen's host; (ii) association in time and space of the vector and the pathogen-infected host; (iii) repeated demonstration of natural infection of the vector and (iv) experimental transmission of the pathogen by the vector.²

For ZIKV transmission in the Americas, criterion 3 has been met only for *Ae. aegypti*, with detection of naturally infected mosquitoes with titers compatible with transmission competence in Mexico (reviewed in ref. 3) and Brazil.⁴ *Ae. albopictus* has also been shown to be capable of laboratory transmission^{5–9} (see also references reviewed in refs 3, 10). Although no field infections have been reported for *Ae. albopictus* in the Americas, they were detected during a 2007

Gabon outbreak (reviewed in ref. 3). In locations where *Ae. aegypti* has been found infected at high rates in the Americas, testing of *Cx. quinquefasciatus*, typically the most common urban tropical mosquito, was unsuccessful aside from three pools collected in Recife, Brazil described by Guedes *et al.*¹ However, the ZIKV RNA levels measured in these Recife pools, reflected in high Ct values (37.6–38.15) representing < 10 infectious units in typical RT-qPCR assays, are incompatible with transmission-competent mosquitoes, which typically have viral titers several orders of magnitude higher^{5–9,11} (see also references reviewed in refs 3, 10). Even naturally infected mosquitoes without viral dissemination to the salivary glands typically have higher titers⁷ (see also references in ref. 10) than reported by Guedes *et al.*¹ Thus, the wild-caught Recife mosquito pools likely contained trace amounts of residual, viremic blood in their guts, undetectable by their colorimetric assay, legs or other dislodged appendages from other infected mosquitoes of different species, or were false-positives.

In laboratory studies of *Cx. quinquefasciatus* other than those of Guedes *et al.*,¹ only one other group has shown transmission of ZIKV by colonized mosquitoes, from an unreported generation number maintained since 2014.¹² At least 15 other studies have found no transmission competence,^{6,7,13,14} (see also references reviewed in ref. 3, 10) even after examining several combinations of geographic strains of mosquito as well as ZIKV, along with different methods of oral exposure. These include other studies from Brazil with colonies established in 2016 and another study from China.⁷ Even after intrathoracic inoculation, generally the most permissive route for arbovirus infection of mosquitoes, *Culex* mosquitoes were found to be refractory to disseminated infection (reviewed in refs 3, 10). Although some of these studies found infection of the midgut without

¹Institute for Human Infections and Immunity, Institute for Translational Research, and Department of Microbiology and Immunology, University of Texas Medical Branch, Galveston, TX, USA; ²Division of Vector-Borne Infectious Disease, Centers for Disease Control and Prevention, Fort Collins, CO, USA; ³Arthropod-Borne and Infectious Diseases Laboratory, Department of Microbiology, Immunology and Pathology, Colorado State University, Ft. Collins, CO, USA; ⁴Institut Pasteur, Department of Virology, Arboviruses and Insect Vectors, Paris, France; ⁵Centro Regional de Investigación en Salud Pública-INSP, Tapachula, Chiapas, Mexico; ⁶Department of Environmental Sciences, Emory University, Atlanta, GA, USA; ⁷Wadsworth Center, New York State Department of Health, Albany, NY, USA; ⁸Laboratório de Mosquitos Transmissores de Hematozoários, Instituto Oswaldo Cruz, Rio de Janeiro, Brazil; ⁹Department of Pathobiological Sciences, University of Wisconsin, Madison, WI, USA; ¹⁰Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Salvador, Brazil; ¹¹Instituto de Saúde Coletiva, Universidade Federal da Bahia, Salvador, Brazil and Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Salvador, Brazil; ¹²Australian Institute of Tropical Health and Medicine, James Cook University, Smithfield, QLD, Australia and ¹³Institute for Human Infections and Department of Pathology, University of Texas Medical Branch, Galveston, TX, USA

Correspondence: SC Weaver

E-mail: sweaver@utmb.edu

Received 3 September 2017; accepted 19 September 2017

dissemination to the saliva, the majority found no indication of any infection after oral exposure. Many of these ZIKV-refractory populations tested are highly competent for West Nile and St. Louis encephalitis flaviviruses, so the specific ZIKV block in these populations would need to be restricted to most but not all *Cx. quinquefasciatus* populations to explain the results of Guedes *et al.*¹ In addition, *Cx. quinquefasciatus* from Recife challenged in another study with the same BRPE243/2015 ZIKV strain used by Guedes *et al.*,¹ as well as with two other ZIKV strains, were consistently refractory to oral infection.¹⁵

The discrepancy between the negative results from so many published studies and the questionable findings of Guedes *et al.*¹ should engender caution in interpretation and conclusions reported by media and public health authorities unless they are verified by more robust results including detection of genuinely transmission-competent mosquitoes in nature. Until further data are collected and other groups can replicate the Recife findings, it is important that public education and interventions remain focused on the conclusion supported by the vast majority of studies: *Ae. aegypti* is the only mosquito species for which we have strong evidence of ZIKV transmission in the Americas.

This submission represents the views solely of the authors and does not constitute those of the Centers for Disease Control and Prevention or the United States Government.

- 1 Guedes DR, Paiva MH, Donato MM *et al.* Zika virus replication in the mosquito *Culex quinquefasciatus* in Brazil. *Emerg Microbes Infect* 2017; **6**: e69.
- 2 Barnett HC. The incrimination of arthropods as vectors of disease. *Proc 11th Int Congr Entomol* 1960; **2**: 341–345.
- 3 Vasilakis N, Weaver SC. Flavivirus transmission focusing on Zika. *Curr Opin Virol* 2017; **22**: 30–35.

- 4 Ferreira-de-Brito A, Ribeiro IP, Miranda RM *et al.* First detection of natural infection of *Aedes aegypti* with Zika virus in Brazil and throughout South America. *Mem Inst Oswaldo Cruz* 2016; **111**: 655–658.
- 5 Azar SR, Roundy CM, Rossi SL *et al.* Differential vector competency of *Aedes albopictus* populations from the Americas for Zika virus. *Am J Trop Med Hygiene* 2017; **97**: 330–339.
- 6 Duchemin JB, Mee PT, Lynch SE *et al.* Zika vector transmission risk in temperate Australia: a vector competence study. *Virology* 2017; **14**: 108.
- 7 Liu Z, Zhou T, Lai Z *et al.* Competence of *Aedes aegypti*, *Ae. albopictus*, and *Culex quinquefasciatus* mosquitoes as Zika virus vectors, China. *Emerg Infect Dis* 2017; **23**: 1085–1091.
- 8 Ciota AT, Bialosuknia SM, Zink SD *et al.* Effects of Zika virus strain and *Aedes* mosquito species on vector competence. *Emerg Infect Dis* 2017; **23**: 1110–1117.
- 9 Di Luca M, Severini F, Toma L *et al.* Experimental studies of susceptibility of Italian *Aedes albopictus* to Zika virus. *Euro Surveill* 2016; **21**: 6–9.
- 10 Lourenco-de-Oliveira R, Failloux AB. Lessons learned on Zika virus vectors. *PLoS Negl Trop Dis* 2017; **11**: e0005511.
- 11 Li MI, Wong PS, Ng LC *et al.* Oral susceptibility of Singapore *Aedes (Stegomyia) aegypti* (Linnaeus) to Zika virus. *PLoS Negl Trop Dis* 2012; **6**: e1792.
- 12 Guo XX, Li CX, Deng YQ *et al.* *Culex pipiens quinquefasciatus*: a potential vector to transmit Zika virus. *Emerg Microbes Infect* 2016; **5**: e102.
- 13 Kenney JL, Romo H, Duggal NK *et al.* Transmission incompetence of *Culex quinquefasciatus* and *Culex pipiens pipiens* from North America for Zika virus. *Am J Trop Med Hygiene* 2017; **96**: 1235–1240.
- 14 Dodson BL, Rasgon JL. Vector competence of *Anopheles* and *Culex* mosquitoes for Zika virus. *PeerJ* 2017; **5**: e3096.
- 15 Fernandes RS, Campos SS, Ribeiro PS *et al.* *Culex quinquefasciatus* from areas with the highest incidence of microcephaly associated with Zika virus infections in the Northeast Region of Brazil are refractory to the virus. *Mem Inst Oswaldo Cruz* 2017; **112**: 577–579.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017