

Reverse Epidemiology: An Experimental Framework to Drive Leishmania Biomarker Discovery in situ by Functional Genetic Screening Using Relevant Animal Models

Laura Piel, Pascale Pescher, Gerald F Späth

▶ To cite this version:

Laura Piel, Pascale Pescher, Gerald F Späth. Reverse Epidemiology: An Experimental Framework to Drive Leishmania Biomarker Discovery in situ by Functional Genetic Screening Using Relevant Animal Models. Frontiers in Cellular and Infection Microbiology, 2018, 8, pp.325. 10.3389/fcimb.2018.00325. pasteur-01974302

HAL Id: pasteur-01974302 https://pasteur.hal.science/pasteur-01974302

Submitted on 8 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Reverse Epidemiology: An Experimental Framework to Drive *Leishmania* Biomarker Discovery *in situ* by Functional Genetic Screening Using Relevant Animal Models

Laura Piel^{1,2}, Pascale Pescher¹ and Gerald F. Späth^{1*}

¹ Institut Pasteur, Unité de Parasitologie Moléculaire et Signalisation, INSERM U1201, Paris, France, ² Université Paris Diderot, Sorbonne Paris Cité, Paris, France

OPEN ACCESS

Edited by:

Javier Moreno, Instituto de Salud Carlos III, Spain

Reviewed by:

Elisa Azuara-Liceaga, Universidad Autónoma de la Ciudad de México, Mexico Bellisa Freitas Barbosa, Federal University of Uberlandia, Brazil

> *Correspondence: Gerald F. Späth gerald.spaeth@pasteur.fr

Specialty section:

This article was submitted to Parasite and Host, a section of the journal Frontiers in Cellular and Infection Microbiology

Received: 18 June 2018 Accepted: 27 August 2018 Published: 19 September 2018

Citation:

Piel L, Pescher P and Späth GF (2018) Reverse Epidemiology: An Experimental Framework to Drive Leishmania Biomarker Discovery in situ by Functional Genetic Screening Using Relevant Animal Models. Front. Cell. Infect. Microbiol. 8:325. doi: 10.3389/fcimb.2018.00325 Leishmania biomarker discovery remains an important challenge that needs to be revisited in light of our increasing knowledge on parasite-specific biology, notably its genome instability. In the absence of classical transcriptional regulation in these early-branching eukaryotes, fluctuations in transcript abundance can be generated by gene and chromosome amplifications, which have been linked to parasite phenotypic variability with respect to virulence, tissue tropism, and drug resistance. Conducting in vitro evolutionary experiments to study mechanisms of Leishmania environmental adaptation, we recently validated the link between parasite genetic amplification and fitness gain, thus defining gene and chromosome copy number variations (CNVs) as important Leishmania biomarkers. These experiments also demonstrated that long-term Leishmania culture adaptation can strongly interfere with epidemiologically relevant, genetic signals, which challenges current protocols for biomarker discovery, all of which rely on in vitro expansion of clinical isolates. Here we propose an experimental framework independent of long-term culture termed "reverse" epidemiology, which applies established protocols for functional genetic screening of cosmid-transfected parasites in animal models for the identification of clinically relevant genetic loci that then inform targeted field studies for their validation as Leishmania biomarkers.

Keywords: Leishmania, biomarker discovery, reverse epidemiology, cosmid screen, functional genetics

INTRODUCTION

Biomarkers are defined as biological characteristics that are objective and quantifiable indicators for responses to therapeutic interventions, or normal and pathogenic biological processes (Biomarkers Definitions Working Group 2001, 2001). With respect to *Leishmania* infection, we can distinguish direct biomarkers that are applied to determine parasite species and prevalence (e.g., parasite-specific proteins, lipids, transcripts, genetic loci), and indirect biomarkers that correspond to different correlates of the host anti-microbial response [e.g., adenosine deaminase (ADA) or cytokines such as IL-10 or TNF] (Kip et al., 2015).

Direct *Leishmania* biomarkers can have either purely diagnostic value (e.g., kinetoplast (k) DNA, ribosomal small sub-unit (SSU) RNA, HSP70 locus, carbohydrate antigens), or prognostic value allowing for the prediction of treatment outcome or disease evolution (e.g., dissemination

in cutaneaous leishmaniasis or development of post-kala-azar dermal leishmaniasis in visceral leishmaniasis). However, despite their potentially important impact on clinical management of leishmaniasis, only few biomarker candidates with potential prognostic value are described, most of which are linked to drug resistance (Vanaerschot et al., 2012; Torres et al., 2013; Hefnawy et al., 2017; Ponte-Sucre et al., 2017). The absence of this class of markers is explained by various biological and technical constraints, some of which are linked to *Leishmania* genome instability that limits biomarker discovery and needs to be considered in ongoing and future biomarkers discovery campaigns.

In the absence of classical transcriptional regulation, Leishmania often regulates transcript and protein abundance by chromosome and gene copy number variations (CNVs) (Dumetz et al., 2017; Prieto Barja et al., 2017), which can drive environmental adaptation (Leprohon et al., 2009; Downing et al., 2011; Rogers et al., 2011; Brotherton et al., 2013; Mukherjee et al., 2013; Ubeda et al., 2014; Zhang et al., 2014; Laffitte et al., 2016). Our recent demonstration that karyotypic fluctuations and haplotype selection allow for fitness gain in culture reveals the importance of Leishmania genome plasticity in short-term evolutionary adaptation (Prieto Barja et al., 2017). Conceivably, the highly dynamic genomic changes occurring during culture adaptation challenge past and current protocols in Leishmania biomarker discovery, which rely on adaptation and massexpansion of field isolates in culture prior to analysis, often resulting in loss of epidemiologically relevant, genetic signals. Here, by drawing from the current literature, we propose an alternative strategy independent of long-term culture that is based on functional genetic screening in relevant animal models. Our review provides an overview on past functional screening results and their documented success in revealing genomic loci that are under environmental selection, and advocates for Leishmania biomarker discovery by combining cosmid selection and subsequent clinical validation, an experimental framework we termed "reverse" epidemiology. In the following we summarize studies that developed and applied cosmid-based approaches to identify new Leishmania factors linked to parasite pathogenicity, tropism and drug resistance, and discuss the potential epidemiological relevance of these factors where clinical data were available.

COSMID-BASED FUNCTIONAL GENETIC SCREENING IN *LEISHMANIA*

Various genetic methods have been successfully applied in the past to identify *Leishmania* genes or genetic markers that are associated with disease outcome or clinical manifestation, including whole genome sequencing (WGS) of isolates (Downing et al., 2011; Rogers et al., 2011; Leprohon et al., 2015), random amplification of polymorphic DNA (RAPD) (Bhattacharyya et al., 1993; Schönian et al., 1996; Mkada-Driss et al., 2014), or assessment of amplified fragment length polymorphisms (AFLP) (Kumar et al., 2009, 2010a; Odiwuor et al., 2011; Jaber et al., 2018). Likewise, cosmid-based functional screens have been applied to discover clinically relevant loci. This approach is based on the genetic transfer of a given cellular phenotype (e.g., drug resistance) from a donor strain to a recipient strain via transfection of a cosmid library. While currently established WGS protocols for *Leishmania* biomarker discovery have been applied on clinical isolates maintained in long-term culture, causing potentially important bias, cosmidbased approaches can directly reveal clinically relevant genotypephenotype relationships, especially when applied *in situ* in infected animals. Even though this functional genetic approach represents a powerful tool, this technology has not been applied in a systematic way at larger scale to drive biomarker discovery.

The preparation and application of a cosmid library is a complex procedure, where genomic DNA fragments of an appropriate size are cloned into purified cosmid DNA and packaged into phages for efficient bacterial transduction, which allows for amplification of the library and assessment of its genomic coverage prior to transfection into parasites by electroporation. The generation of a first series of *Leishmania* shuttle cosmid vectors and the validation of a protocol that allows for genetic complementation and functional screening in these parasites using genomic cosmid libraires was established in 1993 by Beverley and collaborators (Ryan et al., 1993a) followed by Kelly and collaborators in 1994 (Kelly et al., 1994). Subsequently, this protocol was applied in various studies for the identification of *Leishmania* pathogenicity and drug resistance genes.

Cosmid-Based Identification of Novel Leishmania Pathogenicity Factors

Key for Leishmania infectivity is the capacity of procyclic promastigotes to undergo differentiation into infectious metacyclic promastigotes able to resist to complement lysis encountered inside the mammal host following parasite transmission (Sacks and Perkins, 1984; Franke et al., 1985; Howard et al., 1987). Parasite resistance has been largely attributed to the surface glycolipid lipophosphoglycan (LPG), a major Leishmania virulence factor essential for L. major promastigote virulence (Späth et al., 2000, 2003a), that undergoes important modifications during metacyclogenesis (Sacks et al., 1990; Mcconville et al., 1992; Sacks, 2001). LPG biosynthetic genes and their virulence functions have been genetically identified combining cosmid screens with functional null mutant analysis and virulence assessment in macrophages and mice. LPG deficient mutants were generated by chemical mutagenesis, isolated by their failure to agglutinate in the presence of lectin (King and Turco, 1988), transfected with a cosmid library prepared from L. donovani, and screened for restoration of LPG expression using either lectin- or antibody-based agglutination assays revealing the two first LPG biosynthetic genes, a galactofuranose transferase encoded by the gene *lpg1* (Ryan et al., 1993b), and an UdP galactose transporter encoded by lpg2 (Descoteaux et al., 1995). The virulence functions of both genes were confirmed in subsequent studies in L. major lpg1 and lpg2 null mutants (Späth et al., 2000, 2003b). Few years later, by combining cosmid library transfection and antibody panning, Dobson et al. identifed genes encoding arabinosyl- and galactosyltransferases that mediate developmental modifications of LPG during metacyclogenesis (Dobson et al., 2003a,b).

Cosmid-based functional screening has also been applied to gain insight into pathways that govern complement resistance in promastigotes revealing genes that are likely linked to metacyclogenesis. Based on the observation that decrease in resistance to complement lysis is a consequence of long-term maintenance in culture (Lincoln et al., 2004), Dahlin-Laborde et al. used genomic DNA from animal-derived Leishmania infantum (chagasi) promastigotes to construct a cosmid library that was transfected into long-term cultured parasites. The transfected parasites were subjected to complement lysis allowing for the selection of seven different cosmids that conferred increased complement resistance albeit at lower levels compared to short-term cultured control parasites. In-depth analysis of two cosmids revealed genomic fragments of L. infantum chromosome 36 (Dahlin-Laborde et al., 2005), with two subregions encoding, respectively, 5 and 13 genes shown to be critical for the phenotype, including an ADP-ribosylation factorlike protein and an ATP-dependent RNA helicase (Dahlin-Laborde et al., 2008). Cosmid screens were further applied by the Matlashewski team to identify virulence and visceralization factors using libraries prepared with genomic DNA from L. donovani transfected in L. major promastigotes. Transfectants expressing the heterologous library were inoculated into mice by tail vein or footpad injections and cosmids were recovered from parasites that established infection in spleen (type I), skin (type III), or both (type II). Subsequent analysis of individual ORFs by transgenic expression and infection validated an ORF encoding for an unknown protein and a 4.4 kb miniexon gene array on chromosome 36 (Zhang and Matlashewski, 2004). Unlike in L. major, overexpression of the miniexon region in L. braziliensis led to complete virulence attenuation in a hamster model (de Toledo et al., 2009), suggesting species-specific functions of this array. This is further supported by the genetic divergence of this array between new world and old world dermotropic species (Fernandes et al., 1994), which is used as a diagnostic signal for parasite genotyping (Serin et al., 2005; Ovalle-Bracho et al., 2016).

A final example documenting the power of cosmid-based approaches in identifying putative Leishmania virulence factors is represented by a complementation screen conducted using a cosmid library derived from an attenuated HSP100 null mutant that spontaneously recovered infectivity and/or pathogenicity in mice, likely by the amplification of a compensatory locus (Reiling et al., 2006). A screen conducted in mice using cosmidtransfected HSP100 null mutants and subsequent validation experiments revealed P46 as a new virulence factor (Reiling et al., 2010). A follow-up study by Bifeld et al applied a phylogenetic approach on 20 clinical isolates comparing P46 amino acid sequences thus establishing a strong correlation between P46 isoforms and their geographical origin. Transgenic parasites over-expressing three different P46 isoforms in a L. major lab strain were co-injected in BALB/c and C57BL/6 mice. Selection of different isoforms according to the mouse strain suggested that the P46 genetic polymorphism may be linked to parasite adaptation to genetically distinct, region-specific host reservoirs (Bifeld et al., 2015; Table 1).

Cosmid-Based Identification of *Leishmania* Drug Resistance Genes

Since 1999, screening of cosmid libraries has been used as a gain-of-function strategy to identify drug resistance or drug tolerance genes (reviewed in Clos and Choudhury, 2006). Beverley and collaborators established the first proof-of-principle of this approach culturing cosmid transfected *L. major* parasites under pressure of the drugs methotrexate and tubercidin, which resulted in the selection of the known resistance genes *DHFR-TS*, *PTR1*, and *TOR* (Cotrim et al., 1999). The same study identified a new gene encoding a 63 kDa hypothetical protein located on chromosome 31 termed tubercidin-resistant protein (TRP) that is conserved in *Leishmania* and co-localizes in the endoplasmic reticulum in stationary phase promastigotes (Aoki et al., 2016).

Functional complementation has also been a powerful tool for the identification of transporters that can alter drug efficacy. The biopterin transporter bt1, previously named ORF G (Kundig et al., 1999), and the miltefosine (MIL) transporter LdMT (Perez-Victoria et al., 2003) were identified using Leishmania tarentolae transfected with a heterologous L. mexicana cosmid library selected under methotrexate pressure (showing that bt1 can confer resistance), and L. donovani MIL resistant parasites transfected with a L. donovani wild-type cosmid library subjected to MIL selection (showing that a non-mutated LdMT can restore susceptibility). Likewise, the cosmid approach was applied to screen for genes mediating resistance to two inhibitors of ergosterol biosynthesis, terbinafine, and itraconazole, which resulted in the selection of nine different cosmids, some of which conferred cross-resistance to both drugs, and the identification of squalene synthase 1 (SQS1) as an itraconazole resistance gene (Cotrim et al., 1999).

This approach has been recently applied to directly identify clinically relevant drug resistance loci by heterologous screening. Clos and collaborators prepared cosmid libraries from antimony SbIII/SbV resistant or SbIII sensitive/SbV resistant L. braziliensis field isolates that were transfected into SbIII sensitive/SbV resistant promastigotes. Culture under drug pressure selected for cosmids carrying a genomic fragment of chromosome 20, which also conferred drug resistance when transfected into L. infantum (Nühs et al., 2014). A competition assay with full-length or truncated derivatives of the cosmid insert validated ARM58 as a SbIII resistance gene. A more recent study performed by the same group with cosmid-transfected L. infantum extended this finding to the neighboring genes and defined a cluster of three genes, ARM58, ARM56 (previously named ARM58rel), and HSP23 at the telomere of the chromosome 34 that confer increased resistance of intracellular amastigotes against SbV (Tejera Nevado et al., 2016). Using a L. infantum cosmid library, the same team revealed a protein termed P299 that conferred increased resistance of intracellular amastigotes to MIL and reduced promastigote sensitivity to MIL and SbIII, but not pentamidin (Choudhury et al., 2008). Another gene-today annotated as cysteine leucine-rich protein (CLrP, LinJ.34.0570)was revealed causing antimony resistance in L. tarentolae transfected with a cosmid library prepared from arsenite and SbIII resistant parasites (Brochu et al., 2004), and in L. infantum axenic amastigotes (Genest et al., 2008). Brochu et al. also

Gene	ID	chr*	Product	Function	Strain/Isolate	References	Validation**
lpg1	LmjF.25.0010	25	Beta galactofuranosyl transferase	Galactofuranosyl transferase implicated in LPG biosynthesis	L. donovani L1S2D	Ryan et al., 1993b	Yes (Späth et al., 2000)
lpg2	LmjF.34.3120	34	Lipophosphoglycan biosynthetic protein 2	Transmembrane <i>L. donovani</i> 1S/Cl2D Descoteatransporter activity		Descoteaux et al., 1995	Yes (Späth et al., 2003b)
sca2	LmjF.02.0180	2	Phosphoglycan beta 1,2 arabinosyltransferase	Galactosyltransferase <i>L. major</i> FV1 Dobson et al., 2 activity		Dobson et al., 2003a	No
sca1	LmjF.02.0220	2	Phosphoglycan beta 1,2 arabinosyltransferase	Galactosyltransferase activity	L. major FV1	Dobson et al., 2003a	No
scg1	LmjF.07.1170	7	Phosphoglycan beta 1,3 galactosyltransferase 1	Galactosyltransferase	<i>L. major</i> FV1	Dobson et al., 2003b	No
scg2	LmjF.21.0010	21	Phosphoglycan beta 1,3 galactosyltransferase 2	Galactosyltransferase activity	L. major FV1	Dobson et al., 2003b	No
scg3	LmjF.02.0010	2	Phosphoglycan beta 1,3 galactosyltransferase 3	Galactosyltransferase	L. major FV1	Dobson et al., 2003b	No
scg4	LmjF.36.0010	36	phosphoglycan beta 1,3 galactosyltransferase 4	Galactosyltransferase	L. major FV1	Dobson et al., 2003b	No
scg5	LmjF.31.3190	31	phosphoglycan beta 1,3 galactosyltransferase 5	Galactosyltransferase activity	L. major FV1	Dobson et al., 2003b	No
scg6	LmjF.25.2460	25	Phosphoglycan beta 1,3 galactosyltransferase 6	Galactosyltransferase activity	L. major FV1	Dobson et al., 2003b	No
Miniexon		36	Miniexon		L. donovani 1S/Cl2D	Zhang and Matlashewski, 2004	No
P46	LmjF.33.3060	33	46 kD virulence factor	Unknown	L. major 5ASKH	Reiling et al., 2010	No
	LmJF36.0790 - LmjF.36.0840	36	Specific genes involved in phenotype not identified	Not applicable	L. infantum (chagasi)	Dahlin-Laborde et al., 2005	No
	LmJF36.0840 - LmjF.36.0900	36	Specific genes involved in phenotype not identified	Not applicable	L. infantum (Chagasi)	Dahlin-Laborde et al., 2008	No
	LmJF36.3090 - LmjF.36.3210	36	Specific genes involved in phenotype not identified	Not applicable	L. infantum (chagasi)	Dahlin-Laborde et al., 2008	No

*chr, chromosome; **validation refers to loss of function studies establishing a direct link between the gene and parasite pathogenicity.

reported members of the HSP70 protein family as important genes contributing to antimony tolerance, supporting recent phylogenetic evidence that HSP70 family members may allow parasite environmental adaptation with potential important consequences for drug susceptibility (Drini et al., 2016).

Recent work by the Ouellette team coupled cosmid selection and next generation sequencing for drug resistance and drug target gene discovery, proposing a high-throughput capable screening strategy the authors referred to as Cos-Seq (Gazanion et al., 2016). Screening cosmid transfected L. infantum against SbIII, amphotericin B, MIL, paramomycin or pentamidin revealed 64 enriched loci, including 12 common to at least two anti-leishmanial drugs, suggesting the existence of multi-drug resistance genes. This study validated 6 known and uncovered 7 new resistance genes in promastigotes, including two new genes causing methotrexate resistance both encoding for phosphatase 2C-like proteins (LinJ.34.2310 and LinJ.34.2320), one hypothetical protein with leucinerich repeats causing both pentamidin and paromomycin resistance (LinJ.06.1010), a serine/threonine phosphatase causing SbIII resistance (LinJ.12.0610), and phospholipid-translocating ATPase (LinJ30.2270) and C-8 sterol isomerase (LinJ.29.2250) that were revealed screening for MIL resistance (Table 2).

THE FRAMEWORK OF "REVERSE" EPIDEMIOLOGY

The examples described above are testimony to the success of cosmid-based, functional screening approaches to discover genetic loci in Leishmania that are linked to parasite virulence, tissue tropism, and drug resistance. However, even though these loci may represent potential biomarkers with important prognostic value, there are no dedicated, concerted efforts for their validation in clinically relevant settings. One exception includes CLrP, whose increased abundance on RNA and protein levels were correlated with increased Sb resistance in field isolates, albeit only a small number of isolates were used in these studies (Kumar et al., 2010b; Das et al., 2015). For other loci, clinical validation of the functional screening results can be ambiguous, with for example the MRPA and PTR1 genes of the H-locus having been either strictly, partially, or not correlated to Sb resistance in different epidemiological studies (Decuypere et al., 2005, 2012; Mittal et al., 2007; Mukherjee et al., 2007; Mukhopadhyay et al., 2011). Such divergent results may be explained by the polyclonal structure of parasite field isolates and their geographic adaptation, with different resistance mechanisms being selected in genetically distinct

TABLE 2	Genes identified b	v cosmid-based	approach	linked to	l eishmania	drua	resistance o	r suscei	otibility	J.
		y 00011110 00000	approuori	1111001101	Loioi II mai ma	arug	10010101100 0	00000	public	1.

Gene	ID	Product	Function	Drug	Strain*	Validation**	References	
TRP	LmjF.31.2010	Turbicidin-resistant protein	ER protein	Tubercidin	L. major	No	Cotrim et al., 1999	
bt1	LmxM.34.5150	Biopterin transporter	Folate/biopterin transport	Methotrexate	L. mexicana	No	Kundig et al., 1999	
SQS1	LmjF.31.2940	Squalene synthase	Ergosterol biosynth	Itraconazole	L. major	No	Cotrim et al., 1999	
ARM58	LbrM20.0210; LinJ.34.0220	58 kDa antimony resistance marker	Response to drug	SbIII	L. braziliensis; L. infantum	No	Nühs et al., 2014; Tejera Nevado et al., 2016	
ARM56	LinJ.34.0210	56 kDa antimony resistance marker	Response to drug	SbIII	L. braziliensis; L. infantum	No	Nühs et al., 2014; Tejera Nevado et al., 2016	
HSP23	LinJ.34.0230	HSP 23	Response to drug		L. infantum	No	Tejera Nevado et al., 2016	
P299	LinJ.08.0630	P299	Response to drug	Miltefosine/ SbIII	L. infantum	Partially Downing et al., 2011; Jeddi et al., 2014	Choudhury et al., 2008	
CLrP	LinJ.34.0570	Cysteine leucine rich protein	Response to drug	SbIII	L. infantum	Partially (Kumar et al., 2010b; Das et al., 2015)	Genest et al., 2008	
LdMT	LdBPK_131590.1	Miltefosine transporter	Phospholipid-translocating ATPase	Miltefosine	L. donovani	Partially (Coelho et al., 2012; Mondelaers et al., 2016; Shaw et al., 2016; Srivastava et al., 2017)	Perez-Victoria et al., 2003	
	LinJ.34.2310	Phosphatase 2C-like proteins	Catalytic activity	Methotrexate	L. infantum	No	Gazanion et al., 2016	
	LinJ.34.2320 Phosphatase 2C-like proteins		Catalytic activity	Methotrexate L. infantum		No	Gazanion et al., 2016	
	LinJ.06.1010	Leucine Rich Repeat, putative	Protein binding	Pentamidin/ paromomycin	L. infantum	No	Gazanion et al., 2016	
	LinJ.12.0610	Serine/threonine phosphatase	Hydrolase activity and ion binding	SbIII	L. infantum	No	Gazanion et al., 2016	
	LinJ30.2270	Phospholipid- translocating ATPase	Transmembrane transporter	Miltefosine	L. infantum	No	Gazanion et al., 2016	
	LinJ.29.2250	C-8 sterol isomerase	Isomerase	Miltefosine	L. infantum	No	Gazanion et al., 2016	

*Strain used for the generation of the cosmid library; **biological validation in field isolates

isolates (Decuypere et al., 2012). This possibility is supported by our recent demonstration that genetic mosaicism in an individual L. donovani strain can drive polyclonal adaptation, suggesting that different resistance mechanisms may co-exist in sub-populations of any given isolate (Prieto Barja et al., 2017). Such intra-strain specific, polyclonal fitness gain is further supported by the cosmid selection of different genetic loci in response to the same selection pressure applied on a single parasite population in vitro or during animal infection (Cotrim et al., 1999; Dahlin-Laborde et al., 2005; Gazanion et al., 2016). Indeed, such clonal phenotypic variability in a given parasite isolate has been recently documented in L. amazonensis, with important differences in culture proliferation and pathogenic potential observed in untransfected sub-clones or parasites transfected with individual cosmids selected in vivo for increased parasite infectivity (Espiau et al., 2017). Finally, other genes associated with drug resistance or susceptibility identified in cosmid screens failed to be validated in clinical studies such as LdMT, whose mutations were correlated to MIL resistance in promastigotes in culture but could not be associated with MIL resistance or treatment failure in the field (Bhandari et al., 2012). Likewise, PRP1 that has been implicated *in vitro* in resistance to pentamidine with reported cross-resistance to SbIII, did not show increased expression in Sb resistant field isolates (Decuypere et al., 2005, 2012).

Drawing from these examples we propose an experimental framework for the discovery of biomarker candidates by combining functional genetic screens in relevant animal models to reveal loci of interest, which then are validated by dedicated clinical and epidemiological investigations (Figure 1). In this approach, a cosmid library is prepared from parasites freshly derived from clinical isolates that show a phenotype of interest (donor strain). The gene(s) that express this phenotype are identified by transfecting a relevant recipient Leishmania strain and recovery of cosmids from transfectants that gained the phenotype under investigation. These genes can then be validated as biomarkers by quantitative PCR analysis directly applied on clinical samples. Thus, in contrast to classical biomarker discovery, where epidemiological field studies establish a correlation between a clinical phenotype and a genetic locus that then is validated in vitro or in animal studies, the epidemiological protocol we propose is in reverse



hamster infection (in the presence of drug in our example). The selected gene(s) of interested (GOI) will be identified by next generation sequencing (NGS). Correlating the identified genes with the clinical phenotype in dedicated epidemiological studies will then validate the new biomarker.

from lab-based studies back to the field. Even though this approach has its drawbacks (e.g., clinical manifestations caused by gene inactivation or gene deletion cannot be revealed), it provides several interesting advantages that immediately overcome important bottlenecks in *Leishmania* biomarker discovery. First, it is independent of long-term culture that can have an important impact on the parasite genome thus interfering with epidemiologically relevant information. Second, the screening is performed *in situ* in infected animals under environmental constraints that correspond to the clinical setting, thus allowing for the selection of physiologically highly relevant loci. Third, large amounts of parasite can be recovered from different tissues of the infected animals, which can be subjected to direct and even single cell sequencing, thus informing on mechanisms of polyclonal adaptation that may be relevant to the field. Finally, this approach will overcome ethical concerns associated with applying direct genome sequencing on human tissue samples as the cosmid-identified loci will be studied in clinical samples by simple qPCR analysis.

In conclusion, our reverse epidemiology approach exploits genetic amplification for biomarker discovery and thus mimics the very mechanism that has been linked to *Leishmania* genomic adaptation and fitness gain in the field and in culture (Dumetz et al., 2017; Prieto Barja et al., 2017). Cosmid-based functional genetic screening *in situ* linked to clinical validation thus represents a powerful framework that can fill an important gap in the currently rather desolate state of *Leishmania* biomarker discovery, which is challenged by the absence of robust protocols for direct tissue sequencing of parasites in human clinical samples, and the genetic bias caused by parasite long-term culture applied in current epidemiological investigations.

REFERENCES

- Aoki, J. I., Coelho, A. C., Muxel, S. M., Zampieri, R. A., Sanchez, E. M., Nerland, A. H., et al. (2016). Characterization of a novel endoplasmic reticulum protein involved in tubercidin resistance in *Leishmania* major. *PLoS Negl. Trop. Dis.* 10:e0004972. doi: 10.1371/journal.pntd.0004972
- Bhandari, V., Kulshrestha, A., Deep, D. K., Stark, O., Prajapati, V. K., Ramesh, V., et al. (2012). Drug susceptibility in *Leishmania* isolates following miltefosine treatment in cases of visceral leishmaniasis and post kala-azar dermal leishmaniasis. *PLoS Negl. Trop. Dis.* 6:e1657. doi: 10.1371/journal.pntd.0001657
- Bhattacharyya, R., Singh, R., Hazra, T. K., and Majumder, H. K. (1993). Application of polymerase chain reaction with specific and arbitrary primers to identification and differentiation of *Leishmania* parasites. *FEMS Microbiol. Lett.* 114, 99–104. doi: 10.1111/j.1574-6968.1993.tb06557.x
- Bifeld, E., Chrobak, M., Zander, D., Schleicher, U., Schönian G, and Clos, J. (2015). Geographical sequence variation in the *Leishmania* major virulence factor P46. *Infect. Genet. Evol.* 30, 195–205. doi: 10.1016/j.meegid.2014.12.029
- Biomarkers Definitions Working Group 2001 (2001). Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin. Pharmacol. Ther.* 69, 89–95. doi: 10.1067/mcp.2001.113989
- Brochu, C., Haimeur, A., and Ouellette, M. (2004). The heat shock protein HSP70 and heat shock cognate protein HSC70 contribute to antimony tolerance in the protozoan parasite *Leishmania*. *Cell Stress Chaper*. 9, 294–303. doi: 10.1379/CSC-15R1.1
- Brotherton, M. C., Bourassa, S., Leprohon, P., Legare, D., Poirier, G. G., Droit, A., et al. (2013). Proteomic and genomic analyses of antimony resistant *Leishmania infantum* mutant. *PLoS ONE* 8:e81899. doi: 10.1371/journal.pone.0081899
- Choudhury, K., Zander, D., Kube, M., Reinhardt, R., and Clos, J. (2008). Identification of a *Leishmania infantum* gene mediating resistance to miltefosine and SbIII. *Int. J. Parasitol.* 38, 1411–1423. doi: 10.1016/j.ijpara.2008.03.005
- Clos, J., and Choudhury, K. (2006). Functional cloning as a means to identify Leishmania genes involved in drug resistance. Mini Rev. Med. Chem. 6, 123–129. doi: 10.2174/138955706775476028
- Coelho, A. C., Boisvert, S., Mukherjee, A., Leprohon, P., Corbeil, J., and Ouellette, M. (2012). Multiple mutations in heterogeneous miltefosineresistant *Leishmania* major population as determined by whole genome sequencing. *PLoS Negl. Trop. Dis.* 6:e1512. doi: 10.1371/journal.pntd.0001512
- Cotrim, P. C., Garrity, L. K., and Beverley, S. M. (1999). Isolation of genes mediating resistance to inhibitors of nucleoside and ergosterol metabolism in *Leishmania* by overexpression/selection. *J. Biol. Chem.* 274, 37723–37730. doi: 10.1074/jbc.274.53.37723
- Dahlin-Laborde, R. R., Scolaro, E. J., Romine, N. M., Ramer-Tait, A. E., Lei, S. M., and Beetham, J. K. (2008). Characterization of DNA sequences that confer complement resistance in *Leishmania* chagasi. *Ann. N. Y. Acad. Sci.* 1149, 347–351. doi: 10.1196/annals.1428.099

AUTHOR CONTRIBUTIONS

LP wrote the chapter on the use of cosmid libraries regarding virulence and tropism, PP wrote the chapter on the use of cosmid libraries regarding drug resistance, GS corrected the manuscript and wrote introduction and the last chapter detailing the experimental framework.

FUNDING

This work was supported by funds from the Institute Pasteur International Direction strategic fund to the LeiSHield project and the Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases (Grant no. ANR-10-LABX-62-IBEID). LP was supported by the Ph.D. fellowship from the Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation.

- Dahlin-Laborde, R. R., Yu, T. P., and Beetham, J. K. (2005). Genetic complementation to identify DNA elements that influence complement resistance in *Leishmania* chagasi. *J. Parasitol.* 91, 1058–1063. doi: 10.1645/GE-477R.1
- Das, S., Shah, P., Tandon, R., Yadav, N. K., Sahasrabuddhe, A. A., Sundar, S., et al. (2015). Over-expression of cysteine leucine rich protein is related to SAG resistance in clinical isolates of *Leishmania donovani*. *PLoS Negl. Trop. Dis.* 9:e0003992. doi: 10.1371/journal.pntd.0003992
- de Toledo, J. S., Junqueira Dos Santos, A. F., Rodrigues De Moura, T., Antoniazi, S. A., Brodskyn, C., Indiani De Oliveira, C., et al. (2009). *Leishmania* (Viannia) braziliensis transfectants overexpressing the miniexon gene lose virulence in vivo. Parasitol. Int. 58, 45–50. doi: 10.1016/j.parint.2008.09.006
- Decuypere, S., Rijal, S., Yardley, V., De Doncker, S., Laurent, T., Khanal, B., et al. (2005). Gene expression analysis of the mechanism of natural Sb(V) resistance in *Leishmania donovani* isolates from Nepal. *Antimicrob. Agents Chemother.* 49, 4616–4621. doi: 10.1128/AAC.49.11.4616-4621.2005
- Decuypere, S., Vanaerschot, M., Brunker, K., Imamura, H., Muller, S., Khanal, B., et al. (2012). Molecular mechanisms of drug resistance in natural *Leishmania* populations vary with genetic background. *PLoS Negl. Trop. Dis.* 6:e1514. doi: 10.1371/journal.pntd.0001514
- Descoteaux, A., Luo, Y., Turco, S. J., and Beverley, S. M. (1995). A specialized pathway affecting virulence glycoconjugates of *Leishmania*. Science 269, 1869–1872. doi: 10.1126/science.7569927
- Dobson, D. E., Mengeling, B. J., Cilmi, S., Hickerson, S., Turco, S. J., and Beverley, S. M. (2003a). Identification of genes encoding arabinosyltransferases (SCA) mediating developmental modifications of lipophosphoglycan required for sand fly transmission of *Leishmania* major. J. Biol. Chem. 278, 28840–28848. doi: 10.1074/jbc.M302728200
- Dobson, D. E., Scholtes, L. D., Valdez, K. E., Sullivan, D. R., Mengeling, B. J., Cilmi, S., et al. (2003b). Functional identification of galactosyltransferases (SCGs) required for species-specific modifications of the lipophosphoglycan adhesin controlling *Leishmania* major-sand fly interactions. J. Biol. Chem. 278, 15523–15531. doi: 10.1074/jbc.M301568200
- Downing, T., Imamura, H., Decuypere, S., Clark, T. G., Coombs, G. H., Cotton, J. A., et al. (2011). Whole genome sequencing of multiple *Leishmania donovani* clinical isolates provides insights into population structure and mechanisms of drug resistance. *Genome Res.* 21, 2143–2156. doi: 10.1101/gr.12 3430.111
- Drini, S., Criscuolo, A., Lechat, P., Imamura, H., Skalicky, T., Rachidi, N., et al. (2016). Species- and strain-specific adaptation of the HSP70 super family in pathogenic trypanosomatids. *Genome Biol. Evol.* 8, 1980–1995. doi: 10.1093/gbe/evw140
- Dumetz, F., Imamura, H., Sanders, M., Seblova, V., Myskova, J., Pescher, P., et al. (2017). Modulation of aneuploidy in *Leishmania donovani* during adaptation to different *in vitro* and *in vivo* environments and its impact on gene expression. *MBio* 8:e00599-17. doi: 10.1128/mBio.00599-17

- Espiau, B., Vilhena, V., Cuvillier, A., Barral, A., and Merlin, G. (2017). Phenotypic diversity and selection maintain *Leishmania amazonensis* infectivity in BALB/c mouse model. *Mem. Inst. Oswaldo Cruz* 112, 44–52. doi: 10.1590/0074-02760160280
- Fernandes, O., Murthy, V. K., Kurath, U., Degrave, W. M., and Campbell, D. A. (1994). Mini-exon gene variation in human pathogenic *Leishmania* species. *Mol. Biochem. Parasitol.* 66, 261–271. doi: 10.1016/0166-6851(94)90153-8
- Franke, E. D., Mcgreevy, P. B., Katz, S. P., and Sacks, D. L. (1985). Growth cycledependent generation of complement-resistant *Leishmania* promastigotes. *J. Immunol.* 134, 2713–2718.
- Gazanion, E., Fernandez-Prada, C., Papadopoulou, B., and Ouellette, M. (2016). Cos-Seq for high-throughput identification of drug target and resistance mechanisms in the protozoan parasite *Leishmania*. *Proc. Natl. Acad. Sci. U.S.A.* 113, E3012–E3021. doi: 10.1073/pnas.1520693113
- Genest, P. A., Haimeur, A., Legare, D., Sereno, D., Roy, G., Messier, N., et al. (2008). A protein of the leucine-rich repeats (LRRs) superfamily is implicated in antimony resistance in *Leishmania infantum* amastigotes. *Mol. Biochem. Parasitol.* 158, 95–99. doi: 10.1016/j.molbiopara.2007.11.008
- Hefnawy, A., Berg, M., Dujardin, J. C., and De Muylder, G. (2017). Exploiting knowledge on *Leishmania* drug resistance to support the quest for new drugs. *Trends Parasitol.* 33, 162–174. doi: 10.1016/j.pt.2016.11.003
- Howard, M. K., Sayers, G., and Miles, M. A. (1987). Leishmania donovani metacyclic promastigotes: transformation in vitro, lectin agglutination, complement resistance, and infectivity. Exp. Parasitol. 64, 147–156. doi: 10.1016/0014-4894(87)90138-X
- Jaber, H. T., Hailu, A., Pratlong, F., Lami, P., Bastien, P., and Jaffe, C. L. (2018). Analysis of genetic polymorphisms and tropism in East African *Leishmania donovani* by amplified fragment length polymorphism and kDNA minicircle sequencing. *Infect. Genet. Evol.* 65, 80–90. doi: 10.1016/j.meegid.2018.07.016
- Jeddi, F., Mary, C., Aoun, K., Harrat, Z., Bouratbine, A., Faraut, F., et al. (2014). Heterogeneity of molecular resistance patterns in antimony-resistant field isolates of *Leishmania* species from the western Mediterranean area. *Antimicrob. Agents Chemother.* 58, 4866–4874. doi: 10.1128/AAC.02521-13
- Kelly, J. M., Das, P., and Tomas, A. M. (1994). An approach to functional complementation by introduction of large DNA fragments into *Trypanosoma cruzi* and *Leishmania donovani* using a cosmid shuttle vector. *Mol. Biochem. Parasitol.* 65, 51–62. doi: 10.1016/0166-6851(94)90114-7
- King, D. L., and Turco, S. J. (1988). A ricin agglutinin-resistant clone of *Leishmania donovani* deficient in lipophosphoglycan. *Mol. Biochem. Parasitol.* 28, 285–293. doi: 10.1016/0166-6851(88)90013-8
- Kip, A. E., Balasegaram, M., Beijnen, J. H., Schellens, J. H., De Vries, P. J., and Dorlo, T. P. (2015). Systematic review of biomarkers to monitor therapeutic response in leishmaniasis. *Antimicrob. Agents Chemother.* 59, 1–14. doi: 10.1128/AAC.04298-14
- Kumar, A., Boggula, V. R., Misra, P., Sundar, S., Shasany, A. K., and Dube, A. (2010a). Amplified fragment length polymorphism (AFLP) analysis is useful for distinguishing *Leishmania* species of visceral and cutaneous forms. *Acta Trop.* 113, 202–206. doi: 10.1016/j.actatropica.2009.10.006
- Kumar, A., Boggula, V. R., Sundar, S., Shasany, A. K., and Dube, A. (2009). Identification of genetic markers in sodium antimony gluconate (SAG) sensitive and resistant Indian clinical isolates of *Leishmania donovani* through amplified fragment length polymorphism (AFLP). *Acta Trop.* 110, 80–85. doi: 10.1016/j.actatropica.2009.01.005
- Kumar, A., Sisodia, B., Misra, P., Sundar, S., Shasany, A. K., and Dube, A. (2010b). Proteome mapping of overexpressed membrane-enriched and cytosolic proteins in sodium antimony gluconate (SAG) resistant clinical isolate of *Leishmania donovani. Br. J. Clin. Pharmacol.* 70, 609–617. doi: 10.1111/j.1365-2125.2010.03716.x
- Kundig, C., Haimeur, A., Legare, D., Papadopoulou, B., and Ouellette, M. (1999). Increased transport of pteridines compensates for mutations in the high affinity folate transporter and contributes to methotrexate resistance in the protozoan parasite *Leishmania* tarentolae. *EMBO J.* 18, 2342–2351. doi: 10.1093/emboj/18.9.2342
- Laffitte, M. N., Leprohon, P., Papadopoulou, B., and Ouellette, M. (2016). Plasticity of the *Leishmania* genome leading to gene copy number variations and drug resistance. *F1000 Res.* 5:2350. doi: 10.12688/f1000research.9218.1
- Leprohon, P., Fernandez-Prada, C., Gazanion, E., Monte-Neto, R., and Ouellette, M. (2015). Drug resistance analysis by next generation

sequencing in Leishmania. Int. J. Parasitol. Drugs Drug Resist. 5, 26-35. doi: 10.1016/j.ijpddr.2014.09.005

- Leprohon, P., Légaré, Raymond, F., Madore, E., Hardiman, G., Corbeil, J., et al. (2009). Gene expression modulation is associated with gene amplification, supernumerary chromosomes and chromosome loss in antimony-resistant *Leishmania infantum*. *Nucleic Acids Res.* 37, 1387–1399. doi: 10.1093/nar/gkn1069
- Lincoln, L. M., Ozaki, M., Donelson, J. E., and Beetham, J. K. (2004). Genetic complementation of *Leishmania* deficient in PSA (GP46) restores their resistance to lysis by complement. *Mol. Biochem. Parasitol.* 137, 185–189. doi: 10.1016/j.molbiopara.2004.05.004
- Mcconville, M. J., Turco, S. J., Ferguson, M. A., and Sacks, D. L. (1992). Developmental modification of lipophosphoglycan during the differentiation of *Leishmania* major promastigotes to an infectious stage. *EMBO J.* 11, 3593–3600.
- Mittal, M. K., Rai, S., Ashutosh, Ravinder, Gupta, S., Sundar, S., et al. (2007). Characterization of natural antimony resistance in *Leishmania donovani* isolates. *Am. J. Trop. Med. Hyg.* 76, 681–688. Available online at: https://www. ajtmh.org
- Mkada-Driss, I., Lahmadi, R., Chakroun, A. S., Talbi, C., Guerbouj, S., Driss, M., et al. (2014). Screening and characterization of RAPD markers in viscerotropic *Leishmania* parasites. *PLoS ONE* 9:e109773. doi: 10.1371/journal.pone.0109773
- Mondelaers, A., Sanchez-Cañete MP, Hendrickx, S., Eberhardt, E., Garcia-Hernandez, R., Lachaud, L., et al. (2016). Genomic and molecular characterization of miltefosine resistance in *Leishmania infantum* strains with either natural or acquired resistance through experimental selection of intracellular amastigotes. *PLoS ONE* 11:e0154101. doi: 10.1371/journal.pone.0154101
- Mukherjee, A., Boisvert, S., Monte-Neto, R. L., Coelho, A. C., Raymond, F., Mukhopadhyay, R., et al. (2013). Telomeric gene deletion and intrachromosomal amplification in antimony-resistant *Leishmania*. *Mol. Microbiol.* 88, 189–202. doi: 10.1111/mmi.12178
- Mukherjee, A., Padmanabhan, P. K., Singh, S., Roy, G., Girard, I., Chatterjee, M., et al. (2007). Role of ABC transporter MRPA, gamma-glutamylcysteine synthetase and ornithine decarboxylase in natural antimony-resistant isolates of *Leishmania donovani*. J. Antimicrob. Chemother. 59, 204–211. doi: 10.1093/jac/dkl494
- Mukhopadhyay, R., Mukherjee, S., Mukherjee, B., Naskar, K., Mondal, D., Decuypere, S., et al. (2011). Characterisation of antimony-resistant *Leishmania donovani* isolates: biochemical and biophysical studies and interaction with host cells. *Int. J. Parasitol.* 41, 1311–1321. doi: 10.1016/j.ijpara.2011.07.013
- Nühs, A., Schafer, C., Zander, D., Trube, L., Tejera Nevado, P., Schmidt, S., et al. (2014). A novel marker, ARM58, confers antimony resistance to *Leishmania* spp. *Int. J. Parasitol. Drugs Drug Resist.* 4, 37–47. doi: 10.1016/j.ijpddr.2013.11.004
- Odiwuor, S., Vuylsteke, M., De Doncker, S., Maes, I., Mbuchi, M., Dujardin, J. C., et al. (2011). *Leishmania* AFLP: paving the way towards improved molecular assays and markers of diversity. *Infect. Genet. Evol.* 11, 960–967. doi: 10.1016/j.meegid.2011.03.008
- Ovalle-Bracho, C., Díaz-Toro, Y. R., and Muvdi-Arenas, S. (2016). Polymerase chain reaction-miniexon: a promising diagnostic method for mucocutaneous leishmaniasis. *Int. J. Dermatol.* 55, 531–539. doi: 10.1111/ijd.12910
- Perez-Victoria, F. J., Gamarro, F., Ouellette, M., and Castanys, S. (2003). Functional cloning of the miltefosine transporter. a novel P-type phospholipid translocase from *Leishmania* involved in drug resistance. *J. Biol. Chem.* 278, 49965–49971. doi: 10.1074/jbc.M308352200
- Ponte-Sucre, A., Gamarro, F., Dujardin, J. C., Barrett, M. P., López-Vélez, R., García-Hernández R., et al. (2017). Drug resistance and treatment failure in leishmaniasis: a 21st century challenge. *PLoS Negl. Trop. Dis.* 11:e0006052. doi: 10.1371/journal.pntd.0006052
- Prieto Barja, P., Pescher, P., Bussotti, G., Dumetz, F., Imamura, H., Kedra, D., et al. (2017). Haplotype selection as an adaptive mechanism in the protozoan pathogen *Leishmania donovani*. *Nat. Ecol. Evol.* 1, 1961–1969. doi: 10.1038/s41559-017-0361-x
- Reiling, L., Chrobak, M., Schmetz, C., and Clos, J. (2010). Overexpression of a single *Leishmania* major gene enhances parasite infectivity *in vivo* and *in vitro*. *Mol. Microbiol.* 76, 1175–1190. doi: 10.1111/j.1365-2958.2010. 07130.x

- Reiling, L., Jacobs, T., Kroemer, M., Gaworski, I., Graefe, S., and Clos, J. (2006). Spontaneous recovery of pathogenicity by *Leishmania* major hsp100-/- alters the immune response in mice. *Infect. Immun.* 74, 6027–6036. doi: 10.1128/IAI.00773-05
- Rogers, M. B., Hilley, J. D., Dickens, N. J., Wilkes, J., Bates, P. A., Depledge, D. P., et al. (2011). Chromosome and gene copy number variation allow major structural change between species and strains of *Leishmania. Genome Res.* 21, 2129–2142. doi: 10.1101/gr.122945.111
- Ryan, K. A., Dasgupta, S., and Beverley, S. M. (1993a). Shuttle cosmid vectors for the trypanosomatid parasite *Leishmania*. *Gene* 131, 145–150.
- Ryan, K. A., Garraway, L. A., Descoteaux, A., Turco, S. J., and Beverley, S. M. (1993b). Isolation of virulence genes directing surface glycosylphosphatidylinositol synthesis by functional complementation of *Leishmania*. *Proc. Natl. Acad. Sci. U.S.A.* 90, 8609–8613.
- Sacks, D. L. (2001). Leishmania-sand fly interactions controlling species-specific vector competence. Cell. Microbiol. 3, 189–196. doi: 10.1046/j.1462-5822.2001.00115.x
- Sacks, D. L., Brodin, T. N., and Turco, S. J. (1990). Developmental modification of the lipophosphoglycan from *Leishmania* major promastigotes during metacyclogenesis. *Mol. Biochem. Parasitol.* 42, 225–233. doi: 10.1016/0166-6851(90)90165-I
- Sacks, D. L., and Perkins, P. V. (1984). Identification of an infective stage of *Leishmania* promastigotes. *Science* 223, 1417–1419. doi: 10.1126/science.6701528
- Schönian G., Schweynoch, C., Zlateva, K., Oskam, L., Kroon, N., Graser, Y., et al. (1996). Identification and determination of the relationships of species and strains within the genus *Leishmania* using single primers in the polymerase chain reaction. *Mol. Biochem. Parasitol.* 77, 19–29. doi: 10.1016/0166-6851(96)02572-8
- Serin, M. S., Daglioglu, K., Bagirova, M., Allahverdiyev, A., Uzun, S., Vural, Z., et al. (2005). Rapid diagnosis and genotyping of *Leishmania* isolates from cutaneous and visceral leishmaniasis by microcapillary cultivation and polymerase chain reaction-restriction fragment length polymorphism of miniexon region. *Diagn. Microbiol. Infect. Dis.* 53, 209–214. doi: 10.1016/j.diagmicrobio.2005. 05.007
- Shaw, C. D., Lonchamp, J., Downing, T., Imamura, H., Freeman, T. M., Cotton, J. A., et al. (2016). *In vitro* selection of miltefosine resistance in promastigotes of *Leishmania donovani* from Nepal: genomic and metabolomic characterization. *Mol. Microbiol.* 99, 1134–1148. doi: 10.1111/mmi.13291
- Späth, G. F., Epstein, L., Leader, B., Singer, S. M., Avila, H. A., Turco, S. J., et al. (2000). Lipophosphoglycan is a virulence factor distinct from related glycoconjugates in the protozoan parasite *Leishmania* major. *Proc. Natl. Acad. Sci. U.S.A.* 97, 9258–9263. doi: 10.1073/pnas.160257897

- Späth, G. F., Garraway, L. A., Turco, S. J., and Beverley, S. M. (2003a). The role(s) of lipophosphoglycan (LPG) in the establishment of *Leishmania* major infections in mammalian hosts. *Proc. Natl. Acad. Sci. U.S.A.* 100, 9536–9541. doi: 10.1073/pnas.1530604100
- Späth, G. F., Lye, L. F., Segawa, H., Sacks, D. L., Turco, S. J., and Beverley, S. M. (2003b). Persistence without pathology in phosphoglycan-deficient *Leishmania* major. *Science* 301, 1241–1243. doi: 10.1126/science.1087499
- Srivastava, S., Mishra, J., Gupta, A. K., Singh, A., Shankar, P., and Singh, S. (2017). Laboratory confirmed miltefosine resistant cases of visceral leishmaniasis from India. *Parasit. Vectors* 10:49. doi: 10.1186/s13071-017-1969-z
- Tejera Nevado, P., Bifeld, E., Hohn, K., and Clos, J. (2016). A telomeric cluster of antimony resistance genes on chromosome 34 of *Leishmania infantum*. *Antimicrob. Agents Chemother*. 60, 5262–5275. doi: 10.1128/AAC.00544-16
- Torres, D. C., Ribeiro-Alves, M., Romero, G. A., Dávila, A. M., and Cupolillo, E. (2013). Assessment of drug resistance related genes as candidate markers for treatment outcome prediction of cutaneous leishmaniasis in Brazil. *Acta Trop.* 126, 132–141. doi: 10.1016/j.actatropica.2013.02.002
- Ubeda, J. M., Raymond, F., Mukherjee, A., Plourde, M., Gingras, H., Roy, G., et al. (2014). Genome-wide stochastic adaptive DNA amplification at direct and inverted DNA repeats in the parasite *Leishmania*. *PLoS Biol*. 12:e1001868. doi: 10.1371/journal.pbio.1001868
- Vanaerschot, M., Decuypere, S., Downing, T., Imamura, H., Stark, O., De Doncker, S., et al. (2012). Genetic markers for SSG resistance in *Leishmania donovani* and SSG treatment failure in visceral leishmaniasis patients of the Indian subcontinent. J. Infect. Dis. 206, 752–755. doi: 10.1093/infdis/jis424
- Zhang, W. W., and Matlashewski, G. (2004). In vivo selection for Leishmania donovani miniexon genes that increase virulence in Leishmania major. Mol. Microbiol. 54, 1051–1062. doi: 10.1111/j.1365-2958.2004.04327.x
- Zhang, W. W., Ramasamy, G., Mccall, L. I., Haydock, A., Ranasinghe, S., Abeygunasekara, P., et al. (2014). Genetic analysis of *Leishmania donovani* tropism using a naturally attenuated cutaneous strain. *PLoS Pathog.* 10:e1004244. doi: 10.1371/journal.ppat.1004244

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Piel, Pescher and Späth. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.