



# Quantitative genetics of *Aedes aegypti* vector competence for dengue viruses: towards a new paradigm?

Louis A Lambrechts

## ► To cite this version:

Louis A Lambrechts. Quantitative genetics of *Aedes aegypti* vector competence for dengue viruses: towards a new paradigm?. Trends in Parasitology, 2011, 27 (3), pp.111-114. 10.1016/j.pt.2010.12.001 . pasteur-02011024

**HAL Id: pasteur-02011024**

**<https://pasteur.hal.science/pasteur-02011024>**

Submitted on 8 Feb 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Quantitative genetics of *Aedes aegypti* vector**  
2 **competence for dengue viruses: towards a new**  
3 **paradigm?**

4

5

6 **Louis Lambrechts**

7 Department of Virology, Institut Pasteur, 25 rue du Docteur Roux, 75724 Paris Cedex 15,  
8 France.

9

10 *Corresponding author:* Lambrechts, L. (louis.lambrechts@pasteur.fr)

11

## 1   **Abstract**

2

3   Like many other host-pathogen interactions, the vector competence of *Aedes aegypti* for  
4   dengue viruses appears to be determined by genotype-by-genotype interactions, whereby  
5   the outcome of infection depends on the specific combination of mosquito and virus  
6   genotypes. This may complicate efforts to dissect the genetic basis of vector competence  
7   in nature because it obscures mapping between genotype and phenotype and brings into  
8   question the notion of universal mosquito resistance or susceptibility. On the other hand,  
9   it offers novel opportunities to better define compatible vector-pathogen associations  
10   based on integration of both vector and pathogen genomics, which should eventually  
11   improve understanding of pathogen transmission dynamics and the risk of vector-borne  
12   disease emergence.

## 1   **Natural patterns of vector-pathogen specificity**

2

3   Elucidating the genetic determinants underlying patterns of compatibility between insect  
4   vectors and the pathogens they transmit is a major goal of vector biology because they  
5   largely contribute to vector competence (see Glossary) and thus to the public health threat  
6   represented by a given vector-pathogen pair [1]. It is striking, for instance, that parasite  
7   species causing human malaria are exclusively transmitted by *Anopheles* mosquitoes [2],  
8   whereas the vast majority of vectors of arthropod-borne viruses (arboviruses) belong to  
9   the *Culex* and *Aedes* mosquito genera [3]. *An. gambiae*, for example, is the major African  
10   vector of the deadliest human malaria parasite, *Plasmodium falciparum*, but is  
11   incompetent to most arboviruses, with the notable exception of O'nyong-nyong virus [2].  
12   Likewise, *Ae. aegypti* is an efficient vector of yellow fever, dengue, and chikungunya  
13   viruses but does not transmit *P. falciparum* [3]. Vector-pathogen specificity is not only  
14   observed at the genus level, but also at the species level. For example, *An. freeborni* is a  
15   competent vector of the rodent parasite *P. yoelii*, but does not support complete  
16   development of the primate parasite *P. knowlesi*. Conversely, *An. dirus* is a competent  
17   vector of *P. knowlesi*, but is naturally refractory to *P. yoelii* [4,5]. Genetic specificity of  
18   compatibility has also been shown at the intraspecific level between *An. gambiae* and *P.*  
19   *falciparum* [6,7]. Until recently, however, genetic specificity of compatibility at the  
20   intraspecific level had not been formally demonstrated between arboviruses and  
21   mosquitoes.

22

23

## 1    **G x G interactions between dengue viruses and *Aedes aegypti***

2

3    A recent study [8] provided empirical evidence for genotype-by-genotype (G x G)  
4    interactions between dengue viruses and their major vector species worldwide, *Ae.*  
5    *aegypti*. In a reciprocal cross-infection design of three field-derived *Ae. aegypti* pedigrees  
6    and three low-passage dengue-1 virus isolates, several vector competence indices were  
7    dependent on specific combinations of mosquito genotypes and virus isolates (Figure 1).  
8    Because critical environmental factors such as maternal effects and viral titer in blood  
9    meals were carefully controlled in the experimental design, interactions between  
10   mosquito pedigrees and virus isolates could be interpreted as evidence for G x G  
11   interactions. This study showed that individual variation in susceptibility to dengue  
12   viruses among *Ae. aegypti* mosquitoes derives, in part, from interactions between the  
13   mosquito's and the viral genomes. Such genetic interactions between two distinct  
14   organisms can be viewed as intergenomic epistasis [9].

15

16   The finding of G x G interactions between dengue viruses and *Ae. aegypti* is no surprise  
17   because many natural host-pathogen systems are governed by such specific interactions  
18   [7,10–15]. The occurrence of G x G interactions helps to explain the variation previously  
19   observed among *Ae. aegypti* populations in their vector competence for different dengue  
20   virus strains [16,17]. Although stochastic processes also play a role in dengue virus  
21   evolution [18], G x G interactions combined with the fine-scale genetic structure of *Ae.*  
22   *aegypti* populations [19–21] could lead to spatial structuring of viral populations through  
23   adaptation to their local vector populations [8,22].

## 1    **Relative nature of *Ae. aegypti* susceptibility to dengue viruses**

2

3    An important implication of G x G interactions between *Ae. aegypti* and dengue viruses is  
4    that they bring into question the concept of a resistant or susceptible vector genotype in  
5    natural systems. Indeed, no single *Ae. aegypti* pedigree was most resistant or susceptible  
6    to all three virus isolates, and reciprocally no single virus isolate was most successful in  
7    all mosquito pedigrees (Figure 1). Thus, a resistance allele or genotype against one virus  
8    strain may well confer susceptibility against another virus strain. This situation differs  
9    from plant-pathogen systems that are governed by typical gene-for-gene interactions. In  
10   these systems, one resistance allele in the plant host confers resistance to all pathogens  
11   expressing the corresponding ‘avirulence’ factor, whereas a pathogen ‘virulence’ allele  
12   confers universal infectivity [23–25]. Under the gene-for-gene model, a cost of infectivity  
13   to the pathogen is generally required to maintain genetic polymorphism [26–28]. G x G  
14   interactions detected thus far between *Ae. aegypti* and dengue viruses are more consistent  
15   with a matching-allele model of infection genetics, whereby successful infection requires  
16   an exact genetic match between multiple host and pathogen loci [29,30]. Under the  
17   matching-allele model, inspired from self/non-self recognition mechanisms in animals  
18   [31], universal infectivity is not possible and genetic polymorphism can be maintained by  
19   negative frequency-dependent selection [32,33]. In this context, the resistant or  
20   susceptible status of an allele or genotype is frequency-dependent and can only be  
21   inferred at the population level [34]. Strict coevolution, however, is unlikely to occur in  
22   the case of interactions between dengue viruses and *Ae. aegypti* because of the low

1 encounter rate and small fitness cost of infection that probably limit virus-driven  
2 selection of mosquitoes.  
3  
4 That the resistant or susceptible status of a mosquito depends on pathogen genetic  
5 identity will complicate efforts to unravel the genetic basis of *Ae. aegypti* vector  
6 competence for dengue viruses. Several quantitative trait loci (QTL) underlying variation  
7 in *Ae. aegypti* vector competence for dengue-2 virus have been identified [35–37]. The  
8 specific genes and polymorphisms involved are yet to be elucidated. The existence of  
9 significant G x G interactions, however, implies that conclusions from a particular  
10 combination of mapping mosquito population and virus strain will not necessarily apply  
11 in another pair. As was concluded by a meta-analysis of a large set of published QTL  
12 studies [38], it is likely that the genetic architecture of *Ae. aegypti* susceptibility to  
13 dengue viruses will differ among different combinations of vector and virus genotypes. In  
14 other words, sets of QTL and epistatic interactions that explain variation in vector  
15 competence will likely change among different combinations of mapping mosquito  
16 populations and/or virus strains.

### 19 **Investigating the genetic basis of vector-virus specificity**

20  
21 The pessimistic view that G x G interactions will represent a significant hurdle to  
22 elucidate the genetic basis of *Ae. aegypti* vector competence for dengue viruses is  
23 counter-balanced by the new perspectives that they offer. G x G interactions mean that

phenotypic variation in vector competence is influenced not only by the independent, additive effects of the vector and virus genotypes, but also by a genetic component that is specific to the particular vector-virus combination [9]. It is precisely the dissection of this specific component that represents a promising research avenue because it likely includes the genetic determinants of vector-virus specific compatibility, which is a major unresolved question in vector biology. Understanding the genetic basis of vector-pathogen specificity has important implications for the prevention of vector-borne disease, as illustrated by recent cases of vector-borne disease emergence associated with a change in vector-virus specificity. Examples include Venezuelan equine encephalitis [39], West Nile fever [40], and chikungunya [41,42]. Although one should not forget non-genetic contributions to variation in vector competence [43], including environmental influence on G x G interactions [44], genetic polymorphisms underlying vector-virus specificity may ultimately be used as markers to assess the risk of vector-borne disease emergence and to better understand virus transmission dynamics through time and space.

Merging vector and pathogen genetics requires a new approach that transcends the traditional disciplines of medical entomology, virology, parasitology, population and quantitative genetics [45]. This methodological shift shares a common goal with what is now called systems biology; i.e., to model and discover emergent properties of complex systems that cannot be predicted from examination of individual system components, but are essential for understanding the system as a whole [46]. The advent of the post-genomic era in vector biology has provided a wealth of genetic tools and resources [47],



1 which, together with theoretical developments in quantitative genetics [48], constitute a  
2 solid starting point for dissecting the genetic basis of complex traits associated with  
3 vector-pathogen interactions. Following the example of renewed appreciation for intra-  
4 genome epistasis to understand the structure and function of genetic pathways [49], there  
5 are unprecedented opportunities to study gene interactions between two organisms in a  
6 quantitative and comprehensive manner. Challenges ahead include the need to develop  
7 experimental and analytical tools that allow the functional effect of genetic  
8 polymorphisms in two (sometimes very different) genomes to be examined  
9 simultaneously. For instance, *Ae. aegypti* has a diploid DNA genome of an estimated 1.3  
10 Gb [50], whereas dengue viruses have a haploid, positive-strand RNA genome of less  
11 than 11 kb. Another daunting challenge is to elaborate ways to overcome the  
12 overwhelming number of possible genotype combinations. Genetic analysis of pairwise  
13 (or even third- or higher-order) gene interactions will need to be based on strong  
14 hypotheses generated from functional information. Systems modeling will help to narrow  
15 down the list of possible interactions to test empirically those that are predicted to be the  
16 most influential. Ultimately, the perspective will expand to include other interacting  
17 genomes, such as those of the microbiota and endosymbionts whose influence on vector  
18 competence is becoming increasingly apparent [51–54].

## 1   **Conclusions**

2

3   Evidence for G x G interactions between dengue viruses and *Ae. aegypti* calls for a more  
4   integrated view of the genetics underlying this vector-pathogen association. As for other  
5   phenotypes that derive from host-pathogen interactions, variation in vector competence  
6   appears to result, in part, from the interplay between two genomes. Thus, focusing on one  
7   genome without considering the other will inevitably lead to an incomplete picture of the  
8   system. The challenge of future research will be to tackle the shared, comprehensive  
9   nature of the genetic basis of vector competence. Development of high-throughput  
10   functional genomics and systems approaches to biology offer unprecedented  
11   opportunities to accomplish these tasks. This new paradigm in vector biology should  
12   eventually improve our ability to define the risk of vector-borne disease emergence and  
13   understand pathogen transmission dynamics.

14

## 1   **Acknowledgements**

2

3   I thank all the participants of the Keystone Meeting “Molecular Targets for Control of  
4   Vector-Borne Diseases: Bridging Lab and Field Research” held in April 2010 in Copper  
5   Mountain, Colorado, for stimulating discussions. I am grateful to Thomas W. Scott,  
6   Carole Escartin and three anonymous reviewers for helpful comments on an earlier  
7   version of the manuscript. I received support from the French Agence Nationale pour la  
8   Recherche (grant ANR-09-RPDOC-007-01).

## 1   **References**

2

- 3   1     Beerntsen, B.T. *et al.* (2000) Genetics of mosquito vector competence. *Microbiol.*  
4         *Mol. Biol. Rev.* 64, 115–137
- 5   2     Billingsley, P.F. and Sinden, R.E. (1997) Determinants of malaria-mosquito  
6         specificity. *Parasitol. Today* 13, 297–301
- 7   3     Leake, C.J. (1992). Arbovirus-mosquito interactions and vector specificity.  
8         *Parasitol. Today* 8, 123–128
- 9   4     Rosenberg, R. (1985) Inability of *Plasmodium knowlesi* sporozoites to invade  
10        *Anopheles freeborni* salivary glands. *Am. J. Trop. Med. Hyg.* 34, 687–691
- 11   5     Vaughan, J.A. *et al.* (1994) Sporogonic development of *Plasmodium yoelii* in five  
12        anopheline species. *J. Parasitol* 80, 674–681
- 13   6     Harris, C. *et al.* (2010) Polymorphisms in *Anopheles gambiae* immune genes  
14        associated with natural resistance to *Plasmodium falciparum*. *PLoS Pathog.* 6,  
15        e1001112
- 16   7     Lambrechts, L. *et al.* (2005) Host genotype by parasite genotype interactions  
17        underlying the resistance of anopheline mosquitoes to *Plasmodium falciparum*.  
18        *Malar. J.* 4, 3
- 19   8     Lambrechts, L. *et al.* (2009) Genetic specificity and potential for local adaptation  
20        between dengue viruses and mosquito vectors. *BMC Evol. Biol.* 9, 160
- 21   9     Lambrechts, L. (2010) Dissecting the genetic architecture of host-pathogen  
22        specificity. *PLoS Pathog.* 6, e1001019

1 10 Carius, H.J. *et al.* (2001) Genetic variation in a host-parasite association: potential  
2 for coevolution and frequency-dependent selection. *Evolution* 55, 1136–1145.

3 11 de Roode, J.C. and Altizer, S. (2009) Host-parasite genetic interactions and  
4 virulence-transmission relationships in natural populations of Monarch butterflies.  
5 *Evolution* 64, 502–514

6 12 Salvaudon, L. *et al.* (2007) Genotype-specific interactions and the trade-off  
7 between host and parasite fitness. *BMC Evol. Biol.* 7, 189

8 13 Schmid-Hempel, P. *et al.* (1999) Dynamic and genetic consequences of variation  
9 in horizontal transmission for a microparasitic infection. *Evolution* 53, 426–434

10 14 Schulenburg, H. and Ewbank, J.J. (2004) Diversity and specificity in the  
11 interaction between *Caenorhabditis elegans* and the pathogen *Serratia*  
12 *marcescens*. *BMC Evol. Biol.* 4, 49

13 15 Webster, J.P. and Woolhouse, M.E.J. (1998) Selection and strain specificity of  
14 compatibility between snail intermediate hosts and their parasitic schistosomes.  
15 *Evolution* 52, 1627–1634

16 16 Armstrong, P.M. and Rico-Hesse, R. (2001) Differential susceptibility of *Aedes*  
17 *aegypti* to infection by the American and Southeast Asian genotypes of dengue  
18 type 2 virus. *Vector Borne Zoonotic Dis.* 1, 159–168

19 17 Rosen, L. *et al.* (1985) Comparative susceptibility of mosquito species and strains  
20 to oral and parenteral infection with dengue and Japanese encephalitis viruses.  
21 *Am. J. Trop. Med. Hyg.* 34, 603–615

22 18 Holmes, E.C. and Twiddy, S.S. (2003) The origin, emergence and evolutionary  
23 genetics of dengue virus. *Infect. Genet. Evol.* 3, 19–28

1 19 Apostol, B.L. *et al.* (1996) Population genetics with RAPD-PCR markers: the  
2 breeding structure of *Aedes aegypti* in Puerto Rico. *Heredity* 76, 325–334

3 20 Bosio, C.F. *et al.* (2005) Genetic structure of *Aedes aegypti* populations in  
4 Thailand using mitochondrial DNA. *Am. J. Trop. Med. Hyg.* 72, 434–442

5 21 Hemme, R.R. *et al.* (2010) Influence of urban landscapes on population dynamics  
6 in a short-distance migrant mosquito: evidence for the dengue vector *Aedes*  
7 *aegypti*. *PLoS Negl. Trop. Dis.* 4, e634

8 22 Chevillon, C. and Failloux, A.B. (2003) Questions on viral population biology to  
9 complete dengue puzzle. *Trends Microbiol.* 11, 415–421

10 23 Flor, H.H. (1956) The complementary genetic systems in flax and flax rust. *Adv.*  
11 *Genet.* 8, 29–54

12 24 Flor, H.H. (1971) Current status of the gene-for-gene concept. *Annu. Rev.*  
13 *Phytopathol.* 9, 275–296

14 25 Thompson, J.N. and Burdon, J.J. (1992) Gene-for-gene coevolution between  
15 plants and parasites. *Nature* 360, 121–126

16 26 Frank, S.A. (1993) Coevolutionary genetics of plants and pathogens. *Evol. Ecol.*  
17 7, 45–75

18 27 Sasaki, A. (2000) Host-parasite coevolution in a multilocus gene-for-gene system.  
19 *Proc. Biol. Sci.* 267, 2183–2188

20 28 Segarra, J. (2005) Stable polymorphisms in a two-locus gene-for-gene system.  
21 *Phytopathology* 95, 728–736

22 29 Agrawal, A. and Lively, C.M. (2002) Infection genetics: gene-for-gene versus  
23 matching-alleles models and all points in between. *Evol. Ecol. Res.* 4, 79–90

1 30 Frank, S.A. (1993) Specificity versus detectable polymorphism in host-parasite  
2 genetics. *Proc. Biol. Sci.* 254, 191–197

3 31 Grosberg, R.K. and Hart, M.W. (2000) Mate selection and the evolution of highly  
4 polymorphic self/nonself recognition genes. *Science* 289, 2111–2114

5 32 Hamilton, W.D. (1980) Sex versus non-sex versus parasite. *Oikos* 35, 282–290

6 33 Hamilton, W.D. *et al.* (1990) Sexual reproduction as an adaptation to resist  
7 parasites (a review). *Proc. Natl. Acad. Sci. U S A* 87, 3566–3573

8 34 Theron, A. and Coustau, C. (2005) Are *Biomphalaria* snails resistant to  
9 *Schistosoma mansoni*? *J. Helminthol.* 79, 187–191

10 35 Bennett, K.E. *et al.* (2005) Quantitative trait loci that control dengue-2 virus  
11 dissemination in the mosquito *Aedes aegypti*. *Genetics* 170, 185–194

12 36 Bosio, C.F. *et al.* (2000) Quantitative trait loci that control vector competence for  
13 dengue-2 virus in the mosquito *Aedes aegypti*. *Genetics* 156, 687–698

14 37 Gomez-Machorro, C. *et al.* (2004) Quantitative trait loci affecting dengue midgut  
15 infection barriers in an advanced intercross line of *Aedes aegypti*. *Insect. Mol.*  
16 *Biol.* 13, 637–648

17 38 Wilfert, L. and Schmid-Hempel, P. (2008) The genetic architecture of  
18 susceptibility to parasites. *BMC Evol. Biol.* 8, 187

19 39 Brault, A.C. *et al.* (2004) Venezuelan equine encephalitis emergence: enhanced  
20 vector infection from a single amino acid substitution in the envelope  
21 glycoprotein. *Proc. Natl. Acad. Sci. U S A* 101, 11344–11349

1 40 Moudy, R.M. *et al.* (2007) A newly emergent genotype of West Nile virus is  
2 transmitted earlier and more efficiently by *Culex* mosquitoes. *Am. J. Trop. Med.*  
3 *Hyg.* 77, 365–370

4 41 Tsetsarkin, K.A. *et al.* (2007) A single mutation in chikungunya virus affects  
5 vector specificity and epidemic potential. *PLoS Pathog.* 3, e201

6 42 Vazeille, M. *et al.* (2007) Two Chikungunya isolates from the outbreak of La  
7 Reunion (Indian Ocean) exhibit different patterns of infection in the mosquito,  
8 *Aedes albopictus*. *PLoS ONE* 2, e1168

9 43 Black, W.C. *et al.* (2002) Flavivirus susceptibility in *Aedes aegypti*. *Arch. Med.*  
10 *Res.* 33, 379–388

11 44 Thomas, M.B. and Blanford, S. (2003) Thermal biology in insect-parasite  
12 interactions. *Trends Ecol. Evol.* 18, 344–350

13 45 Lambrechts, L. *et al.* (2009) Shifting priorities in vector biology to improve  
14 control of vector-borne disease. *Trop. Med. Int. Health.* 14, 1505–1514

15 46 Schneider, D.J. and Collmer, A. (2010) Studying plant-pathogen interactions in  
16 the genomics era: beyond molecular Koch's postulates to systems biology. *Annu.*  
17 *Rev. Phytopathol.* 48, 457–479

18 47 Lawson, D. *et al.* (2009) VectorBase: a data resource for invertebrate vector  
19 genomics. *Nucleic Acids Res.* 37, D583–587

20 48 Yang, J. *et al.* (2008) Mapping interspecific genetic architecture in a host-parasite  
21 interaction system. *Genetics* 178, 1737–1743

22 49 Phillips, P.C. (2008) Epistasis--the essential role of gene interactions in the  
23 structure and evolution of genetic systems. *Nat. Rev. Genet.* 9, 855–867



1    50    Nene, V. *et al.* (2007) Genome sequence of *Aedes aegypti*, a major arbovirus  
2        vector. *Science* 316, 1718–1723

3    51    Bian, G. *et al.* (2010) The endosymbiotic bacterium *Wolbachia* induces resistance  
4        to dengue virus in *Aedes aegypti*. *PLoS Pathog.* 6, e1000833

5    52    Dong, Y. *et al.* (2009) Implication of the mosquito midgut microbiota in the  
6        defense against malaria parasites. *PLoS Pathog.* 5, e1000423

7    53    Moreira, L.A. *et al.* (2009) A *Wolbachia* symbiont in *Aedes aegypti* limits  
8        infection with dengue, Chikungunya, and *Plasmodium*. *Cell* 139, 1268–1278

9    54    Xi, Z. *et al.* (2008) The *Aedes aegypti* toll pathway controls dengue virus  
10       infection. *PLoS Pathog.* 4, e1000098

11

12

## 1    **Glossary**

2

3    **Epistasis:** Complex genetic interactions arising when the effects of alleles at one locus  
4    depend on the presence of a specific allele at another locus.

5

6    **Genetic architecture:** Number, effect, location, and interactions of genes underlying  
7    phenotypic variation.

8

9    **Genotype-by-genotype (G x G) interaction:** In a host-pathogen system, describes the  
10    intrinsic specificity of compatibility among host and pathogen genotypes. It can be  
11    measured as the statistical deviation from the additive combination of host and pathogen  
12    genotypes in their effects on the infection phenotype.

13

14    **Quantitative trait locus (QTL):** Genomic region that explains part of the phenotypic  
15    variation in a continuous trait.

16

17    **Vector competence:** Intrinsic ability of an arthropod to become infected, allow  
18    replication, and ultimately transmit a pathogen. It is genetically determined but also  
19    influenced by environmental factors.

20


21

1 **Figure Legend**

2

3 **Figure 1.** Genotype-by-genotype interactions between dengue viruses and *Aedes aegypti*.

4 The matrices show experimental measurements of three different vector competence  
5 indices for all pairwise combinations of three dengue virus isolates and three mosquito  
6 pedigrees. Phenotypic values are converted to shades of grey according to the scale bar  
7 indicated below. Each vector competence index depends significantly on the specific  
8 combination of vector genotype and virus isolate. Modified, with permission, from Ref.  
9 [8].



10