



Durability of immunogenicity and strain coverage of MenBvac, a meningococcal vaccine based on outer membrane vesicles: Lessons of the Normandy campaign

François Caron, Julien Sevestre, Eva Hong, Valérie Delbos, Aude Terrade, Eric Mallet, Ala-Eddine Deghmane, Ludovic Lemée, Muhamed-Kheir Taha, François Caron

► To cite this version:

François Caron, Julien Sevestre, Eva Hong, Valérie Delbos, Aude Terrade, et al.. Durability of immunogenicity and strain coverage of MenBvac, a meningococcal vaccine based on outer membrane vesicles: Lessons of the Normandy campaign. *Vaccine*, 2017, 35 (32), pp.4029-4033. 10.1016/j.vaccine.2017.05.065 . pasteur-02013259

HAL Id: pasteur-02013259

<https://pasteur.hal.science/pasteur-02013259>

Submitted on 30 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Durability of immunogenicity and strain coverage of MenBvac, a meningococcal vaccine based on outer membrane vesicles: lessons of the Normandy campaign

Julien Sevestre^{a,b,c,1}, Eva Hong^{c,1}, Valérie Delbos^{a,b}, Aude Terrade^c, Eric Mallet^{a,b}, Ala-Eddine Deghmane^c, Ludovic Lemée^{a,b}, Muhamed-Kheir Taha^{c,2}, François Caron^{a,b,2}

^aCentre Hospitalier Universitaire, Rouen, France

^bUniversité de Rouen, France

^cInstitut Pasteur, Paris, France

¹Both authors contributed equally to this work.

²Both authors contributed equally to this work as senior authors

Corresponding author Muhamed-Kheir Taha

Email: mktaha@pasteur.fr

Running title: Response of OMV-containing meningococcal vaccines

Durability of immunogenicity and strain coverage of MenBvac, a meningococcal vaccine
based on outer membrane vesicles: lessons of the Normandy campaign

Julien Sevestre^{a,b,c,1}, Eva Hong^{c,1}, Valérie Delbos^{a,b}, Aude Terrade^c, Eric Mallet^{a,b}, Ala-Eddine
Deghmane^c, Ludovic Lemée^{a,b}, Muhamed-Kheir Taha^{c,2}, François Caron^{a,b,2}

^aCentre Hospitalier Universitaire, Rouen, France

^bUniversité de Rouen, France

^cInstitut Pasteur, Paris, France

¹Both authors contributed equally to this work.

²Both authors contributed equally to this work as senior authors

Corresponding author Muhamed-Kheir Taha

Email: mktaha@pasteur.fr

Running title: Response of OMV-containing meningococcal vaccines

Abstract

Objectives: MenBvac® is an outer membrane vesicle (OMV)-based meningococcal vaccine. From 2006 to 2012, it was used to control a clonal B outbreak in Normandy (France). We aimed to analyse the durability of the response against the epidemic strain and coverage beyond the vaccine strain. These data should help to optimize the use of OMV-containing vaccines, such as the new 4CMenB/Bexsero® recombinant vaccine.

Methods: Serum bactericidal activity (SBA) was measured in two cohorts of children who received their first dose of MenBvac® at 1-5 years of age and accepted to provide a blood sample either one or four years after a 2+1+1 schedule. All sera were tested against the outbreak strain. Sera from responder subjects were also tested against 12 additional B or C strains which were chosen to entirely, partially, or not at all match the two variable regions (VR1 and VR2) of the PorA vaccine strain.

Results: Only 47.9% and 31.3% of subjects showed an SBA titre consistent with protection one and four years, respectively, after the last boost. Protective SBA titres were observed in all sera against B or C strains that entirely matched P1.7,16, and was high (75-100%) for all but one strain that partially matched VR1 or VR2. Extrapolating our data to the OMV component of 4CMenB/Bexsero® suggests that 14.5% of the current B strains would be covered based on PorA matching to the OMV component of 4CMenB/Bexsero® (regardless of the coverage of the three other vaccine components).

Conclusions: Our data confirm that OMV-based vaccines elicit short-lasting SBA titres and may require repeated booster injections. However, strain coverage may be greater than expected.

Keywords: *Neisseria meningitidis* serogroup B; serum bactericidal activity; outbreak; persistence of vaccine response; invasive meningococcal disease; PorA; vaccination

The Authors declare no conflict of interest.

Highlights

- Meningococcal outbreaks can be controlled by OMV-based vaccines, such as MenBvac
- Serum bactericidal activity of sera from vaccinated subjects was evaluated
- Serum bactericidal activity rapidly declined
- Meningococcal strains expressing partially matching PorA were covered
- OMV-containing vaccines may offer broader protection than originally expected.

Introduction

Capsular polysaccharide vaccines have been successfully developed against *Neisseria meningitidis* of serogroup A, C, W, and Y. Such an approach is not possible for serogroup B strains due to the homology between the group B capsule and the human neural cellular adhesion molecule, leading to poor immunogenicity and a potential risk of autoimmunity. This is particularly problematical because most invasive meningococcal disease in developed countries is caused by serogroup B strains. Since the 1980s, tailor-made meningococcal B vaccines, based on outer membrane vesicles (OMV), have been developed in response to outbreaks, with campaigns in South Africa, Cuba, Norway, Brazil, Chile, and New Zealand [1]. OMV vaccines elicit an immune response that is mainly directed against the protein porin A (PorA) of the outer membrane which includes two variable regions, VR1 and VR2, with a large diversity for both of them. The response is [generally](#) “strain specific” and is not cross-reactive with divergent PorA variants. The last application of an OMV alone-based meningococcal B vaccine was reported in Normandy, France, from 2006 to 2012, where a clonal meningococcal outbreak, due to a strain of serotype 14, sero-subtype P1.7,16 and clonal complex 32 (B:14:P1.7,16/cc32), was controlled using MenBvac®, a Norwegian vaccine designed in the 1980s against a strain expressing the same PorA variant (B:15:P1.7,16/cc32) [2].

Since then, multi-component protein-based vaccines that offer broad coverage against group B *N. meningitidis* have been developed. Such new vaccines do not necessarily make former OMV meningococcal vaccines obsolete [3]. The new multi-component protein-based vaccines do not cover all circulating serogroup B isolates. Indeed, 4CMenB/Bexsero®, the only meningococcal B vaccine available in Europe, [has been first evaluated to](#) cover 73–87% of the B strains circulating in different European countries [4]. [More recent data have predicted a 66% coverage in the UK](#) [5]. These data imply that a tailor-made OMV vaccine

might be requested in the future to control an outbreak due to an uncovered strain. Moreover, OMV have adjuvant properties [6]. They have therefore been included in 4CMenB/Bexsero® (using the preparation developed for MenZB, the OMV vaccine used in New Zealand) to increase both the immunogenicity and persistence of the immune response against the other vaccine components. Also, OMV vaccines may provide cross protection against strains that do not express perfectly matching PorA proteins [7-9]. Importantly, antibodies generated by OMV are independent of those targeting the capsule; thus OMV should work against meningococcal strains from different serogroups, providing that they partially or entirely share the PorA of the vaccine strain. Finally, animal experimentation, as well as several clinical studies, suggest that the Norwegian OMV-based vaccine reduces meningococcal carriage and may hence confer herd protection [10, 11].

During the Normandy outbreak, we performed two immunogenicity studies [11, 12] to evaluate different schedules of MenBvac®. Several volunteers accepted to provide blood five years after the first dose to evaluate the durability of the response and the spectrum of coverage by measuring serum bactericidal titres (SBA), not only against the epidemic strain, but also against other B and non-B strains which are currently endemic in the country.

Methods

Serum bactericidal activity

SBA assays were performed using human serum, devoid of meningococcal lysis activity, as a complement source (hSBA) for serogroup B isolates. Baby rabbit complement (rSBA) was used for serogroup C isolates as previously described [2, 12]. The serum bactericidal activity titres corresponded to the inverse of the final serum dilution causing 50% killing of the inoculum after 1 h of incubation. A protective immune response was defined by an hSBA titre ≥ 4 or a rSBA titre ≥ 8 , as these titres have been suggested as a surrogate for protection [13, 14].

Titres were compared using the percentage of subjects achieving the threshold for protection and by calculating the geometric means of titres (GMT). Data were analyzed using the Chi-squared test, Student's *t*-test, and analysis of variance (ANOVA). A *P* value of < 0.05 was considered to be statistically significant. The Bonferroni correction was applied when the Chi-squared test involved several comparisons.

Durability of immunogenicity against the targeted epidemic strain

The SBA titres against the epidemic clone were analyzed for the two previously assembled cohorts, the first in 2006 from subjects living in the Dieppe area and the second in 2009 from subjects living in the Neufchâtel-en-Bray area [2, 12]. Briefly, children were aged 1-5 years at the time of the first MenBvac® dose. They were selected by random sampling from all subjects of the same age-range of the area eligible to receive MenBvac®. Both cohorts had received a 2+1+1 schedule. The time of administration of the second boost (i.e., the last dose) differed slightly between the two cohorts due to a production shortage of the vaccine that required several adaptations of the initially planned scheme [2]. Thus, subjects received MenBvac® at D0, W6, M8, and M23 in the Dieppe area, and at D0, W6, M8, and M36 in the

Neufchâtel-en-Bray area. Previously enrolled subjects were asked to provide a blood sample four years after the last boost (i.e., at M71) for the Dieppe area [2] and one year after the last boost (i.e., at M50) for the Neufchâtel-en-Bray area [12]. The extension of the former study was approved for both cohorts by the regional ethics committee (Comité de Protection des Personnes Nord-Ouest-1) and written informed consent was obtained from parents or legal guardians of every participant.

Strain coverage

We evaluated the range of strain coverage by MenBvac® by testing a subset of frozen serum samples (collected six weeks after the second boost) of 12 subjects from the Neufchâtel-en-Bray area cohort. The sera came from subjects who responded against strain LNP21362 (B:14:P1.7,16/cc32), which was previously used as a reference strain for SBA assays for the Normandy outbreak [2]. These subjects did not show protective SBA titres before vaccination (D0). Sera were tested against the reference strain (LNP21362), as well as 13 other strains of serogroup B or C harbouring different PorA variable regions (14 SBA tests being the maximal possible due to the low volume of residual serum available for each of these sera). The other 13 strains were selected from B- and C-subgroup strains analysed in 2015 by the National Reference Centre (NRC) for meningococci. In France, almost all cases of invasive meningococcal disease are biologically confirmed by culture and/or PCR at the NRC. The NRC performs complete genotyping of all strains by multi-sequence locus typing (clonal complex) of PorA and FetA [15]. The 13 strains were selected to represent the different combinations of variable regions, VR1 and VR2, partially matching PorA P1.7,16, that are currently circulating in the country. When the same PorA profile was shared by several strains belonging to different clonal complexes, one strain per clonal complex was selected. Human complement was used for the SBA assay of serogroup B isolates. Baby rabbit complement

was used for serogroup C isolates due to the intrinsic meningococcal lysis against these serogroup C strains in all human sera that were used as complement sources.

Finally, the results were extrapolated to the 2015 NRC data to determine the percentage of invasive meningococcal strains currently endemic in the country that completely or partially match the OMV component of MenBvac® (P1.7,16) or 4CMenB/Bexsero® (P1.7-2, 4).

Results

Durability of immunogenicity against the targeted epidemic strain

Table 1 shows the long-term SBA titres for the two cohorts (last line) and compares them to the results previously reported for each scheme [2, 11].

For the Dieppe area cohort, 115 children among the 243 initially included (47%) agreed to participate in the final phase of the study four years after the second boost, amongst which only 31.3% (95% CI 22.8 to 39.8) had an hSBA titre consistent with protection, whereas the geometric mean hSBA was only 3.3 (95% CI 2.8-3.9). These data are in accordance with the marked decline of hSBA titres previously observed 15 months after the first boost, for which only 56% of subjects showed protective titres.

For the Neufchâtel-en-Bray area cohort, 96 children among the 213 initially included (45%) agreed to participate in the final phase of the study one year after the second boost, amongst which only 47.9% (95% CI 37.9 to 57.9) showed protective hSBA titres. This percentage, as well as the geometric mean hSBA value (4.4; 95% CI 3.5-5.5), were not significantly different from those previously achieved one year after the first boost (i.e., at M20).

Strain coverage

The characteristics of the 14 strains studied and the hSBA or rSBA titres achieved against each of them in the sera from 12 (10 for serogroup C due to a limited amount of sera) high responders to MenBvac are presented Table 2.

Among the 14 strains, two (one of serogroup B used as a control and one of serogroup C) completely matched the PorA of MenBvac (P1.7,16) and both belonged to cc32. Nine strains (seven of serogroup B and two of serogroup C) shared either VR1 or VR2 with the vaccine strain or were variants of the P1.7 (VR1) or P1.16 (VR2) family. The last three serogroup B

222 strains had PorA which completely differed from that of the strain vaccine (P1.22, 9, P1.19-2,
223 13-1 and P1.19-1, 15-11). Ten of the 14 strains (71%) belonged to major hyperinvasive clonal
224 complexes (cc11, cc32, cc41/44, cc269).

225 As expected, hSBA titres against the serogroup B isolate (LNP21362) that matched the
226 vaccine strain (P1.7,16) were significantly higher than the threshold of 4 for all 12 sera tested,
227 with a lower limit of the 95% CI > 4.

228 In contrast, the lower limit of the 95% CI of the GMT was lower than the threshold of 4 for
229 the SBA titres against the three isolates (LNP27143, LNP27050 and LNP27000) that did not
230 at all match PorA. One of these strains (LNP27000) did not show coverage in any tested
231 serum. However, 25% and 42% of the volunteers had hSBA titres of 4 and 8 against the other
232 two strains LNP27143 and LNP27050, respectively. Additionally, strain LNP27200, a non-
233 cc32 strain with a partial PorA match (P.21,16-36), also showed a low GMT, with 25% of
234 tested sera displaying hSBA titres of 4.

235 The GMT and the lower limits of the 95% CI were above 4 for the remaining six B strains.
236 The SBA titres were high in all 12 tested sera for three of the six strains (LNP27338,
237 LNP27087 and LNP27386 all belonged to the cc32) and were not significantly different from
238 those achieved against the reference strain (LNP21362). The hSBA titres differed greatly
239 (depending on the strain) for the three remaining strains (LNP27372, LNP27010 and
240 LNP27114). However, the lower limit of the 95% CI remained higher than 4, although the
241 GMT were significantly lower than those obtained against the reference strain LNP21362.

242 We also studied three strains of serogroup C, amongst which one entirely matched PorA of
243 MenBvac (LNP25514) and two partially matched (LNP25553 and LNP26251). The rSBA
244 threshold of 8 was achieved against the three C strains in all tested sera. A hierarchy of

responses was however observed, with the rSBA titres being significantly higher against the strain which entirely matched the vaccine PorA than the other two serogroup C strains.

The percentages of invasive meningococcal strains predicted to be covered by either MenBvac® or 4CMenB/Bexsero® based on their PorA variable regions, and regardless their serogroups, are shown in Table 3. For MenBvac®, only 2.6% of the current French strains totally matched PorA. This putative coverage increased to 16.9% when considering both complete and partially matching (i.e. strains with family 7 VR1 and/or strains with family 16 VR2). For 4CMenB/Bexsero®, the respective percentages were 8.1% and 14.5%.

Discussion

This study was performed in an area of Normandy recently affected by a clonal meningococcal B outbreak that was controlled using MenBvac, an OMV vaccine [2]. The two major results of this study were: (i) the short durability of protection against the targeted strain, despite a four dose vaccination scheme and (ii) an extension of the strain coverage to other B and non-B strains that at least partially shared the PorA of the vaccine strain.

The short persistence of protection was observed even after a second booster dose, although the absence of samples from six weeks after the second booster dose in the Dieppe cohort limits the extrapolation of this conclusion. Several previous studies on OMV vaccines have also reported the rapid decline of SBA titres elicited against the targeted strain [16]. This is not necessarily a major drawback, as such “tailor-made” vaccines are aimed to control an outbreak rather than to offer persistent protection against sporadic events. For the control of a clonal outbreak, two doses of a targeted OMV vaccine given at a 4-6 week interval can be sufficient to “put the fire out” as reported during the first application of MenBvac® in Norway [17]. However, because clonal B outbreaks might naturally persist for up to two decades, obtaining immunity for several years in an affected area is probably pertinent. The four-dose MenBvac® regimen applied during a two-year period in Normandy resulted in limited persistence of the SBA titres against the outbreak strain (B:14:P1.7,16/cc32) which was observed in less than one third of vaccinated 1-5 year-old children 5 years after the first dose. Such poor results do not necessarily exclude a better response in adolescents and young adults, as shown for other OMV vaccines [18]. The short observed persistence of bactericidal antibodies against B:14:P1.7,16/cc32 (even in the Dieppe area, epicentre of the outbreak) may be due to the rapid decline of the response to the vaccine and the lack of natural immunity favoured by a very low rate of pharyngeal carriage of the epidemic strain in the population, as reported by a carriage study performed at the height of the outbreak [10].

295 The broader strain coverage of MenBvac® is relevant in meningococcal vaccinology as the
296 4CMenB/Bexsero® vaccine includes an OMV component. OMV-based vaccines elicit an
297 immune response that is mainly directed against the PorA outer membrane protein. Due to the
298 high variability of PorA among isolates, this type of vaccine is classically considered strain-
299 specific [19]. However, several previous studies with MenBvac® [8, 9, 18, 20] and other
300 OMV vaccines [7, 9, 18, 20] have reported protection beyond the vaccine strain. Rosenqvist
301 *et al.* have shown that MenBvac® induced in adult volunteers significant SBA response
302 against variants of the vaccine strain lacking some outer membrane proteins [8]. The
303 Norwegian Institute of Public Health team had performed a study in adult volunteers
304 comparing the immunogenicity of MenBvac® (P1.7,16 strain) and MeNZB® (P1.7-2,4
305 strain) showing that SBA titres were better against the homologous strain than with the
306 heterologous strain [9, 20]. Tappero *et al* have reported that both MenBvac (P1.7,16 strain)
307 and the Finlay Institute OMV vaccine (P1.15 strain) did not induce significant response in
308 infants while children and adults showed SBA titres against a heterologous strain but to a
309 lesser extent than to the vaccine homologous strain[18]. Boutriau *et al* have analyzed a
310 bivalent OMV vaccine designed to cover a large proportion of B strains circulating in Europe
311 that induced increased SBA titres against the two vaccine strains as well as against three
312 heterologous strains but this vaccine did not reach the market [7]. Our study provides
313 additional data, suggesting that strain coverage can be larger than initially expected.
314 Completely or partially matching one of the two variable regions of PorA (VR1 and VR2)
315 may lead to coverage by the MenBvac® irrespective of the serogroup. However, SBA titres
316 were significantly higher when both VR1 and VR2 matched the vaccine, suggesting
317 synergism of serum bactericidal activity against these variable regions. Further studies may be
318 needed to explore how the alterations of VR regions affect serum bactericidal activity.
319 Moreover, other minor proteins in the OMV can confer protection, as suggested by the SBA

320 titres against the strain LNP27050 that did not match variable regions of PorA P1.7,16. This
321 part of the study had however three limitations. (i) Due to low volumes of residual serum
322 available, it was impossible for a given strain to test all subvariants in terms of VRs; all sub-
323 variants of P1.7,16 may not be covered and testing several other strains may be required. (ii)
324 It was also impossible to evaluate the protection at baseline against the tested strains.
325 However, we have previously reported low level of protective SBA titres against the stain
326 with P1.7,16 before the first dose of MenBvac®) [12].
327 (iii) The broader of coverage has been studied only for MenBvac®; it is not excluded that
328 results may differ for other OMV based vaccines.

329 The strain coverage by 4CMenB/Bexsero® has been extensively studied for the three proteins
330 NadA, fHbp, and NHBA, but much less for its fourth component, corresponding to PorA
331 P1.7-2, 4 from MeNZB® [21]. Previous reports have suggested that the VR1 region (7-2) of
332 PorA was poorly recognised by a monoclonal antibody against subtype P1.7 due to an internal
333 deletion, leading to masking of this region [22]. However, our study shows that polyclonal
334 anti-PorA antibodies, obtained after vaccination in several subjects, achieved SBA titres
335 against the two strains (LNP27010 and LNP27114) that only had VR1 7-2. The potential
336 protection against non-B strains, as suggested by our data, is of interest, as this is expected to
337 enlarge strain converge. Such protection was suggested in reports on the use of 4CmenB®
338 against serogroups W and X [23, 24]. The subjects of our study were not vaccinated against
339 serogroup C meningococci when MenBvac® vaccination started. Indeed, the
340 recommendation for vaccination against MenC only started in France in 2010. The uptake of
341 the vaccine remains very low in France and vaccine coverage in 1-4 year olds was only 50%
342 by the end of 2013 [25].

343 Based on the 2015 database from the NRC, 14.5% of the current serogroup B strains are
344 predicted to be covered by the OMV component of 4CMenB/Bexsero® (regardless of the

coverage by the three other components of the vaccine). The Meningococcal Antigen Typing system (MATS) predicted 85% overage of the serogroup B isolates in France. Importantly, among the 15% (n = 30) of uncovered strains, 13% (n = 4) could be covered on the basis of partial matching with PorA [4]. Thus, strain coverage could be higher than that predicted solely by MATS.

In conclusion, the results of the current study show that OMV-containing vaccines induce immunity, although of short duration, far below that of the vaccine strain. This argues for adapting the OMV component from new meningococcal vaccines to improve the coverage of local strains, particularly in the event of outbreaks that may not be covered by the other components of the vaccines.

Acknowledgements

This work was supported the Institut Pasteur and Rouen University Hospital. The Authors declare no conflicts of interest. JS receives a fellowship from the Conseil Régional de Haute Normandie. We thank Marie-Noelle Ungeheuer (Institut Pasteur-ICAReB) for supply of human serum samples for complement sources

References

- [1] Sadarangani M, Pollard AJ. Serogroup B meningococcal vaccines-an unfinished story. *Lancet Infect Dis.* 2010;10:112-24.
- [2] Caron F, du Chatelet IP, Leroy JP, Ruckly C, Blanchard M, Bohic N, et al. From tailor-made to ready-to-wear meningococcal B vaccines: longitudinal study of a clonal meningococcal B outbreak. *Lancet Infect Dis.* 2011;11:455-63.
- [3] Stephens DS. Outer-membrane-vesicle vaccines: old but not forgotten. *Lancet Infect Dis.* 2011;11:421-2.
- [4] Vogel U, Taha MK, Vazquez JA, Findlow J, Claus H, Stefanelli P, et al. Predicted strain coverage of a meningococcal multicomponent vaccine (4CMenB) in Europe: a qualitative and quantitative assessment. *Lancet Infect Dis.* 2013;13:416-25.
- [5] Medini D, Stella M, Wassil J. MATS: Global coverage estimates for 4CMenB, a novel multicomponent meningococcal B vaccine. *Vaccine.* 2015;33:2629-36.
- [6] Nagaputra JC, Rollier CS, Sadarangani M, Hoe JC, Mehta OH, Norheim G, et al. *Neisseria meningitidis* native outer membrane vesicles containing different lipopolysaccharide glycoforms as adjuvants for meningococcal and nonmeningococcal antigens. *Clin Vaccine Immunol.* 2014;21:234-42.
- [7] Boutriau D, Poolman J, Borrow R, Findlow J, Domingo JD, Puig-Barbera J, et al. Immunogenicity and safety of three doses of a bivalent (B:4:p1.19,15 and B:4:p1.7-2,4) meningococcal outer membrane vesicle vaccine in healthy adolescents. *Clin Vaccine Immunol.* 2007;14:65-73.
- [8] Rosenqvist E, Hoiby EA, Wedege E, Bryn K, Kolberg J, Klem A, et al. Human antibody responses to meningococcal outer membrane antigens after three doses of the Norwegian group B meningococcal vaccine. *Infect Immun.* 1995;63:4642-52.

407 [9] Wedege E, Bolstad K, Aase A, Herstad TK, McCallum L, Rosenqvist E, et al. Functional
408 and specific antibody responses in adult volunteers in new zealand who were given one of
409 two different meningococcal serogroup B outer membrane vesicle vaccines. Clin Vaccine
410 Immunol. 2007;14:830-8.

411 [10] Delbos V, Lemee L, Benichou J, Berthelot G, Taha MK, Caron F. Meningococcal
412 carriage during a clonal meningococcal B outbreak in France. Eur J Clin Microbiol Infect Dis.
413 2013;32:1451-9.

414 [11] Delbos V, Lemee L, Benichou J, Berthelot G, Deghmane AE, Leroy JP, et al. Impact of
415 MenBvac, an outer membrane vesicle (OMV) vaccine, on the meningococcal carriage.
416 Vaccine. 2013;31:4416-20.

417 [12] Caron F, Delbos V, Houivet E, Deghmane AE, Leroy JP, Hong E, et al. Evolution of
418 immune response against *Neisseria meningitidis* B:14:P1.7,16 before and after the outer
419 membrane vesicle vaccine MenBvac. Vaccine. 2012;30:5059–62.

420 [13] McIntosh ED, Broker M, Wassil J, Welsch JA, Borrow R. Serum bactericidal antibody
421 assays - The role of complement in infection and immunity. Vaccine. 2015;33:4414-21.

422 [14] Borrow R, Balmer P, Miller E. Meningococcal surrogates of protection--serum
423 bactericidal antibody activity. Vaccine. 2005;23:2222-7.

424 [15] Jolley KA, Chan MS, Maiden MC. mlstdbNet - distributed multi-locus sequence typing
425 (MLST) databases. BMC Bioinformatics. 2004;5:86.

426 [16] McQuaid F, Snape MD, John TM, Kelly S, Robinson H, Yu LM, et al. Persistence of
427 specific bactericidal antibodies at 5 years of age after vaccination against serogroup B
428 meningococcus in infancy and at 40 months. CMAJ. 2015;187:E215-23.

429 [17] Bjune G, Hoiby EA, Gronnesby JK, Arnesen O, Fredriksen JH, Halstensen A, et al.
430 Effect of outer membrane vesicle vaccine against group B meningococcal disease in Norway.
431 Lancet. 1991;338:1093-6.

- [18] Tappero JW, Lagos R, Ballesteros AM, Plikaytis B, Williams D, Dykes J, et al. Immunogenicity of 2 serogroup B outer-membrane protein meningococcal vaccines: a randomized controlled trial in Chile. *Jama*. 1999;281:1520-7.
- [19] Holst J, Feiring B, Naess LM, Norheim G, Kristiansen P, Hoiby EA, et al. The concept of "tailor-made", protein-based, outer membrane vesicle vaccines against meningococcal disease. *Vaccine*. 2005;23:2202-5.
- [20] Thornton V, Lennon D, Rasanathan K, O'Hallahan J, Oster P, Stewart J, et al. Safety and immunogenicity of New Zealand strain meningococcal serogroup B OMV vaccine in healthy adults: beginning of epidemic control. *Vaccine*. 2006;24:1395-400.
- [21] Sexton K, Lennon D, Oster P, Crengle S, Martin D, Mulholland K, et al. The New Zealand Meningococcal Vaccine Strategy: a tailor-made vaccine to combat a devastating epidemic. *N Z Med J*. 2004;117:U1015.
- [22] Wedege E, Dalseg R, Caugant DA, Poolman JT, Froholm LO. Expression of an inaccessible P1.7 subtype epitope on meningococcal class 1 proteins. *J Med Microbiol*. 1993;38:23-8.
- [23] Ladhani SN, Giuliani MM, Biolchi A, Pizza M, Beebeejaun K, Lucidarme J, et al. Effectiveness of Meningococcal B Vaccine against Endemic Hypervirulent *Neisseria meningitidis* W Strain, England. *Emerg Infect Dis*. 2016;22:309-11.
- [24] Hong E, Giuliani MM, Deghmane AE, Comanducci M, Brunelli B, Dull P, et al. Could the multicomponent meningococcal serogroup B vaccine (4CMenB) control *Neisseria meningitidis* capsular group X outbreaks in Africa? *Vaccine*. 2013;31:1113-6.
- [25] Barret A-S, Parent du Châtelet I, Deghmane A-E, Lepoutre A, Fonteneau L, Maine C, et al. Invasive meningococcal disease in France 2012. *Bull Epidemiol Hbdo*. 2014:25-31.

Table 1
Immunogenicity of MenBvac® vaccine and persistence of the response against B:14:P1.7,16 (strain LNP21362) in two cohorts of children of 1-5 years of age at first dose and previously analyzed at short term.

Dieppe cohort MenBvac at D0-W6-M8-M23				Neufchâtel-en-Bray cohort MenBvac at D0-W6-M8-M36			
Time of sampling	Number	% of subjects with hSBA ≥ 4 [95% CI]	GMT of hSBA titres [95% CI]	Time of sampling	Number	% of subjects with hSBA ≥ 4 [95% CI]	GMT of hSBA titers [95% CI]
-	-	-	-	D0*	213	10.8 (6.6-15)	2.3 (2.1-2.5)
-	-	-	-	M3	172	41.3 (33.9-48.7)	3.3 (2.9-3.7)
M8*	243	37.0 (30.9-39.8)	2.8 (2.6-2.9)	M8*	159	25.5 (19-32.6)	2.8 (2.5-3.1)
M9.5	235	87.7 (83.5-91.9)	11.3 (9.5-13.4)	M9.5	126	84.1 (77.7-90.5)	10.9 (8.7-13.6)
-	-	-	-	M20	126	39.7 (31.2-48.2)	4.1 (3.4-5)
M23*	193	56.3 (49.2-63.3)	3.9 (3.5-4.5)	-	-	-	-
-	-	-	-	M36*	117	21.4 (14-21.4)	2.7 (2.4-3.1)
-	-	-	-	M37.5	99	86.9 (80.3-93.5)	17.5 (13.2-23.2)
-	-	-	-	M50	96	47.9 (37.9-57.9)	4.4 (3.5-5.5)
M71	115	31.3 (22.8-39.8)	3.28 (2.8-3.9)	-	-	-	-

*Sample collected before dose.
D=day. W=week. M=month.

Table 2 Immunogenicity of MenBvac® vaccine against strains of serogroups B or C in 12 responder subjects. Results are expressed in comparison to reference strains that completely match PorA of the vaccine strain.

Study strain number (LNP)	Genotyping					SBA			
	Group	PorA VR1	PorA VR2	cc	FetA	Range of SBA titres (Proportions of sera with protective titres)	GMT of hSBA titres [95% CI]	GMT [95% CI]	p-value
Serogroup B strains; SBA using human complement (hSBA); protective response for hSBA ≥ 4									
Complete match with the vaccine strain									
LNP21362	B	7	16	32	F3-3	16-128 (100%)	107.6	73.5- 157.6	Ref*
Partial match with the vaccine strain									
LNP27010	B	7-2	4	41/44	F1-5	2-16 (92%)	6.4	4.3-9.4	0.0005
LNP27087	B	7	16-26	32	F3-3	32-128 (100%)	114	88.4- 147.1	>0.99
LNP27114	B	7-2	4	162	F5-9	2-128 (75%)	15.1	5.9-39	0.002
LNP27338	B	7-2	16-26	32	F3-3	128-128 (100%)	128	128-128	>0.99
LNP27372	B	7-1	1	32	F3-3	2-128 (92%)	24	9.7-59.5	0.00195
LNP27200	B	21	16-36	865	F5-8	2-4 (25%)	2.4	1.9-2.9	0.0005
LNP27386	B	21	16	41/44	F1-7	128-128 (100%)	128	128-128	>0.99
No match with the vaccine strain									
LNP27000	B	19-1	15-11	269	-	2-2 (0%)	2	2-2	0.0005
LNP27050	B	19-2	13-1	461	-	2-8(42%)	4	2.1-7.5	<0.0001
LNP27143	B	22	9	269	F1-5	2-8 (25%)	2.7	1.9-3.8	0.0005
Serogroup C strains; SBA using rabbit complement (rSBA); protective response for rSBA ≥ 8									
Complete match with the vaccine strain									
LNP25514	C	7	16	32	F3-3	16-512 (100%)	64	26.7- 153.2	Ref**
Partial match with the vaccine strain									
LNP25553	C	17	16-3	11	F1-1	16-32 (100%)	18.4	14.9-22.7	0.016
LNP26251	C	21-7	16	1157	F5-36	64-512 (100%)	161.3	59.7- 435.8	0.019

* Significance level $p < 0.005$ after Bonferroni correction for 10 comparisons

** Significance level $p < 0.025$ after Bonferroni correction for 2 comparisons

Table 3

Percentage of current invasive meningococcal strains in France that are expected be covered by either MenBvac® or 4CMenB/Bexsero® based on their PorA domain (for 4CMenB/Bexsero® the coverage is largely enhanced by the three other components of the vaccine).

2015 French <i>N. meningitidis</i> isolates with complete typing available (n=344)						
	Matching with the P1.7,16 PorA antigen of MenBvac			Matching with the P1.7-2,4 PorA antigen of 4CMenB/Bexsero		
	Complete	Incomplete	Null	Complete	Incomplete	Null
B strains (n=170)	8 (4.7%)	45 (26.5%)	117 (68.8%)	28 (16.5%)	21 (12.4%)	121 (71.1%)
Strains of other groups (n=174)	1 (0.6%)	4 (2.3%)	169 (97.1%)	0 (0%)	1 (0.6%)	173 (99.4%)
Total (n=344)	9 (2.6%)	49 (14.3%)	286(83.1)	28 (8.1%)	22 (6.4%)	294 (85.5%)