

Table 2. Distribution of *S. enterica* subsp. *enterica* and *houtenae* serovars in 171 patients aged ≤ 15 years admitted to the emergency room of the University Hospital of Pointe-à-Pitre

Stools			Blood			Total*		
Serovar	N	%	Serovar	N	%	Serovar	N	%
Panama	54	35.1	Panama	31	53.4	Panama	57	33.3
Enteritidis	23	14.9	Arechavelata	17	29.3	Arechavelata	28	16.4
Arechavelata	18	11.7	4,5,12:i:-	2	3.4	Enteritidis	23	13.5
4,5,12:i:-	15	9.7	Enteritidis	2	3.4	4,5,12:i:-	15	8.8
43:z4,z32:-:†	9	5.8	Newport	2	3.4	43:z4,z32:-:†	9	5.3
Newport	7	4.5	Typhi	2	3.4	Newport	7	4.1
Infantis	5	3.2	Rubislaw	1	1.7	Rubislaw	7	4.1
Rubislaw	5	3.2	Schwarzengrund	1	1.7	Infantis	5	2.9
Typhimurium	4	2.6				Typhimurium	4	2.3
4,12:i:-	2	1.3				Typhi	3	1.8
43:z4,z23:-:†	1	0.6				4,12:i:-	2	1.2
Agona	1	0.6				43:z4,z23:-:†	1	0.6
Braenderup	1	0.6				Agona	1	0.6
Havana	1	0.6				Braenderup	1	0.6
Javiana	1	0.6				Havana	1	0.6
Miami	1	0.6				Javiana	1	0.6
Mississippi	1	0.6				Miami	1	0.6
Montevideo	1	0.6				Mississippi	1	0.6
Pomona	1	0.6				Montevideo	1	0.6
Poona	1	0.6				Pomona	1	0.6
Schwarzengrund	1	0.6				Poona	1	0.6
Typhi	1	0.6				Schwarzengrund	1	0.6
Total	154	100	Total	58	100	Total	171	100

* If more than one isolate with the same serovar and antimicrobial resistance phenotype was recovered from the same patient, only the first was retained.

† *S. enterica* subsp. *houtenae*