

1
2
3 **Impaired phagocytosis directs human monocyte activation in response to fungal derived**
4 **β -glucan particles**

5
6 Giorgio Camilli¹, Elif Eren², David L. Williams³, Vishukumar Aimanianda⁴, Etienne
7 Meunier², and Jessica Quintin^{1#}
8

9
10 ¹ Immunology of Fungal Infections, Department of Mycology, Institut Pasteur, Paris, France.

11 ² Institut de Pharmacologie et Biologie Structurale, IPBS, Université de Toulouse, CNRS,
12 UPS, Toulouse, France.

13 ³ Department of Surgery, Center of Excellence in Inflammation, Infectious Disease and
14 Immunity, James H. Quillen College of Medicine, East Tennessee State University, Johnson
15 City, Tennessee, USA.

16 ⁴ Unité des Aspergillus Department of Mycology, Institut Pasteur, Paris, France.
17

18
19 Keywords: Monocytes, phagocytosis, fungi, β -glucan, inflammation
20

21 # ***Corresponding author :***

22 Dr. Jessica Quintin
23 Immunology of Fungal Infections
24 Mycology Department
25 Bâtiment Metchnikoff (67)
26 Institut Pasteur
27 25, rue du Docteur Roux
28 75015 Paris, France
29 Tel: +33 (0)1 44 38 94 54
30 Fax: +33 (0)1 40 61 31 15
31 Mail: jessica.quintin@pasteur.fr
32
33

34 **Abbreviations**

35 AI : Alkali Insoluble
36 AS : Alkali Soluble
37 ASC : Apoptosis-associated Speck-like Protein Containing a CARD
38 CARD : Caspase recruitment domain
39 CD : Cluster of differentiation
40 CR : Complement receptor
41 Cyt D : Cytochalasin D
42 DCs : Dendritic cells
43 DPI : Diphenyleneiodonium
44 GAPDH : Glyceraldehyde-3-phosphate dehydrogenase
45 HK : heat-killed
46 HK-C.a. : heat-killed *Candida albicans*
47 I κ B α : Inhibitor of NF- κ B, alpha
48 IL: Interleukin
49 ITAM : Immunoreceptor tyrosine-based Activation Motif
50 LPS : Lipopolysaccharide
51 MALT : Mucosa-associated lymphoid tissue lymphoma translocation protein
52 NADPH : Nicotinamide adenine dinucleotide phosphate hydrate
53 NF- κ B : nuclear factor kappa-light-chain-enhancer of activated B cells
54 NLRP3: NACHT, LRR and PYD domains-containing proteins 3
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

PBMC : Peripheral blood mononuclear cell
PI3K : Phosphatidylinositol 3-kinase
PRR : Pattern recognition receptor
ROS : Reactive oxygen species
SHIP : SH2-containing inositol phosphatase
Syk : Spleen tyrosine kinase
TLR : Toll-like receptor
TNF α : Tumor necrosis factor alpha α

Abstract

Recognition of the fungal cell wall carbohydrate β -glucan by the host receptor Dectin-1 elicits broad immunomodulatory responses, such as phagocytosis and activation of oxidative burst. These responses are essential for engulfing and killing fungal pathogens. Phagocytic monocytes are key mediators of these early host inflammatory responses to infection. Remarkably, whether phagocytosis of fungal β -glucan leads to an inflammatory response in human monocytes remains to be established. Here, we show that phagocytosis of heat-killed *Candida albicans* is essential to trigger inflammation and cytokine release. By contrast, inhibition of actin-dependent phagocytosis of particulate (1-3,1-6)- β -glucan induces a strong inflammatory signature. Sustained monocyte activation, induced by fungal β -glucan particles upon actin cytoskeleton disruption, relies on Dectin-1 and results in the classical caspase-1 inflammasome formation through NLRP3, generation of an oxidative burst, NF- κ B activation, and increased inflammatory cytokine release. PI3K and NADPH oxidase were crucial for both cytokine secretion and ROS generation, whereas Syk signaling mediated only cytokine production. Our results highlight the mechanism by which phagocytosis tightly controls the activation of phagocytes by fungal pathogens and strongly suggest that actin cytoskeleton dynamics are an essential determinant of the host's susceptibility or resistance to invasive fungal infections.

Introduction

Prior to the mid-1990s, innate immunity was considered to only be required to 'ingest and kill' microorganisms during the course of an infection, providing the host with time to trigger an effective and specific adaptive immune response. However, the discovery of Toll-like receptors led to a revolution in our understanding of the specific role of innate immunity and how the host immune system is activated upon microbial recognition. Several major classes of pathogen-recognition receptors (PRR) have now been described, each with specificity for conserved pathogen structures [1]. A single fungal species, such as *Candida albicans*, is recognized by multiple PRR, including Toll-like receptor (TLR)1, TLR2, TLR4, TLR6, TLR9, mannose receptor (MR), Dectin-1, Dectin-2, galectin-3, and the lectin domain of complement receptor 3 (CR3) (reviewed in [2]).

Recognition of the most abundant fungal cell wall polysaccharide, β -glucan, by Dectin-1 plays an essential role in the host response to fungal infection [3, 4]. Dectin-1 signaling activates antimicrobial innate immune responses, including phagocytosis and the production of reactive oxygen species (ROS) in myeloid cells [5, 6]. Interestingly, Dectin-1 activation requires a phagocytic synapse, mediated by the binding of particulate glucans, and not by small soluble β -glucan molecules [7]. This synapse allows the phosphorylation of the hemITAM (hemi-immunoreceptor tyrosine-based activation motif) by excluding regulatory phosphatases [7]. Notably, however, the use of either yeast or β -glucan with low purity in a majority of the studies has contributed to conflicting conclusions regarding the role of β -glucan in inducing an inflammatory cytokine response. For example, it has been reported that zymosan, a β -glucan-enriched crude preparation of the *Saccharomyces cerevisiae* fungal cell wall, potentiate cytokine secretion via TLR2 and Dectin-1 signaling [8]. By contrast,

1
2
3 microglial cells, RAW cells (a macrophage cell line) or thioglycollate-elicited macrophages
4
5 fail to secrete cytokines following exposure to pure particulate β -glucan [9, 10]. Walachowski
6
7 and colleagues reported that enriching the cell wall fraction of β -glucan from 15% to 75%
8
9 removes TLR ligands while preserving Dectin-1 activity in murine macrophages [11].
10
11 Furthermore, the most enriched extracts trigger very low nuclear factor kappa-light-chain-
12
13 enhancer of activated B cells (NF- κ B) activity and cytokine production [11].
14
15

16
17
18 The majority of these studies investigating the mechanisms of enriched or pure fungal β -
19
20 glucan-induced inflammatory cytokine response rely upon mouse macrophages and dendritic
21
22 cells (DCs). Strikingly, variation in the reported role of β -glucan in triggering inflammation
23
24 also relies on cell programming. For example, using pure β -glucan particles, depleted
25
26 zymosan, or curdlan, several studies describe that GM-CSF treated mouse macrophages and
27
28 dendritic cells, but not M-CSF treated macrophages, produce cytokines in response to glucan
29
30 stimulation [12-14]. Pure glucan activation of mouse DCs is Dectin-1 and spleen tyrosine
31
32 kinase (Syk) dependent [15, 16] and is terminated by internalization of Dectin-1. Thus, the
33
34 blockade of phagocytosis permits continued receptor signaling at the cell surface [13, 14].
35
36 Hence, despite numerous reports highlighting β -glucan as a key molecule for host phagocytic
37
38 cell activation, crucial information regarding the specific response of human mononuclear
39
40 phagocytes to this molecule is still lacking.
41
42
43
44
45

46
47 Monocytes have long been considered as developmental intermediates between bone marrow
48
49 progenitors and differentiated tissue macrophages. However, some DCs and tissue
50
51 macrophages do not originate from monocytes in non-inflammatory and homeostatic
52
53 conditions. [17, 18]. Furthermore, monocytes perform specific inflammatory functions during
54
55 infection, such as producing inflammatory mediators, tumor necrosis factor α (TNF α), nitric
56
57
58
59
60

1
2
3 oxide and reactive oxygen species [17, 18] and represent key mediators of early host
4 responses to microbes, including fungi [19]. Moreover, the generation of innate immune
5 memory upon challenge with fungal molecules (i.e. β -glucan) highlights the important role of
6 monocytes in regulation of the host responses to infections and has also gained attention for
7 its potential therapeutic implications [20, 21].
8
9
10
11
12

13
14
15 Here, we performed a systematic investigation of the role of phagocytosis on fungal β -glucan-
16 induced inflammation in human monocytes. Our results demonstrate that actin cytoskeleton
17 disruption induced inflammatory cytokine release and enhanced ROS production upon
18 stimulation with a branched *C. albicans* (1-3)- β -glucan. We demonstrate that the synergistic
19 effect on cytokine secretion and ROS generation observed during actin disruption and β -
20 glucan stimulation was dependent upon Dectin-1, phosphatidylinositol 3-kinase (PI3K), and
21 nicotinamide adenine dinucleotide phosphate hydrate (NADPH) oxidase. Conversely, the
22 role of Syk signaling was limited to cytokine production. We further found that the NACHT,
23 LRR, and PYD domains-containing proteins 3 (NLRP3) inflammasome senses phagocytosis
24 inhibition-associated β -glucan-mediated activation to induce caspase-1 cleavage and
25 interleukin (IL)1 β release in human monocytes. These findings highlights the specific
26 mechanism by which phagocytosis tightly controls the activation of primary human
27 monocytes and suggest that actin cytoskeleton dynamics may drive host's susceptibility or
28 resistance to invasive fungal infections.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Phagocytosis and actin dynamics differently impact monocyte responses to *C. albicans* and β -glucans P

Initially, we hypothesized that engulfment of fungal pathogens by human phagocytes is not only a killing and scavenging process but also a potent mechanism of inflammation regulation. To test this hypothesis, we assessed the role of phagocytosis in the immune response triggered by monocytes upon encounter with heat-inactivated/killed (HK) *Candida albicans*. We used HK *C. albicans* to avoid any active delay in the phagocytic process triggered by live and virulent *C. albicans* [22]. Following exposure of primary human monocytes isolated from blood of healthy donors, we observed that HK *C. albicans* induced the release of TNF α , IL6, IL1 β , and IL10 (Figure 1A). This inflammatory response was abrogated when phagocytosis was impaired by the actin polymerization inhibitor, cytochalasin D (CytD)(Figure 1A and SI Figure 1). The inflammatory response was also induced following exposure to HK *Saccharomyces cerevisiae* and inhibited when phagocytosis was blocked (Figure 1B).

The cell wall of yeast, such as *C. albicans* and *S. cerevisiae*, is composed of two main layers [23]. The outer layer is highly enriched with *O*- and *N*-linked glyco/mannoproteins whereas the inner layer contains the skeletal polysaccharides chitin and branched- β -1-3 glucan that confer strength and cell shape. The outer cell wall layer is attached to this framework predominantly by glycosylphosphatidylinositol (GPI)-remnants that are linked to the skeleton through a more flexible β -1-6 glucan [23]. Alkali treatment of the yeast allows separation of the different cell wall carbohydrates. No inflammatory cytokine release was observed following stimulation of human monocytes with the alkali soluble (AS) fraction of *S.*

1
2
3 *cerevisiae*, containing mannoproteins and β -(1-6)-glucan, regardless of the presence or
4 absence of CytD (Figure 1C). Incubation of monocytes with the Alkali Insoluble (AI) fraction
5 containing branched β -(1-3)-glucan, chitin and β -(1-6)-glucan neither did trigger an
6 inflammatory response. Interestingly however, impairing phagocytosis before stimulation
7 with the AI fraction led to strong cytokine release in the supernatant (Figure 1C). Within this
8 fraction, we observed that branched β -(1-3)-glucan (associated with less than 5% chitin) was
9 mimicking the AI fraction whereas β -(1-6)-glucan alone (from AS fraction) were not able to
10 replicate the inflammatory effect observed with Cyt D (Figure 1C). To determine the structure
11 required to trigger the release of cytokine by monocytes when phagocytosis is blocked, we
12 used pure insoluble branched β -(1-3)-glucan from *S. cerevisiae* and *C. albicans*. Both
13 branched β -(1-3)-glucans induced monocyte activation when their engulfment was impaired
14 by Cyt D treatment in a dose dependent manner (Figure 1D; SI Figure 1). The increase in
15 cytokine release was more potent at lower concentrations with branched *C. albicans* β -(1-3)-
16 glucan (Figure 1D). As previously reported for mouse DCs and macrophages [12, 13, 15, 16],
17 we observed that large curdlan particles (insoluble linear β -(1-3)-glucan) stimulated human
18 monocytes in an actin-dependent manner, but laminarin (soluble branched β -(1-3,1-6)-
19 glucan)) did not activate monocytes, even when phagocytosis was inhibited (Figure 1E). To
20 rule out potential aggregations interference (SI Figure 2), mild sonication of branched *C.*
21 *albicans* β -(1-3)-glucan was performed which had no effect on monocyte activation triggered
22 in presence of Cyt D (Figure 1F). Increasing the concentration of branched *C. albicans* β -(1-
23 3)-glucan, or of AI fraction together with Cyt D, showed similarly increased cytokine
24 expression (Figure 1G). By contrast, increasing the concentration of the AS fraction did not
25 induce cytokine expression even in the presence of Cyt D. Of note, combining the AS fraction
26 and the AI fraction had no inhibitory effect on the release of cytokines mediated by AI
27 (Figure 1G). Similarly, increasing the concentration of HK *C. albicans* augmented the initial
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 inflammatory response, but blocking phagocytosis still inhibited the release of cytokines. As
4
5 actin polymerization inhibition led to an increase in cytokine expression which we confirmed
6
7 by using latrunculin B (Figure 1H), we next wanted to test the impact of an actin filament
8
9 polymerizing and stabilizing drug. We observed that jasplakinolide had no effect on HK *C.*
10
11 *albicans*-mediated inflammation or on the absence of response with *C. albicans*-branched β -
12
13 (1-3)-glucan (Figure 1H). Finally, whereas blocking actin polymerization (Cyt D or
14
15 Lantrunculin B), clathrin-mediated endocytosis using chlorpromazine, or acidification of the
16
17 phagolysosome with chloroquine inhibited cytokine production upon HK *C. albicans*
18
19 stimulation, only the inhibition of actin polymerization (Cyt D or Lantrunculin B) was able to
20
21 activate the release of cytokines by human monocytes incubated with *C. albicans* β -(1-3)-
22
23 glucan (Figure 1H).
24
25
26
27
28

29 **IL-1 β release is mediated by the canonical NLRP3 inflammasome in phagocytosis-** 30 31 **impaired monocytes I** 32 33 34

35 As impaired phagocytosis triggered the secretion of IL1 β by human monocytes upon
36
37 branched *C. albicans* β -(1-3)-glucan exposure, we wanted to determine how IL1 β is activated.
38
39 The bioactivity of IL1 β is controlled by activation of the inflammasome [24]. The inactive
40
41 precursor molecule, pro-IL1 β , must be converted to the mature form to become functional
42
43 [24]. Processing of pro-IL1 β is mediated by the enzyme caspase-1, which is activated by
44
45 multi-protein inflammasome complexes such as NLRP3. In macrophages and DCs, activation
46
47 of the NLRP3 inflammasome requires two separate signals [24]. However, in human
48
49 monocytes, lipopolysaccharide (LPS) from Gram-negative bacteria is sufficient to trigger
50
51 caspase-1-dependent IL1 β maturation and secretion [25]. This activity in monocytes is clearly
52
53 distinct from the two-step activation observed in macrophages and DCs [26]. As such, it has
54
55
56
57
58
59
60

1
2
3 become evident that in monocytes LPS is directly sensed by caspase-11 or its human
4 orthologs caspase-4 and caspase-5 in the cytosol to trigger cell death or activate the NLRP3
5 inflammasome [27, 28]. In addition, human caspase-4 and caspase-5 regulate the one-step
6 non-canonical inflammasome activation in human monocytes [29]. Finally in human
7 monocytes, extracellular LPS triggers an alternative inflammasome that relies upon TLR4 -
8 TRIF - RIPK1 - (FADD) – caspase-8 signaling upstream of NLRP3-ASC-caspase-1 [25].
9 Interestingly, both caspase-1 and -8 are associated with linear- β -glucan/Dectin-1 signaling in
10 human DC [30, 31]. Of note, although not associated with fungi or β -glucan, the one-step
11 non-canonical inflammasome activation by LPS is Syk-dependent, a crucial kinase required
12 for β -glucan/Dectin-1 signaling [29].
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Thus, we wanted to determine which of these caspases mediated mature IL1 β release
27 following concomitantly phagocytosis inhibition and *C. albicans* β -(1-3)-glucan exposure in
28 human monocytes. We observed that the pan-caspase inhibitor Z-VAD blocked the release of
29 IL1 β by phagocytosis-impaired human monocytes upon *C. albicans* β -(1-3)-glucan exposure,
30 suggesting caspases mediate IL1 β processing (Figure 2A). Only the caspase-1 inhibitor Z-
31 YVAD, but not caspase-8 Z-IETD nor caspase-4 and -5 Z-LEVD inhibitors inhibited IL1 β
32 release (Figure 2A). Thus, when phagocytosis was impaired, IL1 β release was mediated by
33 the protease caspase-1 following exposure of human monocyte to *C. albicans* β -(1-3)-glucan.
34 Immunoblot experiments confirmed that concomitant Cyt D/ β -glucan treatment promoted
35 both caspase-1 activation and IL1 β processing and release in monocytes (Figure 2B), thus
36 confirming the involvement of an inflammasome in this process. A feature of the canonical
37 inflammasome is its ability to induce ASC polymerization for efficient IL1 β processing after
38 receptor activation [24]. This structure is called a speck and can be imaged by fluorescence
39 [32]. Impairing actin polymerization prior to *C. albicans*-branched β -(1-3)-glucan exposure
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 strongly activated the formation of ASC specks, a phenomenon only mildly induced when
4 actin polymerization was intact (Figure 2C and D). The NLRP3 inflammasome promotes both
5 caspase-1 and -8-mediated IL1 β processing and release in response to β -glucans and fungi
6 [33]. To determine whether NLRP3 could be the mediator of Cyt D/ β -glucan-induced IL1 β
7 processing and release, we used the potassium channel inhibitor glibenclamide, observing that
8 a potassium signal was required for β -glucan-triggered IL1 β release (Figure 2E). Similar
9 results were obtained when using MCC950, a specific NLRP3 inhibitor, in which IL1 β
10 release was strongly diminished (Figure 2E). Intriguingly, although the caspase recruitment
11 domain (CARD)9 - Mucosa-associated lymphoid tissue lymphoma translocation protein
12 (MALT)1 pathway promotes pro-inflammatory cytokine transcription [34], the paracaspase
13 MALT1 inhibitor did not significantly alter mature IL1 β release from human monocytes
14 stimulated with Cyt D/ β -glucan (Figure 2E).
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **Dectin-1, Syk, and PI3K pathways control β -glucan signaling in human phagocytosis-** 32 **impaired monocytes D** 33 34 35 36 37

38 Signalling of β -glucan is mediated by a limited number of surface PRRs on mononuclear
39 phagocytes [35]. As cytokine release by human monocytes, inhibited for phagocytosis, was
40 observed following exposure to *C. albicans* β -(1-3)-glucan we therefore investigated which
41 receptors and signaling pathways were involved.
42
43
44
45

46 Following antibody-mediated inhibition of TLR2 or CR3 receptors, we observed no effect on
47 the release of cytokines whereas blockade of Dectin-1 completely inhibited cytokine secretion
48 mediated by monocytes whose phagocytosis is impaired and stimulated with *C. albicans* β -(1-
49 3)-glucan (Figure 3A). In addition, only a partial inhibition of IL6 cytokine release was
50 observed using a combination of antibodies against CD11b and CD18 subunits of CR3,
51
52
53
54
55
56
57
58
59
60

1
2
3 described as more effective in comparison with blocking the CD11b chain alone [36] (SI
4
5 Figure 3).

6
7
8
9 Dectin-1 activates cellular responses via the Syk/CARD9 [37]. Inhibiting Syk kinase in our
10
11 human primary monocytes system diminished the amount of cytokines produced upon
12
13 stimulation with *C. albicans*-branched β -(1-3)-glucan, suggesting that the stimulation of
14
15 phagocytosis-impaired monocytes by Dectin-1 is Syk kinase-dependent (Figure 3B).
16
17 Phosphoinositide-3 kinase (PI3K) is also a downstream mediator of the Dectin-1 signaling
18
19 pathway [38]. Interestingly, inactivation of PI3K, using wortmannin, had a strong inhibitory
20
21 effect on cytokine production of β -(1-3)-glucan/Cyt D treated monocytes, suggesting that
22
23 PI3K is not only working downstream of Syk but in a concomitant pathway as well (Figure
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
PI3K is not only working downstream of Syk but in a concomitant pathway as well (Figure
3C).

31 We next assessed the activation of the crucial transcription factor NF- κ B by β -(1-3)-glucan
32
33 exposure of human monocytes impaired for phagocytosis. Impairing actin polymerization
34
35 prior to *C. albicans* β -(1-3)-glucan exposure strongly induced phosphorylation of the inhibitor
36
37 of NF- κ B alpha (I κ B α) protein (Figure 4A, B). Phosphorylation was less evident when actin
38
39 polymerization was intact in monocytes treated with by *C. albicans*-branched β -(1-3)-glucan
40
41 (Figure 4A, B). Of interest, the strong phosphorylation of I κ B α observed in monocytes
42
43 treated with CytD and stimulated by *C. albicans* β -(1-3)-glucan correlated with the
44
45 degradation of the protein (Figure 4A, B). Finally, both phosphorylation and degradation of
46
47 I κ B α correlated with the activation and translocation of the canonical NF- κ B transcription
48
49 factor into the nuclei (Figure 4C).
50
51
52
53
54
55
56
57
58
59
60

Differential role of Syk and PI3K in the inflammatory response triggered by β -glucans in monocytes

The immune response mediated by mononuclear phagocytes upon fungal recognition and engulfment is not solely restricted to induction of inflammatory cytokines. Indeed, pattern-recognition receptors in innate immunity trigger reactive oxygen (ROS) species for intracellular killing of fungi [39]. Therefore, we assessed the production of ROS in our monocytes. Stimulation of monocytes with *C. albicans* β -(1-3)-glucan triggered ROS production compared to the basal level detected in resting cells (Figure 5A, D: vehicle). Similar ROS induction was observed when actin polymerization was blocked in untreated monocytes (Figure 5A, D). However, a strong and synergistic induction of ROS was observed when phagocytosis-impaired monocytes were stimulated with *C. albicans* β -(1-3)-glucan. This induction was NADPH dependent as pharmacological inhibition of oxidase activation with either diphenyleneiodonium (DPI) (Figure 5A) or apocynin (SI Figure 4) reduced the production of ROS in a dose-dependent manner. Inhibition of oxidase activation also reduced the secretion of TNF α , IL6, and IL1 β by phagocytosis-impaired monocytes stimulated with *C. albicans* β -(1-3)-glucan (Figure 5B and SI Figure 4).

These results suggest that NADPH-induced ROS are important mediators of the unique inflammatory cytokine signature of phagocytosis-impaired monocytes stimulated with *C. albicans*-branched β -(1-3)-glucan. We, therefore, investigated whether the signaling pathway leading to ROS production shares common mediators with the regulation of cytokine production. ROS generation was observed in monocytes in response to *C. albicans* β -(1-3)-glucan and was blocked by inhibiting either Dectin-1, Syk, or PI3K (Figure 5C, D: DMSO treated). By contrast, inhibition of either Dectin-1 or PI3K, but not Syk, abolished the potent

1
2 synergistic ROS release observed in phagocytosis-impaired monocytes stimulated with *C.*
3 *albicans* β -(1-3)-glucan (Figure 5C, D: Cyt D). This result was surprising given that, unlike
4
5 the concomittent regulation of cytokine release, we observed that oxidative burst, triggered by
6
7 a fungal encounter, depended on PI3K and not Syk.
8
9

10
11
12
13 As we observed that inflammatory cytokine release in phagocytosis-impaired monocytes
14
15 stimulated with *C. albicans* β -(1-3)-glucan was Syk-dependent (Figure 3B), we tested Syk
16
17 involvement in inflammasome and NF- κ B activation. Inhibition of Syk abolished the nuclear
18
19 translocation of NF- κ B transcription factor in phagocytosis-impaired monocytes stimulated
20
21 with *C. albicans* β -(1-3)-glucan (Figure 6A, B). Similarly, inhibition of Syk activity strongly
22
23 reduced caspase-1 activation and the release of active IL-1 β and caspase-1 in the supernatant
24
25 (Figure 6C). Therefore, Syk activity is required for inflammasome and NF- κ B activation, and
26
27 cytokine release.
28
29
30
31
32

33 **Discussion**

34
35
36
37 The surveillance and elimination of fungal pathogens relies heavily on the sentinel behavior
38
39 of phagocytic cells of the innate immune system. As such, innate mononuclear phagocytes
40
41 must recognize, uptake, and kill fungal pathogens. Herein, we demonstrate that phagocytosis
42
43 of fungal cells or fungal cell wall componenets by human primary monocytes also regulates
44
45 the inflammatory responses mediated by the host. We demonstrated that the inflammatory HK
46
47 *C. albicans*-dependent response of monocytes requires phagocytosis and engulfment of the
48
49 yeast. By contrast, ingestion of insoluble branched *C. albicans* β -(1-3)-glucan is inert and
50
51 cytokine secretion from human monocytes is induced only when β -(1-3)-glucan remains in
52
53 the extracellular space. Activation of phagocytosis-impaired-human monocytes by branched
54
55
56
57
58
59
60

1
2
3 *C. albicans* β -(1-3)-glucan through Dectin-1 , Syk and PI3K results in cytokines release and
4
5 NF- κ B activation. In addition we observed NLRP3/ASC/caspase-1 inflammasome formation,
6
7 leading to IL-1 β processing and release. Interestingly our results highlight that the increased
8
9 β -glucan-induced oxidative burst in phagocytosis-impaired monocytes depends on Dectin1
10
11 and PI3K but is independent of the canonical Syk pathway.
12
13

14
15
16 Blocking actin polymerization, hence phagocytosis, converts *C. albicans*-branched β -(1-3)-
17
18 glucan into a potent activator of innate responses via the Dectin-1 pathway. This observation
19
20 is reminiscent of the work of Hernanz-Falc3n and colleagues in which the blockade of glucan
21
22 particles internalization trigger the response of dendritic cells and more specifically mitogen-
23
24 activated protein kinase (MAPK) activation [13]. One might therefore question the role of
25
26 phosphatase in the regulation mediated by actin polymerization. One possible candidate could
27
28 be SH-2 containing inositol 5' polyphosphatase 1 (SHIP1) that has recently been
29
30 demonstrated to interact with the Dectin-1 hemITAM and selectively modulates reactive
31
32 oxygen species production in dendritic cells in response to *C. albicans* [40]. However, the
33
34 study of Blanco-Men3ndez and colleagues showed no effect of the SH-2 containing inositol
35
36 5' polyphosphatase 1 (SHIP1) on the regulation of the secreted cytokines, which was a
37
38 hallmark of our study.
39
40
41
42
43

44
45 Our observation that the insoluble and linear β -glucan curdlan is able to stimulate the release
46
47 of cytokines from human primary monocytes, in an actin-dependent mechanism is consistent
48
49 with other studies in murine macrophages and dendritic cells [13-16]. Indeed, the release of
50
51 cytokines is inhibited when phagocytosis is impaired, a phenomenon reminiscent of what we
52
53 observed with HK *C. albicans*. However, whereas *C. albicans* yeast are evenly distributed
54
55 among the monocytes population and phagocytosed in control situation when actin
56
57
58
59
60

1
2
3 polymerization is functional, (SI Figure 1), very large curdlan aggregates are sporadically
4 distributed and never phagocytosed (SI Figure 1). Indeed, even if monocytes migrate towards
5 the curdlan aggregate, they only surround and bind the β -(1-3)-glucan curdlan particles
6 without phagocytosis (SI Figure 1 and 24 hours time point, data not shown). However, when
7 actin polymerization is blocked, migration of monocyte is constrained and recognition of β -
8 (1-3)-glucan by Dectin-1 is inhibited. This lack of dectin-1 engagement with β -glucans,
9 reported as well by Hernanz-Falcon et al. and Rosas et al. in mice [13, 14], is the likely
10 explanation for the impeded cytokine release. Dectin-1 engagement might, however, still
11 occur with HK *C. albicans* or branched β -(1-3)-glucan isolated from *C. albicans* or *S.*
12 *cerevisiae* as the organisms/particles are more evenly distributed and easily accessible to
13 monocytes (SI Figure 1). Interestingly, whereas Dectin-1 is likely activated by either HK *C.*
14 *albicans* or branched β -(1-3)-glucan, HK *C. albicans* does not elicit cytokine release when
15 phagocytosis is impaired. This observation is reminiscent of the work of Filler et al., who
16 demonstrated that endothelial cell phagocytosis of live *C. albicans* was necessary for
17 endothelial cell stimulation to occur [41]. Strikingly however, and in contrast with our
18 observation in human monocytes, they also determined that killed *C. albicans* did not induce
19 endothelial cell activation, even though killed *C. albicans* are phagocytosed by endothelial
20 cells [41].
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 Our data show that impaired uptake of *C. albicans*-branched β -(1-3)-glucan particles results
45 in high release of IL1 β associated with NLRP3/ASC/caspase-1 inflammasome formation.
46 This observation suggest that NLRP3 could sense the overstimulation of the Dectin-1 receptor
47 by particulate agonists. Intriguingly, inhibition of the para-Caspase activity of MALT1 did not
48 significantly affect the maturation of IL1 β although the CARD9-MALT1 pathway was
49 described to promote pro-inflammatory cytokine transcription. Whether another adaptor from
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the Dectin-1 pathway is involved or if MALT1 plays a role independent from its catalytic
4 domains remains to be investigated. However, MALT1 also expresses a Caspase-Activating
5 Recruitment Domain (CARD) essential for the CBM (CARD-BCL-MALT) tripartite complex
6 formation CARD9-BCL10-MALT1 and signal transduction after Dectin-1 engagement. One
7 should not exclude that upon prolonged interactions between these three partners, the
8 (CARD-BCL-MALT) CBM complex could induce a stress detected by the NLRP3 sensor.
9
10
11
12
13
14
15
16
17

18 The opposite effect on monocyte cytokine responses between whole *Candida albicans* and β -
19 glucan particles is intriguing and raises the question of a putative negative regulation of the
20 phagocytic-dependent cytokine secretion by whole *Candida albicans*. As such, one might
21 hypothesize that mannan in the cell wall of *Candida albicans* acts as a negative regulator of
22 inflammation. Previous studies have shown that mannan contributes to the resistance of *C.*
23 *albicans* to complement activation and complement component 3 (C3) binding by *C. albicans*
24 [42]. However, adding mannans within the alkali-soluble fraction, to the potent β -glucan
25 inducer in the alkali-insoluble fraction did not inhibit the release of cytokines induced by
26 phagocytosis-impaired monocytes. We can, therefore, exclude that alkali soluble fractions of
27 yeast contain inhibitors of the β -glucan mediated inflammation. However, alkali-treatment is
28 very stringent, so we cannot yet exclude that surface proteins on *Candida albicans* are
29 responsible for the mirror effect observed.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 The discrepancy we observed between the different size of branched β -glucan is also of
47 interest. Such discrepancy in size has already been observed in the signaling mediated by
48 Dectin-1 during which the signal is only transduced when two or more receptors cluster
49 together to form a full ITAM motif. This event only occurs with particulate β -glucans, which
50 cluster the receptor in synapse-like structures from which the regulatory tyrosine
51
52
53
54
55
56
57
58
59
60

1
2
3 phosphatases CD45 and CD148 are excluded [7]. Similarly, during the submission of this
4
5 manuscript, Matthew J. Elders and colleagues observed that Dectin-1-mediated immune
6
7 responses in human dendritic cells is highly dependent on β -glucan particle size and the high
8
9 production of IL1 β . Interestingly blocking internalization allows the production of IL1 β [43].
10
11 Our present study now suggests that not only the stoichiometry, but also the capacity to be
12
13 phagocytosed, plays a role in the signaling. Very recently, a study on neutrophils has
14
15 highlighted the differential responses obtained when the pathogen encountered is of small size
16
17 –and phagocytosed-, or of bigger size –hence not phagocytosed [44]. More specifically and
18
19 elegantly, the study demonstrates that the ability to sense the differential localization of ROS
20
21 allows neutrophils to adjust the inflammatory response by modulating their own recruitment
22
23 and cooperation to effectively clear microbes of different size. Whether a similar mechanism
24
25 is involved in monocytes is still questionable. But such a hypothesis might be supported by
26
27 the observation that human peripheral blood mononuclear cells (PBMCs) use distinctly
28
29 Dectin-1 or CR3 receptor and overlapping, but distinct, signaling pathways for the oxidative
30
31 burst in response to challenge by different physical forms of β -glucan [36].
32
33
34
35
36

37
38 It is also possible that the source of the cell wall fragment may also be of importance. In this
39
40 study we focused on the yeast form of *Candida albicans* and its cell wall component β -
41
42 glucan. But it has been demonstrated that with *Aspergillus fumigatus*, another important
43
44 human fungal pathogen, in contrast to conidia, hyphal fragments induce NLRP3
45
46 inflammasome assembly, caspase-1 activation and IL1 β release from a human monocyte cell
47
48 line [45]. *C. albicans* hyphal β -glucan, but not of yeast, can stimulate the release of IL1 β by
49
50 monocytes-derived-macrophages [46]. The ability of *C. albicans* to switch from a unicellular
51
52 yeast form into a filamentous hyphal form has been presented as essential for activation of the
53
54 NLRP3 inflammasome, as *C. albicans* mutants incapable of forming hyphae were defective in
55
56
57
58
59
60

1
2
3 their ability to induce murine macrophage IL1 β secretion [47]. By contrast, Mukaremera and
4 colleagues demonstrated that *C. albicans* hyphae stimulate proportionally lower levels of
5 certain cytokines from human PBMCs per unit of cell surface area than yeast cells, but did not
6 suppress cytokine responses when copresented with yeast cells. This latter observation is
7 intriguing as germ tubes with an attached parent yeast cell does not induce a response that
8 would normally be associated with free yeast cells [48].
9
10
11
12
13
14
15
16
17

18 Our study also raises the question of the importance of phagocytosis in the inflammatory
19 response mediated in situations where phagocytosis is naturally impaired. As such, *Candida*
20 *albicans* is known to produce potent biofilms rendering it tolerant to antifungal treatment
21 [49]. In such biofilms, innate immune cells can hardly migrate [50] rendering phagocytosis
22 very unlikely. However, soluble matrix β -(1-3)-glucans are accessible [49]. One might
23 hypothesize that the combined detection of non-phagocytosed soluble β -(1-3)-glucans
24 together with the non-phagocytosis of whole *Candida albicans* cells participates in the non-
25 effective immune responses triggered towards *Candida albicans* biofilm.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Materials and methods

Cell isolation

Human PBMCs were separated on Ficoll-Paque (GE Healthcare) by density centrifugation of heparinized blood from healthy donors ("Etablissement Français du Sang" EFS, Paris, France, habilitation HS-2015-25101). Monocytes were purified by negative selection using the Pan Monocyte Isolation Kit (Miltenyi Biotec, Bergisch Gladbach, Germany). The purity evaluated by flow cytometry was routinely >90%. Cells were seeded at 5×10^5 cells/ml in RPMI 1640, supplemented with 2 mM GlutaMax, 50 $\mu\text{g/ml}$ gentamicin and 1 mM sodium pyruvate (Gibco, Invitrogen, Carlsbad, CA, USA). Cells were maintained in a humidified atmosphere of 5% CO_2 at 37°C.

Stimulation experiments

When indicated 1×10^5 monocytes were pretreated with DMSO (SIGMA), cytochalasin D (1 $\mu\text{g/ml}$, C8273, SIGMA), latrunculin B (2,5 μM , 428020, EMD Millipore), jasplakinolide (0,1 μM , 420127, EMD Millipore), chlorpromazine (1 $\mu\text{g/ml}$, C8138, SIGMA) or chloroquine (25 μM , C6628, SIGMA) for 1 hour. Different stimuli were then added to the cells: heat-inactivated *Candida albicans* (UC820 strain, 30 min at 95 degrees, $10^5/\text{ml}$ unless otherwise stated), heat-inactivated *Saccharomyces cerevisiae* (BY471 strain, 30 min at 95 degrees, $10^6/\text{ml}$ unless otherwise stated), purified β -glucan from *Candida albicans* (1 $\mu\text{g/ml}$ unless otherwise stated; depyrogenated was provided by Dr. David L. Williams, East Tennessee State University) (β -glucan from *Candida albicans* was purified as described [46]); purified β -glucan from *Saccharomyces cerevisiae* (1 or 10 $\mu\text{g/ml}$, G5011, SIGMA); Laminarin from *Laminaria digitata* (1 or 10 $\mu\text{g/ml}$, L9634, SIGMA); curdlan from *Alcaligenes faecalis* (1 or 10 $\mu\text{g/ml}$, Wako); alkali-soluble and insoluble fractions from *Saccharomyces cerevisiae* (1, 10

1
2
3 or 50 µg/ml); β-(1-6)-glucan from *Saccharomyces cerevisiae* (1 or 10 µg/ml); β-(1-3)/(1-6)-
4
5 glucan from *Saccharomyces cerevisiae* (1 or 10 µg/ml). Preparations from the cell wall of
6
7 *Saccharomyces cerevisiae* (BY4741 strain) were provided by Vishukumar Aimanianda,
8
9 Institut Pasteur, Paris, France. Alkali-insoluble fractions was prepared and β-(1-6)-glucan was
10
11 purified from from *S. cerevisiae* (BY4741 strain) as described earlier [51]. To obtain alkali-
12
13 soluble fraction, in brief, the excess of NaBH₄ in the supernatant after NaOH-NaBH₄
14
15 treatment of the sediment (cell wall fraction; [51]) was destroyed using 2% acetic acid,
16
17 followed by dialysis against water for until neutrality and freeze drying. The β-(1-3)-glucan
18
19 (approximately 4% branched by β-(1-6)-linkage; [52]) was purified from alkali-insoluble
20
21 fraction using the protocol described earlier [53]. The stimuli were pre-incubated for 1h at
22
23 room temperature with (10µg/ml) polymyxin B (92283, SIGMA).
24
25
26
27
28

29 For inhibition, before adding DMSO or cytochalasin D, monocytes were preincubated for 30
30
31 min with: 20 µM Ac-YVAD-cmk (caspase-1 inhibitor, inh-yvad, InvivoGen); 20 µM z-VAD-
32
33 fmk (pan-caspase inhibitor, G7231, Promega); 20 µM z-IETD-fmk (caspase-8 inhibitor, 1148,
34
35 BioVision); 10 µM z-LEVD-fmk (caspase-4/5 inhibitor, ALX-260-142, Alexis); 20 µM z-
36
37 VRPR-fmk (MALT1 inhibitor, ALX-260-166, Alexis); 1 µM MCC950 (NLRP-3 inhibitor,
38
39 CP-456773, SIGMA); 125 µM glibenclamide (inhibitor of potassium efflux, G0639,
40
41 SIGMA); 1-2 µM Syk inhibitor (574711, EMD Millipore); 100 nM (wortmannin, tlr1-wtm,
42
43 InvivoGen); 5 µg/ml anti-Dectin-1 Ab (ab82888, abcam) or 5 µg/ml IgG1 Isotype control
44
45 (401402, Biolegend); 10 µg/ml anti-TLR2 Ab (16-9922-82, eBioscience) or 10 µg/ml IgG2a
46
47 Isotype control (16-4724-82, eBioscience); 10 µg/ml anti-CR3 Ab (AF1730, R&D SYSTEM)
48
49 or 10 µg/ml normal goat IgG control (AB-108-C, R&D SYSTEM); 1-5-10 µM DPI (NADPH
50
51 oxidase inhibitor, D2926, SIGMA); 0,1 mM Apocynin (NADPH oxidase inhibitor, 178385,
52
53
54
55
56
57
58
59
60

1
2
3 SIGMA). The CR3 receptor was also blocked using a combination of antibodies against both
4 the CD11b and CD18 subunits (SI Figure 3) as previously described [36]: LM2/1, a mouse
5 anti-human IgG1 monoclonal antibody to the I domain of CD11b chain of CR3 (10 µg/mL;
6 eBioscience); VIM12, a mouse monoclonal IgG1 anti-human antibody to the lectin domain of
7 CD11b chain of CR3 (10 µg/mL; invitrogen); and IB4, a mouse monoclonal IgG2a anti-
8 human antibody to the CD18 chain of CR3 (10 µg/mL; Calbiochem). Isotype controls, mouse
9 IgG1 (clone P3; eBioscience) and mouse IgG2a (clone eBM2A; eBioscience) were used at the
10 same concentration as the blocking antibodies. Positive controls for inhibitors that did not
11 show any effect in our system are depicted in SI Figure 7.
12
13
14
15
16
17
18
19
20
21
22
23

24 **Cytotoxicity assay**

25
26 To exclude any impact of cytotoxicity of the different inhibitory compounds used in this
27 study we measured lactate dehydrogenase (LDH) activity released from the cytosol of
28 damaged cells into the supernatant (SI Figure 5). LDH release was measured using the
29 Cytotoxicity Detection kit (Roche, Mannheim, Germany) according to the manufacturer's
30 instructions. Monocytes were pretreated with the indicated inhibitors for 30 min (see
31 “Stimulation experiments” for all the concentrations used) and exposed to CytD (1 µg/ml) for
32 1h. Monocytes were then stimulated with 1 µg/ml β-glucan C.a. for 24h before cell-free
33 culture supernatants were collected and incubated with LDH assay solution at 25°C for 30
34 min. The optical density values were analyzed at 490 nm by subtracting the reference value at
35 620 nm. The experiment was repeated at least three times and the results were expressed as a
36 percentage of the maximum LDH release, obtained by lysing the cells in 1% Triton X-100.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Immunoblot

Total cellular protein extracts were obtained by lysing 2×10^6 cells for 30 min at 4°C in lysis buffer (Pierce RIPA Buffer) in the presence of protease and phosphatase inhibitors (ROCHE). Samples were centrifuged for 15 minutes at 4°C, 15000 RPM and supernatants collected in new tubes. Cell culture supernatants were collected and precipitated in Trichloroacetate (TCA) 10% (v/v) for 1 hour on ice. Proteins were centrifuged for 30 min/4°C at 15000 RPM and pellets washed with ice-cold acetone. Samples were centrifuged for 15 min, 4°C, 15000 rpms. Proteins from cell extracts and supernatants were suspended in Laemli Buffer 1X and boiled at 95°C for 10 min. Proteins from cell extracts and supernatants were either mixed using a ratio 1:1 or individually kept. 30µL of each sample were allowed to migrate by SDS-PAGE and transferred with the iBlot apparatus (Life Technologies). Immunoblotting was performed by use of the following primary antibodies: anti-phospho-IKB (5A5), anti-IKB (44D4), anti-NLRP3 (D4D8T), anti-Caspase-1 (D7F10), anti-IL1β (3A6) (all from Cell Signalling). Anti-mouse or anti-rabbit IgG, HRP-linked antibodies (GE Healthcare) were used as secondary antibodies, and protein bands were visualized by use of ECL Prime (GE Healthcare) and a myECL imager (Thermo Fisher Scientific). Proteins were normalized for Glyceraldehyde-3-phosphate dehydrogenase (GAPDH, Santa Cruz Biotechnology) by use of ImageJ software (NIH, Bethesda, MD, USA).

Cytokine assay

TNFα, IL6 and IL1β levels on monocytes were determined by enzyme-linked immunosorbent assays (ELISA) on 24h monocyte cell culture supernatants according to manufacturer's instructions (R&D Systems). For IL10, supernatants from human cell cultures were collected and cytokine detection was performed using the ELISA MAX™ Standard Set (BioLegend), according to the manufacturer's instructions.

Reactive oxygen species measurements

For measurement of reactive oxygen species production, a luminol-enhanced chemiluminescence assay was used. 1×10^5 monocytes, resuspended in Hank's Balanced Salt Solution (HBSS, GIBCO), were seeded in a white 96-well assay plate (Corning) and pre-treated for 30 min with the vehicle or the inhibitors indicated. Cells were then treated for 1h with DMSO or cytochalasin D diluted in Hank's Balanced Salt Solution (HBSS). The β -glucan (1 $\mu\text{g/ml}$) and Luminol (50 μM) were added, and light emission was measured every 2 min for a period of 120 min using the Synergy H1 microplate reader (Biotek). The light emission levels are expressed as RLU (Relative Light Units = luminescence given by the luminometer). All measurements were performed at least in duplicate, depending on the number of cells available.

Fluorescence

Treated monocytes were fixed in 4% formaldehyde solution for 10 min at 37°C, then washed with phosphate buffered saline (PBS) and excess of formaldehyde solution was quenched for 10 min with 0.1M of Glycine. Cells were permeabilized with Saponin 0.2%/BSA 3% for 20 min at 4°C. Antibodies against p65 (1/800, L8F6, Cell Signalling), ASC (1/500, sc-22514-R, clone N.15, SantaCruz Biotechnologies) are incubated at 4°C in Saponin/BSA solution overnight. The day after, samples were incubated with secondary antibodies anti-mouse-488 and anti-rabbit-568 (Invitrogen). Hoechst (Invitrogen) is used at 1/5000 for nucleus staining. Coverslips were mounted using Vectashield mounting medium (Vectorlabs) and sealed. Cells were imaged using a spinning disk microscope with a 60X oil objective. Images are analyzed using Image J software.

Internalization assay

HK *C. albicans* were labeled with fluorescein isothiocyanate (FITC; SIGMA) for 1h in the dark and thoroughly washed with phosphate buffered saline (PBS) before incubating with the effectors. For β -glucan-FITC labelling, in brief, glucan preparations were suspended in FITC solution prepared in carbonate buffer (pH 9.6) (glucan:FITC concentration ratio of 5:1), incubated overnight at 4°C, washed with carbonate buffer (3x), with PBS (1x) and then suspended in PBS. Monocytes were then incubated with HK fungi or β -glucan for 2h. For quenching of not internalized fluorescein isothiocyanate (FITC)-particles, cells were treated for 3 min with trypan blue (0.2%; Molecular Probes, Life Technologies). Internalization was assessed with an EVOS FL Cell Imaging system (Thermo Fisher Scientific).

Statistics

In vitro monocyte experiments were performed at least 3 times, and the results were analyzed using a Wilcoxon matched-pairs signed-rank test (unless otherwise stated). Synergy was defined when ROS level produced upon the concomitant Cyt D and β -glucan treatment was significantly greater than the addition of ROS levels produced with Cyt D and β -glucan alone. A P value below 0.05 was considered statistically significant. All data were analyzed using GraphPad Prism software version 5.0 (La Jolla, CA, USA). Data are shown as means \pm standard errors of the means.

Acknowledgments

We thank Dr Molly Ingersoll for critical reading of the manuscript. We thank Rachel Lavenir for her technical help thorough the study. G.C. and J.Q. were supported by the Institut Carnot Pasteur MI grant ANR 11-CARN 017-01 and the ANR JCJC grant ANR-16-CE15-0014-01 (to J.Q.). E.E. and E.M. were supported by the FRM « Amorçage Jeunes Equipes » grant

AJE20151034460 and ATIP-Avenir Grant (to E.M.). This work was also supported, in part, by National Institutes of Health grants GM119197, GM53522 and GM083016 to D.L.W.

Conflict of Interest

The authors declare no financial or commercial conflict of interest

References

- [1] **Akira, S., Uematsu, S. and Takeuchi, O.**, Pathogen recognition and innate immunity. *Cell* 2006. **124**: 783-801.
- [2] **Gow, N. A., Netea, M. G., Munro, C. A., Ferwerda, G., Bates, S., Mora-Montes, H. M., Walker, L. et al.**, Immune recognition of *Candida albicans* beta-glucan by dectin-1. *J Infect Dis* 2007. **196**: 1565-1571.
- [3] **Taylor, P. R., Tsoni, S. V., Willment, J. A., Dennehy, K. M., Rosas, M., Findon, H., Haynes, K. et al.**, Dectin-1 is required for beta-glucan recognition and control of fungal infection. *Nat Immunol* 2007. **8**: 31-38.
- [4] **Gantner, B. N., Simmons, R. M. and Underhill, D. M.**, Dectin-1 mediates macrophage recognition of *Candida albicans* yeast but not filaments. *Embo J* 2005. **24**: 1277-1286.
- [5] **Herre, J., Marshall, A. S., Caron, E., Edwards, A. D., Williams, D. L., Schweighoffer, E., Tybulewicz, V. et al.**, Dectin-1 uses novel mechanisms for yeast phagocytosis in macrophages. *Blood* 2004. **104**: 4038-4045.
- [6] **Underhill, D. M., Rossnagle, E., Lowell, C. A. and Simmons, R. M.**, Dectin-1 activates Syk tyrosine kinase in a dynamic subset of macrophages for reactive oxygen production. *Blood* 2005. **106**: 2543-2550.
- [7] **Goodridge, H. S., Reyes, C. N., Becker, C. A., Katsumoto, T. R., Ma, J., Wolf, A. J., Bose, N. et al.**, Activation of the innate immune receptor Dectin-1 upon formation of a 'phagocytic synapse'. *Nature* 2011. **472**: 471-475.
- [8] **Brown, G. D., Herre, J., Williams, D. L., Willment, J. A., Marshall, A. S. and Gordon, S.**, Dectin-1 mediates the biological effects of beta-glucans. *J Exp Med* 2003. **197**: 1119-1124.
- [9] **Shah, V. B., Huang, Y., Keshwara, R., Ozment-Skelton, T., Williams, D. L. and Keshwara, L.**, Beta-glucan activates microglia without inducing cytokine production in Dectin-1-dependent manner. *J Immunol* 2008. **180**: 2777-2785.
- [10] **Dennehy, K. M., Ferwerda, G., Faro-Trindade, I., Pyz, E., Willment, J. A., Taylor, P. R., Kerrigan, A. et al.**, Syk kinase is required for collaborative cytokine production induced through Dectin-1 and Toll-like receptors. *Eur J Immunol* 2008. **38**: 500-506.
- [11] **Walachowski, S., Tabouret, G. and Foucras, G.**, Triggering Dectin-1-Pathway Alone Is Not Sufficient to Induce Cytokine Production by Murine Macrophages. *PloS one* 2016. **11**: e0148464.
- [12] **Goodridge, H. S., Shimada, T., Wolf, A. J., Hsu, Y. M., Becker, C. A., Lin, X. and Underhill, D. M.**, Differential use of CARD9 by dectin-1 in macrophages and dendritic cells. *J Immunol* 2009. **182**: 1146-1154.

- 1
2
3 [13] **Hernanz-Falcon, P., Joffre, O., Williams, D. L. and Reis e Sousa, C.,** Internalization of Dectin-1 terminates induction of inflammatory responses. *Eur J Immunol* 2009. **39**: 507-513.
- 4
5
6 [14] **Rosas, M., Liddiard, K., Kimberg, M., Faro-Trindade, I., McDonald, J. U., Williams, D. L., Brown, G. D. et al.,** The induction of inflammation by dectin-1 in vivo is dependent on myeloid cell programming and the progression of phagocytosis. *J Immunol* 2008. **181**: 3549-3557.
- 7
8
9
10 [15] **LeibundGut-Landmann, S., Gross, O., Robinson, M. J., Osorio, F., Slack, E. C., Tsoni, S. V., Schweighoffer, E. et al.,** Syk- and CARD9-dependent coupling of innate immunity to the induction of T helper cells that produce interleukin 17. *Nat Immunol* 2007. **8**: 630-638.
- 11
12
13
14 [16] **Yoshitomi, H., Sakaguchi, N., Kobayashi, K., Brown, G. D., Tagami, T., Sakihama, T., Hirota, K. et al.,** A role for fungal {beta}-glucans and their receptor Dectin-1 in the induction of autoimmune arthritis in genetically susceptible mice. *J Exp Med* 2005. **201**: 949-960.
- 15
16
17
18 [17] **Auffray, C., Sieweke, M. H. and Geissmann, F.,** Blood monocytes: development, heterogeneity, and relationship with dendritic cells. *Annu Rev Immunol* 2009. **27**: 669-692.
- 19
20
21
22 [18] **Ginhoux, F. and Jung, S.,** Monocytes and macrophages: developmental pathways and tissue homeostasis. *Nat Rev Immunol* 2014. **14**: 392-404.
- 23
24
25 [19] **Shi, C. and Pamer, E. G.,** Monocyte recruitment during infection and inflammation. *Nat Rev Immunol* 2011. **11**: 762-774.
- 26
27
28 [20] **Saeed, S., Quintin, J., Kerstens, H. H., Rao, N. A., Aghajani-refah, A., Matarese, F., Cheng, S. C. et al.,** Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science* 2014. **345**: 1251086.
- 29
30
31 [21] **van der Meer, J. W., Joosten, L. A., Riksen, N. and Netea, M. G.,** Trained immunity: A smart way to enhance innate immune defence. *Mol Immunol* 2015. **68**: 40-44.
- 32
33
34 [22] **Bain, J. M., Louw, J., Lewis, L. E., Okai, B., Walls, C. A., Ballou, E. R., Walker, L. A. et al.,** *Candida albicans* hypha formation and mannan masking of beta-glucan inhibit macrophage phagosome maturation. *MBio* 2014. **5**: e01874.
- 35
36
37 [23] **Gow, N. A., van de Veerdonk, F. L., Brown, A. and Netea, M. G.,** *Candida albicans* morphogenesis and the host defence: discriminating invasion from colonization. *Nat Rev Microbiol* 2012. **10**: 112-122.
- 38
39
40 [24] **Broz, P. and Dixit, V. M.,** Inflammasomes: mechanism of assembly, regulation and signalling. *Nat Rev Immunol* 2016. **16**: 407-420.
- 41
42
43 [25] **Gaidt, M. M., Ebert, T. S., Chauhan, D., Schmidt, T., Schmid-Burgk, J. L., Rapino, F., Robertson, A. A. et al.,** Human Monocytes Engage an Alternative Inflammasome Pathway. *Immunity* 2016. **44**: 833-846.
- 44
45
46 [26] **Hogquist, K. A., Nett, M. A., Unanue, E. R. and Chaplin, D. D.,** Interleukin 1 is processed and released during apoptosis. *Proc Natl Acad Sci U S A* 1991. **88**: 8485-8489.
- 47
48
49 [27] **Kayagaki, N., Warming, S., Lamkanfi, M., Vande Walle, L., Louie, S., Dong, J., Newton, K. et al.,** Non-canonical inflammasome activation targets caspase-11. *Nature* 2011. **479**: 117-121.
- 50
51
52 [28] **Shi, J., Zhao, Y., Wang, Y., Gao, W., Ding, J., Li, P., Hu, L. et al.,** Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature* 2014. **514**: 187-192.
- 53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- [29] **Vigano, E., Diamond, C. E., Spreafico, R., Balachander, A., Sobota, R. M. and Mortellaro, A.**, Human caspase-4 and caspase-5 regulate the one-step non-canonical inflammasome activation in monocytes. *Nat Commun* 2015. **6**: 8761.
- [30] **Gringhuis, S. I., Kaptein, T. M., Wevers, B. A., Theelen, B., van der Vlist, M., Boekhout, T. and Geijtenbeek, T. B.**, Dectin-1 is an extracellular pathogen sensor for the induction and processing of IL-1beta via a noncanonical caspase-8 inflammasome. *Nat Immunol* 2012. **13**: 246-254.
- [31] **Ganesan, S., Rathinam, V. A., Bossaller, L., Army, K., Kaiser, W. J., Mocarski, E. S., Dillon, C. P. et al.**, Caspase-8 modulates dectin-1 and complement receptor 3-driven IL-1beta production in response to beta-glucans and the fungal pathogen, *Candida albicans*. *J Immunol* 2014. **193**: 2519-2530.
- [32] **Tzeng, T. C., Schattgen, S., Monks, B., Wang, D., Cerny, A., Latz, E., Fitzgerald, K. et al.**, A Fluorescent Reporter Mouse for Inflammasome Assembly Demonstrates an Important Role for Cell-Bound and Free ASC Specks during In Vivo Infection. *Cell Rep* 2016. **16**: 571-582.
- [33] **Gurung, P., Burton, A. and Kanneganti, T. D.**, NLRP3 inflammasome plays a redundant role with caspase 8 to promote IL-1beta-mediated osteomyelitis. *Proc Natl Acad Sci U S A* 2016. **113**: 4452-4457.
- [34] **Schmitt, A., Grondona, P., Maier, T., Brandle, M., Schonfeld, C., Jager, G., Kosnopfel, C. et al.**, MALT1 Protease Activity Controls the Expression of Inflammatory Genes in Keratinocytes upon Zymosan Stimulation. *The Journal of investigative dermatology* 2016. **136**: 788-797.
- [35] **Netea, M. G., Brown, G. D., Kullberg, B. J. and Gow, N. A.**, An integrated model of the recognition of *Candida albicans* by the innate immune system. *Nat Rev Microbiol* 2008. **6**: 67-78.
- [36] **Bose, N., Wurst, L. R., Chan, A. S., Dudney, C. M., LeRoux, M. L., Danielson, M. E., Will, P. M. et al.**, Differential regulation of oxidative burst by distinct beta-glucan-binding receptors and signaling pathways in human peripheral blood mononuclear cells. *Glycobiology* 2014. **24**: 379-391.
- [37] **Drummond, R. A., Saijo, S., Iwakura, Y. and Brown, G. D.**, The role of Syk/CARD9 coupled C-type lectins in antifungal immunity. *Eur J Immunol* 2011. **41**: 276-281.
- [38] **Shah, V. B., Ozment-Skelton, T. R., Williams, D. L. and Keshvara, L.**, Vav1 and PI3K are required for phagocytosis of beta-glucan and subsequent superoxide generation by microglia. *Mol Immunol* 2009. **46**: 1845-1853.
- [39] **Kawai, T. and Akira, S.**, The roles of TLRs, RLRs and NLRs in pathogen recognition. *Int Immunol* 2009. **21**: 317-337.
- [40] **Blanco-Menendez, N., del Fresno, C., Fernandes, S., Calvo, E., Conde-Garrosa, R., Kerr, W. G. and Sancho, D.**, SHIP-1 Couples to the Dectin-1 hemITAM and Selectively Modulates Reactive Oxygen Species Production in Dendritic Cells in Response to *Candida albicans*. *J Immunol* 2015. **195**: 4466-4478.
- [41] **Filler, S. G., Pfunder, A. S., Spellberg, B. J., Spellberg, J. P. and Edwards, J. E., Jr.**, *Candida albicans* stimulates cytokine production and leukocyte adhesion molecule expression by endothelial cells. *Infect Immun* 1996. **64**: 2609-2617.
- [42] **Boxx, G. M., Kozel, T. R., Nishiya, C. T. and Zhang, M. X.**, Influence of mannan and glucan on complement activation and C3 binding by *Candida albicans*. *Infect Immun* 2010. **78**: 1250-1259.
- [43] **Elder, M. J., Webster, S. J., Chee, R., Williams, D. L., Hill Gaston, J. S. and Goodall, J. C.**, β -Glucan Size Controls Dectin-1-Mediated Immune Responses in

- 1
2
3 Human Dendritic Cells by Regulating IL-1 β Production. *Frontiers in immunology*
4 2017. **8**.
- 5 [44] **Warnatsch, A., Tsourouktsoglou, T. D., Branzk, N., Wang, Q., Reincke, S.,**
6 **Herbst, S., Gutierrez, M. et al.**, Reactive Oxygen Species Localization Programs
7 Inflammation to Clear Microbes of Different Size. *Immunity* 2017. **46**: 421-432.
- 8 [45] **Said-Sadier, N., Padilla, E., Langsley, G. and Ojcius, D. M.**, Aspergillus
9 fumigatus stimulates the NLRP3 inflammasome through a pathway requiring
10 ROS production and the Syk tyrosine kinase. *PloS one* 2010. **5**: e10008.
- 11 [46] **Lowman, D. W., Greene, R. R., Bearden, D. W., Kruppa, M. D., Pottier, M.,**
12 **Monteiro, M. A., Soldatov, D. V. et al.**, Novel structural features in Candida
13 albicans hyphal glucan provide a basis for differential innate immune recognition
14 of hyphae versus yeast. *J Biol Chem* 2014. **289**: 3432-3443.
- 15 [47] **Joly, S., Ma, N., Sadler, J. J., Soll, D. R., Cassel, S. L. and Sutterwala, F. S.**, Cutting
16 edge: Candida albicans hyphae formation triggers activation of the Nlrp3
17 inflammasome. *J Immunol* 2009. **183**: 3578-3581.
- 18 [48] **Mukaremera, L., Lee, K. K., Mora-Montes, H. M. and Gow, N. A. R.**, Candida
19 albicans Yeast, Pseudohyphal, and Hyphal Morphogenesis Differentially Affects
20 Immune Recognition. *Frontiers in immunology* 2017. **8**: 629.
- 21 [49] **Bonhomme, J. and d'Enfert, C.**, Candida albicans biofilms: building a
22 heterogeneous, drug-tolerant environment. *Curr Opin Microbiol* 2013. **16**: 398-
23 403.
- 24 [50] **Alonso, M. F., Gow, N. A., Erwig, L. P. and Bain, J. M.**, Macrophage Migration Is
25 Impaired within Candida albicans Biofilms. *J. Fungi* 2017. **3**: 31.
- 26 [51] **Aimanianda, V., Clavaud, C., Simenel, C., Fontaine, T., Delepierre, M. and**
27 **Latge, J. P.**, Cell wall beta-(1,6)-glucan of Saccharomyces cerevisiae: structural
28 characterization and in situ synthesis. *J Biol Chem* 2009. **284**: 13401-13412.
- 29 [52] **Aimanianda, V., Simenel, C., Garnaud, C., Clavaud, C., Tada, R., Barbin, L.,**
30 **Mouyna, I. et al.**, The Dual Activity Responsible for the Elongation and Branching
31 of beta-(1,3)-Glucan in the Fungal Cell Wall. *MBio* 2017. **8**.
- 32 [53] **Stephen-Victor, E., Karnam, A., Fontaine, T., Beauvais, A., Das, M., Hegde, P.,**
33 **Prakhar, P. et al.**, Aspergillus fumigatus cell wall alpha-(1,3)-glucan stimulates
34 regulatory T cell polarization by inducing PD-L1 expression on human dendritic
35 cells. *J Infect Dis* 2017.
- 36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legends

Figure 1. Effects of phagocytosis impairment on the response of human monocytes towards whole *Candida albicans* and the cell wall fraction composed of branched β -glucan. (A-B) Human monocytes were left untreated (ctrl) or stimulated with either heat-killed *Candida albicans* (HK-C.a., A) or heat-killed *Saccharomyces cerevisiae* (HK-S.c., B) in the presence (Cyt D) or in the absence (DMSO) of cytochalasin D. (C) Cells were stimulated with the Alkali-Soluble (AS) and Insoluble (AI) fractions, β -(1-6)-glucan and β -(1-3)/(1-6) glucan fractions from the cell wall of *Saccharomyces cerevisiae* after being treated with DMSO or Cyt D. (D) Data show the stimulation experiments of monocytes with two doses (1 or 10 $\mu\text{g/ml}$) of branched β -(1-3)-glucan isolated from *C. albicans* (β -glucan C.a.) or from *S. cerevisiae* (β -glucan S.c.) in the presence or in the absence of Cyt D. (E) Data show the stimulation experiments of monocytes with two doses (1 or 10 $\mu\text{g/ml}$) of insoluble linear β -(1-3)-glucan (curdlan) or of soluble branched β -(1-3)-glucan (laminarin) in the presence or in the absence of Cyt D. (F) The particulate branched β -(1-3)-glucan isolated from the cell wall of *C. albicans* was sonicated (Sonifier cell Disruptor B-30. Output control 4, Duty cycle 40%; 2 times 10 cycle with 5min on ice in between the two sets). Monocytes were then exposed to either sonicated or not-sonicated β -glucan in the presence or in the absence of Cyt D. (G) Cells pre-treated with DMSO or Cyt D were stimulated for 24h with the indicated concentrations of β -glucan from *C. albicans* (β -glucan C.a.), HK-C.a., AS and AI fractions from S.c., or the combination of AS+AI (10+10 $\mu\text{g/ml}$). (H) Monocytes were pre-treated with either DMSO (control), or with cytochalasin D (Cyt D) or latrunculin B (Lat B) (actin polymerization inhibitors), or with jasplakinolide (Jsp, actin polymerization inducer), or with chlorpromazine (CLP, clathrin-mediated endocytic inhibitor) or the lysosomal inhibitor cloroquine (CLQ). Cells were then stimulated with either branched β -(1-3)-glucan isolated

1
2
3 from *C. albicans* (β -glucan C.a.) or with heat-killed *Candida albicans* (HK-C.a.). For all the
4
5 experiments, culture supernatants were collected after 24h and concentration of secreted
6
7 TNF α , IL6, IL1 β and IL10 was determined by ELISA. Data were reproducible for all the four
8
9 cytokines tested even if histograms are not shown. Graphs show the mean \pm SEM of at least
10
11 three independent experiments. For (A, H), n = 8; for (B-G), n = 6; *p < 0.05, **p < 0.01.
12
13 Wilcoxon matched-pairs signed-rank test.
14
15
16
17

18 **Figure 2.** The role of the NLRP3-ASC-caspase-1 inflammasome in β -glucan induced IL1 β .
19
20 (A and E) Monocytes were pre-treated with either z-VAD-fmk (pan-caspase inhibitor), z-
21
22 YVAD-fmk (caspase-1 inhibitor), z-IETD-fmk (caspase-8 inhibitor), z-LEVD-fmk (caspase-
23
24 4/5 inhibitor), MCC950 (NLRP3 inhibitor), glibenclamide (inhibitor of potassium (K⁺)
25
26 efflux) or z-VRPR-fmk (MALT1 inhibitor) before treatment with either DMSO or Cyt D for
27
28 1h. Cells were then stimulated with branched β -(1-3)-glucan isolated from *C. albicans* (β -
29
30 glucan C.a.), culture supernatants were collected after 24h and secretion of IL1 β was
31
32 determined by ELISA. Graphs show the mean \pm SEM of four independent experiments (n=8).
33
34 **P < 0.001 (B) Monocytes were stimulated for 6h with or without β -glucan C.a. in the
35
36 presence or absence of Cyt D, as indicated. Caspase-1 (pro-form and cleaved p20), NLRP3
37
38 and IL1 β (pro- and mature IL1 β) expression was assessed by western blot. Arrows indicate
39
40 the specific band for NLRP3 and the mature and active form of IL1 β and caspase-1,
41
42 respectively. GAPDH was used as the loading control. Blots are representative of three
43
44 independent experiments. (C) Immunofluorescence microscopy of ASC inflammasomes.
45
46 Representative images of monocytes stimulated for 6h with or without β -glucan C.a. in the
47
48 presence or absence of Cyt D and stained with Hoechst in blue (DNA) and ASC in red.
49
50 Original magnification. 60x, Scale Bar: 50 μ m. (D) Quantification of the percentage ASC
51
52
53
54
55
56
57
58
59
60

1
2
3 specks from (C) (3x200 cells analysed with Image J). *P < 0.05 and **P < 0.01; 2-tailed
4
5 Student's t test.
6
7
8

9 **Figure 3.** The role of the Dectin-1/Syk/PI3K pathway in the overproduction of cytokines by
10 monocytes impaired for phagocytosis and stimulated with β -glucan. Monocytes were pre-
11 treated with anti-Dectin-1 Ab, anti-TLR2 Ab, anti-CR3 Ab or isotype control Abs in (A). Syk
12 inhibitor in (B), or PI3K inhibitor (WRT; wortmannin) in (C), were used on monocytes as
13 indicated. After the first incubation with inhibitors, cells were treated either with DMSO or
14 Cyt D and finally stimulated with branched β -(1-3)-glucan isolated from *C. albicans* (β -
15 glucan C.a.). Culture supernatants were collected after 24h and secretion of TNF α , IL6 and
16 IL1 β was determined by ELISA. Graphs show the mean \pm SEM of at least three independent
17 experiments. For (A), n = 6; for (B) and (C), n = 9; *p < 0.05, **p < 0.01. Wilcoxon matched-
18 pairs signed-rank test.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34


35 **Figure 4.** I κ B α phosphorylation, I κ B α degradation and p65 nuclear translocation in
36 cytochalasin D/ β -glucan stimulated monocytes. (A) Western blot analysis of phospho-I κ B α
37 and I κ B α in cell lysates from monocytes treated for 6h with or without β -glucan C.a. in the
38 presence or absence of Cyt D. GAPDH was used as a loading control. (B) Graphs represent
39 the densitometric analysis of the blots. Immunoreactive bands were normalized to GAPDH.
40 The bars represent mean values \pm SEM of the analysis of 4 independent experiments. (C)
41 Cells were treated as in (A) and nuclear translocation of p65 NF- κ B was visualized by
42 immunostaining with a specific antibody recognizing p65 and an alexa-fluor 488-conjugated
43 secondary antibody (green). For reference the nuclei are stained Hoescht (blue). One
44 representative image of three independent experiments is shown. (D) Quantification of nuclei
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 with p65 translocation (3x200 nuclei analysed with Image J). Original magnification. 60x,
4
5 scale bar: 50 μ M. *P < 0.05 and **P < 0.01; 2-tailed Student's t test.
6
7
8

9 **Figure 5.** Analysis of ROS production in human monocytes with impaired phagocytosis
10 mechanism upon β -glucan stimulation. Monocytes were pre-treated for 30 min with (A) the
11 indicated concentrations of DPI (NADPH oxidase inhibitor) before being treated with either
12 DMSO or Cyt D for 1h. Cells were then left untreated or stimulated with particulate β -glucan
13 C.a.. Luminol was added, and Reactive oxygen species (ROS) production was monitored by
14 chemiluminescence during time (2h) and expressed as relative light units (RLU). Graphs
15 show mean \pm SEM corresponding to up to three independent experiments (n=3); each
16 experimental condition was assessed in triplicate. (B) Monocytes were pre-treated with DPI,
17 as indicated. After 30 min of incubation, cells were treated either with DMSO or Cyt D and,
18 after an additional 1h, stimulated with β -glucan C.a.. Culture supernatants were collected after
19 24h and secretion of TNF α , IL6 and IL1 β was determined by ELISA. Graphs show the mean
20 + SEM of at least three independent experiments (n = 8); **p < 0.01. Wilcoxon matched-pairs
21 signed-rank test. (C-D) For inhibition, monocytes were pre-incubated for 30 min with the
22 vehicle (control), Syk inhibitor (2 μ M), anti-Dectin-1 Ab, or PI3K inhibitor (wortmannin) and
23 then treated as in (A). ROS production was monitored by chemiluminescence. Graphs show
24 mean + SEM corresponding to three independent experiments (n=5); each experimental
25 condition was assessed in duplicate or triplicate depending on the number of cells available.
26 (D) RLU data are plotted as histograms of mean + SEM of the maximal ROS production
27 corresponding to three independent experiments (n = 5). Significant synergy is represented as
28 # p<0,05, ## p<0,01. *P < 0.05, **P < 0.01 and ***P < 0.001; one-way ANOVA.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure 6.** Syk activity assessment for inflammasome, and NF- κ B activation in phagocytosis-
4 impaired-monocyte stimulated with *C. albicans*-branched β -(1-3)-glucan. (A) Monocytes
5 treated for 6h with β -glucan C.a. in the presence or absence of Cyt D and of the Syk inhibitor
6 (2 μ M). Nuclear translocation of p65 NF- κ B was visualized by immunostaining with a
7 specific antibody recognizing p65 and a alexa-fluor 488-conjugated secondary antibody
8 (green). For reference the nuclei are stained Hoescht (blue). Original magnification. 60x, scale
9 bar: 50 μ M Image is representative of three independent experiments. (B) Quantification of
10 nuclei with p65 translocation (4x50 nuclei analysed with Image J). (C) Cells were treated as
11 in (A). Caspase-1 (pro-form and cleaved p20), and IL1 β (pro- and mature IL1 β) expressions
12 were assessed by western blot. Mature IL1 β and caspase-1 were also detected in cell culture
13 supernatants. GAPDH was used as the loading control. Blots are representative of three
14 independent experiments.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **Figure 7.** Schematic model illustrating potential signaling pathways triggered by the binding
32 of β -glucan to dectin-1 in human monocytes with normal (left) or impaired phagocytic (right)
33 function. Attenuation of dectin-1 signaling is associated with the engulfment of the
34 receptor/ligand complex. By contrast, blocking of actin-dependent phagocytosis led to
35 increased cell activation. Black lines depict the interactions that normally occur upon
36 phagocytosis and β -glucan/dectin-1 interaction, whereas red lines indicate the proposed
37 signaling events that enhance the strength and efficacy of signal propagation.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60







Figure 2






Figure 4




Figure 5






Figure 6




Figure 7

Supporting Information

Supporting Information Figure 1. Internalization of FITC-labelled heat-killed *C. albicans* or β -glucans from *C. albicans* or from *S. cerevisiae* or curdlan, by monocytes in the presence or in the absence of cytochalasin D. Extracellular fluorescence is left unquenched or quenched by trypan blue. Original magnification, x20.

Supporting Information Figure 2. Differential Interference Contrast (DIC) of pure monocytes incubated with or without *C. albicans*-branched β -(1,3)-glucan in the presence or absence of Cytochalasin D (CytD). No aggregates of β -glucan can be observed with or without CytD. It is rather a network of external filaments of β -(1,3)-glucan that can be observed in the right bottom left, when phagocytosis is impaired. Original magnification, x20.

Supporting Information Figure 3. Monocytes were pre-treated with a cocktail of 3 anti-CR3 antibodies (3x α CR3) or isotype controls as described in Materials and Methods. After the first 30 min incubation with the anti-CR3 antibodies cocktail, cells were treated for 1h either with DMSO or Cyt D and finally stimulated with branched β -(1,3)-glucan isolated from *C. albicans* (β -glucan C.a.). Culture supernatants were collected after 24h and secretion of TNF- α , IL-6 and IL-1 β was determined by ELISA. Graphs show the mean \pm SEM of at least three independent experiments, n = 10; *p < 0.05.

Supporting Information Figure 4. Analysis of Cyt D plus β -glucan C.a.-induced oxidative burst and cytokine production in the presence of NADPH oxidases inhibitor apocynin. (A) Monocytes were pre-treated for 30 min with the vehicle (control) or apocynin (Apo; NADPH oxidase inhibitor) before being treated with either DMSO or Cyt D for 1h. Cells were then left


1
2
3 untreated or stimulated with particulate β -glucan C.a.. Luminol was added, ROS production
4
5 was monitored by chemiluminescence during time (2h) and expressed as relative light units.
6
7
8 Graphs show mean \pm SEM corresponding to up to three independent experiments. For each
9
10 condition, values were normalized to the average reading of the untreated control
11
12 (DMSO/Untreated/Vehicle). (B) Cells were treated as in (A), culture supernatants were
13
14 collected after 24h and secretion of TNF- α , IL-6 and IL-1 β was determined by ELISA.
15
16
17 Graphs show the mean \pm SEM of three independent experiments (n = 6). *p < 0.05; **p <
18
19 0.01.
20
21
22
23

24 **Supporting Information Figure 5.** Lactate dehydrogenase cytotoxicity assay for monocytes
25
26 after exposure to several inhibitors. Cells were pre-incubated for 30 min with the indicated
27
28 inhibitors before stimulation for 24h with Cyt D (1 μ g/ml) + β -glucan C.a. (1 μ g/ml). The
29
30 results were normalized to the maximum LDH release and expressed as fold change
31
32 compared to control cells. Graphs show the mean \pm SEM of at least three independent
33
34 experiments. **p < 0.01 and ***p < 0.001 versus control (none); one-way ANOVA.
35
36
37
38
39


40 **Supporting Information Figure 6.** (A) Monocytes were pre-treated for 30 min with either z-
41
42 LEVD-fmk (caspase-4/5 inhibitor) or z-YVAD-fmk (caspase-1 inhibitor) before treatment
43
44 with LPS (10 ng/ml) (LPS-EB Ultrapure, InvivoGen) for 24h. In (B), cells were pre-treated
45
46 with z-VRPR-fmk (MALT1 inhibitor) before stimulation with HK-C.a. (10^5 /ml) for 24h.
47
48 Culture supernatants were collected and secretion of IL-1 β was determined by ELISA.
49
50
51 Graphs show the mean \pm SEM of four independent experiments (n=7). *P < 0.05 and **P <
52
53 0.01. (C) macrophages were stimulated with z-IETD-fmk (caspase-8 inhibitor) or LPS
54
55 (200ng/mL) alone or in combination. IL-1 β release and cell death were assessed by ELISA
56
57 and LDH cytotoxicity assay kit (Roche, Mannheim, Germany), respectively. When indicated,
58
59
60

1
2
3 the cells were pre-treated with 40 μ M of necrostatin-1 (SIGMA) 30minutes prior the
4 stimulation to block the z-IETD+LPS-induced necroptotic cell death. Graphs show the mean
5
6
7 \pm SEM of 3 independent experiments (n=4 amount 3).
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56






1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56







1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

