



HAL
open science

Low prevalence of gamma-secretase complex gene mutations in a large cohort of predominantly Caucasian patients with Hidradenitis Suppurativa

Sabine Duchatelet, Snaigune Miskinyte, Maia Delage, Marie-Noelle Ungeheuer, Thi Lam, Farida Benhadou, Allard R.J. V. Vossen, Errol P Prens, Olivier Cogrel, Marie Beylot-Barry, et al.

► To cite this version:

Sabine Duchatelet, Snaigune Miskinyte, Maia Delage, Marie-Noelle Ungeheuer, Thi Lam, et al.. Low prevalence of gamma-secretase complex gene mutations in a large cohort of predominantly Caucasian patients with Hidradenitis Suppurativa. *Journal of Investigative Dermatology*, 2020, 10.1016/j.jid.2019.10.025 . pasteur-02547297

HAL Id: pasteur-02547297

<https://pasteur.hal.science/pasteur-02547297>

Submitted on 19 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Journal Pre-proof

Low prevalence of *gamma-secretase complex* gene mutations in a large cohort of predominantly Caucasian patients with Hidradenitis Suppurativa

Sabine Duchatelet, Snaigune Miskinyte, Maia Delage, Marie-Noëlle Ungeheuer, Thi Lam, Farida Benhadou, Véronique Del Marmol, Allard R.J. V. Vossen, Errol P. Prens, Olivier Cogrel, Marie Beylot-Barry, Céline Girard, Julien Vidil, Olivier Join-Lambert, Mélanie Parisot, Patrick Nitschké, Sylvain Hanein, Sylvie Fraitag, Hessel H. Van der Zee, Didier Bessis, Giovanni Damiani, Andrea Altomare, Yi-Hua Liao, Georgios Nikolakis, Christos C. Zouboulis, Aude Nassif, Alain Hovnanian

PII: S0022-202X(20)30224-4

DOI: <https://doi.org/10.1016/j.jid.2019.10.025>

Reference: JID 2330

To appear in: *The Journal of Investigative Dermatology*

Received Date: 29 July 2019

Revised Date: 23 September 2019

Accepted Date: 1 October 2019

Please cite this article as: Duchatelet S, Miskinyte S, Delage M, Ungeheuer M-N, Lam T, Benhadou F, Del Marmol V, Vossen ARJV, Prens EP, Cogrel O, Beylot-Barry M, Girard C, Vidil J, Join-Lambert O, Parisot M, Nitschké P, Hanein S, Fraitag S, Van der Zee HH, Bessis D, Damiani G, Altomare A, Liao Y-H, Nikolakis G, Zouboulis CC, Nassif A, Hovnanian A, Low prevalence of *gamma-secretase complex* gene mutations in a large cohort of predominantly Caucasian patients with Hidradenitis Suppurativa, *The Journal of Investigative Dermatology* (2020), doi: <https://doi.org/10.1016/j.jid.2019.10.025>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 The Authors. Published by Elsevier, Inc. on behalf of the Society for Investigative Dermatology.



Title page

Low prevalence of *gamma-secretase complex* gene mutations in a large cohort of predominantly Caucasian patients with Hidradenitis Suppurativa

Author List

Sabine Duchatelet^{1,2}, Snaigune Miskinyte^{1,2}, Maia Delage^{3,4}, Marie-Noëlle Ungeheuer⁵, Thi Lam³, Farida Benhadou^{4,6}, Véronique Del Marmol^{4,6}, Allard R. J. V. Vossen^{4,7}, Errol P. Prens^{4,7}, Olivier Cogrel⁸, Marie Beylot-Barry⁸, Céline Girard⁹, Julien Vidil³, Olivier Join-Lambert^{4,10,11,12}, Mélanie Parisot^{2,13}, Patrick Nitschke^{2,14}, Sylvain Hanein^{2,15}, Sylvie Fraitag¹⁶, Hessel H. Van der Zee^{4,7}, Didier Bessis⁹, Giovanni Damiani^{4,17,18}, Andrea Altomare^{17,18}, Yi-Hua Liao¹⁹, Georgios Nikolakis^{4,20}, Christos C. Zouboulis^{4,20}, Aude Nassif^{3,4}, Alain Hovnanian^{1,2,4,21}

Affiliations

¹Laboratory of Genetic Skin Diseases, INSERM U1163 Imagine Institute, Paris, France; ²Paris University, Paris, France; ³Medical Center, Pasteur Institute, Paris, France; ⁴European Hidradenitis Suppurativa Foundation e.V., Dessau, Germany; ⁵Clinical Investigation and Access to Research Bioresources (ICAReB) platform, Institut Pasteur, Paris, France; ⁶Department of Dermatology, Hopital Erasme - Université Libre de Bruxelles, Belgium; ⁷Department of Dermatology, Erasmus University Medical Center, Rotterdam, the Netherlands; ⁸Department of Dermatology, Centre Hospitalier Universitaire de Bordeaux, Bordeaux, France; ⁹Department of Dermatology, Saint-Eloi Hospital and Montpellier University Hospital, Montpellier, France; ¹⁰Microbiology Department, Necker-Enfants Malades Hospital, APHP, Paris, France; ¹¹Physiopathologie des Infections Systémiques, Necker-Enfants Malades Hospital, APHP, Paris, France; ¹²Microbiology Department, CHU de Caen, Normandy University, Caen, France; ¹³Genomics Core Facility, Imagine Institute-

Structure Fédérative de Recherche Necker, INSERM U1163 and INSERM US24/CNRS UMS3633, Paris, France; ¹⁴Bioinformatics Core Facility, INSERM U1163 Imagine Institute, Paris, France; ¹⁵Translational Genetic, Imagine Institute, UMR 1163, INSERM, Paris, France; ¹⁶Department of Pathology Necker–Enfants Malades Hospital, Paris, France ; ¹⁷I.R.C.C.S. Istituto Ortopedico Galeazzi, Milan, Italy; ¹⁸Department of Biomedical, Surgical and Dental Sciences, University of Milan, Milan, Italy; ¹⁹Department of Dermatology, National Taiwan University Hospital, Taipei, Taiwan; ²⁰Departments of Dermatology, Venereology, Allergology and Immunology, Dessau Medical Center, Brandenburg Medical School Theodor Fontane, Dessau, Germany; ²¹Department of Genetics, Necker–Enfants Malades Hospital, Assistance Publique des Hôpitaux de Paris, (AP-HP), Paris, France.

Corresponding Author:

Alain Hovnanian

Imagine Institute, Laboratory of Genetic skin diseases: from disease mechanism to therapy,
24 boulevard du Montparnasse, 75015 Paris

Phone: +33 (0)1 42 75 42 89

Fax : +33 (0)1 42 75 42 17

Email: alain.hovnanian@inserm.fr

ORCID:

Sabine Duchatelet: 0000-0003-2537-0385

Snaigune Miskinyte: 0000-0001-5865-3117

Maia Delage: 0000-0003-4310-7214

Marie-Noëlle Ungeheuer: 0000-0002-1639-9613

Thi Lam: 0000-0002-4659-3011

Farida Benhadou: 0000-0002-4533-8297

Véronique Del Marmol: 0000-0002-4359-0667

Allard R. J. V. Vossen: 0000-0002-1448-5972

Errol P. Prens: 0000-0002-8158-660X

Olivier Cogrel: 0000-0002-8878-8351

Marie Beylot-Barry: 0000-0001-6150-1229

Céline Girard: 0000-0002-9666-6354

Julien Vidil: 0000-0002-9963-5449

Olivier Join-Lambert: 0000-0002-4888-8630

Mélanie Parisot: 0000-0003-4312-2035

Patrick Nitschké: 0000-0002-2094-3298

Sylvain Hanein: 0000-0001-5817-6330

Sylvie Fraitag: 0000-0002-2258-2051

Hessel H. Van der Zee: 0000-0002-2874-7726

Didier Bessis: 0000-0002-3815-5417

Giovanni Damiani: 0000-0002-2390-6505

Andrea Altomare: 0000-0003-4171-9399

Yi-Hua Liao: 0000-0002-4434-3780

Georgios Nikolakis: 0000-0002-0920-9092

Christos C. Zouboulis: 0000-0003-1646-2608

Aude Nassif: 0000-0002-2407-0834

Alain Hovnanian: 0000-0003-3412-7512

Short title: *gamma-secretase complex* gene mutations in a HS cohort

Keywords: Hidradenitis Suppurativa, Dowling-Degos disease, gamma-secretase complex, targeted NGS, Genetics, *NCSTN*, *PSENEN*, *PSENI*, Crohn's disease, spondyloarthropathy

Abbreviations: HS, Hidradenitis Suppurativa; DDD, Dowling-Degos disease; GSC, gamma-secretase complex; PG, pyoderma gangrenosum; PASH, pyoderma gangrenosum, acne and suppurative hidradenitis; PAPASH, pyogenic arthritis, acne, pyoderma gangrenosum and suppurative hidradenitis; NGS, Next Generation Sequencing.

Letter text**To the Editor**

Hidradenitis suppurativa (HS) is a chronic dermatosis characterized by nodules and abscesses in apocrine gland-bearing sites. Mutations in three *gamma-secretase complex* (*GSC*) genes have been identified in autosomal dominant forms of HS (Frew et al., 2017) (Tables S1-S2). *NCSTN*, *PSENI* and *PSENE1* mutations have been reported in 41 unrelated HS patients, one Chinese HS family and 21 unrelated HS patients with or without Dowling-Degos disease (DDD) respectively (Frew et al., 2017). The functional consequences of these mutations are yet to be elucidated. In the absence of segregation analysis and functional studies supporting their causality, it is possible that some reported variants are benign (Table S2).

Three studies investigated *GSC* gene mutations in small patient cohorts (Table S1). One study explored *NCSTN*, *PSENI* and *PSENE1* in 48 British patients (Pink et al., 2012). A second study analyzed the six *GSC* genes (*NCSTN*, *PSENI*, *PSENE1*, *PSENE2*, *APHIA* and *APHIB*) in 20 patients from South Wales (Ingram et al., 2013). A third study investigated *NCSTN* in 95 Caucasian patients (Liu et al., 2016).

In contrast to our previous studies which focused on multiplex kindreds or in patients with syndromic HS and led to the identification of 4 *NCSTN* mutations (Duchatelet et al., 2019, Duchatelet et al., 2015, Miskinyte et al., 2012, Sonbol et al., 2018), the current study aimed at determining the prevalence of *GSC* gene mutations among a large cohort involving 169 unrelated mostly Caucasian patients, with isolated or syndromic HS. This study was approved by the institutional review board (CPP Ile-de-France Paris). Subjects provided written informed consent and consented to the publication of their images. Cohort characteristics are detailed in Tables 1 and S3. We performed mutational analysis of the six *GSC* genes using a targeted NGS gene panel, which also included *PSTPIP1* involved in

PASH (pyoderma gangrenosum (PG), acne, HS) and PAPASH (pyogenic arthritis, acne, PG, HS) syndromes, *FGFR2* and *POFUT1* possibly implicated in HS (Calderon-Castrat et al., 2016, Gonzalez-Villanueva et al., 2018, Higgins et al., 2017).

We identified two unreported *NCSTN* (NM_015331) mutations, p.Gly61Val (c.182G>T) and p.Gly576Val (c.1727G>T), in two unrelated familial cases (Patients 111 and 150), predicted to be damaging by computational tools (Table S4). The first mutation co-segregates with HS in the family (Figures 1 and S1). The proband's daughter carrying the familial mutation had an atypical presentation with multiple comedones outside HS predilection sites, suggesting the involvement of modifier genes, epigenetic and/or environmental factors. Co-segregation of the second mutation with the disease was not studied, as no additional family member was available. We also identified a previously reported *NCTSN* variant c.996+7G>A in a sporadic patient (Patient 2) (Pink et al., 2012). However, this variant was inherited from the unaffected father and the study of the subject's keratinocytes mRNA did not detect *NCSTN* splicing nor expression anomaly, suggesting it is a non-deleterious change. In a familial HS case (Patient 30), we identified a previously unreported *PSENI* (NM_000021) mutation p.Ser390Glufs*20 (c.1167_1168insGA), inherited from the mother with Crohn's disease and spondylarthropathy (Figures 1 and S1). We found *PSENE1* (NM_172341) mutations in 5 HS patients with DDD (Figures 1 and S1). In patient 157, we identified the previously reported c.168T>G p.Tyr56* Jewish Ashkenazi founder mutation, inherited from the mother with Jewish Ashkenazi ancestry. Three *PSENE1* mutations were not previously described: c.304T>A p.*102Argext*50 (Patient 158), c.166+2T>C (Patients 159 and 160) and c.66dup p.Phe23Valfs*98 (Patient 161). Haplotype analysis for patients 159 and 160 did not reveal a common haplotype, arguing against a

founder mutation (Figure 1). The c.166+2T>C mutation is predicted to abolish the splicing donor site (Table S4). No causal mutation was identified in the remaining 160 patients.

Functional studies on primary keratinocytes from *PSENE1*-mutated patients 157 and 158 showed a 2-fold reduction in *PSENE1* transcript levels in patient 157 compared with controls and a 25% reduction in patient 158 (Figure S2). Western-blot analysis revealed no mutated PEN2 protein in both patients, a low level of wild-type protein (~13%) in patient 157, but a normal level of wild-type protein in patient 158 (Figure S3). We could show impaired NCSTN maturation and stability of PSEN1 protein fragments and a reduction in *HES1* (Notch target) transcript levels in patient 157 compared with controls (Figures S2-S3), suggesting impaired GSC function and Notch signaling. In contrast, patient 158 displayed no detectable abnormality at the protein nor RNA levels (Figures S2-S3), although previous studies demonstrated the critical role of the length and sequence of the C-terminus of PEN2 for GSC activity (Prokop et al., 2005). Previous studies demonstrated that some *NCSTN* mutations did not impair GSC activity nor Notch signaling (Zhang and Sisodia, 2015), that the *PSEN1* p.Pro242Leufs*11 mutation may lead to enhanced Notch signaling in zebrafish (Newman et al., 2014) and that the total number of mature GSC was unaltered in fibroblast solubilized membrane fractions harboring *NCSTN* p.Glu333_Gln367del or *PSENE1* p.Phe23Valfs*98 mutations (Pink et al., 2016) (Table S2).

To date, Notch was used as a hallmark for GSC function, hypothesizing that Notch signaling was involved in HS pathogenesis. However, these findings suggest that the disease mechanisms are complex and that these mutations, together with the p.*102Argext*50 *PSENE1* mutation, may act through the alteration of a GSC catalytic activity independent mechanism, and/or impaired Notch-independent downstream signaling. Alternatively, mutations may become deleterious only in particular cells or conditions (temperature, pH or salt concentrations). Therefore, the cells which were investigated (fibroblasts or epidermal

keratinocytes) may not be appropriate and other cell types such as follicular keratinocytes or sebocytes could be more suitable to investigate HS pathophysiology. Further experiments to elucidate the functional consequences of these mutations on GSC activity and downstream pathways are required.

In summary, in this and previous studies, we have examined *GSC* gene mutations in a total of 188 unrelated HS patients and identified mutations in 8 out of 169 patients in the current study and in 4 out of 19 patients in our former studies. Our findings confirm that *GSC* gene mutations are uncommon in the largest HS cohort investigated so far (12 mutations/188 patients i.e. 6.4 %), further document *PSENE1* mutations in HS patients with DDD, report a European case of HS due to a *PSENI* mutation and expand the spectrum of *NCSTN*-associated clinical manifestations.

Data Availability Statement

Targeted Next Generation Sequencing datasets related to this article are hosted at the NCBI Sequence Read Archive under the accession code PRJNA552677.

Conflict of Interest

Professor del Marmol has received speaker and/or board fees from BMS, Abvie, and Sanofi not related to this publication. The remaining authors state no conflict of interest.

Acknowledgments

We are grateful to the patients and their relatives for participating in this study and to AFRH (French Association for Research on HS). We acknowledge the use of the bioresources of the Necker Imagine DNA biobank (BB-033-00065). We thank Cécile Masson for bioinformatics assistance, Amina Ait-Saadi and CRT-CC (Clinical Core of the Center for Translational Sciences) for regulatory support. This work was supported by the Fondation pour la Recherche Médicale (ROXANNE project, LMV20100519581) and the Société Française de Dermatologie. The Department of Dermatology, Hopital Erasme - Université Libre de Bruxelles, Belgium; the Department of Dermatology, Erasmus University Medical Center, Rotterdam, the Netherlands; the Department of Dermatology, Centre Hospitalier Universitaire de Bordeaux, Bordeaux, France; the I.R.C.C.S. Istituto Ortopedico Galeazzi, Milan, Italy; and the Departments of Dermatology, Venereology, Allergology and Immunology, Dessau Medical Center, Brandenburg Medical School Theodor Fontane, Dessau, Germany are health care units of the European Reference Network for Rare and Complex Skin Diseases (ERN Skin).

CRedit statement

Conceptualization: SD, AN, AH

Data Curation: not applicable

Formal Analysis: SD

Funding Acquisition: OJL, AN, AH

Investigation: SD, SM

Methodology: SD, SH

Project Administration: AN, AH

Resources: MD, MNU, TL, FB, VDM, ARJVV, EPP, OC, MBB, CG, JV, OJL, MP, PN, SH,
SF, HHVDZ, DB, GD, AA, YHL, GN, CCZ, AN

Software: not applicable

Supervision: AN, AH

Validation: SD, AN

Visualization: SD

Writing - Original Draft: SD

Writing - Review & Editing: all authors

References

- Calderon-Castrat X, Bancalari-Diaz D, Roman-Curto C, Romo-Melgar A, Amoros-Cerdan D, Alcaraz-Mas LA, et al. PSTPIP1 gene mutation in a pyoderma gangrenosum, acne and suppurative hidradenitis (PASH) syndrome. *The British journal of dermatology* 2016;175(1):194-8.
- Duchatelet S, Join-Lambert O, Delage M, Miskinyte S, Guet-Revillet H, Lam T, et al. Remission of chronic acne fulminans and severe hidradenitis suppurativa with targeted antibiotherapy. *JAAD case reports* 2019;5(6):525-8.
- Duchatelet S, Miskinyte S, Join-Lambert O, Ungeheuer MN, Frances C, Nassif A, et al. First nicastrin mutation in PASH (pyoderma gangrenosum, acne and suppurative hidradenitis) syndrome. *The British journal of dermatology* 2015;173(2):610-2.
- Frew JW, Vekic DA, Woods J, Cains GD. A systematic review and critical evaluation of reported pathogenic sequence variants in hidradenitis suppurativa. *The British journal of dermatology* 2017;177(4):987-98.
- Gonzalez-Villanueva I, Gutierrez M, Hispan P, Betlloch I, Pascual JC. Novel POFUT1 mutation associated with hidradenitis suppurativa-Dowling-Degos disease firm up a role for Notch signalling in the pathogenesis of this disorder. *The British journal of dermatology* 2018;178(4):984-6.
- Higgins R, Pink A, Hunger R, Yawalkar N, Navarini AA. Generalized Comedones, Acne, and Hidradenitis Suppurativa in a Patient with an FGFR2 Missense Mutation. *Frontiers in medicine* 2017;4:16.
- Ingram JR, Wood M, John B, Butler R, Anstey AV. Absence of pathogenic gamma-secretase mutations in a South Wales cohort of familial and sporadic hidradenitis suppurativa (acne inversa). *The British journal of dermatology* 2013;168(4):874-6.

- Liu M, Davis JW, Idler KB, Mostafa NM, Okun MM, Waring JF. Genetic analysis of NCSTN for potential association with hidradenitis suppurativa in familial and nonfamilial patients. *The British journal of dermatology* 2016;175(2):414-6.
- Miskinyte S, Nassif A, Merabtene F, Ungeheuer MN, Join-Lambert O, Jais JP, et al. Nicastrin mutations in French families with hidradenitis suppurativa. *The Journal of investigative dermatology* 2012;132(6):1728-30.
- Newman M, Wilson L, Verdile G, Lim A, Khan I, Moussavi Nik SH, et al. Differential, dominant activation and inhibition of Notch signalling and APP cleavage by truncations of PSEN1 in human disease. *Human molecular genetics* 2014;23(3):602-17.
- Pink AE, Dafou D, Desai N, Holmes O, Hobbs C, Smith CH, et al. Hidradenitis suppurativa: haploinsufficiency of gamma-secretase components does not affect gamma-secretase enzyme activity in vitro. *The British journal of dermatology* 2016;175(3):632-5.
- Pink AE, Simpson MA, Desai N, Dafou D, Hills A, Mortimer P, et al. Mutations in the gamma-secretase genes NCSTN, PSENEN, and PSEN1 underlie rare forms of hidradenitis suppurativa (acne inversa). *The Journal of investigative dermatology* 2012;132(10):2459-61.
- Prokop S, Haass C, Steiner H. Length and overall sequence of the PEN-2 C-terminal domain determines its function in the stabilization of presenilin fragments. *Journal of neurochemistry* 2005;94(1):57-62.
- Sonbol H, Duchatelet S, Miskinyte S, Bonsang B, Hovnanian A, Misery L. PASH syndrome (pyoderma gangrenosum, acne and hidradenitis suppurativa): a disease with genetic heterogeneity. *The British journal of dermatology* 2018;178(1):e17-e8.

Zhang X, Sisodia SS. Acne inversa caused by missense mutations in NCSTN is not fully compatible with impairments in Notch signaling. *The Journal of investigative dermatology* 2015;135(2):618-20.

Journal Pre-proof

Table 1. Patient characteristics

Sex:	
Male	74 (44%)
Female	95 (56%)
Ethnicity:	
Caucasian	107 (63%)
African	24 (14%)
Asian	3 (2%)
Other and mixed	23 (14%)
Unknown	12 (7%)
Form :	
Familial	86 (51%)
Sporadic	48 (28%)
False sporadic ¹	27 (16%)
Unknown	8 (5%)
Hurley stage severity score :	
1	62 (37%)
2	44 (26%)
3	56 (33%)
Unknown	7 (4%)
Body mass index (mean) :	26.4 (range 17.0-46.0)
<18.5	1 (<1%)
18.5-24.9	67 (40%)
25-29.9	54 (32%)
≥30	36 (21%)
Unknown	11 (6%)
Smoking history	115 (68%)
Age of HS onset (mean)	20 (range 1-55 years)
Syndromic Form :	13(8%)
PASH	5 (3%)
PAPASH	1 (<1%)
PASS	2 (1%)
HS + Dowling-Degos disease	5 (3%)
Comorbidities	
severe acne	19 (11%)
acne	34 (20%)
dissecting cellulitis of the scalp	6 (4%)
inflammatory bowel disease	18 (11%)
inflammatory rheumatologic diseases	15 (9%)
atopy, asthma, eczema	41 (24%)
psoriasis	2 (1%)
diabetes	14 (8%)
Adamantiades-Behçet's disease	1 (<1%)
Familial Mediterranean fever	1 (<1%)

¹“False sporadic” cases had in fact first or second-degree relatives who presented either a few abscesses/cysts localized to HS predilection sites or isolated pilonidal sinus, without completely fulfilling criteria for HS, and/or had inflammatory bowel disease, rheumatologic inflammatory disease or severe acne.


Of note, there was a bias in the recruitment of HS patients in this cohort since the enrollment mainly involved patients with severe forms of HS.

Journal Pre-proof


Figure 1. Genealogical trees of patients harboring mutations in *NCSTN*, *PSENI* or *PSENE1* genes

Arrows indicate the probands. Haplotype reconstruction with informative microsatellites markers (D19S416, D19S425, D19S893, D19S224 and D19S220) on chromosome 19q13.11-q13.13 flanking *PSENE1* was performed for patient 159 and her two affected daughters, as well as for patient 160 and one of his affected sister, as illustrated. No common haplotype for these two families was identified. Patient 160 belongs to a family in which some patients presented only HS and were unavailable for the genetic study. Of note, patient 2 *NCSTN* variant was excluded as there is no co-segregation of this splicing variant with HS and no effects on mRNA expression or splicing, arguing against a causal mutation.


NCSTN
p.Gly61Val




NCSTN
p.Gly576Val



PSEN1
p.Ser390Glufs*20




PSENEN
p.Tyr56*




PSENEN
p.*102Argext*50




PSENEN
c.166+2T>C



PSENEN
p.Phe23Valfs*98



PSENEN
c.166+2>T>C



- ● HS
- ● HS + Dowling-Degos disease
- ○ Individuals without clinical information
- * Individuals clinically and molecularly diagnosed

Markers	Mb	Band
Centromere		
D19S416	34.06	19q13.11
D19S425	35.49	
D19S893	35.57	19q13.12
<i>PSENEN</i>	36.23	
D19S224	36.52	
D19S220	38.43	19q13.13
Telomere		

Supplemental Table S1. Studies investigating gamma-secretase complex gene mutations in HS patients

Reference	Number of unrelated patients studied (Form)	Number of patients with <i>NCSTN</i> mutations (Form)	Number of patients with <i>PSENI</i> mutations (Form)	Number of patients with <i>PSENEP</i> mutations (Form)	Ethnicity
Wang et al., 2010	6 (Familial)	3 (Familial)	1 (Familial)	2 (Familial)	Chinese
Li et al., 2011	2 (1 Familial + 1 Sporadic)	2 (1 Familial + 1 Sporadic)	-	-	Chinese
Liu et al., 2011	2 (Familial)	2 (Familial)	-	-	Chinese
Pink et al., 2011	7 (Familial)	1 (Familial)	0	1 (Familial)	British
Miskinyte et al., 2012	14 (Familial)	3 (Familial)	0	0	French
Pink et al., 2012	48 (20 Familial + 28 Sporadic)	3 (Sporadic)	0	0	British (2) and Afro-Caribbean (1)
Haines et al., 2012	19	1	0	0	Singaporean Chinese in origin
Zhang et al., 2013	2 (Familial)	2 (Familial)	-	-	Chinese
Ingram et al., 2013	20 (12 Familial + 8 Sporadic)	0	0	0	South Wales
Jiao et al., 2013	1 (Familial)	1 (Familial)	-	-	Chinese
Nomura et al., 2013	10 (1 Familial + 9 Sporadic)	1 (Familial)	-	-	Japanese
Ma et al., 2013	1 (Familial)	1 (Familial)	-	-	Chinese
Nomura et al., 2014	1 (Familial)	1 (Familial)	-	-	Japanese
Chen et al., 2015	1 (Familial)	1 (Familial)	-	-	African American
Duchatelet et al., 2015	3 (2 Familial + 1 adopted child)	1 (Adopted child)	0	0	French
Yang et al., 2015	1 (Familial)	1 (Familial)	-	-	Chinese
Panmontha et al., 2015	2 (Familial)	-	-	2 (Familial)	Thai
Xu et al., 2016	2 (Familial)	2 (Familial)	-	-	Chinese
Liu et al., 2016	95 (57 Familial + 38 Sporadic)	2 (1 Familial + 1 Sporadic)	-	-	Australian, Canadian, Czech, Danish, French, German, Greeks, Hungarian, Dutch, Puerto Rican, Swedish, Swiss, Turkish and American
Liu et al., 2016	1 (Sporadic)	-	-	1 (Sporadic)	Chinese
Ratnamala et al., 2016	2 (Familial)	2 (Familial)	-	-	Indian
Xiao et al., 2016	1 (Familial)	1 (Familial)	-	-	Chinese
Faraji Zonooz et al.,	1 (Familial)	1 (Familial)	-	-	Iranian

2016					
Zhang et al., 2016	1 (Familial)	1 (Familial)	-	-	Chinese
Zhou et al., 2016	2 (Familial)	-	-	2 (Familial)	Chinese
Pink et al., 2016	1 (Familial)	1 (Familial)	-	-	British
Ralser et al., 2017	6 (3 Familial + 3 Sporadic)	-	-	6 (3 Familial + 3 Sporadic)	German (3), French (1), Indian (1), Thai (1)
Li et al., 2017	2 (Familial)	-	-	2 (Familial)	Chinese
Sonbol et al., 2018	1 (Sporadic)	0	0	0	French
Pavlovsky et al., 2018	4 (Familial)	-	-	4 (Familial)	Jewish Ashkenazi (4)
Kan et al., 2018	1 (Familial)	-	-	1 (Familial)	Japanese
Wu et al., 2018	1 (Familial)	1 (Familial)	-	-	Chinese
Shi et al., 2018	1 (Familial)	1 (Familial)	-	-	Chinese
Xiao et al., 2018	1 (Sporadic)	1 (Sporadic)	-	-	Chinese
Duchatelet et al., 2019	1 (Familial)	0	0	0	French
He et al., 2019	3 (Familial) ¹	3 (Familial) ¹	-	-	Chinese
Takeichi et al., 2019	1 (Familial)	1 (Familial)	-	-	Japanese

¹In total, 5 familial HS cases harboring *NCSTN* mutations were reported but two of them were previously published in Liu et al., 2011 and Xiao et al., 2016

Supplemental Table S2. Summary of gamma-secretase complex gene mutations reported in HS

Gene	Mutation		Familial segregation	Functional studies	Reference
	Nucleotide	Protein			
NCSTN	c.97G>A	p.Gly33Arg	Yes	None	Takeichi et al., 2019
	c.210_211delAG	p.Val72Tyrf*16 ¹	Yes	None	Liu et al., 2011
	c.218delC	p.Ile73Thrfs*3 ²	Yes	None	Wu et al., 2018
	c.223G>A	p.Val75Ile	Yes	<i>In vitro</i> assay in fibroblasts: no impact on PSEN1 endoproteolysis, Notch processing nor nuclear signaling	Zhang et al., 2013 / Zhang et al., 2015
	c.344_351del ³	p.Thr115Asnfs*20	n/a (adopted child)	None	Duchatelet et al., 2015
	c.349C>T	p.Arg117*	Yes/Yes/nr (familial)	Blood mRNA study: reduced NCSTN expression	Wang et al., 2010/Chen et al., 2015/Liu et al., 2016
	c.477C>A	p.Cys159*	Yes	Skin immunohistochemistry: reduced NCSTN staining; Skin mRNA and protein study: decreased NSCTN expression	Xiao et al., 2016 / He et al., 2019
	c.487delC	p.Gln163Serfs*39	Yes	mRNA study from peripheral blood mononuclear cells: significant reduction of transcript level	Miskinyte et al., 2012
	c.497C>A	p.Ser166*	Yes	None	Ma et al., 2014
	c.553G>A	p.Asp185Asn	nr (sporadic)	<i>In vitro</i> assay in fibroblasts: no impact on PSEN1 endoproteolysis, Notch processing nor nuclear signaling	Pink et al., 2012 / Zhang et al., 2015
	c.582+1delG	p.Cys195Lysfs*16 (exon 6 skipping) + p.Cys195_Gly332del (exons 6-8 skipping)	Yes	Peripheral lymphocytes mRNA study: marked reduction of NCSTN expression, abnormal splicing	Nomura et al., 2013
	c.617C>A	p.Ser206*	Yes	None	Shi et al., 2018

<i>NCSTN</i>	c.632C>G	p.Pro211Arg	Mutation not found in the patient' healthy relatives (sporadic)	<i>In vitro</i> assay in fibroblasts: no impact on PSEN1 endoproteolysis, Notch processing nor nuclear signaling	Li et al., 2011 / Zhang et al., 2015
	c.647A>C	p.Gln216Pro	Yes	<i>In vitro</i> assay in fibroblasts: functionally inactive mutant (on Notch signaling processing and PSEN1 endoproteolysis) but effect on activity negated when the mutant is co-expressed with the wild-type	Zhang et al., 2013 / Zhang et al., 2015
	c.686_687dup ⁴	p.Cys230Profs*32 ⁴	Yes	None	Ratnamala et al., 2016
	c.887A>G	p.Glu296Gly	Yes	None	Xu et al., 2016
	c.944C>T	p.Ala315Val	Yes	None	Zang et al., 2016
	c.978delG	p.Met326Ilefs*31	nr	None	Haines et al., 2012
	c.996+1G>A ⁵	p.Glu333_Gln367del (exon 9 skipping)	nr	Fibroblast mRNA study: abnormal transcript, reduced <i>NCSTN</i> expression; Fibroblast protein study: no mutant protein detected, decreased mature <i>NCSTN</i> in total cell fractions but no abnormality for <i>NCSTN</i> , <i>PSEN1</i> and <i>PEN2</i> expression in solubilized membrane preparations	Pink et al. 2016
	c.996+7G>A ⁶	p.?	nr (sporadic)	lymphoblast mRNA study: reduced <i>NCSTN</i> expression	Pink et al., 2012
	c.1101+1 G>A	No abnormal transcript	Yes	mRNA study: No abnormal transcript detected, reduced <i>NCSTN</i> expression	Pink et al., 2011
	c.1101+10 A>G ⁷	p.?	nr (sporadic)	lymphoblast mRNA study: no aberrant transcripts, normal <i>NCSTN</i> expression	Pink et al., 2012
	c.1229C>T	p.Ala410Val	nr (sporadic)	None	Liu et al., 2016
<i>NCSTN</i>	c.1258C>T	p.Gln420*	Yes	mRNA and protein studies	Jiao et al., 2013/Yang et al., 2015

				from whole skin, epidermis, dermis, keratinocytes or fibroblasts: reduced expression	
	c.1294C>T	p.Arg432*	Yes	Skin immunohistochemistry: reduced NCSTN staining; Skin mRNA and protein study: decreased NSCTN expression	He et al., 2019
	c.1300C>T	p.Arg434*	Yes/Yes	mRNA study from peripheral blood mononuclear cells: normal expression	Miskinyte et al., 2012 / Xu et al., 2016
	c.1352+1G>A	p.?	Yes/Yes	Skin immunohistochemistry: reduced NCSTN staining; Skin mRNA and protein study: decreased NSCTN expression	Liu et al., 2011 / He et al., 2019
	c.1534C>T	p.Gln512*	Yes	Skin immunohistochemistry: reduced NCSTN staining; Skin mRNA and protein study: decreased NSCTN expression	He et al., 2019
	c.1551+1G>A	p.Ala486_Thr517del	Yes	Blood mRNA study: Reduced <i>NCSTN</i> expression; abnormal splicing with exon 13 skipping	Wang et al., 2010
	c.1635C>G ³	p.Tyr545*	Yes	None	Faraji Zonooz et al., 2016
	c.1695T>G	p.Tyr565*	Yes	None	Li et al., 2011
	c.1702C>T	p.Gln568*	Yes (incomplete penetrance)	None	Nomura et al., 2014
	c.1752delG	p.Glu584Aspfs*44	Yes	Blood and skin mRNA study: reduced <i>NCSTN</i> expression; Skin protein study: decreased NSCTN expression; Skin immunohistochemistry: reduced NCSTN staining	Wang et al., 2010 / He et al., 2019
	c.1768 A>G	p.Ser590AlafsX3 and another product	Yes	mRNA study from peripheral blood mononuclear cells: abnormal splicing; no reduction of <i>NCSTN</i> transcript levels	Miskinyte et al., 2012
<i>NCSTN</i>	c.1800_1801delTG ⁸	p.Tyr600* ⁸	Yes	None	Ratnamala et al., 2016

	c.2584-2585delCA	Reduced expression	nr (sporadic)	Reporter assay in HEK293T cells: decreased expression through miR-155	Xiao et al., 2018
<i>PSENI</i>	c.725delC	p.Pro242Leufs*11	Yes	Blood mRNA study: reduced <i>PSENI</i> expression; Zebrafish study : increased Notch signaling	Wang et al., 2010 / Newman et al., 2014
	c.868+16G>T ^{9,10}	p.?	No (familial)	None	Ma et al., 2014
	c.953A>G ⁹	p.Glu318Gly	nr	None	Ingram et al. 2013
<i>PSENE1</i>	c.35T>A ¹¹	p.Leu12*	Yes	None	Ralser et al., 2017
	c.43_56del ¹¹	p.Cys15Profs*101 ¹²	Yes	None	Kan et al., 2018
	c.62-1G>C ¹³	p.Gly22_Tyr56del (exon 3 skipping)	Yes	Exon trapping: exon 3 skipping	Ralser et al., 2017
	c.66delG ¹¹	p.Phe23Leufs*46	Yes/ <i>De novo</i> mutation (sporadic)	None	Wang et al., 2010 / Liu et al., 2016
	c.66_67insG ¹¹	p.Phe23Valfs*98 ¹²	Yes	Blood and fibroblast mRNA study: reduced <i>PSENE1</i> expression; Fibroblast protein study: no mutant protein detected, reduced PEN2 expression in total cell fractions but no abnormality for NCSTN, PSEN1 and PEN2 expression in solubilized membrane preparations	Pink et al.2011 / Pink et al. 2016
	c.84dup	p.Leu29Serfs*92 ¹²	Yes	Leukocyte mRNA study: Increased <i>PSENE1</i> expression	Panmontha et al., 2015
	c.115C>T ¹¹	p.Arg39*	nr (sporadic)	Protein expression in HEK293T cells: no mutant protein	Ralser et al., 2017
	c.166+2T>C ^{11,14}	p.Gly22_Tyr56del (exon 3 skipping)	nr (sporadic)	mRNA study : abnormal splicing	Ralser et al., 2017
	c.167-2A>G ¹¹	p.?	Yes	None	Zhou et al., 2016

<i>PSENEN</i>	c.168T>G ¹¹	p.Tyr56*	Yes	mRNA study from keratinocytes: decreased <i>PSENEN</i> expression; Notch-responsive reporter assay in primary keratinocytes: decreased Luciferase activity	Pavlovsky et al., 2018
	c.194T>G ¹¹	p.Leu65Arg	Yes/Yes	None	Zhou et al., 2016 / Li et al., 2017
	c.229_230insCACCC ¹¹	p.Ile77Thrfs*45 ¹²	Yes	None	Li et al., 2017 (Zhang et al., unpublished data)
	c.279delC	p.Phe94Serfs*51 ¹²	Yes	None	Wang et al., 2010

Mutations are numbered according to the nomenclature of the Human Genome Variation Society, such that +1 is the A of the start codon (ATG) of the cDNA sequences of *NCSTN* (Genbank accession number NM_015331), *PSENI* (NM_000021), and *PSENEN* (NM_172341).

n/a: not applicable; nr: not reported

¹This mutation was initially reported as p.Thr70fsX18

²This mutation was initially reported as p.Pro73Leufs*15

³PASH syndrome

⁴This mutation was initially reported as c.687insCC p.Cys230Profs*31

⁵This mutation was initially reported as c.1125+1G>A

⁶This variant is unlikely to be causal according to our findings (absence of co-segregation with the disease and no effect on *NCSTN* splicing or expression)

⁷This variant was reported to be unlikely pathogenic based on mRNA study results

⁸This mutation was initially reported as c.1799delTG p.Leu600X

⁹These variants were reported to be unlikely pathogenic (absence of co-segregation with the disease, observed in healthy controls, predicted to be benign)

¹⁰This variant was initially reported as c.837+16G>T

¹¹HS associated with Dowling-Degos disease or hyperpigmentation

¹²Mutations predicting an altered and elongated protein

¹³Patients with this mutation presented Dowling-Degos disease associated or not with HS

¹⁴This mutation was initially reported as g.1412T>C

Supplemental Table S3. Clinical information for the 169 HS patients from the cohort

Patient number	Form	Sex	Geographic origin	Age	BMI	Smoking history	Age at onset	Hurley stage	Comorbidities
1	Familial	F	EU	40	20.2	Yes	18	1	atopy
2	False sporadic	M	EU	27	20.9	Yes	18	1	acne, atopy
3	Familial	M	EU	28	20.6	Yes	13	1	DCS, eczema
4	False sporadic	F	EU	51	29.0	Yes	25	1	atopy
5	Familial	F	EU	46	30.2	Yes	27	1	atopy
6	Familial	F	EU	30	34.9	Yes	13	1	acne, atopy
7	Familial	F	EU	47	26.7	Yes	15	1	severe acne, atopy, AS
8	Familial	F	AF	26	35.9	Yes	15	1	atopy, SAPHO syndrome
9	Familial	F	AF	27	21.9	No	7	1	atopy
10	Sporadic	F	EU	27	29.0	Yes	19	1	acne, atopy
11	Familial	F	EU	49	26.8	Yes	19	1	acne, atopy
12	Familial	F	EU	38	27.0	Yes	19	1	acne, atopy
13	Familial	F	EU	31	26.0	Yes	13	2	diabetes, atopy
14	Familial	M	AF	37	21.6	Yes	18	2	acne, atopy
15	Familial	M	AF	30	21.8	No	18	2	acne
16	Sporadic	F	EU	47	23.9	Yes	24	2	none
17	Sporadic	M	EU	47	31.4	Yes	24	2	severe acne, atopy
18	Sporadic	F	EU	35	28.1	Yes	10	2	acne, atopy
19	False sporadic	M	EU	35	25.0	Yes	24	2	acne, atopy
20	Familial	M	EU	26	22.0	Yes	12	2	acne, atopy
21	Sporadic	M	EU	31	21.0	Yes	24	2	acne, atopy
22	Sporadic	M	EU	49	20.6	Yes	12	2	severe acne, atopy
23	Familial	M	EU	28	27.8	Yes	17	2	severe acne, atopy
24	Familial	M	AF	58	27.5	Yes	18	2	diabetes, atopy, acne
25	Sporadic	M	EU	32	20.1	Yes	17	2	none
26	Familial	F	EU	34	35.0	Yes	16	2	SAPHO syndrome, atopy
27	Familial	F	EU	33	25.3	Yes	14	2	fibromyalgia

28	Familial	F	EU	39	31.2	No	13	2	Gougerot-Sjogren syndrome, atopy
29	Familial	F	EU	43	20.1	Yes	21	2	chronic diarrhea
30	Familial	M	EU	25	32.0	Yes	17	1	Crohn's disease, atopy
31	Familial	F	EU	48	33.0	Yes	29	1	craniopharyngioma and adrenal insufficiency, diabetes
32	Sporadic	M	EU	24	27.0	No	15	2	none
33	Sporadic	F	EU	31	21.3	Yes	14	1	unlabeled arthralgias, acne
34	Sporadic	M	AF	28	17.0	No	19	3	DCS, eczema
35	Sporadic	M	EU	38	24.5	Yes	22	1	severe acne, atopy
36	Sporadic	F	AF	35	25.0	No	16	3	breast cancer
37	Familial	F	AF	29	28.4	Yes	21	1	atopy, osteomalacia
38	Familial	F	AF	55	20.0	No	16	1	Systemic Lupus
39	Sporadic	M	EU	41	27.0	Yes	22	2	DCS
40	False sporadic	M	EU	19	28.2	No	11	1	acne
41	Sporadic	M	EU- Maghreb	34	22.2	Yes	26	1	Irritable bowel syndrome, asthma
42	False sporadic	M	EU	29	24.9	Yes	25	2	none
43	Sporadic	F	EU	36	28.1	Yes	17	1	AS, severe acne
44	Sporadic	F	AF	38	33.9	Yes	17	2	diabetes
45	False sporadic	F	EU	36	31.6	No	27	3	acne
46	False sporadic	F	EU	49	28.9	Yes	27	2	atopy
47	False sporadic	F	EU	25	19.5	No	19	2	Crohn's disease, acne, atopy
48	False sporadic	F	EU	41	25.2	Yes	30	3	Crohn's disease
49	False sporadic	F	EU	20	22.9	No	14	1	none
50	Familial	F	Saudi Arabia	27	39.3	No	9	3	anodontia
51	Familial	M	Maghreb	29	27.8	Yes	15	2	acne
52	Familial	F	n/a	20	n/a	n/a	16	3	n/a
53	Familial	M	EU	61	26.1	Yes	25	3	none
54	Familial	M	EU	62	20.3	Yes	40	3	none
55	Familial	M	EU	32	28.9	Yes	23	2	severe acne
56	Familial	M	EU	25	29.4	No	21	3	severe acne, atopy

57	Familial	M	EU	35	21.6	No	13	1	severe acne, atopy
58	Familial	M	EU	38	31.8	Yes	22	2	DCS
59	Familial	F	EU	41	31.6	Yes	15	1	Darier's disease
60	Familial	M	Maghreb	27	24.8	Yes	9	1	none
61	Familial	M	EU	30	29.4	Yes	16	2	none
62	Familial	F	EU	n/a	n/a	n/a	n/a	n/a	none
63	Familial	M	AF-EU	40	20.2	Yes	17	2	Crohn's disease, eczema
64	Familial	M	AF	63	31.0	No	33	3	diabetes, acne
65	False sporadic	M	EU	33	23.5	Yes	25	3	severe acne
66	False sporadic	M	EU	42	29.0	Yes	26	3	severe acne, diabetes, eczema
67	Familial	F	EU	44	28.3	Yes	15	3	atopy
68	n/a	M	EU	69	24.2	Yes	52	3	severe acne, DCS
69	Familial	M	EU	71	30.9	Yes	55	2	diabetes, atopy
70	Sporadic	M	EU	38	22.9	Yes	17	1	severe acne
71	Familial	F	EU	36	20.0	Yes	24	2	Crohn's disease
72	false Sporadic	F	n/a	34	n/a	n/a	15	3	plantar keratoderma, multiple sclerosis
73	Familial	M	AF	29	19.4	Yes	18	3	Crohn's disease
74	False sporadic	F	EU	31	20.1	Yes	16	2	AS
75	Sporadic	M	n/a	n/a	n/a	n/a	n/a	n/a	DCS
76	Familial	F	AF	31	23.6	No	16	3	breast cancer
77	Familial	F	AF	24	36.1	No	17	2	epidermolysis bullosa, atopy, eczema
78	Sporadic	M	AF	25	27.5	Yes	17	3	severe acne
79	Familial	M	Maghreb	62	23.9	Yes	34	3	acne
80	False sporadic	F	EU	38	29.5	Yes	12	3	acne
81	False sporadic	M	EU	38	29.5	Yes	31	3	acne, atopy
82	Familial	M	EU	35	22.6	Yes	15	1	acne
83	Familial	M	EU	52	25.7	Yes	17	3	none
84	Familial	F	EU	40	27.7	Yes	15	1	RA, atopy, psoriasis
85	False sporadic	F	EU	50	21.3	Yes	25	3	Crohn's disease, RA, Behçet's disease, APL syndrome

86	Sporadic	F	AF-EU	64	23.4	No	54	2	Crohn's disease, Steinert myotonia, diabetes
87	Familial	F	EU	45	32.5	Yes	15	1	none
88	Familial	F	EU	42	25.3	Yes	18	1	AS
89	n/a	F	Maghreb	32	19.9	Yes	11	1	none
90	False sporadic	F	EU	35	24.4	Yes	20	1	none
91	Familial	F	EU	52	21.0	Yes	30	1	acne
92	Familial	F	EU	37	23.7	Yes	15	1	Ulcerative colitis
93	Familial	M	EU-Maghreb	35	25.5	Yes	23	1	none
94	Familial	F	EU	35	21.0	Yes	20	1	none
95	False sporadic	F	EU	49	27.4	Yes	17	1	acne
96	n/a	F	AF	35	33.5	Yes	16	1	acne
97	Familial	M	Asia	48	32.6	No	15	1	diabetes, acne
98	Familial	F	EU	56	24.7	Yes	18	1	none
99	Familial	M	AF	31	21.9	No	19	2	none
100	Familial	M	EU	55	29.9	Yes	46	2	none
101	False sporadic	F	EU	39	28.7	Yes	28	2	AS
102	Sporadic	F	Maghreb	25	20.6	Yes	15	3	SAPHO syndrome
103	Familial	M	EU	15	28.0	No	13	3	none
104	Familial	M	AF	13	22.6	No	8	3	none
105	Familial	F	EU	30	27.6	Yes	22	3	Crohn's disease
106	Familial	F	AF	19	26.3	No	11	3	adrenal hyperplasia
107	Familial	M	EU	16	21.8	No	13	2	severe acne, craniostenosis
108	Familial	F	EU	25	23.9	No	11	3	none
109	False sporadic	M	EU	46	33.0	Yes	20	3	diabetes
110	Familial	M	EU	29	28.4	n/a	n/a	3	FMF
111	Familial	M	EU	51	32.7	No	17	1	none
112	Sporadic	M	EU	56	36.8	Yes	13	3	diabetes
113	Familial	F	n/a	26	n/a	n/a	n/a	n/a	none
114	Sporadic	F	AF	30	27.5	No	15	3	none
115	Familial	F	EU	40	21.5	No	19	2	Crohn's disease, AS, PG

116	Sporadic	F	n/a	33	n/a	n/a	31	3	none
117	Familial	F	EU-AF	24	23.2	Yes	17	3	severe acne, severe acne
118	Sporadic	M	EU	35	20.0	Yes	16	3	none
119	Sporadic	M	EU	44	21.9	No	12	3	Crohn's disease
120	n/a	F	Maghreb	48	22.0	No	47	3	none
121	Familial	F	AF	31	33.4	Yes	11	3	AS
122	Sporadic	F	AF	23	18.8	No	11	2	Crohn's disease
123	Familial	F	EU	21	23.1	Yes	13	2	Ulcerative colitis
124	Familial	F	Asia	17	20.7	No	16	3	severe acne
125	Sporadic	F	AF	46	23.5	Yes	19	1	none
126	Sporadic	M	n/a	29	30	Yes	20	1	severe acne
127	Sporadic	F	n/a	36	n/a	Yes	n/a	1	n/a
128	Sporadic	F	n/a	33	n/a	No	11	1	none
129	Familial	M	n/a	33	32.4	Yes	n/a	1	none
130	Familial	F	EU	71	27.1	Yes	22	1	none
131	Familial	F	EU	38	39.0	Yes	20	1	Crohn's disease
132	n/a	F	Iran	24	26.2	Yes	6	1	none
133	Familial	M	Maghreb	37	34.3	Yes	21	1	diabetes
134	Sporadic	F	EU	42	35.5	Yes	16	1	none
135	Familial	F	EU	33	28.9	Yes	20	1	none
136	Sporadic	F	Maghreb	31	35.5	No	22	1	none
137	Sporadic	F	AF	42	35.6	No	32	1	none
138	Sporadic	F	EU+AF	23	27.6	Yes	13	1	none
139	Sporadic	F	EU	28	22.1	Yes	22	1	severe acne
140	Sporadic	F	EU	42	27.7	Yes	20	1	none
141	Sporadic	F	EU	28	30.1	Yes	11	1	none
142	n/a	F	EU	33	28.6	Yes	18	1	none
143	Sporadic	F	EU	25	22.3	Yes	19	1	none
144	n/a	M	EU	29	24.8	Yes	15	3	none
145	false Sporadic	F	EU	26	23.2	Yes	15	3	none

146	Sporadic	F	Saudi Arabia	16	43.0	No	1	3	none
147	Sporadic	F	EU	25	21.7	Yes	15	3	none
148	Sporadic	M	EU	44	27.4	Yes	27	3	none
149	false Sporadic	M	Maghreb	71	22.7	Yes	40	3	none
150	Familial	F	EU	30	21.7	No	23	3	none
151	false Sporadic	M	n/a	30	n/a	Yes	26	1	none
152	Sporadic	F	Algeria	58	31.2	Yes	50	3	psoriasis
153	false Sporadic	M	EU	34	26.3	Yes	23	2	none
154	Familial	F	EU	47	25.1	No	22	3	AS
155	Familial	F	Maghreb	50	33.2	No	38	3	diabetes
156	false Sporadic	M	EU	29	23.1	Yes	22	3	Crohn's disease
157	Familial	F	EU	44	21.4	No	24	1	Dowling-Degos
158	n/a	M	EU	60	26.2	No	18	3	Dowling-Degos, diabetes, atopy
159	Familial	F	EU	66	25.4	No	13	n/a	Dowling-Degos
160	Familial	M	EU	54	42.4	n/a	15	n/a	Dowling-Degos
161	Sporadic	M	n/a	53	n/a	n/a	n/a	n/a	Dowling-Degos
162	Familial	F	EU	56	27	Yes	37	3	PG+Acne : PASH syndrome
163	Familial	M	Caribbean	35	19.9	Yes	30	3	PG+Acne : PASH syndrome
164	False sporadic	M	EU-Algeria-Morocco	22	19.6	Yes	14	3	PG+Acne+Spondyloarthritis : PASS syndrome
165	Sporadic	M	Morocco	47	27.2	Yes	n/a	2	PG+Acne : PASH syndrome
166	Sporadic	F	EU	40	20.8	Yes	18	3	PG+Acne+arthritis : PAPASH syndrome
167	Sporadic	M	EU	45	32.4	No	16	2	PG+Acne+Spondyloarthritis : PASS syndrome
168	Familial	M	n/a	22	n/a	n/a	n/a	n/a	PG+Acne : PASH syndrome
169	Sporadic	M	Asia	40	22.1	No	35	2	PG+Acne : PASH syndrome

BMI: Body Mass Index, DCS: Dissecting cellulitis of the scalp, AS: Ankylosing Spondylarthritis, RA: Rheumatoid arthritis, SAPHO: Synovitis, acne, pustulosis palmoplantaris, hyperostosis and osteitis; PG: Pyoderma gangrenosum and FMF: Familial Mediterranean fever, EU: Europe, AF: Africa

Journal Pre-proof

Supplemental Table S4. *In silico* prediction analysis of identified mutations

Gene	Mutation	Grantham score	Polyphen-2	SIFT	MutationTaster	Panther	MutationAssessor	AlignGVGD	SNAP2
<i>NCSTN</i>	p.Gly576Val	109	Probably damaging (score 0.998)	Deleterious (score 0.05)	Disease causing (prob 0.999)	probably damaging (preservation time 1368)	Impact function : medium (FI Score 2.745)	Most likely to interfere with function (Class 65)	effect (score 81)
<i>NCSTN</i>	p.Gly61Val	109	Probably damaging (score 0.999)	Deleterious (score 0.00)	Disease causing (prob 0.999)	probably damaging (preservation time 1628)	Impact function : medium (FI Score : 3.085)	Most likely to interfere with function (Class 65)	effect (score 93)
<i>PSENI</i>	p.Ser390Glufs*20	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
<i>PSENEEN</i>	p.Tyr56*	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
<i>PSENEEN</i>	p.*102Argext*50	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
<i>PSENEEN</i>	p.Phe23Valfs*98	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a


Gene	Mutation	SNP&GO	Condell	FATHMM	Provean	MutPred2	VEST-4	MutPred-LOF	DDIG	ENTPRISE-X
<i>NCSTN</i>	p.Gly576Val	Disease (prob 0.759)	Deleterious (score 0.632)	Pathogenic (score -1.56)	Deleterious (score -4.78)	Functional consequences (score 0.890)	pathogenic (score 0.955)	n/a	n/a	n/a
<i>NCSTN</i>	p.Gly61Val	Disease (prob 0.913)	Deleterious (score 0.656)	Pathogenic (score -5.29)	Deleterious (score -8.50)	Functional consequences (score 0.931)	pathogenic (score 0.997)	n/a	n/a	n/a
<i>PSENI</i>	p.Ser390Glufs*20	n/a	n/a	n/a	n/a	n/a	pathogenic (score 0.971)	No pathogenicity (score 0.445)	disease (score 0.935)	Deleterious (score 0.970)
<i>PSENEEN</i>	p.Tyr56*	n/a	n/a	n/a	n/a	n/a	pathogenic (score 0.887)	Pathogenicity (score 0.548)	disease (score 0.842)	Deleterious (score 0.973)
<i>PSENEEN</i>	p.*102Argext*50	n/a	n/a	n/a	n/a	n/a	benign (score 0.088)	No pathogenicity (score 0.380)	n/a	n/a
<i>PSENEEN</i>	p.Phe23Valfs*98	n/a	n/a	n/a	n/a	n/a	pathogenic (score 0.853)	No pathogenicity (score 0.439)	disease (score 0.883)	Deleterious (score 0.893)

Gene	Mutation	SSF	MaxEnt	NNSPLICE	GeneSplicer	HSF
<i>PSENFEN</i>	c.166+2T>C	Loss of donor site (71.05 ⇒ -)	Loss of donor site (7.39 ⇒ -)	Loss of donor site (0.72 ⇒ -)	Loss of donor site (4.70 ⇒ -)	Loss of donor site (81.46 ⇒ 54.63)

n/a: not applicable

Journal Pre-proof

Patient 158




Patient 30 mother




Patient 111 niece




Patient 157



Jo





Supplemental Figure Legends**Supplemental Figure S1. Histopathological and clinical features of HS patients with identified mutations**

Hematoxylin and eosin staining of affected skin from patient 158 with HS and Dowling-Degos disease carrying a *PSENEN* mutation, showed epidermal hyperplasia with elongated and hyperpigmented rete ridges.

Hyperpigmentation and numerous multipore comedones in the groin in Patient 30's affected mother (harboring *PSEN1* mutation and presenting with HS associated with Crohn's disease and ankylosing spondylitis). Nodules of the groin and the buttocks in Patient 111's affected niece (carrying *NCSTN* mutation). Hyperpigmentation of the intergluteal fold, the groin and the axillary regions in Patient 157 (affected with HS and Dowling-Degos disease, harboring a *PSENEN* mutation). Subjects consented to the publication of their images.

Supplemental Figure S2. Relative *PSENEN* and *HES1* transcripts abundance in *PSENEN*-mutated patients versus control primary human epidermal keratinocytes.

Quantitative RT-PCRs of keratinocyte RNA from Patients 157 (p.Tyr56*) and 158 (p.*102Argext*50) show (a) decreased *PSENEN* expression level in both patients compared to control RNA, suggesting nonsense-mediated mRNA decay and unstable mutant mRNA degradation respectively, and (b) reduction in *HES1* (used as a hallmark for Notch signaling activation) expression level in patient 157. *PGKI* was used as an endogenous control.

Supplemental Figure S3. Immunoblot of PEN2, NCSTN and PSEN1 in *PSENEN*-mutated patients and control primary human epidermal keratinocytes.

Western blot analysis of keratinocyte lysates from control subjects, Patients 157 and 158 probed for PEN2 (a), NCSTN (b) and PSEN1 (c) show decreased PEN2 levels, a significant reduction in mature NCSTN expression and in PSEN1 fragments in Patient 157, but no

abnormality for Patient 158. No mutated PEN2 protein was observed in patients 157 and 158 (size prediction ~7 kDa for p.Tyr96* and ~17 kDa for p.*102Argext*50). Unaltered PSEN1 endoproteolysis was noted in both patients. Actin was used as a loading control.

Journal Pre-proof

Supplementary references

- Barrandon Y, Green H. Three clonal types of keratinocyte with different capacities for multiplication. *Proc Natl Acad Sci U S A* 1987;84(8):2302-6.
- Chen S, Mattei P, You J, Sobreira NL, Hinds GA. gamma-Secretase Mutation in an African American Family With Hidradenitis Suppurativa. *JAMA dermatology* 2015;151(6):668-70.
- Faraji Zonooz M, Sabbagh-Kermani F, Fattahi Z, Fadaee M, Akbari MR, Amiri R, et al. Whole Genome Linkage Analysis Followed by Whole Exome Sequencing Identifies Nicastrin (NCSTN) as a Causative Gene in a Multiplex Family with gamma-Secretase Spectrum of Autoinflammatory Skin Phenotypes. *The Journal of investigative dermatology* 2016;136(6):1283-6.
- Goossens D, Moens LN, Nelis E, Lenaerts AS, Glassee W, Kalbe A, et al. Simultaneous mutation and copy number variation (CNV) detection by multiplex PCR-based GS-FLX sequencing. *Human mutation* 2009;30(3):472-6.
- He Y, Xu H, Li C, Zhang X, Zhou P, Xiao X, et al. Nicastrin/miR-30a-3p/RAB31 Axis Regulates Keratinocyte Differentiation by Impairing EGFR Signaling in Familial Acne Inversa. *The Journal of investigative dermatology* 2019;139(1):124-34.
- Itan Y, Shang L, Boisson B, Ciancanelli MJ, Markle JG, Martinez-Barricarte R, et al. The mutation significance cutoff: gene-level thresholds for variant predictions. *Nature methods* 2016;13(2):109-10.
- Jiao T, Dong H, Jin L, Wang S, Wang J. A novel nicastrin mutation in a large Chinese family with hidradenitis suppurativa. *The British journal of dermatology* 2013;168(5):1141-3.
- Kan T, Takahagi S, Shindo H, Tanaka A, Kawai M, Hide M. A unique clinical phenotype of a patient bearing a newly identified deletion mutation in the PSENEN gene along with the pathogenic serum desmoglein-1 antibody. *Clinical and experimental dermatology* 2018;43(3):329-32.
- Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nature genetics* 2014;46(3):310-5.
- Li C, Li W, Xu H, Zhang X, Su B, Zhang W, et al. PSENEN Mutation Carriers with Co-manifestation of Acne Inversa (AI) and Dowling-Degos Disease (DDD): Is AI or DDD the Subphenotype? *The Journal of investigative dermatology* 2017;137(10):2234-6.
- Li CR, Jiang MJ, Shen DB, Xu HX, Wang HS, Yao X, et al. Two novel mutations of the nicastrin gene in Chinese patients with acne inversa. *The British journal of dermatology* 2011;165(2):415-8.
- Liu Y, Gao M, Lv YM, Yang X, Ren YQ, Jiang T, et al. Confirmation by exome sequencing of the pathogenic role of NCSTN mutations in acne inversa (hidradenitis suppurativa). *The Journal of investigative dermatology* 2011;131(7):1570-2.
- Liu Y, Miao T, Ma J, Shao L, Luo S, Li Y, et al. PSENEN c.66delG in sporadic acne inversa. *European journal of dermatology : EJD* 2016;26(3):298-9.

- Nomura Y, Nomura T, Sakai K, Sasaki K, Ohguchi Y, Mizuno O, et al. A novel splice site mutation in NCSTN underlies a Japanese family with hidradenitis suppurativa. *The British journal of dermatology* 2013;168(1):206-9.
- Nomura Y, Nomura T, Suzuki S, Takeda M, Mizuno O, Ohguchi Y, et al. A novel NCSTN mutation alone may be insufficient for the development of familial hidradenitis suppurativa. *Journal of dermatological science* 2014;74(2):180-2.
- Panmontha W, Rerknimitr P, Yeetong P, Srichomthong C, Suphapeetiporn K, Shotelersuk V. A Frameshift Mutation in PEN-2 Causes Familial Comedones Syndrome. *Dermatology* 2015;231(1):77-81.
- Pink AE, Simpson MA, Brice GW, Smith CH, Desai N, Mortimer PS, et al. PSENEN and NCSTN mutations in familial hidradenitis suppurativa (Acne Inversa). *The Journal of investigative dermatology* 2011;131(7):1568-70.
- Pink AE, Dafou D, Desai N, Holmes O, Hobbs C, Smith CH, et al. Hidradenitis suppurativa: haploinsufficiency of gamma-secretase components does not affect gamma-secretase enzyme activity in vitro. *The British journal of dermatology* 2016;175(3):632-5.
- Ralser DJ, Basmanav FB, Tafazzoli A, Wititsuwannakul J, Delker S, Danda S, et al. Mutations in gamma-secretase subunit-encoding PSENEN underlie Dowling-Degos disease associated with acne inversa. *The Journal of clinical investigation* 2017;127(4):1485-90.
- Ratnamala U, Jhala D, Jain NK, Saiyed NM, Raveendrababu M, Rao MV, et al. Expanding the spectrum of gamma-secretase gene mutation-associated phenotypes: two novel mutations segregating with familial hidradenitis suppurativa (acne inversa) and acne conglobata. *Experimental dermatology* 2016;25(4):314-6.
- Shi TW, Bai N, Zhang JA, Lu F, Chen XB, Kong XD, et al. Mutations in the gamma-secretase genes PSEN1, PSENEN, and NCSTN in a family with acne inversa. *European journal of dermatology : EJD* 2018;28(3):374-6.
- Takeichi T, Matsumoto T, Nomura T, Takeda M, Niwa H, Kono M, et al. A novel NCSTN missense mutation in the signal peptide domain causes hidradenitis suppurativa, which has features characteristic of an autoinflammatory keratinization disease. *The British journal of dermatology* 2019.
- Wang B, Yang W, Wen W, Sun J, Su B, Liu B, et al. Gamma-secretase gene mutations in familial acne inversa. *Science* 2010;330(6007):1065.
- Wu C, Yang J, Zhang S, Li J, Jin H, Zhang X. A novel NCSTN gene mutation in a Chinese family with acne inversa. *Molecular genetics and genomics : MGG* 2018;293(6):1469-75.
- Xiao X, He Y, Li C, Zhang X, Xu H, Wang B. Nicastrin mutations in familial acne inversa impact keratinocyte proliferation and differentiation through the Notch and phosphoinositide 3-kinase/AKT signalling pathways. *The British journal of dermatology* 2016;174(3):522-32.
- Xiao YJ, Yang Y, Liang YH. hsa-miR-155 targeted NCSTN 3'UTR mutation promotes the pathogenesis and development of acne inversa. *International Journal of Clinical and Experimental Pathology* 2018;11(4):1878-89.

- Xu H, Xiao X, Hui Y, Zhang X, He Y, Li C, et al. Phenotype of 53 Chinese individuals with nicastrin gene mutations in association with familial hidradenitis suppurativa (acne inversa). *The British journal of dermatology* 2016;174(4):927-9.
- Yang JQ, Wu XJ, Dou TT, Jiao T, Chen XB, Min M, et al. Haploinsufficiency caused by a nonsense mutation in NCSTN underlying hidradenitis suppurativa in a Chinese family. *Clinical and experimental dermatology* 2015;40(8):916-9.
- Zhang C, Wang L, Chen L, Ren W, Mei A, Chen X, et al. Two novel mutations of the NCSTN gene in Chinese familial acne inverse. *Journal of the European Academy of Dermatology and Venereology : JEADV* 2013;27(12):1571-4.
- Zhang S, Meng J, Jiang M, Zhao J. Characterization of a Novel Mutation in the NCSTN Gene in a Large Chinese Family with Acne Inversa. *Acta dermato-venereologica* 2016;96(3):408-9.
- Zhou C, Wen GD, Soe LM, Xu HJ, Du J, Zhang JZ. Novel Mutations in PSENEN Gene in Two Chinese Acne Inversa Families Manifested as Familial Multiple Comedones and Dowling-Degos Disease. *Chinese medical journal* 2016;129(23):2834-9.
- Zouboulis CC, Del Marmol V, Mrowietz U, Prens EP, Tzellos T, Jemec GB. Hidradenitis Suppurativa/Acne Inversa: Criteria for Diagnosis, Severity Assessment, Classification and Disease Evaluation. *Dermatology* 2015a;231(2):184-90.
- Zouboulis CC, Desai N, Emtestam L, Hunger RE, Ioannides D, Juhasz I, et al. European S1 guideline for the treatment of hidradenitis suppurativa/acne inversa. *Journal of the European Academy of Dermatology and Venereology : JEADV* 2015b;29(4):619-44.

Supplemental Table Legends**Supplemental Table S1. Studies investigating gamma-secretase complex gene mutations in HS patients**

¹In total, 5 familial HS cases harboring *NCSTN* mutations were reported but two of them were previously published in Liu et al., 2011 and Xiao et al., 2016

Supplemental Table S2. Summary of gamma-secretase complex gene mutations reported in HS

Mutations are numbered according to the nomenclature of the Human Genome Variation Society, such that +1 is the A of the start codon (ATG) of the cDNA sequences of *NCSTN* (Genbank accession number NM_015331), *PSENI* (NM_000021), and *PSENE1* (NM_172341).

n/a: not applicable; nr: not reported

¹This mutation was initially reported as p.Thr70fsX18

²This mutation was initially reported as p.Pro73Leufs*15

³PASH syndrome

⁴This mutation was initially reported as c.687insCC p.Cys230Profs*31

⁵This mutation was initially reported as c.1125+1G>A

⁶This variant is unlikely to be causal according to our findings (no co-segregation with the disease and no effect on *NCSTN* splicing or expression)

⁷This variant was reported as unlikely pathogenic based on mRNA study results

⁸This mutation was initially reported as c.1799delTG p.Leu600X

⁹These variants were reported as unlikely pathogenic (no co-segregation with the disease, observed in healthy controls, predicted to be benign)

¹⁰ This variant was initially reported as c.837+16G>T

¹¹HS associated with Dowling-Degos disease or hyperpigmentation

¹²mutations predicting an altered and elongated protein

¹³patients with this mutation presented Dowling-Degos disease associated or not with HS

¹⁴This mutation was initially reported as g.1412T>C

Supplemental Table S3. Clinical information for the 169 HS patients from the cohort

BMI: Body Mass Index, DCS: Dissecting cellulitis of the scalp, AS: Ankylosing Spondylarthritis, RA: Rheumatoid arthritis, SAPHO: Synovitis, acne, pustulosis palmoplantaris, hyperostosis and osteitis; PG: Pyoderma gangrenosum and FMF: Familial Mediterranean fever, EU: Europe, AF: Africa

Table S4. *In silico* prediction analysis of identified mutations

n/a: not applicable

Supplementary Methods

Clinical diagnostic criteria

Diagnosis of HS was made according to European guidelines (Zouboulis et al., 2015a, Zouboulis et al., 2015b) including recurrent or suppurating typical lesions (nodules, abscesses, sinus tracts or hypertrophic scars) in typical localizations (axillae, inguinal folds, anal cleft, sub-mammary regions) occurring more than twice in six months.

Study subjects

The DNA bioresources originated from three protocols dedicated to studies of Hidradenitis suppurativa which were all approved by French Ethical Committee (CPP Ile de France I, CPP Ile de France 3). All protocols were parallelly approved by the French drug agency (AFFSSAPS/ANSM). Written informed consents were obtained from all subjects included. Samples and associated data were managed by the ICAReB platform at Pasteur Institute or by Necker Imagine DNA biobank following all applicable regulations and standards. The probands and their relatives provided a blood sample. Patients 2, 157 and 158 provided punch biopsies of skin. De-identified normal skin from surgical margins was obtained for controls.

Targeted Next Generation Sequencing

A custom SureSelect gene panel was designed using the SureDesign software (Agilent H. sapiens, hg19, GRCh37, February 2009). The target regions covered 751.97 kb (panel v1) or 866.10 kb (panel v2), including coding exons and splice junctions of the 185 genes (panel v1: 2613 exons) or 210 genes (panel v2: 2803 exons) respectively. Illumina compatible precapture barecoded genomic DNA libraries were constructed according to the manufacturer's protocol (Ovation Ultralow, Nugen Technologies). Briefly, genomic DNA (1 - 3µg) was mechanically fragmented by sonication with a Covaris S2 Ultrasonicator, ligated to multiplexing Illumina compatible paired-end adapters, amplified by PCR with indexed (barcoded) primers for sequencing, and hybridized to biotinylated complementary 120-bp

RNA custom-designed capture probes in a solution-based reaction. Hybridization was performed at 65 °C for at least 16 h, followed by paired-end sequencing on the Illumina HiSeq 2500 (130 x130 bases) or on a NovaSeq 6000 (100 x 100 bases) at the Genomics Core Facility at the Imagine Institute.

Data analysis was performed with Paris Descartes University/Imagine Institute's Bioinformatics Core facility. After demultiplexing, sequences were aligned to the reference human genome hg19 using the Burrows-Wheeler Aligner. Downstream processing was carried out with the Genome Analysis Toolkit (GATK), SAMtools and Picard, following documented best practices (<http://www.broadinstitute.org/gatk/guide/topic?name=best-practices>). Variant calls were made with the GATK, Unified Genotyper, freebayes and SAMtools based on the 72nd version of ENSEMBL database. Variants were annotated and analyzed using the in-house PolyDiag software interface which filters out irrelevant and common polymorphisms based on frequencies extracted from public databases (dbSNP, Exome Variant Server and Exome Aggregation Consortium). Consequences of mutations on protein function were predicted using Polyphen2 (<http://genetics.bwh.harvard.edu/pph2>), SIFT (<https://sift.bii.a-star.edu.sg/>) and MutationTaster (www.mutationtaster.org). Mutations were then ranked on the basis of the predicted impact of each variant by combined annotation-dependent depletion (CADD) (Kircher et al., 2014), and compared with the mutation significance cut-off, a gene-level specific cut-off for CADD scores (<http://pec630.rockefeller.edu:8080/MS>) (Itan et al., 2016). To evaluate Copy number variations (CNVs, i.e duplication and large deletion events, for each individual) the relative read count for each targeted region was determined as the ratio of the read count for that region divided by the total absolute read counts of all targeted regions of the design. The ratio of the relative read count of a region in a given individual over the average relative read

counts in other individuals of the run resulted in the estimated copy number for that region in that individual (method adapted from Goossens et al., 2009).

The mean depth of coverage per sample was >300X (except for patients 51, 148, 150, 152 and 155 for whom it was comprised between 267X and 298X) permitting to maximize the detection of variant in regions with less coverage and to enable accurate Copy Number Variant Analysis of the exons of the panel. In average, 99% of the targeted exonic bases were covered at least with 30 independent reads.

Prioritization of the variants was performed thanks to the PolyDiag interface. We excluded known variants with a minor allele frequency $\geq 5\%$ listed in databases or variants previously identified in “in-house” exomes. We subsequently selected for variants affecting splice sites or coding regions (nonsynonymous, nonsense, frameshift, start/stop gain or loss) and predicted to be damaging by *in silico* prediction tools. Additional prediction tools were then used to predict the pathogenicity of candidate missense variants identified (Panther, MutationAssessor, AlignGVGD, SNAP2, SNP&GO, Condel, FATHMM, Provean and MutPred2). Other computational tools were also used (VEST-4, MutPred-LOF, DDIG and ENTPIRE-X). To evaluate the consequence of the splicing-site mutation, we used *in silico* prediction tools (SSF, MaxEnt, NSPLICE, GeneSplicer and HSF).

Microsatellites genotyping

Primers sequences of microsatellite markers (available at the Working Draft of the Human Genome available at UCSC, Human Assembly (GRCh37/hg19)) used for haplotype analysis at *PSENN* locus (19q13.12) are listed below.

D19S416 (AFM304XF5) (5'-3'): F=CCTGTCCCAGAGAGACCCTA,

R=AAGAGAGTGTGCCATTTGCT;

D19S425 (AFMA139WE9) (5'-3'): F=CCACAGGTGTGCATAAAAG,

R=GCCATGTGACTGTAGCAGA;

D19S893 (AFMB004WH1) (5'-3'): F=AATCCTGAGACTGGGGG,

R=TGGTGACACACTGGTGAC;

D19S224 (AFM240VC1) (5'-3'): F=AACACCATTCCTCATCTTCC,

R=CCCAGGCCCTATCTGA;

D19S220 (AFM214YF12) (5'-3'): F=ATGTTTCAGAAAGGCCATGTCATTTG,

R=TCCCTAACGGATACACAGCAACAC.

FAM-labelled amplified fragments were electrophoresed on an Applied Biosystems 3500XL genetic analyser using GeneScan™ 400 HD ROX™ Size Standard (Life technologies) and analyzed using the Gene-Mapper Software 5 (Applied Biosystems, Foster City, CA).

Cell culture

4 mm punch biopsies were explanted to isolate primary epidermal keratinocytes, which were maintained in Green medium on a feeder layer of lethally irradiated 3T3 mouse fibroblasts as described previously (Barrandon and Green, 1987). For experiments, keratinocytes were grown in 0.06 mM CaCl₂ EpiLife medium (Invitrogen) to 70-80% confluence.

RNA isolation from primary human epidermal keratinocytes and RT-PCR

Subconfluent cultured keratinocytes were harvested, RNA was prepared via RNeasy Mini (Qiagen), and cDNA was prepared with random hexamer priming and SuperScript IV reverse transcriptase (LifeTechnologies). PCR was performed in triplicate for each sample with qPCR MesaGreen Mastermix (Eurogentec) on a 7500 Sequence Detection (Applied Biosystems). Six control subjects were used, giving similar results. The experiment was replicated at least three times. PGK1 was used as an endogenous control.

RT-PCR primers

Quantitative PCR was performed with *PSENE1* or *HES1* and normalization to *PGK1*.

PSENE1 (5'-3'): F=ATGAACCTGGAGCGAGTGTC,

R=GTGTAGGCTGGGACAAGGAA;

HES1 (5'-3'): F=CCAAAGACAGCATCTGAGCA, R=AGAATGTCCGCCTTCTCCA;

PGK1 (5'-3'): F=CTGTGGCTTCTGGCATAACCT, R=AATCTGCTTAGCCCGAGTGA.

Journal Pre-proof

Protein isolation from primary human epidermal keratinocytes and immunoblotting

Keratinocytes were lysed in buffer (50 mM pH 8 Tris-HCl, 150 mM NaCl, 1% Nonidet P-40, 5 mM pH 8 EDTA pH8, Complete protease inhibitors [Roche]), incubated on ice for 30 min and sonicated for 2 min. Protein extracts were isolated by centrifugation at 13000 g at 4°C for 20 minutes. 30 µg samples were loaded in Laemmli buffer (62.5 mM pH 6.8 Tris HCl, 5% β-mercaptoethanol, 2% SDS, 10% glycerol, 0.002% bromophenol blue), separated by SDS-PAGE, transferred onto Nitrocellulose membrane (Trans-Blot Turbo Nitrocellulose, BioRad), and incubated with primary and secondary antibodies, with chemiluminescence detection (ECL Plus Western Blotting Substrate, Pierce) using the ChemiDoc Imaging System (BioRad). β-actin served as loading control. Experiments were replicated three times. Two control subjects were used, giving similar results.

Antibodies: Primary antibodies included rabbit polyclonal anti-PEN2 (Enzo ADI-905-736), rabbit polyclonal anti-NCSTN (Sigma N1660) and mouse monoclonal anti-PSEN1 (Chemicon international MAB5232). Secondary antibodies were HRP-linked anti-rabbit or anti-mouse IgG antibodies (Cell Signaling #7074 and #7076).

Statistical analyses: Statistical significance of observed differences was assessed by two-tailed Mann-Whitney U tests. Means with s.e.m. are shown.