

## Legionella Effectors Explored with INSeq: New Functional Insights

Monica Rolando, Carmen Buchrieser

## ▶ To cite this version:

Monica Rolando, Carmen Buchrieser. Legionella Effectors Explored with INSeq: New Functional Insights. Trends in Microbiology, 2018, 26 (3), pp.169-170. 10.1016/j.tim.2018.01.003 . pasteur-02552011

## HAL Id: pasteur-02552011 https://pasteur.hal.science/pasteur-02552011

Submitted on 16 Jun2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Legionella effectors explored with INSeq: new functional insights
2	
3	Monica Rolando <sup>1,2</sup> and Carmen Buchrieser <sup>1,2,*</sup>
4	
5	
6	<sup>1</sup> Institut Pasteur, Biologie des Bactéries Intracellulaires, Paris, France
7	<sup>2</sup> CNRS UMR 3525, Paris, France
8	
9	
10	
11	
12	
13	
14	Keywords
15	Legionella pneumophila, bacterial effectors, transposon insertion sequencing, virulence
16	factors
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	*
29	For correspondence:
30	Carmen Buchrieser
31	Biologie des Bacteries Intracellulaires, Institut Pasteur
32	28, rue du Dr. Koux, $75724$ Paris Cedex 15, France
33	1  el:  +33.1.45.68.83.72
34	E-mail: <i>cbuch(a)pasteur.fr</i>

- 35 Abstract

*Legionella pneumophila* secretes over 300 effector proteins that manipulate host cells. This
multiplicity of effectors hampers the characterization of their individual roles. Shames *et al.*report a new approach to solve the enigma of *Legionella* effector function by using INSeq to
analyse effector functions in the context of infection.

45

46 Understanding how bacterial pathogens modulate host pathways and hence counteract 47 host defence and cause disease is fundamental in our fight against infectious diseases. The 48 intracellular pathogen *Legionella pneumophila*, the causative agent of Legionnaires' disease, 49 is a unique model to study pathogen manipulation of host pathways during infection, with its 50 sophisticated arsenal of molecular weapons that are secreted into the host cell to counteract 51 host defences.

52 L. pneumophila is primarily an environmental bacterium that replicates within 53 protozoan hosts (amoeba). When Legionella-contaminated aerosols are inhaled the bacterium 54 can replicate within alveolar macrophages and cause an atypical pneumonia in susceptible 55 individuals. L. pneumophila utilizes the same processes to parasitize human or protozoan cells 56 as the L. pneumophila life cycle has been shown to be identical in all host cell types 57 examined: the bacterium is internalized in a membrane-bound compartment called the 58 Legionella containing vacuole (LCV) that escapes from lysosomal degradation, with the LCV 59 remodeled through recruitment of host endoplasmic reticulum-derived vesicles into a 60 protective vacuole for bacterial replication [1].

61 All key steps of the establishment of the LCV and the ability to replicate 62 intracellularly are dependent on a functional Dot/Icm type IV secretion system (T4SS). This highly conserved machinery is essential for virulence as it secretes over 300 effector proteins 63 64 into the host cell [2]. Genome sequencing has revealed a considerable plasticity and diversity in the effector repertoires encoded by different L. pneumophila strains, suggesting that this 65 diversity allows adaptation to the different hosts L. pneumophila can parasitize [3]. Uniquely, 66 67 among the over 300 T4SS effectors are an abundance of proteins with high similarity to eukaryotic proteins or proteins encoding eukaryotic-like domains, suggesting that molecular 68 69 mimicry of eukaryotic proteins is a main virulence strategy of L. pneumophila [4]. Thus, the 70 broad spectrum of tools employed for bacterial survival and replication in eukaryotic cells 71 seems to have evolved during co-evolution of L. pneumophila and its protozoan hosts by 72 horizontal gene transfer among domains of life, allowing this bacterium to parasitize a wide 73 variety of hosts.

Although the T4SS is strictly necessary to initiate and sustain intracellular replication of *L. pneumophila*, the contribution of individual effectors remains minor [5]. Indeed, after the first excitement of scientists that *L. pneumophila* may encode as many functions as effectors, it soon became evident that high functional redundancy was present, and that most of the effectors did not show a virulence phenotype in the standard infection models used. Furthermore, although a few effectors analyzed individually have been shown to play a role in
intracellular replication of *L. pneumophila*, an efficient method to systematically assess the
contribution of each effector was missing.

82 Shames and colleagues recently developed a new approach to define the contribution 83 of the many L. pneumophila effector proteins during infection using a high-throughput transposon insertion sequencing technique called INSeq [6]. Guided by the principle of 84 signature tagged mutagenesis, a number of groups have previously developed multiplex 85 screening systems that combined random transposon mutagenesis and next generation 86 87 sequencing (TnSeq) [7]. These comprehensive approaches allow high-resolution functional screening of whole genomes to identify key genes across various conditions *in vitro* or during 88 89 host colonization. However, a limitation of in vivo experiments is that during the establishment of infection a considerable number of bacteria can be killed or removed 90 91 stochastically; thus one cannot distinguish whether the bacteria were eliminated by chance or because they were indeed less fit. The approach of Shames et al. to circumvent such 92 93 bottlenecks was to reduce the complexity of the input pool, establishing an effector mutant pool (EMP) library where 297 of the 315 genes predicted to encode effector proteins were 94 95 represented. Subsequently they used the INseq technique to study the effects of this EMP on virulence phenotypes in amoeba, bone marrow-derived macrophages (BMDMs), and mice. 96 97 This comprehensive approach allowed them to cluster individual mutants in distinct categories depending on their fitness phenotype. Mutants showing a universal defect, a 98 99 mammalian-, lung-, BMDM-, or amoeba-specific virulence defect were identified. Indeed, 100 effectors already known to play a role in L. pneumophila virulence were selected (e.g. mavN 101 [8]), confirming the power of the technique to identify proteins implicated in virulence. 102 Moreover, virulence phenotypes were detected for a large number of uncharacterized effector 103 mutants, opening new vistas on the characterization of secreted L. pneumophila proteins. Due 104 to the unbiased approach, effector proteins that contributed both positively and negatively to 105 bacterial fitness could also be identified.

Another strength of the study was that the effector mutants were analyzed in three different host systems, which opened the possibility of evaluating the contribution of individual effectors in specific host contexts. An example is LegC4, an effector that, when deleted, conferred a competitive advantage on the mutant during *in vivo* murine infections, but not for *ex vivo* cellular infections. This was likely due to a role of LegC4 in stimulating pro-inflammatory responses *in vivo* that drive enhanced host control of the infection.

Furthermore, new metaeffectors were identified. First described by Kubori and 112 colleagues [9], metaffectors are defined as effectors that act on another effector to modulate 113 114 its function in the host cell. The characterization of metaeffectors is an important step in 115 understanding L. pneumophila-host interactions as uncovering their spatiotemporal regulation 116 provides insight on how the bacterium modulates diverse host cell functions at discrete stages 117 of infection. Shames and colleagues showed that Lpg2505 is a metaeffector that regulates the previously characterized effector SidI [10]. Interestingly, as with all metaeffectors identified 118 119 to date, *lpg2505* constitutes an operon with *sidI*, and suppresses SidI toxicity to prevent host damage that would limit L. pneumophila replication efficiency. 120

In summary, INSeq analyses of a defined effector mutant pool in three different infection models allowed virulence phenotypes to be assigned to multiple effectors for which functions were previously unknown. Furthermore, effectors having distinct impact in different host models, or metaeffectors and effectors that act negatively on fitness can be identified. This new approach to identifying previously cryptic effector functions can now be applied across multiple bacterial pathogens and infection systems, opening the door to systematic deconvolution of host-pathogen interactions.

- 128 References
- 129
- Isberg, R.R. et al. (2009) The *Legionella pneumophila* replication vacuole: making a cosy
   niche inside host cells. Nat Rev Microbiol 7 (1), 13-24.
- 132 2. Ensminger, A.W. (2016) *Legionella pneumophila*, armed to the hilt: justifying the largest
  133 arsenal of effectors in the bacterial world. Curr Opin Microbiol 29, 74-80.
- 3. Gomez-Valero, L. and Buchrieser, C. (2013) Genome dynamics in *Legionella*: the basis of versatility and adaptation to intracellular replication. Cold Spring Harb Perspect Med 3 (6), a009993.
- 4. Cazalet, C. et al. (2004) Evidence in the *Legionella pneumophila* genome for exploitation
  of host cell functions and high genome plasticity. Nat Genet 36 (11), 1165-1173.
- 5. O'Connor, T.J. et al. (2011) Minimization of the *Legionella pneumophila* genome reveals
  chromosomal regions involved in host range expansion. Proc Natl Acad Sci U S A 108 (36), 14733-40.
- 6. Shames, S.R. et al. (2017) Multiple *Legionella pneumophila* effector virulence phenotypes
  revealed through high-throughput analysis of targeted mutant libraries. Proc Natl Acad Sci U S A 114 (48), E10446-E10454.
- 7. van Opijnen, T. and Camilli, A. (2013) Transposon insertion sequencing: a new tool for
  systems-level analysis of microorganisms. Nat Rev Microbiol 11 (7), 435-442.
- 8. Portier, E. et al. (2015) *IroT/mavN*, a new iron-regulated gene involved in *Legionella pneumophila* virulence against amoebae and macrophages. Environ Microbiol 17 (4), 1338-50.
- 9. Kubori, T. et al. (2010) *Legionella* metaeffector exploits host proteasome to temporally
   regulate cognate effector. PLoS Pathog 6 (12), e1001216.
- 152 10. Shen, X. et al. (2009) Targeting eEF1A by a *Legionella pneumophila* effector leads to
  153 inhibition of protein synthesis and induction of host stress response. Cell Microbiol 11 (6),
  154 911-926.
- 155