



## On hepatocellular carcinoma in South America and early-age onset of the disease

Pascal Pineau, Eloy Ruiz, Eric Deharo, Stephane Bertani

### ► To cite this version:

Pascal Pineau, Eloy Ruiz, Eric Deharo, Stephane Bertani. On hepatocellular carcinoma in South America and early-age onset of the disease. Clinics and Research in Hepatology and Gastroenterology, 2019, 43 (5), pp.522-526. 10.1016/j.clinre.2018.10.019 . pasteur-02641989

**HAL Id: pasteur-02641989**

**<https://pasteur.hal.science/pasteur-02641989>**

Submitted on 20 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1    **On hepatocellular carcinoma in South America and early-age onset of**  
2    **the disease**

3

4    Pascal Pineau<sup>1\*</sup>, Eloy Ruiz<sup>2</sup>, Eric Deharo<sup>3</sup>, Stéphane Bertani<sup>3\*</sup>

5

6    <sup>1</sup> Institut Pasteur, Unité Organisation Nucléaire et Oncogenèse, INSERM, U  
7    993, Paris, France

8    <sup>2</sup> Instituto Nacional de Enfermedades Neoplásicas, Departamento de Cirugía  
9    en Abdomen, Lima, Peru

10   <sup>3</sup> Université de Toulouse, IRD, UPS, UMR 152 PHARMADEV, Toulouse,  
11   France

12   \*Correspondence: P.P. (email: [pascal.pineau@pasteur.fr](mailto:pascal.pineau@pasteur.fr)) and S.B.  
13   ([stephane.bertani@ird.fr](mailto:stephane.bertani@ird.fr))

14

## **Summary**

Hepatocellular carcinoma (HCC) is one of the most predominant tumor types worldwide, being particularly prevalent in sub-Saharan Africa and East Asia. However, HCC is inexplicably underreported in South America, despite unsettling clinical epidemiological trends of the disease on this continent. Here, we review the current knowledge on HCC presentation in Peru. We emphasize the well documented occurrence of an early-age nosological form of the disease in Andean descent populations. We further discuss the reasons for such clinical presentation, as well as the implications for liver cancer screening, management, and prevention.

## **Keywords**

Liver cancer; Cancer risk factor; Hepatitis B virus (HBV); Indigenous people; Low- and middle-income countries; Global health transition

## **Abbreviations**

AFP, alpha-fetoprotein; CELAC, Community of Latin American and Caribbean States; HBV, Hepatitis B virus; HBsAg, HBV surface antigen; HCC, Hepatocellular carcinoma; HCV, Hepatitis C virus; INEN, National Cancer Institute of Peru

## Introduction

Hepatocellular carcinoma (HCC), the main form of primary liver cancer, is the seventh most common malignancy in incidence and the third leading cause of tumor-related death in the world [1]. Global clinical epidemiology of HCC reported hitherto chiefly delineates a prominent patient profile corresponding grossly to males over 45 years old with chronic liver diseases [2]. The incidence rate of HCC has doubled worldwide during the last two decades, with nearly 85% of the recorded cases and highest rates of disability-adjusted life-years occurring in low- and middle-income countries [3–5]. The largest burdens of HCC are borne in sub-Saharan Africa and East Asia, where the highly endemic chronic infection with hepatitis B virus (HBV) and dietary exposure to mutagenic aflatoxins potentialize one another [6,7]. In contrast, the incidence of HCC observed in more economically developed countries is associated most of all with hepatitis C virus (HCV) infection and heavy alcohol intake that are often associated with comorbid conditions, such as non-alcoholic fatty liver disease, diabetes mellitus, hereditary hemochromatosis or even  $\alpha$ -1 antitrypsin deficiency [6,8–10]. These risk factors trigger overtime liver cirrhosis that progresses in a significant proportion of cases in a hepatocarcinogenic process.

Reviews of the global burden of liver cancer have repetitively overlooked the epidemiology of the disease in South America creating a gap between the existing situation in the field and the overall clinical epidemiology of HCC described in the relevant literature [2,11]. Yet, while the incidence rate of primary liver cancer in South America is considered low to intermediate, the

epidemiological trends and the clinical presentation of HCC on this continent are displaying striking and worrying features as a whole. For example, South America is part of the Community of Latin American and Caribbean States (CELAC), which is the world region with the greatest incidence rise for liver cancer monitored during the last decade [12]. Furthermore, some physicians have reported on the continent, since the seventies, the dual occurrence of an early-onset form of HCC in younger individuals concomitantly with a more conventional older patient population.

#### **Clinical epidemiological peculiarities and historical background**

In the first instance, Donayre and colleagues reported in a cohort of 60 Peruvian HCC patients between 1969 and 1997 a mean age of 45 years old, with as many individuals in their third decade as people in their seventh decade [13]. The gender ratio (Male:Female) was balanced at 1.3, which appears to be another peculiarity of the clinical epidemiology of HCC in South America compared with elsewhere where the disease afflicts significantly many more men than women [14]. Forty-five percent of the patients were seropositive for the HBV surface antigen (HBsAg). Unfortunately, no formal diagnosis of cirrhosis was provided in this study. However, the authors consistently described an extensive hepatomegaly due to the explosive development of HCC in a short timescale as the predominant feature of clinical presentation.

Second, our group of medical practitioners and researchers described the clinical and demographic features of 232 consecutive patients who underwent

liver resection for HCC between 1990 and 2006 at the National Cancer Institute of Peru (INEN) [15]. In this cohort, the data were consistent with the observations made previously by Donayre and colleagues: the gender ratio was 1.5 and the mean and median ages were relatively young with 41.4 and 36 years old, respectively. Forty-four percent of the patients were HBsAg seropositive [HBsAg(+)], and individuals presented with an extended hepatomegaly due to the development of sizeable HCC of 15 cm-diameter on average. Surprisingly, only 16.3% of the patients had cirrhosis, and the serum level of alpha-fetoprotein (AFP) was exceedingly high with a mean value over 100,000 ng/mL.

Building on these studies, and in order to provide further insight into this intriguing clinical epidemiological situation, we assembled the largest cohort to date in South America; 1,541 Peruvian patients who were consecutively diagnosed with HCC at INEN between 1997 and 2010 [16]. A comprehensive analysis of the demographics substantiated that both mean and median ages observed were misleading and resulted, in fact, from a genuine bimodal Gaussian age-based distribution with a first peak at age 25 and a second peak at age 64, each mode integrating 50% of the overall patient population (Bimodality Index: 1.95; Moment of Mixture: 44.8 years old; Skewness Coefficients: +0.4 and -0.6, respectively).

This age-based dispersion delineated two distinct subpopulations of HCC patients with specific clinical features in terms of, *inter alia*, gender balance, tumor size, distant metastasis and recurrence [16]. For example, 71% of the

111 younger patient individuals were HBsAg(+) compared with 22.5% in the older  
112 patient population. As mentioned above, two remarkable hallmarks of HCC  
113 clinical presentation in Peru were the significant sizing of an intrahepatic  
114 tumor and the overall low frequency of associated cirrhosis in only 10% of the  
115 cases examined, this ratio barely reaching 5% in the younger patient  
116 population. Noticeably, AFP serum concentration was dramatically heightened  
117 in the younger patient group compared with the older one, culminating with c.  
118 380,000 ng/mL on average. Taken together, these findings support the idea  
119 that a singular biological process is driving liver tumorigenesis in a fraction of  
120 HCC patients in South America; and the clinical context described herein is  
121 uniquely concerning enough to be an important research topic.

122  
123 Afterward, investigators originating from a broad consortium of regional  
124 countries have recently conducted a multicenter study in six South American  
125 countries including Argentina, Brazil, Colombia, Ecuador, Peru, and Uruguay.  
126 In two different articles analyzing the dataset collected from 1,336 patients  
127 with HCC between 2005 and 2015, they corroborated at an upper regional  
128 level the peculiarities of the disease originally observed in the Peruvian HCC  
129 patient population; unfortunately, though, they did not refer to the works  
130 previously published on this specific issue [17,18]. In their studies, the authors  
131 highlighted the large proportion of HCC diagnosed below age 50 in South  
132 America, notably in Peru, and correlated here again the occurrence of early  
133 age HCC with a high prevalence of HBV and mild pervasiveness of  
134 associated cirrhosis. In a subsequent scientific correspondence, Debes  
135 suggested environmental toxins as a foe on early-age HCC in South America,

pointing out the putative role of dietary aflatoxin exposure as already described in Africa [19,20].

### **Mutation spectrum**

For our part, we undertook in a second phase a molecular analysis of nine HCC-related gene mutation hotspots (*i.e.*, *ARID2*, *AXIN1*, *BRAF*, *CTNNB1*, *NFE2L2*, *H/K/N-RAS*, and *TP53*) in 80 Peruvian HCC patients in furtherance to deepen our understanding of HCC in South America [21]. We demonstrated therein that Peruvian HCC featured a peculiar pattern of somatic mutations. The number of genetic alterations observed within these mutational hotspots was relatively low (0 to 3 per tumor) and the intensity of the mutagenic process rather mild with only two mutation hotspots (*i.e.*, *CTNNB1* and *TP53*) reaching 22% of mutants in both early- and late-onset forms of HCC. The preeminent class of HCC-associated genetic defects was epitomized by short deletions affecting notably the Wnt pathway. To the best of our knowledge, this mutation spectrum is not only unprecedented for HCC but also unique among solid tumors in which indels are usually monitored as a marginal subset of genetic alterations, confirming further the distinctive positioning of HCC in South America [22].

### **Risk factors**

In the same study mentioned above from 2014, we addressed the commentary made *a posteriori* first by Chan and colleagues in 2017 and then by Debes in 2018 on the putative brunt of dietary aflatoxin intoxication in early-age onset of HCC afflicting the patients in South America [17,20].



Among the 80 Peruvian patients analyzed, we found only one carrier of the aflatoxin B1-induced R249S *TP53* gene mutation, who was, for the record, an older individual [21,23]. Together with the evaluation of the dietary aflatoxin risk factor, we also considered the possibility of a traditional misapplication of medicinal plants that could contribute to the development of early-age onset of HCC. Indeed, such self-medication disuse that enhances the risk of cancer has been highlighted for *Aristolochia* plants in Chinese traditional medicine [24]. We thus performed a cross-sectional study among 88 Peruvian patients with liver cancer to document their herbal medicine practices [25]. Most of the plant species cited in the survey were of common use in Peru, not being reported hitherto to have carcinogenic potential. Moreover, we discarded, in another preliminary report, the possibility in Peruvian HCC patients of chronic infection with the liver fluke *Fasciola hepatica*, which is found to be endemic in the Andean highlands and has previously been associated with liver parenchymal insults [26,27].

## **Histology**

Taking into account that a significant fraction of HCC patients from South America do not present with full-fledged cirrhosis, we then decided to comprehensively specify, from a histological point of view, the pathological features of both tumor and non-tumor liver parenchymata of 50 HCC patients from Peru [28]. Interestingly, younger HCC patients presented with virtually no fibrolamellar carcinomas, which is a histotype occurring almost exclusively in non-cirrhotic liver and allegedly earlier than age 40 [29]. In addition, a large share of the liver tumors was steato-hepatitic HCC, a relatively rare variant

that is ordinarily associated with non-alcoholic fatty liver diseases and HCV-related cirrhosis [30]. Above all, our survey emphasized the relatively healthy status of the liver in Peruvian HCC patients. Tumors were arising mostly in liver parenchyma with low to mild degrees of fibrosis; this figure is at odds with the current view on the topic, as HCC in non-cirrhotic, non-fibrotic livers is claimed to represent a small minority of the cases [31]. Similarly, the levels of fatty liver and steatohepatitis (8%), as well as siderosis, were relatively low; this latter observation refuting one of the hypotheses formulated by Ponzetto and colleagues to explain the occurrence of early-age HCC among patients in South America [19]. In addition, the micro-steatotic pattern observed in case of infection with the genotype III of hepatitis Delta virus was conspicuously absent from the series. However, a significant proportion of younger HCC patients presented with a high density of clear cell foci of cellular alteration within the non-tumor liver parenchyma [28]. From a morpho-histological perspective, these hepatic foci are reminiscent of liver lesions observed in rodent models that have been subjected to genotoxic chemicals [32]. The exact burden of these clear cell foci and their potential role in liver carcinogenesis has to be ascertained in HCC patients from South America. Therefore, together with prevalent HBV infection, it is likely that the intercession of hitherto poorly documented environmental, metabolic, or infectious cofactors, such as alternative mycotoxins to aflatoxin B1, diabetes mellitus, or even co-infection with *Helicobacter* spp., which is highly endemic in the region [8,19,20,33].

## **Role of hepatitis B virus**

As HBV is still suspected to be the prominent etiological agent and its prevalence monitored serologically associates with younger HCC patients in South America, we performed an in-depth molecular study of HBV infection in 65 HCC patients from Peru [34]. A narrow majority of individuals (51%) were monitored HBsAg(+) and, thus, considered as genuinely infected at the onset of the disease. Using an ultra-sensitive assay, HBV DNA was, however, detected at a very low viral DNA burden in more than 80% of cases, disclosing hence a substantial rate of occult HBV infections in HCC patients [28,34,35]. A phylogenetic analysis of the viral sequences clustered every isolate within the sub-genotype F1b, which is a clade encountered historically in indigenous people of the Americas, notably in Alaska where the occurrence of early-age HCC has been described as well [36–38]. Intriguingly, HBV DNA sequence variations suggest an age-dependent restriction process, as viral genomes in younger patients displayed significantly higher frequency of mutations at di-pyrimidine sites (*i.e.*, TpT and CpC), which was until that time an unprecedented feature in the HBV genome [34]. These findings sharply contrast with the prevailing paradigm that relates higher HBV DNA loads with early-age HCC development [39]. We made assumptions that, in Native communities of the Americas, HBV-associated hepato-carcinogenesis might depart substantially from that broadly observed in other populations [34].

## **Clinical management**

Altogether, the clinical and biological singularities of HCC and associated comorbidities observed in South America have some implications for the management strategies for screening, detection, diagnosis, and prognosis of

HCC patients from the region, as well as the vigilance of the population at risk. For instance, protocols used to screen for HBV infection, notably occult ones, as well as to detect early HCC development in subjects at risk should be tailored to the local situation. Furthermore, Trevisani and colleagues asserted that non-cirrhotic, non-fibrotic HCC clinically represents a distinctive nosological form of HCC with good amenability to liver resection even in cases of major hepatectomy [31]. This observation was confirmed by our review of the 253 Peruvian HCC patients who consecutively underwent a curative hepatectomy at INEN between 1991 and 2011 [40]. In our hands, the survival outcomes of liver resection observed were in good standing with those recorded with liver transplantation in cirrhotic patients with an early-stage tumor, despite the fact that the greatest part of the interventions performed were major hepatectomies due to the size and topography of the tumors exsected. This should be of great concern to the health policy-makers and group of experts aiming to build a regional consensus for HCC prognosis stratification and treatment guidelines and provide state-of-the-art prescriptions for local physicians and scientists, such as the Latin American Association for the Study of the Liver (LAASL) [41].

### **Concluding remark**

Finally, another uncertainty concerns the magnitude of the phenomenon frequently observed in Peru; as to whether it is restricted to the Peruvian Andean communities or also occurs in other human populations, perhaps to a lesser degree. The observations made by Debes and colleagues confirm our original findings at an upper South American continental level [17,18].

Nevertheless, the occurrence of early-age forms of HCC in Alaskan Native people, 10,000 km distant from the indigenous communities of Peru but infected with the very same HBV clade, leads to speculation about the eventuality of a widespread, particular hepato-carcinogenic process shared among all populations with Americas' indigenous ancestry component [16,34,37,42].

**Disclosure of interest:** The authors declare that they have no competing interest.

**Acknowledgements:** The authors are grateful to Elizabeth Elliott, Brian Gadd, and Joshua Lee Halford for their valuable editorial assistance.

**Funding:** This work was supported by the French National Alliance for Life Sciences and Health [grant number ENV201408]; the French National League Against Cancer, Paris, France; and the French National Research Institute for Sustainable Development.

## References

- [1] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018. doi:10.3322/caac.21492.
- [2] Llovet JM, Zucman Rossi J, Pikarsky E, Sangro B, Schwartz M, Sherman M, et al. Hepatocellular carcinoma. *Nat Rev Dis Primer* 2016;2:16018. doi:10.1038/nrdp.2016.18.
- [3] Bray F, Jemal A, Grey N, Ferlay J, Forman D. Global cancer transitions according to the Human Development Index (2008-2030): a population-based study. *Lancet Oncol* 2012;13:790–801. doi:10.1016/S1470-2045(12)70211-5.
- [4] Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-386. doi:10.1002/ijc.29210.
- [5] Soerjomataram I, Lortet Tieulent J, Parkin DM, Ferlay J, Mathers C, Forman D, et al. Global burden of cancer in 2008: a systematic analysis of disability-adjusted life-years in 12 world regions. *Lancet* 2012;380:1840–50. doi:10.1016/S0140-6736(12)60919-2.
- [6] Perz JF, Armstrong GL, Farrington LA, Hutin YJF, Bell BP. The contributions of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer worldwide. *J Hepatol* 2006;45:529–38. doi:10.1016/j.jhep.2006.05.013.

- 303 [7] Kucukcakan B, Hayrulai Musliu Z. Challenging role of dietary aflatoxin  
304 B1 exposure and hepatitis B infection on risk of hepatocellular  
305 carcinoma. Open Access Maced J Med Sci 2015;3:363–9.  
306 doi:10.3889/oamjms.2015.032.
- 307 [8] Li X, Wang X, Gao P. Diabetes mellitus and risk of hepatocellular  
308 carcinoma. BioMed Res Int 2017;2017:5202684.  
309 doi:10.1155/2017/5202684.
- 310 [9] Singh AK, Kumar R, Pandey AK. Hepatocellular carcinoma: causes,  
311 mechanism of progression and biomarkers. Curr Chem Genomics Transl  
312 Med 2018;12:9–26. doi:10.2174/2213988501812010009.
- 313 [10] Testino G, Leone S, Borro P. Alcohol and hepatocellular carcinoma: a  
314 review and a point of view. World J Gastroenterol 2014;20:15943–54.  
315 doi:10.3748/wjg.v20.i43.15943.
- 316 [11] Bray F, Piñeros M. Cancer patterns, trends and projections in Latin  
317 America and the Caribbean: a global context. Salud Publica Mex  
318 2016;58:104–17.
- 319 [12] Wong MCS, Jiang JY, Goggins WB, Liang M, Fang Y, Fung FDH, et al.  
320 International incidence and mortality trends of liver cancer: a global  
321 profile. Sci Rep 2017;7:45846. doi:10.1038/srep45846.
- 322 [13] Donayre M, Bussalleu A, Berríos J, Cok J. Carcinoma primario de  
323 hígado en el Hospital Nacional Cayetano Heredia (Enero 1969 - Abril  
324 1997). Hallazgos clínicos y de laboratorio. Rev Gastroenterol Peru  
325 1999;19:15–25.
- 326 [14] Yang D, Hanna DL, Usher J, LoCoco J, Chaudhari P, Lenz HJ, et al.

- 327 Impact of sex on the survival of patients with hepatocellular carcinoma: a  
328 Surveillance, Epidemiology, and End Results analysis. *Cancer*  
329 2014;120:3707–16. doi:10.1002/cncr.28912.
- 330 [15] Ruiz E, Sanchez J, Celis J, Payet E, Berrospi F, Chavez I, et al.  
331 Resultados a corto y largo plazo de la resección hepática por  
332 hepatocarcinoma. Análisis de 232 resecciones consecutivas. *Rev*  
333 *Gastroenterol Peru* 2007;27:223–37.
- 334 [16] Bertani S, Pineau P, Loli S, Moura J, Zimic M, Deharo E, et al. An  
335 atypical age-specific pattern of hepatocellular carcinoma in Peru: a  
336 threat for Andean populations. *PLOS One* 2013;8:e67756.  
337 doi:10.1371/journal.pone.0067756.
- 338 [17] Chan AJ, Balderramo D, Kikuchi L, Ballerga EG, Prieto JE, Tapias M, et  
339 al. Early age hepatocellular carcinoma associated with hepatitis B  
340 infection in South America. *Clin Gastroenterol Hepatol* 2017;15:1631–2.  
341 doi:10.1016/j.cgh.2017.05.015.
- 342 [18] Debes JD, Chan AJ, Balderramo D, Kikuchi L, Gonzalez Ballerga E,  
343 Prieto JE, et al. Hepatocellular carcinoma in South America: evaluation  
344 of risk factors, demographics and therapy. *Liver Int* 2018;38:136–43.  
345 doi:10.1111/liv.13502.
- 346 [19] Ponzetto A, Diella FA, Holton J. Aetiology of hepatocellular carcinoma in  
347 South America. *Liver Int* 2018;38:956–7. doi:10.1111/liv.13703.
- 348 [20] Debes JD. Early hepatocellular carcinoma in South America: what is to  
349 blame? *Liver Int* 2018;38:957–8. doi:10.1111/liv.13717.
- 350 [21] Marchio A, Bertani S, Rojas Rojas T, Doimi F, Terris B, Deharo E, et al.



351 A peculiar mutation spectrum emerging from young Peruvian patients  
 352 with hepatocellular carcinoma. PLOS One 2014;9:e114912.  
 353 doi:10.1371/journal.pone.0114912.

354 [22] Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler  
 355 KW. Cancer genome landscapes. Science 2013;339:1546–58.  
 356 doi:10.1126/science.1235122.

357 [23] Mao R, Liu J, Liu G, Jin S, Xue Q, Ma L, et al. Whole genome  
 358 sequencing of matched tumor, adjacent non-tumor tissues and  
 359 corresponding normal blood samples of hepatocellular carcinoma  
 360 patients revealed dynamic changes of the mutations profiles during  
 361 hepatocarcinogenesis. Oncotarget 2017;8:26185–99.  
 362 doi:10.18632/oncotarget.15428.

363 [24] Nortier JL, Martinez MC, Schmeiser HH, Arlt VM, Bieler CA, Petein M, et  
 364 al. Urothelial carcinoma associated with the use of a Chinese herb  
 365 (*Aristolochia fangchi*). N Engl J Med 2000;342:1686–92.  
 366 doi:10.1056/NEJM200006083422301.

367 [25] Rojas Rojas T, Bourdy G, Ruiz E, Cerapio JP, Pineau P, Gardon J, et al.  
 368 Herbal medicine practices of patients with liver cancer in Peru: a  
 369 comprehensive study toward integrative cancer management. Integr  
 370 Cancer Ther 2018;17:52–64. doi:10.1177/1534735416681642.

371 [26] Machicado C, Bertani S, Herrera Velit P, Espinoza J, Ruiz E, Marcos L.  
 372 Negative serology of *Fasciola hepatica* infection in patients with liver  
 373 cancer in Peru: a preliminary report. Rev Soc Bras Med Trop  
 374 2018;51:231–3. doi:10.1590/0037-8682-0180-2017.

- 375 [27] Cabada MM, Morales ML, Webb CM, Yang L, Bravenec CA, Lopez M, et  
376 al. Socioeconomic factors associated with *Fasciola hepatica* infection  
377 among children from 26 communities of the Cusco region of Peru. Am J  
378 Trop Med Hyg 2018. doi:10.4269/ajtmh.18-0372.
- 379 [28] Cano L, Cerapio JP, Ruiz E, Marchio A, Turlin B, Casavilca S, et al.  
380 Liver clear cell foci and viral infection are associated with non-cirrhotic,  
381 non-fibrolamellar hepatocellular carcinoma in young patients from South  
382 America. Sci Rep 2018;8:9945. doi:10.1038/s41598-018-28286-0.
- 383 [29] Kakar S, Burgart LJ, Batts KP, Garcia J, Jain D, Ferrell LD.  
384 Clinicopathologic features and survival in fibrolamellar carcinoma:  
385 comparison with conventional hepatocellular carcinoma with and without  
386 cirrhosis. Mod Pathol 2005;18:1417–23.  
387 doi:10.1038/modpathol.3800449.
- 388 [30] Salomao M, Yu WM, Brown RS, Emond JC, Lefkowitz JH.  
389 Steatohepatic hepatocellular carcinoma (SH-HCC): a distinctive  
390 histological variant of HCC in hepatitis C virus-related cirrhosis with  
391 associated NAFLD/NASH. Am J Surg Pathol 2010;34:1630–6.  
392 doi:10.1097/PAS.0b013e3181f31caa.
- 393 [31] Trevisani F, Frigerio M, Santi V, Grignaschi A, Bernardi M.  
394 Hepatocellular carcinoma in non-cirrhotic liver: a reappraisal. Dig Liver  
395 Dis 2010;42:341–7. doi:10.1016/j.dld.2009.09.002.
- 396 [32] Ribback S, Calvisi DF, Cigliano A, Sailer V, Peters M, Rausch J, et al.  
397 Molecular and metabolic changes in human liver clear cell foci resemble  
398 the alterations occurring in rat hepatocarcinogenesis. J Hepatol

399 2013;58:1147–56. doi:10.1016/j.jhep.2013.01.013.

400 [33] Eusebi LH, Zagari RM, Bazzoli F. Epidemiology of *Helicobacter pylori*  
401 infection. *Helicobacter* 2014;19 Suppl 1:1–5. doi:10.1111/hel.12165.

402 [34] Marchio A, Cerapio JP, Ruiz E, Cano L, Casavilca S, Terris B, et al.  
403 Early-onset liver cancer in South America associates with low hepatitis B  
404 virus DNA burden. *Sci Rep* 2018;8:12031. doi:10.1038/s41598-018-  
405 30229-8.

406 [35] Hollinger FB, Sood G. Occult hepatitis B virus infection: a covert  
407 operation. *J Viral Hepat* 2010;17:1–15. doi:10.1111/j.1365-  
408 2893.2009.01245.x.

409 [36] von Meltzer M, Vásquez S, Sun J, Wendt UC, May A, Gerlich WH, et al.  
410 A new clade of hepatitis B virus subgenotype F1 from Peru with unusual  
411 properties. *Virus Genes* 2008;37:225–30. doi:10.1007/s11262-008-  
412 0261-x.

413 [37] Gounder PP, Bulkow LR, Snowball M, Negus S, Spradling PR,  
414 McMahon BJ. Hepatocellular carcinoma risk in Alaska Native children  
415 and young adults with hepatitis B virus: retrospective cohort analysis. *J*  
416 *Pediatr* 2016;178:206–13. doi:10.1016/j.jpeds.2016.08.017.

417 [38] Livingston SE, Simonetti JP, McMahon BJ, Bulkow LR, Hurlburt KJ,  
418 Homan CE, et al. Hepatitis B virus genotypes in Alaska Native people  
419 with hepatocellular carcinoma: preponderance of genotype F. *J Infect*  
420 *Dis* 2007;195:5–11. doi:10.1086/509894.

421 [39] Chen CJ, Yang HI, Iloeje UH, REVEAL-HBV Study Group. Hepatitis B  
422 virus DNA levels and outcomes in chronic hepatitis B. *Hepatol Baltim Md*

423 2009;49:S72-84. doi:10.1002/hep.22884.

424 [40] Ruiz E, Rojas Rojas T, Berrospi F, Chávez I, Luque C, Cano L, et al.

425 Hepatocellular carcinoma surgery outcomes in the developing world: a

426 20-year retrospective cohort study at the National Cancer Institute of

427 Peru. *Heliyon* 2016;2:e00052. doi:10.1016/j.heliyon.2015.e00052.

428 [41] Méndez Sánchez N, Ridruejo E, Alves de Mattos A, Chávez Tapia NC,

429 Zapata R, Paraná R, et al. Latin American Association for the Study of

430 the Liver (LAASL) clinical practice guidelines: management of

431 hepatocellular carcinoma. *Ann Hepatol* 2014;13 Suppl 1:S4-40.

432 [42] Hayashi S, Khan A, Simons BC, Homan C, Matsui T, Ogawa K, et al. A

433 novel association between core mutations in hepatitis B virus genotype

434 F1b and hepatocellular carcinoma in Alaskan Native People. *Hepatology*

435 2018. doi:10.1002/hep.30111.

436