



HAL
open science

Cressdnaviricota : a Virus Phylum Unifying Seven Families of Rep-Encoding Viruses with Single-Stranded, Circular DNA Genomes

Mart Krupovic, Arvind Varsani, Darius Kazlauskas, Mya Breitbart, Eric A Delwart, Karyna Rosario, Natalya Yutin, Yuri I Wolf, Balázs Harrach, F. Murilo Zerbini, et al.

► To cite this version:

Mart Krupovic, Arvind Varsani, Darius Kazlauskas, Mya Breitbart, Eric A Delwart, et al.. Cressdnaviricota : a Virus Phylum Unifying Seven Families of Rep-Encoding Viruses with Single-Stranded, Circular DNA Genomes. *Journal of Virology*, 2020, 94 (12), pp.e00582-20. 10.1128/JVI.00582-20 . pasteur-02861253

HAL Id: pasteur-02861253

<https://pasteur.hal.science/pasteur-02861253>

Submitted on 8 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 ***Cressdnaviricota*: a virus phylum unifying seven families of Rep-encoding viruses with**
2 **single-stranded, circular DNA genomes**

3
4 Mart Krupovic ^{a#}, Arvind Varsani ^{b,c}, Darius Kazlauskas ^d, Mya Breitbart ^e, Eric Delwart ^{f,g}, Karyna
5 Rosario ^e, Natalya Yutin ^h, Yuri I. Wolf ^h, Balázs Harrach ⁱ, F. Murilo Zerbini ^j, Valerian V. Dolja ^k, Jens
6 H. Kuhn ^l, Eugene V. Koonin ^h

7
8 a – Archaeal Virology Unit, Institut Pasteur, 75015 Paris, France

9 b – The Biodesign Center for Fundamental and Applied Microbiomics, Center for Evolution and
10 Medicine, School of Life Sciences, Arizona State University, Tempe, AZ, USA

11 c – Structural Biology Research Unit, Department of Integrative Biomedical Sciences, University of Cape
12 Town, Cape Town, South Africa

13 d – Institute of Biotechnology, Life Sciences Center, Vilnius University, Saulėtekio av. 7, Vilnius, 10257,
14 Lithuania

15 e – University of South Florida, College of Marine Science, 140 7th Avenue South, St. Petersburg, FL,
16 33701, USA

17 f – Vitalant Research Institute, San Francisco, CA, USA

18 g – UCSF Dept. of Laboratory Medicine, University of California-San Francisco, San Francisco, CA,
19 USA

20 h – National Center for Biotechnology Information, National Library of Medicine, National Institutes of
21 Health, Bethesda, MD, 20894, USA

22 i – Institute for Veterinary Medical Research, Centre for Agricultural Research, Hungária krt. 21,
23 Budapest, 1143, Hungary

24 j – Departamento de Fitopatologia/Bioagro, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil

25 k – Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA

26 l – Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious Diseases,
27 National Institutes of Health, Frederick, Maryland, USA

28
29
30 # – Correspondence

31 E-mail: mart.krupovic@pasteur.fr

32
33
34 **Keywords:** *Bacilladnaviridae*, *Circoviridae*, *Geminiviridae*, *Genomoviridae*, *Nanoviridae*,
35 *Redondoviridae*, *Smacoviridae*, International Committee on Taxonomy of Viruses (ICTV), CRESS DNA
36 viruses, virus classification, virus evolution

37 **Text**

38 Eukaryotic single-stranded (ss) DNA viruses are classified into ten families (Table 1) but many remain
39 unclassified (1, 2). Viruses from seven families (*Bacilladnaviridae*, *Circoviridae*, *Geminiviridae*,
40 *Genomoviridae*, *Nanoviridae*, *Redondoviridae*, and *Smacoviridae*) have small circular genomes, some
41 encompassing only two genes, encoding the replication initiation protein (Rep) and the capsid protein (3-
42 5). The genomes are replicated by the rolling-circle mechanism initiated by the virus-encoded Rep
43 proteins of the HUH endonuclease superfamily (6-10). In eukaryotic ssDNA viruses, the HUH
44 endonuclease domain is followed by a superfamily 3 helicase (S3H) domain, which is not found in
45 prokaryotic ssDNA virus Repls, although homologous two-domain Repls are encoded by several groups of
46 bacterial plasmids (5). Informally, this assemblage of ssDNA viruses has been referred to as “circular,
47 Rep-encoding single-stranded (CRESS) DNA viruses” (4, 8). Viruses from the seven families infect hosts
48 across the eukaryotic domain, including animals (*Circoviridae*), diatoms (*Bacilladnaviridae*), fungi
49 (*Genomoviridae*) and plants (*Nanoviridae* and *Geminiviridae*), whereas members of the *Smacoviridae* and
50 the newly created *Redondoviridae* have been discovered by metagenomics and are suspected to infect
51 animals (3, 11), although association of smacovirids with methanogenic archaea has also been suggested
52 (12).

53
54 Phylogenetic analysis of classified and unclassified CRESS-DNA viruses revealed 13 well supported
55 clades (Figure 1). Six clades correspond to unclassified viruses, denoted CRESSV1 through CRESSV6
56 (13). The phylogeny splits into two large clades: (1) geminivirids, genomovirids, and CRESSV6 viruses;
57 (2) all other CRESS-DNA viruses (Figure 1). These clades are supported by detailed analyses of the
58 conserved catalytic motifs of Rep proteins (5, 14), with Repls of viruses in clade 1 sharing the GRS motif
59 (15, 16), which is not found in Repls of viruses from clade 2. By contrast, Repls of all members of clade 2
60 contain a conserved arginine finger in the helicase domain, absent in geminivirids and genomovirids,
61 supporting the monophyly of the Repls in clade 2.

62
63 Unlike the Repls, the capsid proteins of CRESS-DNA viruses are not orthologous, and apparently were
64 acquired from RNA viruses on multiple, independent occasions (5, 14, 17-19). Thus, the gene encoding
65 the characteristic two-domain Rep is the only universally conserved gene in this virus supergroup and thus
66 can serve as a scaffold for mapping evolutionary events. Accordingly, it has been proposed that Repls be
67 used as a basis for megataxonomic classification of CRESS-DNA viruses (20).

68
69 Recently, the International Committee on Taxonomy of Viruses (ICTV) has extended its taxonomic rank
70 structure to facilitate the classification of the entire spectrum of virus diversity (21, 22). This new
71 taxonomic framework was adopted to formalize the evolutionary relationships among CRESS-DNA
72 viruses. Below we outline these taxonomic changes, which have been ratified by the ICTV in March of
73 2020 and are now part of the official virus taxonomy:

- 74
- 75 • Clade 1: Families *Geminiviridae* and *Genomoviridae* were included into the new order *Geplafuvirales*
76 (gemini/genomo, plants, fungi, and the suffix -virales for order taxa). In anticipation of the future
77 classification of the CRESSV6 virus group, the order *Geplafuvirales* has been assigned into a new
78 class, *Repensiviricetes* (a portmanteaux of Rep-encoding single strand, and the suffix -viricetes for
79 class taxa).
 - 80
81 • Clade 2: Viruses of the families *Bacilladnaviridae*, *Circoviridae*, *Nanoviridae*, *Redondoviridae* and
82 *Smacoviridae*, and unclassified groups CRESSV1–5 are sufficiently distinct from each other,
83 precluding their unification into a single order. The differences extend beyond the separation of the
84 Repls in the phylogenetic tree and include non-homologous capsid proteins, different genome lengths
85 (Table 1) and genome architectures (e.g., nanovirids have multipartite genomes, whereas other viruses
86 have monopartite genomes). Notably, circovirids cluster with CRESSV1–3 viruses, whereas nanovirids

- 87 cluster with CRESSV4–5 viruses, justifying order-rank taxa for these assemblages. Thus, five orders
88 have been created for classification of clade 2 viruses:
- 89 - *Baphyvirales* (to include *Bacilladnaviridae*): after bacillariophytes (diatoms, hosts of viruses in
90 this taxon), and the suffix -virales;
 - 91 - *Circlivirales* (to include *Circoviridae* and, in due course, CRESSV1–3 viruses): circo-like, and the
92 suffix -virales;
 - 93 - *Cremevirales* (to include *Smacoviridae*): CRESS, metagenomics, and the suffix -virales;
 - 94 - *Mulpavirales* (to include *Nanoviridae* and, in due course, CRESSV4 and CRESSV5 viruses): after
95 multipartite genomes of viruses in this taxon, and the suffix -virales;
 - 96 - *Recrevirales* (to include the newly approved *Redondoviridae*): redondoviruses, CRESS, and the
97 suffix -virales.

98
99 Viruses from the established families within the five orders encode order-specific capsid proteins, which
100 are not recognizably similar at the sequence level across the orders. This feature and the Rep phylogeny
101 should help in assigning newly identified viruses to a particular taxon. Notably, Reps of satellite nucleic
102 acids from the family *Alphasatellitidae* (23) and nanovirids form a clade (Figure 1); thus a taxon unifying
103 *Alphasatellitidae* and *Nanoviridae* might be considered in the future. The five orders were further assigned
104 to the class *Arfiviricetes* (arginine finger, and the suffix -viricetes for class taxa). Finally, the two classes
105 *Repensiviricetes* and *Arfiviricetes* were grouped into the phylum *Cressdnaviricota* (a contraction of
106 CRESS DNA, and the suffix -viricota for phylum taxa), thereby unifying all officially recognized families
107 of CRESS-DNA viruses.

110 Funding

111 M.K. was supported by l'Agence Nationale de la Recherche (project ENVIRA, no. ANR-17-CE15-0005-
112 01). E.V.K., Y.I.W. and N.Y. are supported by the Intramural Research Program of the U.S. National
113 Institutes of Health (National Library of Medicine). B.H. was supported by the Hungarian National
114 Research, Development and Innovation Office (NN128309). This work was supported in part through the
115 Laulima Government Solutions, LLC prime contract with the US National Institute of Allergy and
116 Infectious Diseases (NIAID) under Contract No. HHSN272201800013C. J.H.K. performed this work as
117 an employee of Tunnell Government Services (TGS), a subcontractor of Laulima Government Solutions,
118 LLC under Contract No. HHSN272201800013C.

119
120 The views and conclusions contained in this document are those of the authors and should not be
121 interpreted as necessarily representing the official policies, either expressed or implied, of the U.S.
122 Department of Health and Human Services or of the institutions and companies affiliated with the authors.

124 Figure legend

125 **Figure 1.** Unrooted maximum likelihood phylogenetic tree of Rep proteins from CRESS-DNA viruses.
126 Closely related sequence groups are collapsed into triangles, the side lengths of which are proportional to
127 the distances between closest and farthest leaf nodes. The Rep alignment used for the tree reconstruction
128 was taken from Kazlauskas et al. (5, 13) and supplemented with sequences of redondovirids (3). The
129 maximum likelihood phylogenetic tree was constructed using PhyML (24) with automatic selection of the
130 best-fit substitution model for a given alignment. The best model identified by PhyML was RtREV. The
131 branch support was assessed using aBayes implemented in PhyML. The scale bar represents the number
132 of substitutions per site.

133 **References**

- 134 1. Simmonds P, Adams MJ, Benko M, Breitbart M, Brister JR, Carstens EB, Davison AJ, Delwart E,
135 Gorbalenya AE, Harrach B, Hull R, King AM, Koonin EV, Krupovic M, Kuhn JH, Lefkowitz EJ, Nibert
136 ML, Orton R, Roossinck MJ, Sabanadzovic S, Sullivan MB, Suttle CA, Tesh RB, van der Vlugt RA,
137 Varsani A, Zerbini FM. 2017. Consensus statement: Virus taxonomy in the age of metagenomics.
138 Nat Rev Microbiol 15:161-168.
- 139 2. Tisza MJ, Pastrana DV, Welch NL, Stewart B, Peretti A, Starrett GJ, Pang YS, Krishnamurthy SR,
140 Pesavento PA, McDermott DH, Murphy PM, Whited JL, Miller B, Brenchley J, Rosshart SP,
141 Reherrmann B, Doorbar J, Ta'ala BA, Pletnikova O, Troncoso JC, Resnick SM, Bolduc B, Sullivan
142 MB, Varsani A, Segall AM, Buck CB. 2020. Discovery of several thousand highly diverse circular
143 DNA viruses. eLife 9:e51971.
- 144 3. Abbas AA, Taylor LJ, Dothard MI, Leiby JS, Fitzgerald AS, Khatib LA, Collman RG, Bushman FD.
145 2019. *Redondoviridae*, a family of small, circular DNA viruses of the human oro-respiratory tract
146 associated with periodontitis and critical illness. Cell Host Microbe 25:719-729 e4.
- 147 4. Zhao L, Rosario K, Breitbart M, Duffy S. 2019. Eukaryotic circular Rep-encoding single-stranded
148 DNA (CRESS DNA) viruses: Ubiquitous viruses with small genomes and a diverse host range. Adv
149 Virus Res 103:71-133.
- 150 5. Kazlauskas D, Varsani A, Koonin EV, Krupovic M. 2019. Multiple origins of prokaryotic and
151 eukaryotic single-stranded DNA viruses from bacterial and archaeal plasmids. Nat Commun
152 10:3425.
- 153 6. Chandler M, de la Cruz F, Dyda F, Hickman AB, Moncalian G, Ton-Hoang B. 2013. Breaking and
154 joining single-stranded DNA: the HUH endonuclease superfamily. Nat Rev Microbiol 11:525-38.
- 155 7. Krupovic M. 2013. Networks of evolutionary interactions underlying the polyphyletic origin of
156 ssDNA viruses. Curr Opin Virol 3:578-86.
- 157 8. Rosario K, Duffy S, Breitbart M. 2012. A field guide to eukaryotic circular single-stranded DNA
158 viruses: insights gained from metagenomics. Arch Virol 157:1851-71.
- 159 9. Ilyina TV, Koonin EV. 1992. Conserved sequence motifs in the initiator proteins for rolling circle
160 DNA replication encoded by diverse replicons from eubacteria, eucaryotes and archaeobacteria.
161 Nucleic Acids Res 20:3279-85.
- 162 10. Koonin EV, Ilyina TV. 1993. Computer-assisted dissection of rolling circle DNA replication.
163 Biosystems 30:241-68.
- 164 11. Varsani A, Krupovic M. 2018. *Smacoviridae*: a new family of animal-associated single-stranded
165 DNA viruses. Arch Virol 163:2005-2015.
- 166 12. Díez-Villaseñor C, Rodriguez-Valera F. 2019. CRISPR analysis suggests that small circular single-
167 stranded DNA smacoviruses infect Archaea instead of humans. Nat Commun 10:294.
- 168 13. Kazlauskas D, Varsani A, Krupovic M. 2018. Pervasive chimerism in the replication-associated
169 proteins of uncultured single-stranded DNA viruses. Viruses 10:E187.
- 170 14. Kazlauskas D, Dayaram A, Kraberger S, Goldstien S, Varsani A, Krupovic M. 2017. Evolutionary
171 history of ssDNA bacilladnaviruses features horizontal acquisition of the capsid gene from ssRNA
172 nodaviruses. Virology 504:114-121.
- 173 15. Nash TE, Dallas MB, Reyes MI, Buhman GK, Ascencio-Ibanez JT, Hanley-Bowdoin L. 2011.
174 Functional analysis of a novel motif conserved across geminivirus Rep proteins. J Virol 85:1182-
175 92.
- 176 16. Varsani A, Krupovic M. 2017. Sequence-based taxonomic framework for the classification of
177 uncultured single-stranded DNA viruses of the family *Genomoviridae*. Virus Evol 3:vew037.
- 178 17. Diemer GS, Stedman KM. 2012. A novel virus genome discovered in an extreme environment
179 suggests recombination between unrelated groups of RNA and DNA viruses. Biol Direct 7:13.

- 180 18. Krupovic M, Ravantti JJ, Bamford DH. 2009. Geminiviruses: a tale of a plasmid becoming a virus.
181 BMC Evol Biol 9:112.
- 182 19. Roux S, Enault F, Bronner G, Vaultot D, Forterre P, Krupovic M. 2013. Chimeric viruses blur the
183 borders between the major groups of eukaryotic single-stranded DNA viruses. Nat Commun
184 4:2700.
- 185 20. Koonin EV, Dolja VV, Krupovic M, Varsani A, Wolf YI, Yutin N, Zerbini FM, Kuhn JH. 2020. Global
186 organization and proposed megataxonomy of the virus world. Microbiol Mol Biol Rev 84:e00061-
187 19.
- 188 21. Siddell SG, Walker PJ, Lefkowitz EJ, Mushegian AR, Adams MJ, Dutilh BE, Gorbalenya AE, Harrach
189 B, Harrison RL, Junglen S, Knowles NJ, Kropinski AM, Krupovic M, Kuhn JH, Nibert M, Rubino L,
190 Sabanadzovic S, Sanfaçon H, Simmonds P, Varsani A, Zerbini FM, Davison AJ. 2019. Additional
191 changes to taxonomy ratified in a special vote by the International Committee on Taxonomy of
192 Viruses (October 2018). Arch Virol 164:943-946.
- 193 22. Gorbalenya AE, Krupovic M, Mushegian A, Kropinski AM, Siddell S, Varsani A, Adams MJ, Davison
194 AJ, Dutilh BA, Harrach B, Harrison RL, Junglen S, King AMQ, Knowles NJ, Lefkowitz EJ, Nibert ML,
195 Rubino L, Sabanadzovic S, Sanfaçon H, Simmonds P, Walker PJ, Zerbini FM, Kuhn JH. 2020. The
196 new scope of virus taxonomy: partitioning the virosphere into 15 hierarchical ranks. Nat
197 Microbiol 5:668-674.
- 198 23. Briddon RW, Martin DP, Roumagnac P, Navas-Castillo J, Fiallo-Olive E, Moriones E, Lett JM,
199 Zerbini FM, Varsani A. 2018. *Alphasatellitidae*: a new family with two subfamilies for the
200 classification of geminivirus- and nanovirus-associated alphasatellites. Arch Virol 163:2587-2600.
- 201 24. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and
202 methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0.
203 Syst Biol 59:307-21.

204

205 Table 1. Families of ssDNA viruses associated with eukaryotic hosts. Families unified into the phylum
 206 *Cressdnaviricota* are in bold.

Family	Host	Virion ϕ, nm	Genome topology	Genome size, kb	ORFs	Replication protein[#]
<i>Anelloviridae</i>	animals	30–32	circular	2–4	2–4	Rep
<i>Bidnaviridae</i>	animals (insects)	20–24	linear	6–6.5 per segment (2 segments)	6	pPolB
<i>Parvoviridae</i>	animals	23–28	linear	4–6.3	3–4	HUH-Rep
<i>Bacilladnaviridae</i>	algae (diatoms)	33–38	circular	5.5–6	4	HUH-Rep
<i>Circoviridae</i>	animals	15–25	circular	1.7–2.1	2	HUH-Rep
<i>Geminiviridae</i>	plants	22x38	circular	2.5–3 per segment (1–2 segments)	4–8	HUH-Rep
<i>Genomoviridae</i>	fungi, insects	20–22	circular	2–2.4	2	HUH-Rep
<i>Nanoviridae</i>	plants	17–20	circular	0.98–1.1 per segment (6–9 segments)	5–8	HUH-Rep
<i>Redondoviridae</i>	?	?	circular	3.0–3.1	3	HUH-Rep
<i>Smacoviridae</i>	?	?	circular	2.3–2.9	2	HUH-Rep

207 [#]HUH-Rep, Rep of the HUH endonuclease superfamily; pPolB, protein-primed family B DNA
 208 polymerase.