



## **Klebsiella pneumoniae carriage in low-income countries: antimicrobial resistance, genomic diversity and risk factors**

Bich-Tram Huynh, Virginie Passet, Andriniana Rakotondrasoana, Thierno Diallo, Alexandra Kerleguer, Melanie Hennart, Agathe De Lauzanne, Perlinot Herindrainy, Abdoulaye Seck, Raymond Bercion, et al.

### **► To cite this version:**

Bich-Tram Huynh, Virginie Passet, Andriniana Rakotondrasoana, Thierno Diallo, Alexandra Kerleguer, et al.. *Klebsiella pneumoniae carriage in low-income countries: antimicrobial resistance, genomic diversity and risk factors*. *Gut microbes*, 2020, pp.1-13. 10.1080/19490976.2020.1748257 . pasteur-02866178

**HAL Id: pasteur-02866178**

**<https://pasteur.hal.science/pasteur-02866178>**

Submitted on 12 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

***Klebsiella pneumoniae* carriage in low-income countries:  
antimicrobial resistance, genomic diversity and risk factors**

**Bich-Tram HUYNH<sup>1</sup>, Virginie PASSET<sup>2</sup>, Andriniaina RAKOTONDRASOA<sup>3</sup>, Thierno  
DIALLO<sup>4</sup>, Alexandra KERLEGUER<sup>5</sup>, Melanie HENNART<sup>2</sup>, Agathe De LAUZANNE<sup>6</sup>,  
Perlinot HERINDRAINY<sup>7</sup>, Abdoulaye SECK<sup>4</sup>, Raymond BERCION<sup>4</sup>, Laurence  
BORAND<sup>6</sup>, Maria PARDOS DE LA GANDARA<sup>8</sup>, Elisabeth DELAROCQUE-  
ASTAGNEAU<sup>1</sup>, Didier GUILLEMOT<sup>1</sup>, Muriel VRAY<sup>9</sup>, Benoit GARIN<sup>3</sup>, Jean-Marc  
COLLARD<sup>3</sup>, Carla RODRIGUES<sup>2</sup> and Sylvain BRISSE<sup>2 \*</sup>**

1- UMR 1181 "Biostatistics, Biomathematics, Pharmacoepidemiology and Infectious  
Diseases"(B2PhEID). Institut Pasteur, INSERM, University of Versailles Saint-Quentin

2- Institut Pasteur, Biodiversity and Epidemiology of Bacterial Pathogens, Paris, France

3- Experimental bacteriology unit, Institut Pasteur Madagascar

4- Laboratory of medical biology, Institut Pasteur de Dakar

5- Medical Biology Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

6- Epidemiology & Public Health Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

7- Epidemiology & Public Health Unit, Institut Pasteur Madagascar

8- Institut Pasteur, Centre National de Référence des Escherichia coli, Shigella et Salmonella,  
Paris, France

9- Epidemiology & Public Health Unit, Institut Pasteur de Dakar

**\*Corresponding author:** S. Brisse. Institut Pasteur, Biodiversity and Epidemiology of  
Bacterial Pathogens, 28 rue du Docteur Roux, F-75724 Paris, France. E-mail:  
sbrisse@pasteur.fr; Phone +33 1 45 68 83 34.

## **Abstract**

### **Background**

*Klebsiella pneumoniae* (hereafter, *Kp*) is a major public health threat responsible for high levels of multidrug resistant (MDR) human infections. Besides, *Kp* also causes severe infections in the community, especially in Asia and Africa. Although most *Kp* infections are caused by endogenous intestinal carriage, little is known about the prevalence and microbiological characteristics of *Kp* in asymptomatic human carriage, and attached risk factors including environmental sources exposure.

### **Methods**

Here, 911 pregnant women from communities in Madagascar, Cambodia, and Senegal were screened for gut colonization by *Kp*. Characteristics of *Kp* strains (antimicrobial susceptibility, genomic diversity, virulence, and resistance genes) were defined, and associated risk factors were investigated.

### **Results**

*Kp* carriage rate was 55.9%, and *Kp* populations were highly heterogeneous (6 phylogroups, 325 sequence types, Simpson index 99.6%). One third of *Kp* isolates had acquired antimicrobial resistance genes. MDR-*Kp* (11.7% to 39.7%) and extended spectrum beta-lactamase (ESBL)-producing *Kp* (0.7% to 14.7%) varied among countries. Isolates with virulence genes were detected (14.5%). Environmental exposure factors including food, animal contacts, or hospitalization of household members were associated with carriage of *Kp*, antimicrobial resistance and hypervirulence. However, risk factors were country-specific and *Kp* subpopulation-specific.

### **Conclusion**

46 This large-scale multicenter study uncovers the huge diversity of Kp in human gut carriage,  
47 demonstrates that antimicrobial resistance is widespread in communities of three low-income  
48 countries, and underlines the challenges posed by Kp colonization to the control of  
49 antimicrobial resistance.

50 **Key words:** *Klebsiella pneumoniae*, carriage, antibiotic resistance, genomic diversity,  
51 community, low-income countries

## BACKGROUND

*Klebsiella pneumoniae* (Kp) represents an increasing threat to public health. Kp strains that produce extended spectrum beta-lactamases (ESBLs) and carbapenemases often exhibit cross-resistance to other available agents, and pan-resistant *Klebsiella* are increasingly described.<sup>1</sup> In low-income countries (LICs), Kp is responsible for a large part of early neonatal infections and for severe community-acquired infections.<sup>2</sup> The latter can occur in previously healthy persons and are caused by Kp strains that are regarded as hypervirulent (HV-Kp).<sup>1</sup> In LICs, multidrug resistant (MDR)-Kp is one important driver of unfavorable outcome in infections because appropriate therapy is often not available nor affordable. Worryingly, the convergence of MDR and HV phenotypes in single Kp strains is being increasingly reported.<sup>3</sup>

The taxonomy of Kp has been recently updated, highlighting the existence of five different species distributed in seven phylogroups (Kp1 to Kp7): *K. pneumoniae sensu stricto* (phylogroup Kp1), *K. quasipneumoniae* [subsp. *quasipneumoniae* (Kp2) and subsp. *similipneumoniae* (Kp4)], *K. variicola* [subsp. *variicola* (Kp3) and subsp. *tropica* (Kp5)], ‘*K. quasivariicola*’ (Kp6) and *K. africana* (Kp7).<sup>4</sup> In the manuscript, ‘Kp’ abbreviation refers to the *K. pneumoniae* species complex (‘*sensu lato*’, including the five species/seven phylogroups).

Kp is a well-established bacteria of the gut microbiota, and most Kp infections are caused by endogenous strains that previously colonize the host asymptomatically.<sup>5,6</sup> However, Kp carriage has been mostly analyzed within health-care facilities and with a focus on antimicrobial resistant strains. Currently, little is known about naturally occurring human carriage Kp populations, and on the prevalence and diversity of MDR-Kp or HV-Kp among these.

Kp is a ubiquitous bacteria present in animals, environmental sources, and food.<sup>7</sup> Therefore, environmental exposures or behavioral factors may impact individual Kp carriage. Understanding the determinants of Kp carriage may lead to novel strategies to control the spread of this pathogen. To our knowledge, risk factors associated with Kp carriage have not been investigated in the community, nor in LICs.

The objectives of this study were to define the prevalence, antimicrobial resistance, genomic diversity, and virulence potential of Kp isolates carried in pregnant women from three LICs, and to identify risk factors associated with colonization of Kp and its resistant or virulent subpopulations.

## **METHODS**

### ***Study design***

This cross-sectional study was nested within the BIRDY program.<sup>8</sup> Briefly, it consists of a cohort of pregnant women and their newborns in Madagascar, Senegal, and Cambodia with an urban and rural site in each country (Figure S1). Pregnant women enrolled between January 2015 and December 2016 were invited to participate in the present *Klebsiella* carriage study. A maternal stool sample was taken at delivery or, in case of home delivery, shortly after delivery. Data on maternal socio-demographic characteristics, medical history, and exposures, possibly related to Kp and its resistant or virulent subpopulations carriage, including dietary habits, antibiotic consumption, and contacts with animals, were collected.

### ***Klebsiella pneumoniae isolation***

Stool was retrieved using swabs or stool samples, inoculated in a Luria-Bertani broth with amoxicillin (10 mg/L final concentration) and incubated 18 h at 37°C.<sup>9</sup> The enrichment was performed with amoxicillin because Kp isolates carry a constitutively expressed class A beta-

99 lactamase (of *bla*<sub>SHV</sub>, *bla*<sub>OKP</sub> or *bla*<sub>LEN</sub> type) that confers intrinsic resistant to aminopenicillins  
100 (such as ampicillin and amoxicillin), whereas many other bacteria including *E. coli* (the most  
101 abundant potential competitor) are susceptible to amoxicillin. 100 microliters of the  
102 enrichment culture were plated onto selective Simmons citrate with inositol medium agar  
103 plates.<sup>10</sup> Large, yellow, glossy colonies suspected of being *Klebsiella* were identified using  
104 either API20E or mass spectrometry (MALDI-ToF, Bruker). The first isolate identified as  
105 either *Klebsiella pneumoniae* or *Klebsiella variicola* was kept. For 7 carriers, two isolates  
106 genetically distinct based on 7-gene MLST were retained.<sup>11</sup>

### 107 ***Antimicrobial susceptibility testing***

108 The susceptibility of Kp isolates to 16 antimicrobial agents (amoxicillin + clavulanic acid,  
109 ticarcillin + clavulanic acid, piperacillin + tazobactam, cephalothin, cefotaxime, ceftazidime,  
110 cefepime, aztreonam, imipenem, gentamicin, amikacin, tobramycin, nalidixic acid,  
111 ciprofloxacin, tetracycline, and trimethoprim + sulfamethoxazole) was defined using the agar  
112 disk diffusion method and interpreted according to EUCAST 2015 guidelines, except for  
113 cephalothin, nalidixic acid and tetracycline (CLSI M100-S25). ESBL isolates were identified  
114 using the double-disk method.

### 115 ***Genomic sequencing and bioinformatics analyses***

116 NextSeq-500 sequencing of Nextera XT 2×150 nt libraries was performed. Genomic  
117 assemblies (Table S1) were obtained using SPAdes v3.9.<sup>12</sup> Multilocus sequence typing  
118 (MLST) was performed using the BIGSdb-Kp database (<https://bigsdb.pasteur.fr>).<sup>11,13</sup> This  
119 resource was also used for detecting quinolone target genes mutations, and variants of the *wzi*  
120 and intrinsic class A beta-lactamases.<sup>13,14</sup> Novel variants were submitted to BIGSdb (for  
121 *bla*<sub>SHV</sub>, *bla*<sub>OKP</sub>, and *bla*<sub>LEN</sub> variants) and to NCBI beta-lactamases database (*bla*<sub>SHV-206</sub> to  
122 *bla*<sub>SHV-228</sub>). Other beta-lactamases and resistance gene families were detected using Kleborate

(<https://github.com/katholt/Kleborate>). Isolates that carried only *bla*<sub>SHV</sub> (or *bla*<sub>OKP</sub> or *bla*<sub>LEN</sub>), *fosA* and *oqxAB* genes were considered as having an ancestral resistome. Other antimicrobial resistance genes were considered acquired. To evaluate the correlation between the antimicrobial resistance phenotype and genotype, and to depict co-resistance networks, we constructed a correlation matrix for binary variables (in the case of resistance genes: 1, presence; 0, absence; in the case of antimicrobial drugs: 1, resistant/intermediate; 0, susceptible) using the ‘corr.test’ function (Pearson method, which for a pair of binary variables equates to the Phi coefficient) from the ‘corrplot’ R package. Significant correlations were visualized utilizing the ‘corrplot’ function from the same package.

Plasmids were detected using PlasmidFinder

(<https://cge.cbs.dtu.dk/services/PlasmidFinder/>).<sup>15</sup> Kleborate

(<https://github.com/katholt/Kleborate>) was used to search for yersiniabactin, colibactin, aerobactin, and salmochelin clusters, and to predict the capsular types.<sup>16,17</sup> BIGSdb-Kp was used to define the presence of other virulence-associated genes (Table S1).<sup>18</sup> Isolates harboring at least one of the genes *rmpA* and *rmpA2*, and/or at least one complete gene cluster among *iucABCD-iutA* (aerobactin), *iroBCDN* (salmochelin), *ybtAEPQSTUX-fuyA-irp1-irp2* (yersiniabactin), *clbABCDEFGHJKLMNOPQR* (colibactin) and *allABCDRS* (allantoinase) were considered virulent. Hypervirulent Kp were defined as isolates harboring at least one of the genes *rmpA* and *rmpA2*, and/or at least one complete gene cluster among *iucABCD-iutA* (aerobactin) and *iroBCDN* (salmochelin). We defined ‘high-risk clones’ as clones harboring 7-gene MLST sequence types (ST) represented at least 10 times in NCBI genomes (April 2019: 6,258 genomes available) and mentioned in the title or abstract of at least five publications in NCBI PubMed (“Klebsiella” + “pneumoniae” + “STxxx”, April 2019).

#### ***Phylogenetic and statistical analyses***

Genome phylogeny was derived from MASH distances using JolyTree  
(<https://gitlab.pasteur.fr/GIPhy/JolyTree>) and visualized using iTOL (<https://itol.embl.de/>).<sup>19</sup>  
Unique strains were defined with a  $\leq 10$  allelic mismatch cutoff within allelic variation  
recorded using BIGSdb genome comparator function based on the 5,073 protein-coding genes  
of strain Kp616 (GCA\_003076555.1; 90% identity and 90% length coverage).

Statistical analyses were performed using Stata version 15 (Stata Corp., College Station, TX).  
We defined multidrug resistant (MDR) strains as being resistant to at least one antimicrobial  
agent in at least 3 antibacterial categories. Chi square or Fisher exact test was used to compare  
the prevalence of Kp subsets (all, MDR, ESBL) and antibiotic resistance patterns among  
countries. Risk factor analysis was performed separately for each country given the inter-  
country variation of the pregnant women's characteristics. We constructed univariate analysis  
to examine associations between women's environmental exposures, and overall Kp, MDR-,  
ESBL-, and virulent Kp carriage. All variables with a univariate p-value  $\leq 0.2$  were included  
in multivariate analysis. We used a logistic regression model for all, MDR, and ESBL-Kp  
subsets. We conducted backwards stepwise elimination of non-significant parameters  
( $p < 0.05$ ). All multivariate models were adjusted for site (urban/rural). For the seven women  
from whom 2 Kp were isolated, both were kept in the multivariate analysis; sensitivity  
analyses performed with and without these duplicates led to identical results.<sup>4,20–22</sup>

## ***Ethics***

The study was approved by the ethics committees of Madagascar (068-MSANP/CE), Senegal  
(SEN 14-20), Cambodia (132 NEHCR) and was authorized by the Institut Pasteur in Paris.  
Written informed consent was given by all participants.

## **RESULTS**

### ***Kp carriage***

A total of 911 pregnant women from Madagascar (n=423), Cambodia (n=152), and Senegal (n=336) were enrolled. For 874 (95.9%) of these, both epidemiological data and a fecal sample were obtained (Figure S2). Women enrolled in the urban sites represented 32.6%, 55.5%, and 35.2% of the total in Madagascar, Cambodia, and Senegal, respectively. The study population (Table 1) was young (mean 27 years; SD 6.4) and underprivileged, as a large majority (79.3%-88.9%) did not complete secondary school and an important fraction did not have toilets within the housing (Madagascar: 93.1%; Cambodia: 49.3%; and Senegal: 29.9%).

Kp colonization was detected in 489 of 874 (55.9%) women. Carriage rate per country (Figure 1A) was significantly lower in Senegal (40.2%, 95% CI [36.3-46.9];  $p<0.001$ ) compared to Madagascar (64.7%, CI [59.9-69.2]) and Cambodia (66.4%, [58.3-73.7]). Carriage rate was lower in the urban area than in the rural area (Figure 1A) in Senegal (29.5% *versus* 45.4%,  $p=0.006$ ) but not in Madagascar (63.6% *versus* 65.2%,  $p=0.7$ ) and Cambodia (61.5% *versus* 70.4%,  $p=0.3$ ).

#### ***Kp phylogenetic diversity***

Genomic sequence-based phylogenetic relationships (Figure 2) revealed the presence of six deep lineages, corresponding to *K. pneumoniae* phylogroups Kp1 (70.1%), Kp2 (2%), Kp3 (10.7%), Kp4 (15.7%) and Kp5 (2.3%) and a novel phylogroup we named Kp7, described elsewhere as *K. africana*.<sup>4</sup> Kp1 was the most frequent phylogroup in each country (68-73.7%), but phylogroups distribution differed among countries ( $p<0.001$ ) (Figure 1B): Kp4 was the second most common group in Cambodia and Senegal (27.8% and 20.4%, respectively) but not in Madagascar (Kp3: 17.2%; Kp4: 8.4%).

A remarkably high genotypic diversity was found based on MLST. There were 325 STs, a majority of which (242, 74.5%) were represented by a single isolate (Figure S3). The genotype distribution into multiple low-frequency STs was reflected by a very high (99.6%) Simpson

diversity index, even within individual countries or sites (range, 98.7 – 99.7%). High-risk clones (Table S2) represented 70 (14.1%) isolates (Table S3). Among 11 STs with  $\geq 5$  isolates, six were high-risk clones and nine were isolated in two or three countries (Figure 1C). However, based on the number of allelic differences between strains within these clonal groups, only two strains were observed in more than one carrier (Figure 1C; Table S1).

### ***Antimicrobial resistance***

Antimicrobial resistance rates differed among the three countries (Figure 3A; Table S4). Low levels of resistance were observed for imipenem (0.8%) and amikacin (0.4%). Resistance rates to quinolones (15.5%), other aminoglycosides (17.5%), and third-generation cephalosporins (14.4%) were higher ( $p < 0.001$ ) in Cambodia compared to Madagascar (2.7%, 8%, and 10%, respectively) and Senegal (~3% each). ESBL-Kp were almost absent in Senegal (0.7%) but they were frequent in Cambodia (14.4%) and Madagascar (8.4%). MDR-Kp rates were higher in Madagascar (39.7%) compared to Cambodia (25.8%) and Senegal (11.7%;  $p < 0.001$ ). Multidrug-resistance was higher in the urban site compared to the rural site only in Cambodia (42.5% vs 14%,  $p = 0.002$ ; Table S4). In Cambodia, 10.3% of the isolates were resistant to at least 8 categories of antibiotics. As expected, higher resistance levels were observed for older antibiotics (amoxicillin and ticarcillin with clavulanate, tetracycline, and trimethoprim/sulfamethoxazole).

Comparisons among phylogroups showed differences only in Cambodia, with higher resistance rates against gentamicin and tetracycline in Kp1 ( $p = 0.04$  and  $p = 0.006$ , respectively; Table S3).

### ***Antimicrobial resistance genes, plasmids and sublineages***

As expected, *oqxAB*, *fosA*, and the chromosomal beta-lactamase gene were ubiquitous (Table S3; Figure 3B); 117 variants of the latter were found, with *bla<sub>SHV-11</sub>* ( $n = 145$ , 29.7%) being the most frequent (Figure S4). Remarkably, 334 (67.4%) of isolates had no acquired resistance

gene nor any GyrA or ParC alterations, thus corresponding to the ancestral (“wild-type”) resistome type. The median number of acquired resistance genes was 3 (similar in each country), ranging from 9 to 11 among ESBL producers. Of the 37 ESBL enzymes, CTX-M-15 was the most common (n=22, 60%). Resistance genes were strongly associated with phenotypic resistance (Figure 3C), and some gene associations were depicted, reflecting their genetic linkage on common genetic platforms (plasmids, transposons, integrons) and resulting in co-resistance phenotypes. For example, *bla*<sub>OXA-1</sub>, *aac*(6')-Ib-cr, *catB*, and *qnrB* were associated with *bla*<sub>CTX-M-15</sub> gene, whereas *bla*<sub>LAP-2</sub>, *qnrS*, and *aac*(3)-IIId were linked to *bla*<sub>CTX-M-14/-27</sub>; detection rates for both combinations were higher in Cambodia (Table S5).

Plasmid replicons were found in 70.6%, 73.9%, and 55.5% of Kp from Madagascar, Cambodia, and Senegal, respectively. Twenty-two replicon types were detected (Table S2), of which IncFIB(K) (36.1%), Col440I (27.4%), IncFII (26.4%), IncR (15.9%), and IncFIA(HI1) (15.1%) were the most frequent. However, the predominant type in Senegal was Col440I (31.4%), and plasmid distribution differed among countries (p<0.001).

Seventy isolates (14.1%) belonged to widespread genotypes commonly associated with ESBL or carbapenemase production (Table S3). Furthermore, among ESBL-Kp, 19 (51.4%) belonged to previously recognized MDR STs. Although ST307 is emerging as a major carbapenemase-producing sublineage<sup>23,24</sup>, our ST307 carriage isolates did not harbor carbapenemase genes.

### ***Virulence genes and convergence with resistance***

Seventy-two (14.5%) virulent isolates were found (Figure 1), with yersiniabactin being the most frequent virulence factor (62 isolates, 12.5%; Figure S5; TableS5). The regulator of mucoid phenotype genes *rmpA* and *rmpA2* were present in six isolates (1.2%), whereas aerobactin and salmochelin clusters were present in 14 (2.8%) and 11 (2.2%) isolates, respectively; these hypervirulence-associated clusters were observed mostly in Madagascar

and were associated with STs of capsular serotype K2 previously described as hypervirulent<sup>25,26</sup>: ST25, ST65, ST375, and ST380 (Table S3). In contrast, no isolate of the liver abscess-associated serotype K1 ST23 was found.

Virulence and antibiotic resistance elements were generally observed in distinct isolates (Table S3; Figure S6). In particular, aerobactin and salmochelin hypervirulence clusters were associated with ancestral resistome isolates (in Madagascar,  $p=0.05$  and  $p=0.06$ , respectively). However, in one urban Cambodian isolate (SB5663, ST17), a convergence of MDR (*bla*<sub>CTX-M-27</sub>, *bla*<sub>LAP-2</sub>, *dfrA*, *sul2*, *tetA*, *floR*) and hypervirulence (*rmpA*, yersiniabactin, salmochelin/ICEKp1) was observed (Table S3; Figure S6).

#### **Associations between environmental exposures and specific characteristics of Kp**

Risk factors of Kp, MDR, ESBL, and virulent Kp carriage were investigated (Table 2; Table S7). The use of antibiotics during pregnancy was associated with a higher risk of Kp carriage (in Madagascar, adjusted odds ratio (aOR)=2.1;  $p=0.03$ ), as was dry fish consumption (in Cambodia: aOR=2;  $p=0.05$ ) and contact with chicken (in Senegal: aOR=1.9;  $p=0.008$ ). In contrast, hands washing after using toilets (in Madagascar: aOR=0.5;  $p=0.02$ ), but also manipulating animal excrement (in Madagascar, aOR=0.5;  $p=0.008$ ), were associated with a reduced risk of Kp carriage.

Among Kp-colonized women, those who reported washing their hands after using toilets (in Madagascar: aOR=0.4;  $p=0.02$ ), to eat dried meat (in Madagascar: aOR=0.4;  $p=0.01$ ), or to be in contact with chicken (in Senegal: aOR=0.2;  $p=0.01$ ) were less at risk of being MDR-Kp colonized. In contrast, women who reported eating raw vegetables at least once per week (in Madagascar: aOR=2.4;  $p=0.03$ ) were more at risk of being MDR-Kp colonized.

Regarding virulent Kp carriage, hospitalization of a household member in the previous year (in Madagascar: aOR=4.0;  $p=0.001$ ), consumption of rabbit meat (in Madagascar: aOR=3.6;

p=0.01), and use of fecal material as fertilizer (Senegal: 3.7, p=0.02) were risk factors. In contrast, fish consumption was associated with decreased virulent Kp colonization (Senegal: OR=0.2; p=0.02).

No risk factor for MDR and virulent Kp carriage in Cambodia and for ESBL Kp carriage in the 3 countries were identified.

## DISCUSSION

Despite the importance of Kp carriage in the ecology and spread of this pathogen, knowledge on Kp colonization is scarce, especially in LIC communities. In previous carriage studies, other *Enterobacteriaceae* including *E. coli* were considered together with Kp, which represented a minority of studied isolates.<sup>27,28</sup> In addition, previous work focused on ESBL- or carbapenemase-producing *Enterobacteriaceae* carriage, overlooking the natural diversity and ecology of susceptible populations from which resistant strains evolve.

The design of our community-based multicentric study is unique in that the naturally colonizing population of Kp was investigated irrespective of antimicrobial resistance phenotypes, providing insights into the natural ecology of Kp in human carriage. We found that Kp carriage rate per country ranged from 40% to 66%. Kp community carriage studies in high-income countries found carriage rates ranging from 6% to 35%; rates were much higher in one study of Chinese volunteers living in Asian countries.<sup>5–7,29</sup> Although hospitalization is known to increase Kp carriage, our estimates are even higher than the prevalence of Kp carriage reported in hospital settings, typically ranging from 20 to 35%.<sup>5,6,27</sup> Kp abundance in the gut microbiota is low; combining an enrichment step with a *Klebsiella* selective medium may have contributed to recover Kp at higher rates here than in previous studies.<sup>30</sup> Of concern, pregnant women can transmit to their children during delivery or after birth, potentially leading to neonatal sepsis.<sup>31</sup> Our data demonstrate that Kp1 is the most prevalent phylogroup in all sites. Currently, recognized high-risk clones all belong to Kp1, and this phylogroup is the most frequent in

human infections.<sup>26,32</sup> Our results suggest that the clinical predominance of Kp1 might be a direct reflection of the ecological dominance of this group in carriage, rather than to a higher infectious potential compared to other phylogroups. Consistent with this hypothesis, the high rate of Kp4 carriage in Cambodia is mirrored by common Kp4 bloodstream infections in this country.<sup>3</sup>

A striking inter-carrier genetic diversity of Kp was found, with almost three quarters of isolates having a unique ST. Further, genomic sequencing ruled out in all cases except two, the occurrence of recent transmission. Human Kp carriage populations thus appear structured into individual carriers acting as rarely connected islands, favoring evolutionary diversification into a multitude of genotypes.

Kp isolates causing hypervirulent infections belong to recognized sublineages of capsular serotypes K1 and K2, but the source of these infections is poorly known. K1-ST23 was common among Asian healthy carriers.<sup>33</sup> Here, several isolates belonging to hypervirulent K2 sublineages were found, suggesting that the human gut may act as a reservoir of these strains.

Antimicrobial resistance differed among countries, likely reflecting the different levels of antibiotic use, hygiene, and infection control between the 3 countries. The proportion of MDR and ESBL-producing isolates was high in Madagascar and Cambodia. These findings are worrisome, as the gut is a well-known reservoir of infections by these strains.<sup>5,6</sup> Besides, the role of Kp as a source of ESBL genes and other resistance determinants that can be transmitted horizontally to other pathogens is a clear cause of concern.<sup>34</sup> We did not find any carbapenemase-producing Kp, indicating that its spread in the community is still limited in these countries, consistent with low carbapenem consumption.<sup>35</sup>

We were able to investigate for the first time, risk factors associated with natural carriage of Kp and its genetic or phenotypic subpopulations. We found that women who washed their hands after toilets use were less likely to be colonized with Kp or MDR Kp in Madagascar. Hand

washing is effective in the transmission of fecal-oral infection in the community, but is often neither properly done nor practiced at key times.<sup>36</sup> Our finding supports that promotion of a simple hygienic behavior could contribute to the control of the spread of antimicrobial resistance in the community.

Consumption of food or contacts with some animals were found associated with Kp, MDR, or virulent Kp carriage. Raw vegetables or rabbit meat (although rabbit meat consumption was low) were associated with a higher risk of being colonized with MDR-Kp or virulent Kp, respectively, suggesting a possible food origin for these isolates. Bacteria can persist on products that are inadequately cooked prior to consumption. However, at which step of the food supply chain these products are contaminated remains an open question. This study calls for more focused investigations of the presence of Kp on food. We also found that manipulating animal excrement and having animal contacts appear protective of Kp and MDR-Kp colonization. One explanation could lie in negative ecological associations between Kp and other bacteria during animal hosts colonization. Future studies on interactions of Kp with the gut microbiota are warranted.

We acknowledge several limitations to this study. Due to the cross-sectional design, the interpretation of the risk factors must be cautious, in particular due to recall bias. Our sample size was not very large, specifically in Cambodia, which might explain that we found few risk factors associated with MDR or virulent Kp. Although the majority of the samples consisted in a stool collection (85%), endorectal swabbing was performed instead when stool was not available. However, previous studies showed that bacterial communities defined from endorectal swabbing and stool specimens are similar.<sup>37,38</sup> In addition, we used an enrichment step that allows the recovery of Kp isolates even when the amount of starting material is low, which could potentially be the case when using rectal swabs.

Human-to-human contact was not investigated in depth here. Characteristics reflecting a

potential human transmission (number of household members) were collected and were not significantly associated with Kp carriage. We performed our carriage study in pregnant women, who represent a homogeneous and generally healthy population, with limited socioeconomic status bias. However, the immune system is modified during pregnancy, impacting the gut microbiota.<sup>39</sup> Whether pregnancy has an impact on Kp carriage would require dedicated studies. Finally, food or environmental sampling performed in parallel to human screening would have allowed comparing Kp populations in these niches, possibly leading to their association with human carriage, but these studies will be particularly challenging in light of the huge diversity of Kp revealed herein.<sup>40</sup>

## CONCLUSIONS

To conclude, this multicentric study provides the most detailed picture to date of Kp colonization in human LICs communities. One important outcome is the high prevalence of Kp carriage, and high levels of antimicrobial resistance among carriage Kp isolates. Our findings also illustrate the challenges of understanding the transmission dynamics of ubiquitous pathogens. As Kp carriage risk factors were country-dependent, major drivers could not be pointed out, probably reflecting the influence of a complex combination of antibiotic use, hygienic and environmental conditions in the given local context. Consequently, our results suggest that there is “no one size fits all” intervention to tackle the Kp community burden and that public health strategies to control Kp spread may need to be defined based on local specificities.

## **Competing interest**

The authors declare that they have no competing interests.

## **Funding**

This work was supported by the Institut Pasteur programme Action Concertée Inter-Pasteurienne (Grant A-14-2014) and by the French government's Investissement d'Avenir program Laboratoire d'Excellence 'Integrative Biology of Emerging Infectious Diseases' (grant ANR-10-LABX-62-IBEID). Support to the BIRDY project was provided by the Total Foundation, MSD Avenir and the Monaco Department of International Cooperation.

CR was supported financially by the MedVetKlebs project, a component of European Joint Programme One Health EJP, which has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 773830.

The funders had no role in study design, data collection and analysis, interpretation, or writing of the manuscript.

## **Acknowledgements**

We thank all collaborators of the BIRDY program: Andrianonimiadana Lova Maminirina, Andriambololona Mino, Herisoa Feno, Rabearitiana Lydos, Rahelirivao Bodonirina Tanjona, Rakotobe Miora, Rakotoarimanana Feno Manitra Jacob, Randriarilala Heritiana, Randrianirina Frédérique, Volahasina Antsa Tanjona, Andrianirina Zafitsara Zo, Vincent Richard, Awa Ndir, Amy Gassama Sow, Balla Sy, Marguerite Diatta, Pape Samba Dieye, Jean Baptiste Diouf, Joseph Faye, Abibatou Ndiaye, Bouya Ndao, Thida Chon, Veronique Ngo, Arnaud Tarantola, Sok Touch, Patrice Piola, Sophie Goyet, Siyin Lach, Elsa Kermorvant-Duchemin, Michael Padget, Laurence Watier, and Abdou Armya Youssouf.

385 We thank all physicians, laboratory staff, field interviewers, and community workers involved  
386 in the project. We thank Leonardo Panunzi for assistance with genomic analyses of resistance  
387 genes, and the “Plateforme de Microbiologie Mutualisée” of Institut Pasteur for genomic  
388 sequencing. We are grateful to all women participating in the BIRDY program.

## 389    **References:**

- 390    1.    Woodford N, Turton JF, Livermore DM. Multiresistant Gram-negative bacteria: the role  
391        of high-risk clones in the dissemination of antibiotic resistance. *FEMS Microbiol Rev*  
392        2011; 35:736–55.
- 393    2.    Zaidi AK, Thaver D, Ali SA, Khan TA. Pathogens associated with sepsis in newborns  
394        and young infants in developing countries. *Pediatr Infect J* 2009; 28:S10-8.
- 395    3.    Wyres KL, Wick RR, Judd LM, Froumine R, Tokolyi A, Gorrie CL, Lam MMC,  
396        Duchêne S, Jenney A, Holt KE. Distinct evolutionary dynamics of horizontal gene  
397        transfer in drug resistant and virulent clones of *Klebsiella pneumoniae*. *PLoS Genet*  
398        2019; 15:e1008114.
- 399    4.    Rodrigues C, Passet V, Rakotondrasoa A, Diallo TA, Criscuolo A, Brisse S. Description  
400        of *Klebsiella africanensis* sp. nov., *Klebsiella variicola* subsp. *tropicalensis* subsp. nov.  
401        and *Klebsiella variicola* subsp. *variicola* subsp. nov. *Res Microbiol* 2019;
- 402    5.    Gorrie CL, Mirceta M, Wick RR, Edwards DJ, Thomson NR, Strugnell RA, Pratt NF,  
403        Garlick JS, Watson KM, Pilcher DV, et al. Gastrointestinal Carriage Is a Major  
404        Reservoir of *Klebsiella pneumoniae* Infection in Intensive Care Patients. *Clin Infect Dis*  
405        2017; 65:208–15.
- 406    6.    Martin RM, Cao J, Brisse S, Passet V, Wu W, Zhao L, Malani PN, Rao K, Bachman  
407        MA. Molecular Epidemiology of Colonizing and Infecting Isolates of *Klebsiella*  
408        *pneumoniae*. *mSphere* [Internet] 2016; 1. Available from:  
409        <http://www.ncbi.nlm.nih.gov/pubmed/27777984>
- 410    7.    Brisse S, Grimont F, Grimont PA, Schleifer K-H, Stackebrandt E. The Genus *Klebsiella*.  
411        In: Dworkin M, Falkow S, Rosenberg E, editors. *The Prokaryotes. A handbook on the*  
412        *biology of bacteria*. Springer; 2006.
- 413    8.    Huynh B-T, Kermorvant-Duchemin E, Herindrainy P, Padget M, Rakotoarimanana FMJ,  
414        Feno H, Hariniaina-Ratsima E, Rahelirivao T, Ndir A, Goyet S, et al. Bacterial  
415        Infections in Neonates, Madagascar, 2012-2014. *Emerg Infect Dis* 2018; 24:710–7.
- 416    9.    Haeggman S, Löfdahl S, Paauw A, Verhoef J, Brisse S. Diversity and evolution of the  
417        class A chromosomal beta-lactamase gene in *Klebsiella pneumoniae*. *Antimicrob Agents*  
418        *Chemother* 2004; 48:2400–8.
- 419    10.    Van Kregten E, Westerdaal NA, Willers JM. New, simple medium for selective recovery  
420        of *Klebsiella pneumoniae* and *Klebsiella oxytoca* from human feces. *J Clin Microbiol*  
421        1984; 20:936–41.
- 422    11.    Diancourt L, Passet V, Verhoef J, Grimont PAD, Brisse S. Multilocus sequence typing  
423        of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol* 2005; 43:4178–82.
- 424    12.    Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,  
425        Nikolenko SI, Pham S, Prjibelski AD, et al. SPAdes: a new genome assembly algorithm  
426        and its applications to single-cell sequencing. *J Comput Biol* 2012; 19:455–77.

- 427 13. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb  
428 software, the PubMLST.org website and their applications. Wellcome Open Res 2018;  
429 3:124.
- 430 14. Brisse S, Passet V, Haugaard AB, Babosan A, Kassis-Chikhani N, Struve C, Decre D.  
431 wzi Gene sequencing, a rapid method for determination of capsular type for *Klebsiella*  
432 strains. J Clin Microbiol 2013; 51:4073–8.
- 433 15. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller  
434 Aarestrup F, Hasman H. In silico detection and typing of plasmids using PlasmidFinder  
435 and plasmid multilocus sequence typing. Antimicrob Agents Chemother 2014; 58:3895–  
436 903.
- 437 16. Lam MMC, Wyres KL, Judd LM, Wick RR, Jenney A, Brisse S, Holt KE. Tracking key  
438 virulence loci encoding aerobactin and salmochelin siderophore synthesis in *Klebsiella*  
439 pneumoniae. Genome Med 2018; 10:77.
- 440 17. Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, Holt KE.  
441 Identification of *Klebsiella* capsule synthesis loci from whole genome data. Microb  
442 Genomics 2016; 2:e000102.
- 443 18. Bojer MS, Struve C, Ingmer H, Hansen DS, Krogfelt KA. Heat resistance mediated by a  
444 new plasmid encoded Clp ATPase, ClpK, as a possible novel mechanism for nosocomial  
445 persistence of *Klebsiella pneumoniae*. PloS One 2010; 5:e15467.
- 446 19. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy  
447 AM. Mash: fast genome and metagenome distance estimation using MinHash. Genome  
448 Biol 2016; 17:132.
- 449 20. Blin C, Passet V, Touchon M, Rocha EPC, Brisse S. Metabolic diversity of the emerging  
450 pathogenic lineages of *Klebsiella pneumoniae*. Env Microbiol 2017; 19:1881–98.
- 451 21. Brisse S, Verhoef J. Phylogenetic diversity of *Klebsiella pneumoniae* and *Klebsiella*  
452 oxytoca clinical isolates revealed by randomly amplified polymorphic DNA, *gyrA* and  
453 *parC* genes sequencing and automated ribotyping. Int J Syst Evol Microbiol 2001;  
454 51:915–24.
- 455 22. Long SW, Linson SE, Ojeda Saavedra M, Cantu C, Davis JJ, Brettin T, Olsen RJ.  
456 Whole-Genome Sequencing of a Human Clinical Isolate of the Novel Species *Klebsiella*  
457 quasivariicola sp. nov. Genome Announc 2017; 5.
- 458 23. Villa L, Feudi C, Fortini D, Brisse S, Passet V, Bonura C, Endimiani A, Mammina C,  
459 Ocampo AM, Jimenez JN, et al. Diversity, virulence, and antimicrobial resistance of the  
460 KPC-producing *Klebsiella pneumoniae* ST307 clone. Microb Genomics 2017;  
461 3:e000110.
- 462 24. Wyres KL, Hawkey J, Hetland MAK, Fostervold A, Wick RR, Judd LM, Hamidian M,  
463 Howden BP, Löhr IH, Holt KE. Emergence and rapid global dissemination of CTX-M-  
464 15-associated *Klebsiella pneumoniae* strain ST307. J Antimicrob Chemother 2019;  
465 74:577–81.

- 466 25. Bialek-Davenet S, Criscuolo A, Ailloud F, Passet V, Jones L, Delannoy-Vieillard AS,  
467 Garin B, Le Hello S, Arlet G, Nicolas-Chanoine MH, et al. Genomic definition of  
468 hypervirulent and multidrug-resistant *Klebsiella pneumoniae* clonal groups. *Emerg*  
469 *Infect Dis* 2014; 20:1812–20.
- 470 26. Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, Dance D, Jenney A,  
471 Connor TR, Hsu LY, Severin J, et al. Genomic analysis of diversity, population  
472 structure, virulence, and antimicrobial resistance in *Klebsiella pneumoniae*, an urgent  
473 threat to public health. *Proc Natl Acad Sci U A* 2015; 112:E3574-81.
- 474 27. Luvsansharav UO, Hirai I, Nakata A, Imura K, Yamauchi K, Niki M, Komalamisra C,  
475 Kusolsuk T, Yamamoto Y. Prevalence of and risk factors associated with faecal carriage  
476 of CTX-M beta-lactamase-producing Enterobacteriaceae in rural Thai communities. *J*  
477 *Antimicrob Chemother* 2012; 67:1769–74.
- 478 28. Chereau F, Herindrainy P, Garin B, Huynh BT, Randrianirina F, Padget M, Piola P,  
479 Guillemot D, Delarocque-Astagneau E. Colonization of extended-spectrum-beta-  
480 lactamase- and NDM-1-producing Enterobacteriaceae among pregnant women in the  
481 community in a low-income country: a potential reservoir for transmission of  
482 multiresistant Enterobacteriaceae to neonates. *Antimicrob Agents Chemother* 2015;  
483 59:3652–5.
- 484 29. Lin Y-T, Siu LK, Lin J-C, Chen T-L, Tseng C-P, Yeh K-M, Chang F-Y, Fung C-P.  
485 Seroepidemiology of *Klebsiella pneumoniae* colonizing the intestinal tract of healthy  
486 chinese and overseas chinese adults in Asian countries. *BMC Microbiol* 2012; 12:13.
- 487 30. Conlan S, Kong HH, Segre JA. Species-level analysis of DNA sequence data from the  
488 NIH Human Microbiome Project. *PLoS One* 2012; 7:e47075.
- 489 31. Chan GJ, Lee AC, Baqui AH, Tan J, Black RE. Risk of early-onset neonatal infection  
490 with maternal infection or colonization: a global systematic review and meta-analysis.  
491 *PLoS Med* 2013; 10:e1001502.
- 492 32. Brisse S, van Himbergen T, Kusters K, Verhoef J. Development of a rapid identification  
493 method for *Klebsiella pneumoniae* phylogenetic groups and analysis of 420 clinical  
494 isolates. *Clin Microbiol Infect* 2004; 10:942–5.
- 495 33. Chung DR, Lee H, Park MH, Jung SI, Chang HH, Kim YS, Son JS, Moon C, Kwon KT,  
496 Ryu SY, et al. Fecal carriage of serotype K1 *Klebsiella pneumoniae* ST23 strains closely  
497 related to liver abscess isolates in Koreans living in Korea. *Eur J Clin Microbiol Infect*  
498 *Dis* 2012; 31:481–6.
- 499 34. Wyres KL, Holt KE. *Klebsiella pneumoniae* as a key trafficker of drug resistance genes  
500 from environmental to clinically important bacteria. *Curr Opin Microbiol* 2018; 45:131–  
501 9.
- 502 35. Klein EY, Van Boeckel TP, Martinez EM, Pant S, Gandra S, Levin SA, Goossens H,  
503 Laxminarayan R. Global increase and geographic convergence in antibiotic consumption  
504 between 2000 and 2015. *Proc Natl Acad Sci U S A* 2018; 115:E3463–70.

- 505 36. Halder AK, Tronchet C, Akhter S, Bhuiya A, Johnston R, Luby SP. Observed hand  
506 cleanliness and other measures of handwashing behavior in rural Bangladesh. BMC  
507 Public Health 2010; 10:545.
- 508 37. Bassis CM, Moore NM, Lolans K, Seekatz AM, Weinstein RA, Young VB, Hayden  
509 MK, CDC Prevention Epicenters Program. Comparison of stool versus rectal swab  
510 samples and storage conditions on bacterial community profiles. BMC Microbiol 2017;  
511 17:78–78.
- 512 38. Reyman M, van Houten MA, Arp K, Sanders EAM, Bogaert D. Rectal swabs are a  
513 reliable proxy for faecal samples in infant gut microbiota research based on 16S-rRNA  
514 sequencing. Sci Rep 2019; 9:16072–16072.
- 515 39. Koren O, Goodrich JK, Cullender TC, Spor A, Laitinen K, Kling Bäckhed H, Gonzalez  
516 A, Werner JJ, Angenent LT, Knight R, et al. Host Remodeling of the Gut Microbiome  
517 and Metabolic Changes during Pregnancy. Cell 2012; 150:470–80.
- 518 40. Hanage WP. Two Health or Not Two Health? That Is the Question. mBio 2019; 10.

519

## Figures

### Figure 1. Rates of Kp carriage and genotype diversity.

**A.** Prevalence of Kp carriage across countries and sites. The difference of prevalence between urban and rural sites was statistically significant in Senegal. CI, confidence interval. **B.** Phylogroup distribution in the three countries. Phylogroup Kp6 was not found; phylogroup Kp7 was only found in one isolate from Senegal. **C.** Frequency and geographic distribution of the most common carriage STs. 1% frequency corresponds to 5 isolates. Urban bar sectors are dotted with white. Within single STs, genome-wide variation ruled-out recent transmission among individual carriers, except in two instances (< 10 allelic mismatches out of 5,073 gene loci) in Madagascar (ST37: urban-rural triplet; ST48: rural pair) represented by oval shapes inside the bar sectors. High-risk clones are represented by an asterisk above the bars.

### Figure 2. Phylogenetic relationships.

The scale bar corresponds to 0.01 substitutions per site. iTOL (<https://itol.embl.de/>) was used to visualize country of origin (circles at branch tips), phylogroups (background color in first circle comprising the isolates names), the ancestral or acquired character of isolates' resistome (second circle), ESBL phenotype (third circle), and virulence genes as indicated in front of each external circle.

### Figure 3. Antimicrobial resistance.

**A.** Rates (%) of phenotypic non-susceptibility. ESBL, extended-spectrum beta-lactamases detected by the double-disk synergy test (DDST); IMI, imipenem; AMK, amikacin; GEN, gentamicin; CIP, ciprofloxacin; SXT, trimethoprim-sulfamethoxazole. **B.** Prevalence (%) of resistance genes. **C.** Correlation matrix between antimicrobial resistance genotype and

542 phenotype (Pearson method). Blank squares represent correlations without statistical  
543 significance ( $p > 0.05$ ). Positive correlation is depicted by blue circles, whereas red circles  
544 represent significant negative correlation. The size and strength of color represent the  
545 numerical value of the Phi correlation coefficient. Black boxes represent genes commonly  
546 found together in the same strain. The plot was created with the ‘corrplot’ R package.  
547 Antimicrobial agents: CEP, cephalothin; TOB, tobramycin; GEN, gentamicin; SXT,  
548 trimethoprim-sulfamethoxazole; TET, tetracycline; CAZ, ceftazidime; ATM, aztreonam;  
549 CTX, cefotaxime; FEP, cefepime; DDST, double-disk synergy test; CIP, ciprofloxacin; NAL,  
550 nalidixic acid; TZP, piperacillin-tazobactam; AMC, amoxicillin-clavulanic acid; TCC,  
551 ticarcillin; IMI, imipenem; AMK, amikacin.

552 **Tables**

553 **Table 1.** Characteristics of the inclusion population

|                                          | Madagascar         | Cambodia   | Senegal    | p-value |
|------------------------------------------|--------------------|------------|------------|---------|
|                                          | N=405              | N=146      | N=323      |         |
|                                          | n (%) or mean (SD) |            |            |         |
| <b>Urban Site</b>                        | 132 (32.6)         | 81 (55.5)  | 105 (35.2) | <0.001  |
| <b>Age (years)</b>                       | 25.9 (6.4)         | 27 (5.7)   | 28 (6.71)  | 0.08    |
| <b>Gestity</b>                           |                    |            |            |         |
| 1                                        | 121 (29.9)         | 48 (32.9)  | 70 (21.7)  | 0.04    |
| >1                                       | 284 (70.1)         | 119 (67.1) | 253 (78.3) |         |
| <b>Education</b>                         |                    |            |            |         |
| Absence/primary school                   | 90 (22.2)          | 78 (53.4)  | 227 (70.3) | <0.001  |
| Partial secondary school                 | 231 (57.0)         | 50 (34.2)  | 62 (19.2)  |         |
| Complete secondary or higher             | 84 (20.7)          | 18 (12.3)  | 34 (11.1)  |         |
| <b>Occupancy</b>                         |                    |            |            |         |
| Unemployed                               | 263 (64.9)         | 50 (34.3)  | 258 (79.9) | <0.001  |
| Manual job                               | 129 (31.9)         | 90 (61.6)  | 55 (17.0)  |         |
| Clerical job                             | 13 (3.2)           | 6 (4.1)    | 10 (3.1)   |         |
| <b>House characteristics</b>             |                    |            |            |         |
| Individual housing                       | 135 (33.3)         | 93 (64.1)  | 210 (65.0) | <0.001  |
| Housing within a compound                | 133 (32.8)         | 30 (20.7)  | 73 (22.6)  |         |
| Accommodation shared with other families | 137 (33.8)         | 22 (15.2)  | 40 (12.4)  |         |
| <b>Electricity</b>                       | 299 (73.8)         | 145 (99.3) | 307 (95.6) | <0.001  |
| <b>Toilets</b>                           |                    |            |            |         |
| Inside the house                         | 28 (6.9)           | 74 (50.6)  | 225 (70.2) | <0.001  |
| Outside the house                        | 377 (93.1)         | 72 (49.3)  | 96 (29.9)  |         |
| <b>Pregnancy follow-up</b>               |                    |            |            |         |
| Skilled health-care workers              | 384 (97.2)         | 142 (97.3) | 322 (99.7) | 0.02    |

571

**Table 2.** Environmental exposures and colonization with Kp, MDR KP and virulent Kp

|                                         | Kpn +           | Kpn -           | Multivariate analysis* |       |                                       | MDR Kp       | non MDR Kp | Multivariate analysis* |      |                                              | Virulent Kp | non virulent Kp | Multivariate analysis* |       |
|-----------------------------------------|-----------------|-----------------|------------------------|-------|---------------------------------------|--------------|------------|------------------------|------|----------------------------------------------|-------------|-----------------|------------------------|-------|
|                                         | N=262<br>(n, %) | N=143<br>(n, %) | aOR<br>[95%CI]         | p     |                                       | N=104        | N=158      | aOR                    | p    |                                              | N=34        | N=228           | aOR                    | p     |
| <b>Madagascar</b>                       |                 |                 |                        |       |                                       |              |            |                        |      |                                              |             |                 |                        |       |
| <b>Antibiotherapy during pregnancy</b>  | 42 (16.0)       | 15(10.5)        | 2.1<br>[1.1-4.1]       | 0.03  | <b>Hand washing after toilets use</b> | 65(70.7)     | 107(81.1)  | 0.4<br>[0.2-0.9]       | 0.02 | <b>Rabbit meat consumption</b> §§            | 7(20.6)     | 15(6.6)         | 3.6<br>[1.3-9.8]       | 0.01  |
| <b>Hand washing after toilets use</b>   | 172(76.9)       | 115(86.4)       | 0.5<br>[0.3-0.9]       | 0.02  | <b>Dry meat consumption</b> §§        | 12(11.9)     | 32(20.9)   | 0.4<br>[0.2-0.8]       | 0.01 | <b>Hospitalization of a household member</b> | 7(20.6)     | 13(5.8)         | 4.0<br>[1.4-11.2]      | 0.009 |
| <b>Manipulation of animal excrement</b> | 35(13.4)        | 31(21.7)        | 0.5<br>[0.2-0.8]       | 0.008 | <b>Raw vegetables consumption</b> §   | 91<br>(88.4) | 128 (82.1) | 2.4<br>[1.1-5.5]       | 0.03 |                                              |             |                 |                        |       |
| <b>Cambodia</b>                         |                 |                 |                        |       |                                       |              |            |                        |      |                                              |             |                 |                        |       |
|                                         | N=97            | N=49            | aOR                    | p     |                                       |              |            |                        |      |                                              |             |                 |                        |       |
| <b>Dry fish consumption §</b>           | 51(52.6)        | 17(34.7)        | 2.0<br>[1.0-4.1]       | 0.05  |                                       |              |            |                        |      |                                              |             |                 |                        |       |
| <b>Senegal</b>                          |                 |                 |                        |       |                                       |              |            |                        |      |                                              |             |                 |                        |       |
|                                         | N=137           | N=193           | aOR                    | p     |                                       | N=16         | N=121      | aOR                    | p    |                                              | N=29        | N=108           | aOR                    | p     |
| <b>Contact with chicken</b>             | 82(59.9)        | 78(40.4)        | 1.9<br>[1.2-3.1]       | 0.008 | <b>Contact with chicken</b>           | 5(31.3)      | 77(63.6)   | 0.2<br>[0.06-0.7]      | 0.01 | <b>Fish consumption §</b>                    | 23 (79.3)   | 101(93.5.)      | 0.2<br>[0.1-0.8]       | 0.02  |
|                                         |                 |                 |                        |       |                                       |              |            |                        |      | <b>Use of fecal material as fertilizer</b>   | 8(27.6)     | 11(10.2)        | 3.7<br>[1.7-11.0]      | 0.02  |

. \* All multivariate analyses are adjusted on site. Only significant parameters are shown

§ Consumption frequency was categorized as “at least once per week” versus “less than once per week”

§§ Consumption frequency was categorized as “ever” versus “never”

## **Supplementary data**

**Table S1.** *Klebsiella* isolates and genome sequence information.

**Table S2.** High-risk clones.

**Table S3.** Characteristics of the 496 *Klebsiella* isolates: genotypes, antimicrobial susceptibility, capsular types, resistance genes, virulence genes, metal tolerance gene clusters and plasmid replicons.

**Table S4.** Frequencies of antimicrobial resistance phenotypes among *K. pneumoniae* isolates, by country and site.

**Table S5.** Frequencies of antimicrobial resistance genes among *K. pneumoniae* isolates, by country and site.

**Table S6.** Frequencies of virulence-associated genes among *K. pneumoniae* isolates, by country and site.

**Table S7.** Environmental exposures and colonization with Kp, MDR-Kp and virulent Kp among pregnant women.

**Figure S1.** Geographical maps with study sites.

**Figure S2.** Flow charts of inclusion of pregnant women and Kp isolates.

**Figure S3.** MLST sequence type (ST) diversity.

**Figure S4.** Diversity of the beta-lactamases-encoding genes (*bla<sub>SHV</sub>*, *bla<sub>OKP</sub>*, *bla<sub>LEN</sub>*).

**Figure S5.** Rates of virulence clusters or genes, by country of isolation.

**Figure S6.** Details of presence per isolate of antimicrobial resistance phenotype, and of genes coding for antimicrobial resistance, virulence or plasmid replicons and metal tolerance (MT).

Isolates are ordered vertically according to the same phylogeny (left) as in the main

phylogeny Figure. ST, sequence type. Note that cluster *kfuABC* was conserved in non-Kp1 isolates but not within Kp1.

**Table S1. Klebsiella isolates and their genomic characteristics**

| Isolate | Species               | Phylo-group | ST   | Country  | Site  | Isolation year | from same donor | Unique strain # | Sequencing coverage | No. Contigs | Assembly size (nt) | N50    | Accession Number |
|---------|-----------------------|-------------|------|----------|-------|----------------|-----------------|-----------------|---------------------|-------------|--------------------|--------|------------------|
| SB5624  | Klebsiella pneumoniae | Kp1         | 661  | Cambodia | Rural | 2016           |                 |                 | 86                  | 72          | 5340922            | 316248 | ERS3557085       |
| SB5625  | Klebsiella pneumoniae | Kp1         | 2643 | Cambodia | Urban | 2016           |                 |                 | 79                  | 44          | 5107551            | 292488 | ERS3557086       |
| SB5627  | Klebsiella pneumoniae | Kp1         | 2683 | Cambodia | Urban | 2016           |                 |                 | 78                  | 78          | 5482965            | 243024 | ERS3557087       |
| SB5629  | Klebsiella pneumoniae | Kp1         | 111  | Cambodia | Rural | 2016           |                 |                 | 49                  | 163         | 5402739            | 129791 | ERS3557088       |
| SB5632  | Klebsiella pneumoniae | Kp1         | 15   | Cambodia | Rural | 2016           |                 |                 | 65                  | 99          | 5407397            | 195332 | ERS3557091       |
| SB5634  | Klebsiella pneumoniae | Kp1         | 1035 | Cambodia | Urban | 2016           |                 |                 | 65                  | 120         | 5151356            | 116192 | ERS3556883       |
| SB5635  | Klebsiella pneumoniae | Kp1         | 133  | Cambodia | Rural | 2016           |                 |                 | 91                  | 63          | 5371762            | 217541 | ERS3557093       |
| SB5637  | Klebsiella pneumoniae | Kp1         | 2648 | Cambodia | Urban | 2016           |                 |                 | 79                  | 92          | 5529200            | 199112 | ERS3557095       |
| SB5639  | Klebsiella pneumoniae | Kp1         | 883  | Cambodia | Urban | 2016           |                 |                 | 85                  | 92          | 5446274            | 206771 | ERS3557096       |
| SB5640  | Klebsiella pneumoniae | Kp1         | 661  | Cambodia | Urban | 2016           |                 |                 | 80                  | 70          | 5402508            | 280597 | ERS3557097       |
| SB5643  | Klebsiella pneumoniae | Kp1         | 37   | Cambodia | Urban | 2016           |                 |                 | 83                  | 70          | 5399978            | 204189 | ERS3556780       |
| SB5646  | Klebsiella pneumoniae | Kp1         | 1574 | Cambodia | Urban | 2016           |                 |                 | 69                  | 55          | 5269781            | 250108 | ERS3557102       |
| SB5647  | Klebsiella pneumoniae | Kp1         | 2651 | Cambodia | Urban | 2016           |                 |                 | 76                  | 64          | 5385958            | 312099 | ERS3556894       |
| SB5649  | Klebsiella pneumoniae | Kp1         | 2652 | Cambodia | Rural | 2016           |                 |                 | 78                  | 54          | 5244578            | 224777 | ERS3557103       |
| SB5651  | Klebsiella pneumoniae | Kp1         | 2654 | Cambodia | Urban | 2016           |                 |                 | 79                  | 66          | 5271961            | 293441 | ERS3556914       |
| SB5652  | Klebsiella pneumoniae | Kp1         | 307  | Cambodia | Urban | 2016           |                 |                 | 71                  | 93          | 5436326            | 171620 | ERS3556931       |
| SB5653  | Klebsiella pneumoniae | Kp1         | 307  | Cambodia | Urban | 2016           |                 |                 | 78                  | 87          | 5420162            | 214814 | ERS3556932       |

|        |                       |     |      |          |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|----------|-------|------|----|-----|---------|--------|------------|
| SB5654 | Klebsiella pneumoniae | Kp1 | 2655 | Cambodia | Rural | 2016 | 84 | 107 | 5361896 | 248906 | ERS3556822 |
| SB5655 | Klebsiella pneumoniae | Kp1 | 1145 | Cambodia | Rural | 2016 | 78 | 90  | 5346312 | 240496 | ERS3557105 |
| SB5657 | Klebsiella pneumoniae | Kp1 | 17   | Cambodia | Urban | 2016 | 82 | 47  | 5062967 | 271406 | ERS3557106 |
| SB5658 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Rural | 2016 | 68 | 82  | 5543682 | 207586 | ERS3557107 |
| SB5659 | Klebsiella pneumoniae | Kp1 | 76   | Cambodia | Rural | 2016 | 70 | 86  | 5504433 | 240821 | ERS3557108 |
| SB5660 | Klebsiella pneumoniae | Kp1 | 17   | Cambodia | Rural | 2016 | 90 | 32  | 5070557 | 297443 | ERS3557109 |
| SB5662 | Klebsiella pneumoniae | Kp1 | 394  | Cambodia | Urban | 2016 | 90 | 52  | 5407929 | 219809 | ERS3556919 |
| SB5663 | Klebsiella pneumoniae | Kp1 | 17   | Cambodia | Urban | 2016 | 91 | 91  | 5543635 | 264515 | ERS3556903 |
| SB5664 | Klebsiella pneumoniae | Kp1 | 105  | Cambodia | Urban | 2016 | 90 | 39  | 5163741 | 295616 | ERS3556781 |
| SB5666 | Klebsiella pneumoniae | Kp1 | 200  | Cambodia | Rural | 2016 | 91 | 48  | 5214665 | 289243 | ERS3557112 |
| SB5668 | Klebsiella pneumoniae | Kp1 | 2706 | Cambodia | Rural | 2016 | 92 | 44  | 5160021 | 292030 | ERS3557113 |
| SB5672 | Klebsiella pneumoniae | Kp1 | 2709 | Cambodia | Rural | 2016 | 89 | 114 | 5155751 | 148981 | ERS3556943 |
| SB5673 | Klebsiella pneumoniae | Kp1 | 2710 | Cambodia | Rural | 2016 | 91 | 70  | 5460208 | 317648 | ERS3557117 |
| SB5675 | Klebsiella pneumoniae | Kp1 | 1782 | Cambodia | Rural | 2016 | 89 | 74  | 5376631 | 193102 | ERS3556915 |
| SB5676 | Klebsiella pneumoniae | Kp1 | 2711 | Cambodia | Rural | 2016 | 91 | 84  | 5125733 | 160797 | ERS3556868 |
| SB5680 | Klebsiella pneumoniae | Kp1 | 2713 | Cambodia | Rural | 2016 | 69 | 98  | 5359302 | 172998 | ERS3556917 |
| SB5681 | Klebsiella pneumoniae | Kp1 | 1662 | Cambodia | Rural | 2016 | 74 | 62  | 5219300 | 233701 | ERS3557121 |
| SB5682 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Rural | 2016 | 88 | 56  | 5381492 | 245190 | ERS3557122 |
| SB5683 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Rural | 2016 | 90 | 59  | 5415330 | 240098 | ERS3557123 |
| SB5684 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Rural | 2016 | 80 | 64  | 5549627 | 204833 | ERS3557124 |

|        |                       |     |      |          |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|----------|-------|------|----|-----|---------|--------|------------|
| SB5686 | Klebsiella pneumoniae | Kp1 | 2690 | Cambodia | Urban | 2016 | 96 | 64  | 5150093 | 252822 | ERS3556846 |
| SB5687 | Klebsiella pneumoniae | Kp1 | 2691 | Cambodia | Rural | 2016 | 96 | 50  | 5174574 | 268291 | ERS3557126 |
| SB5688 | Klebsiella pneumoniae | Kp1 | 1662 | Cambodia | Rural | 2016 | 96 | 48  | 5226977 | 314174 | ERS3557127 |
| SB5690 | Klebsiella pneumoniae | Kp1 | ND   | Cambodia | Rural | 2016 | 94 | 54  | 5282590 | 306163 | ERS3557128 |
| SB5692 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Urban | 2016 | 92 | 54  | 5555920 | 283787 | ERS3557130 |
| SB5693 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Urban | 2016 | 91 | 55  | 5557640 | 285763 | ERS3557131 |
| SB5694 | Klebsiella pneumoniae | Kp1 | 2692 | Cambodia | Rural | 2016 | 92 | 97  | 5437713 | 148731 | ERS3556803 |
| SB5695 | Klebsiella pneumoniae | Kp1 | 34   | Cambodia | Urban | 2016 | 94 | 81  | 5183897 | 181968 | ERS3556900 |
| SB5696 | Klebsiella pneumoniae | Kp1 | 2693 | Cambodia | Urban | 2016 | 94 | 77  | 5321489 | 205650 | ERS3557132 |
| SB5698 | Klebsiella pneumoniae | Kp1 | 2504 | Cambodia | Urban | 2016 | 94 | 54  | 5314104 | 293448 | ERS3556895 |
| SB5699 | Klebsiella pneumoniae | Kp1 | 111  | Cambodia | Rural | 2016 | 93 | 48  | 5193407 | 339667 | ERS3557134 |
| SB5701 | Klebsiella pneumoniae | Kp1 | 2694 | Cambodia | Rural | 2016 | 95 | 50  | 5230149 | 298062 | ERS3556847 |
| SB5702 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Rural | 2016 | 95 | 73  | 5434817 | 258434 | ERS3556908 |
| SB5703 | Klebsiella pneumoniae | Kp1 | 35   | Cambodia | Urban | 2016 | 95 | 73  | 5442485 | 285425 | ERS3556909 |
| SB5705 | Klebsiella pneumoniae | Kp1 | 17   | Cambodia | Urban | 2016 | 95 | 52  | 5333660 | 310773 | ERS3556918 |
| SB5706 | Klebsiella pneumoniae | Kp1 | 2695 | Cambodia | Rural | 2016 | 95 | 74  | 5142988 | 258274 | ERS3557136 |
| SB5707 | Klebsiella pneumoniae | Kp1 | 307  | Cambodia | Urban | 2016 | 93 | 57  | 5316377 | 307481 | ERS3556933 |
| SB5708 | Klebsiella pneumoniae | Kp1 | 307  | Cambodia | Rural | 2016 | 92 | 63  | 5307325 | 286380 | ERS3556934 |
| SB5712 | Klebsiella pneumoniae | Kp1 | 111  | Cambodia | Rural | 2016 | 96 | 40  | 5207036 | 359321 | ERS3557140 |
| SB5713 | Klebsiella pneumoniae | Kp1 | 48   | Cambodia | Urban | 2016 | 94 | 112 | 5379213 | 179168 | ERS3556935 |

|        |                                                    |     |      |          |       |      |    |     |         |        |            |
|--------|----------------------------------------------------|-----|------|----------|-------|------|----|-----|---------|--------|------------|
| SB5714 | Klebsiella pneumoniae                              | Kp1 | 1315 | Cambodia | Urban | 2016 | 94 | 102 | 5348903 | 152725 | ERS3556920 |
| SB5715 | Klebsiella pneumoniae                              | Kp1 | 76   | Cambodia | Urban | 2016 | 93 | 57  | 5321869 | 313319 | ERS3556782 |
| SB5716 | Klebsiella pneumoniae                              | Kp1 | 469  | Cambodia | Urban | 2016 | 94 | 48  | 5138345 | 263428 | ERS3556901 |
| SB5720 | Klebsiella pneumoniae                              | Kp1 | 36   | Cambodia | Urban | 2016 | 93 | 47  | 5211442 | 318748 | ERS3557142 |
| SB5721 | Klebsiella pneumoniae                              | Kp1 | 774  | Cambodia | Rural | 2016 | 94 | 40  | 5104795 | 347891 | ERS3557143 |
| SB5722 | Klebsiella pneumoniae                              | Kp1 | 2701 | Cambodia | Urban | 2016 | 94 | 49  | 5195511 | 281539 | ERS3557144 |
| SB5723 | Klebsiella pneumoniae                              | Kp1 | 631  | Cambodia | Urban | 2016 | 94 | 63  | 5314776 | 281221 | ERS3556916 |
| SB5724 | Klebsiella pneumoniae                              | Kp1 | 2702 | Cambodia | Urban | 2016 | 90 | 80  | 5480956 | 225763 | ERS3556869 |
| SB5725 | Klebsiella pneumoniae                              | Kp1 | 2703 | Cambodia | Rural | 2016 | 93 | 65  | 5398757 | 299290 | ERS3557145 |
| SB5670 | Klebsiella variicola subsp. variicola              | Kp3 | 2645 | Cambodia | Rural | 2016 | 88 | 50  | 5454754 | 225824 | ERS3557115 |
| SB5678 | Klebsiella variicola subsp. variicola              | Kp3 | 2645 | Cambodia | Rural | 2016 | 89 | 41  | 5446243 | 264614 | ERS3557120 |
| SB5691 | Klebsiella variicola subsp. variicola              | Kp3 | 2645 | Cambodia | Rural | 2016 | 94 | 41  | 5489856 | 314983 | ERS3557129 |
| SB5697 | Klebsiella variicola subsp. variicola              | Kp3 | 2645 | Cambodia | Urban | 2016 | 95 | 36  | 5464940 | 384747 | ERS3557133 |
| SB5626 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2644 | Cambodia | Urban | 2016 | 65 | 74  | 5282775 | 264726 | ERS3556929 |
| SB5630 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1770 | Cambodia | Urban | 2016 | 72 | 193 | 5493483 | 148255 | ERS3557089 |
| SB5631 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2684 | Cambodia | Rural | 2016 | 79 | 38  | 5120402 | 368084 | ERS3557090 |
| SB5633 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2685 | Cambodia | Urban | 2016 | 82 | 36  | 5072270 | 383723 | ERS3557092 |
| SB5636 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2686 | Cambodia | Urban | 2016 | 84 | 30  | 5089222 | 384645 | ERS3557094 |
| SB5638 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1584 | Cambodia | Rural | 2016 | 93 | 109 | 5429245 | 175660 | ERS3556799 |
| SB5641 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2689 | Cambodia | Rural | 2016 | 84 | 28  | 5044032 | 499401 | ERS3557098 |

|        |                                                    |     |      |          |       |      |    |     |         |        |            |
|--------|----------------------------------------------------|-----|------|----------|-------|------|----|-----|---------|--------|------------|
| SB5642 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2689 | Cambodia | Rural | 2016 | 84 | 29  | 5043810 | 390857 | ERS3557099 |
| SB5644 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2649 | Cambodia | Rural | 2016 | 60 | 114 | 5365452 | 226173 | ERS3557100 |
| SB5645 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2650 | Cambodia | Rural | 2016 | 73 | 40  | 5092929 | 379605 | ERS3557101 |
| SB5648 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1308 | Cambodia | Rural | 2016 | 59 | 126 | 5448005 | 180080 | ERS3556867 |
| SB5650 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2653 | Cambodia | Rural | 2016 | 73 | 35  | 5057265 | 379105 | ERS3557104 |
| SB5656 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2489 | Cambodia | Rural | 2016 | 81 | 121 | 5136202 | 144535 | ERS3556942 |
| SB5661 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2704 | Cambodia | Rural | 2016 | 91 | 41  | 5132256 | 270896 | ERS3557110 |
| SB5665 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2705 | Cambodia | Urban | 2016 | 89 | 34  | 5137797 | 371558 | ERS3557111 |
| SB5669 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2707 | Cambodia | Rural | 2016 | 89 | 47  | 5209783 | 394292 | ERS3557114 |
| SB5671 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2708 | Cambodia | Rural | 2016 | 89 | 32  | 5195937 | 342629 | ERS3557116 |
| SB5674 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2653 | Cambodia | Rural | 2016 | 90 | 39  | 5199048 | 370553 | ERS3557118 |
| SB5677 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2687 | Cambodia | Rural | 2016 | 61 | 89  | 5500390 | 248746 | ERS3557119 |
| SB5679 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2688 | Cambodia | Rural | 2016 | 87 | 106 | 5219193 | 237135 | ERS3556944 |
| SB5685 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2696 | Cambodia | Urban | 2016 | 96 | 32  | 5267118 | 580124 | ERS3557125 |
| SB5700 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1584 | Cambodia | Rural | 2016 | 92 | 66  | 5196530 | 242909 | ERS3557135 |
| SB5709 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2697 | Cambodia | Rural | 2016 | 97 | 30  | 5001814 | 371361 | ERS3557137 |
| SB5710 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2698 | Cambodia | Rural | 2016 | 95 | 45  | 5078634 | 356527 | ERS3557138 |
| SB5711 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2699 | Cambodia | Rural | 2016 | 95 | 31  | 5179150 | 382603 | ERS3557139 |
| SB5718 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1697 | Cambodia | Rural | 2016 | 95 | 86  | 5167614 | 172755 | ERS3556848 |
| SB5719 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2700 | Cambodia | Rural | 2016 | 95 | 29  | 5085967 | 378994 | ERS3557141 |

|        |                       |     |      |            |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5372 | Klebsiella pneumoniae | Kp1 | 8    | Madagascar | Urban | 2015 | 90 | 76  | 5178652 | 206796 | ERS3556948 |
| SB5376 | Klebsiella pneumoniae | Kp1 | 1621 | Madagascar | Urban | 2015 | 90 | 55  | 5330747 | 338002 | ERS3556951 |
| SB5378 | Klebsiella pneumoniae | Kp1 | 1621 | Madagascar | Urban | 2015 | 78 | 64  | 5323070 | 288601 | ERS3556953 |
| SB5380 | Klebsiella pneumoniae | Kp1 | 70   | Madagascar | Urban | 2015 | 86 | 57  | 5476561 | 305068 | ERS3556954 |
| SB5382 | Klebsiella pneumoniae | Kp1 | 1621 | Madagascar | Urban | 2015 | 83 | 59  | 5319369 | 288601 | ERS3556956 |
| SB5383 | Klebsiella pneumoniae | Kp1 | 491  | Madagascar | Urban | 2015 | 90 | 115 | 5243061 | 141273 | ERS3556957 |
| SB5384 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Urban | 2015 | 87 | 87  | 5436056 | 148274 | ERS3556958 |
| SB5385 | Klebsiella pneumoniae | Kp1 | 1873 | Madagascar | Urban | 2015 | 87 | 55  | 5222041 | 208934 | ERS3556959 |
| SB5389 | Klebsiella pneumoniae | Kp1 | 873  | Madagascar | Urban | 2015 | 89 | 65  | 5416819 | 257302 | ERS3556961 |
| SB5390 | Klebsiella pneumoniae | Kp1 | 111  | Madagascar | Urban | 2015 | 50 | 70  | 5429342 | 326979 | ERS3556886 |
| SB5391 | Klebsiella pneumoniae | Kp1 | 1    | Madagascar | Urban | 2015 | 92 | 54  | 5334032 | 262838 | ERS3556887 |
| SB5392 | Klebsiella pneumoniae | Kp1 | 2467 | Madagascar | Urban | 2015 | 92 | 61  | 5191536 | 329548 | ERS3556962 |
| SB5393 | Klebsiella pneumoniae | Kp1 | 873  | Madagascar | Urban | 2015 | 91 | 47  | 5444209 | 355310 | ERS3556963 |
| SB5395 | Klebsiella pneumoniae | Kp1 | 2177 | Madagascar | Urban | 2015 | 91 | 36  | 5020730 | 367844 | ERS3556964 |
| SB5397 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Urban | 2015 | 90 | 113 | 5368646 | 160219 | ERS3556793 |
| SB5398 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Urban | 2015 | 82 | 66  | 5420415 | 271077 | ERS3556830 |
| SB5399 | Klebsiella pneumoniae | Kp1 | 629  | Madagascar | Urban | ND   | 80 | 94  | 5371811 | 205169 | ERS3556853 |
| SB5401 | Klebsiella pneumoniae | Kp1 | 2467 | Madagascar | Urban | 2015 | 86 | 58  | 5180397 | 295639 | ERS3556967 |
| SB5403 | Klebsiella pneumoniae | Kp1 | 2471 | Madagascar | Urban | 2016 | 66 | 191 | 5367052 | 89127  | ERS3556815 |
| SB5404 | Klebsiella pneumoniae | Kp1 | 45   | Madagascar | Urban | 2016 | 64 | 79  | 5396683 | 203125 | ERS3556969 |

|        |                       |     |      |            |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5406 | Klebsiella pneumoniae | Kp1 | 2473 | Madagascar | Urban | 2016 | 65 | 174 | 5165061 | 70752  | ERS3556970 |
| SB5407 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Urban | 2016 | 75 | 98  | 5449031 | 136675 | ERS3556971 |
| SB5408 | Klebsiella pneumoniae | Kp1 | 2474 | Madagascar | Urban | 2016 | 40 | 157 | 5460594 | 73383  | ERS3556972 |
| SB5411 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Urban | 2016 | 89 | 114 | 5333761 | 142933 | ERS3556832 |
| SB5412 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Urban | 2016 | 84 | 142 | 5345078 | 92641  | ERS3556854 |
| SB5415 | Klebsiella pneumoniae | Kp1 | 441  | Madagascar | Urban | 2016 | 69 | 102 | 5457548 | 208158 | ERS3556977 |
| SB5417 | Klebsiella pneumoniae | Kp1 | 2480 | Madagascar | Urban | 2016 | 81 | 77  | 5418208 | 241277 | ERS3556833 |
| SB5418 | Klebsiella pneumoniae | Kp1 | 2481 | Madagascar | Urban | 2016 | 72 | 97  | 5241241 | 172192 | ERS3556834 |
| SB5419 | Klebsiella pneumoniae | Kp1 | 3331 | Madagascar | Urban | 2016 | 81 | 98  | 5072035 | 128862 | ERS3556876 |
| SB5421 | Klebsiella pneumoniae | Kp1 | 17   | Madagascar | Urban | 2016 | 85 | 61  | 5424932 | 293693 | ERS3556910 |
| SB5422 | Klebsiella pneumoniae | Kp1 | 964  | Madagascar | Urban | 2016 | 85 | 83  | 5420021 | 183254 | ERS3556816 |
| SB5423 | Klebsiella pneumoniae | Kp1 | 3100 | Madagascar | Urban | 2016 | 91 | 130 | 5287089 | 89139  | ERS3556855 |
| SB5424 | Klebsiella pneumoniae | Kp1 | 2482 | Madagascar | Urban | 2016 | 92 | 133 | 5084066 | 89186  | ERS3556980 |
| SB5425 | Klebsiella pneumoniae | Kp1 | 2482 | Madagascar | Urban | 2016 | 88 | 122 | 5079152 | 92130  | ERS3556981 |
| SB5426 | Klebsiella pneumoniae | Kp1 | 25   | Madagascar | Urban | 2016 | 89 | 84  | 5541863 | 192330 | ERS3556982 |
| SB5427 | Klebsiella pneumoniae | Kp1 | 1787 | Madagascar | Urban | 2016 | 73 | 71  | 5404088 | 324273 | ERS3556923 |
| SB5429 | Klebsiella pneumoniae | Kp1 | 48   | Madagascar | Urban | 2016 | 60 | 76  | 5512363 | 175690 | ERS3556930 |
| SB5432 | Klebsiella pneumoniae | Kp1 | 1661 | Madagascar | Urban | 2016 | 72 | 45  | 5087071 | 265120 | ERS3556986 |
| SB5434 | Klebsiella pneumoniae | Kp1 | 2484 | Madagascar | Urban | 2016 | 73 | 166 | 5230428 | 78973  | ERS3556817 |
| SB5435 | Klebsiella pneumoniae | Kp1 | 872  | Madagascar | Urban | ND   | 74 | 126 | 5250432 | 123643 | ERS3556776 |

|        |                       |     |      |            |       |      |          |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----------|----|-----|---------|--------|------------|
| SB5436 | Klebsiella pneumoniae | Kp1 | 2485 | Madagascar | Urban | 2016 |          | 92 | 71  | 5369420 | 203248 | ERS3556987 |
| SB5438 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Urban | 2016 | strain B | 93 | 40  | 5230669 | 284494 | ERS3556989 |
| SB5440 | Klebsiella pneumoniae | Kp1 | 25   | Madagascar | Urban | 2016 |          | 92 | 64  | 5549052 | 211951 | ERS3556990 |
| SB5441 | Klebsiella pneumoniae | Kp1 | 1537 | Madagascar | Urban | 2016 |          | 93 | 41  | 5152938 | 310877 | ERS3556991 |
| SB5442 | Klebsiella pneumoniae | Kp1 | 101  | Madagascar | Urban | 2016 |          | 91 | 95  | 5406482 | 250934 | ERS3556836 |
| SB5443 | Klebsiella pneumoniae | Kp1 | 2487 | Madagascar | Urban | 2016 |          | 93 | 94  | 5267782 | 141582 | ERS3556777 |
| SB5446 | Klebsiella pneumoniae | Kp1 | 770  | Madagascar | Urban | 2016 |          | 85 | 41  | 5056940 | 282476 | ERS3556994 |
| SB5447 | Klebsiella pneumoniae | Kp1 | 2938 | Madagascar | Rural | 2015 |          | 84 | 101 | 5314141 | 176731 | ERS3556995 |
| SB5450 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Rural | 2015 |          | 93 | 88  | 5430447 | 148274 | ERS3556997 |
| SB5451 | Klebsiella pneumoniae | Kp1 | 36   | Madagascar | Rural | 2015 |          | 83 | 82  | 5005670 | 133698 | ERS3556998 |
| SB5452 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Rural | 2015 |          | 92 | 83  | 5445429 | 150520 | ERS3556999 |
| SB5453 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Rural | 2015 |          | 71 | 115 | 5444588 | 123903 | ERS3557000 |
| SB5456 | Klebsiella pneumoniae | Kp1 | 36   | Madagascar | Rural | ND   |          | 76 | 153 | 5479592 | 167910 | ERS3556837 |
| SB5460 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Rural | 2015 |          | 66 | 109 | 5518271 | 160927 | ERS3557003 |
| SB5461 | Klebsiella pneumoniae | Kp1 | 20   | Madagascar | Rural | 2015 |          | 57 | 74  | 5405677 | 225887 | ERS3557004 |
| SB5462 | Klebsiella pneumoniae | Kp1 | 45   | Madagascar | Rural | 2015 |          | 89 | 34  | 5161179 | 357889 | ERS3557005 |
| SB5463 | Klebsiella pneumoniae | Kp1 | 491  | Madagascar | Rural | 2015 |          | 93 | 106 | 5239597 | 141269 | ERS3557006 |
| SB5464 | Klebsiella pneumoniae | Kp1 | 1303 | Madagascar | Rural | 2015 |          | 90 | 85  | 5018683 | 149691 | ERS3556936 |
| SB5466 | Klebsiella pneumoniae | Kp1 | 2632 | Madagascar | Rural | 2015 |          | 68 | 116 | 5217543 | 173144 | ERS3557007 |
| SB5467 | Klebsiella pneumoniae | Kp1 | 2715 | Madagascar | Rural | 2015 |          | 88 | 56  | 5162213 | 203033 | ERS3556800 |

|        |                       |     |      |            |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5468 | Klebsiella pneumoniae | Kp1 | 2177 | Madagascar | Rural | 2015 | 83 | 53  | 5015760 | 261489 | ERS3557008 |
| SB5470 | Klebsiella pneumoniae | Kp1 | 1978 | Madagascar | Rural | 2015 | 91 | 55  | 5264571 | 269053 | ERS3556888 |
| SB5471 | Klebsiella pneumoniae | Kp1 | 36   | Madagascar | Rural | 2015 | 91 | 47  | 5228028 | 318470 | ERS3557009 |
| SB5472 | Klebsiella pneumoniae | Kp1 | 1198 | Madagascar | Rural | 2015 | 62 | 81  | 5347913 | 264893 | ERS3556838 |
| SB5473 | Klebsiella pneumoniae | Kp1 | 2473 | Madagascar | Rural | 2015 | 91 | 149 | 5136430 | 83330  | ERS3557010 |
| SB5478 | Klebsiella pneumoniae | Kp1 | 2504 | Madagascar | Rural | 2015 | 76 | 66  | 5247596 | 260223 | ERS3557011 |
| SB5479 | Klebsiella pneumoniae | Kp1 | 22   | Madagascar | Rural | 2015 | 68 | 120 | 5405462 | 169554 | ERS3556839 |
| SB5480 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Rural | 2015 | 70 | 94  | 5304577 | 173154 | ERS3556937 |
| SB5481 | Klebsiella pneumoniae | Kp1 | 2505 | Madagascar | Rural | 2015 | 64 | 102 | 5479015 | 166616 | ERS3556788 |
| SB5482 | Klebsiella pneumoniae | Kp1 | 2506 | Madagascar | Rural | 2015 | 73 | 76  | 5260234 | 227988 | ERS3556889 |
| SB5483 | Klebsiella pneumoniae | Kp1 | 399  | Madagascar | Rural | 2015 | 73 | 99  | 5496447 | 174799 | ERS3556840 |
| SB5486 | Klebsiella pneumoniae | Kp1 | 1966 | Madagascar | Rural | 2016 | 90 | 53  | 5215468 | 277324 | ERS3557013 |
| SB5489 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Rural | 2016 | 91 | 99  | 5559122 | 136788 | ERS3557016 |
| SB5491 | Klebsiella pneumoniae | Kp1 | 491  | Madagascar | Rural | 2016 | 79 | 163 | 5275772 | 88615  | ERS3557017 |
| SB5492 | Klebsiella pneumoniae | Kp1 | 491  | Madagascar | Rural | 2016 | 82 | 158 | 5258941 | 90365  | ERS3557018 |
| SB5494 | Klebsiella pneumoniae | Kp1 | 831  | Madagascar | Rural | 2016 | 82 | 76  | 5426956 | 204301 | ERS3556856 |
| SB5497 | Klebsiella pneumoniae | Kp1 | 432  | Madagascar | Rural | 2016 | 88 | 41  | 5038579 | 269791 | ERS3557022 |
| SB5498 | Klebsiella pneumoniae | Kp1 | 2552 | Madagascar | Rural | 2016 | 88 | 47  | 5204652 | 277765 | ERS3557023 |
| SB5499 | Klebsiella pneumoniae | Kp1 | 22   | Madagascar | Rural | 2016 | 87 | 86  | 5304984 | 226309 | ERS3556849 |
| SB5500 | Klebsiella pneumoniae | Kp1 | 873  | Madagascar | Rural | 2016 | 88 | 46  | 5302640 | 296215 | ERS3557024 |

|        |                       |     |      |            |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5501 | Klebsiella pneumoniae | Kp1 | 45   | Madagascar | Urban | 2016 | 91 | 48  | 5183010 | 207611 | ERS3557146 |
| SB5502 | Klebsiella pneumoniae | Kp1 | 8    | Madagascar | Rural | 2016 | 77 | 53  | 5175960 | 230593 | ERS3557025 |
| SB5506 | Klebsiella pneumoniae | Kp1 | 2554 | Madagascar | Rural | 2016 | 76 | 54  | 5259561 | 293683 | ERS3557027 |
| SB5507 | Klebsiella pneumoniae | Kp1 | 2555 | Madagascar | Rural | ND   | 78 | 65  | 5212584 | 169101 | ERS3557028 |
| SB5508 | Klebsiella pneumoniae | Kp1 | 2938 | Madagascar | Rural | 2016 | 70 | 92  | 5323442 | 183578 | ERS3557029 |
| SB5509 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Rural | 2016 | 83 | 125 | 5384152 | 123057 | ERS3556938 |
| SB5510 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Rural | 2016 | 84 | 82  | 5429075 | 163610 | ERS3557030 |
| SB5511 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Rural | 2016 | 86 | 97  | 5444987 | 148885 | ERS3557031 |
| SB5512 | Klebsiella pneumoniae | Kp1 | 2408 | Madagascar | Rural | 2016 | 87 | 90  | 5300346 | 194594 | ERS3557032 |
| SB5513 | Klebsiella pneumoniae | Kp1 | 17   | Madagascar | Rural | 2016 | 90 | 79  | 5180162 | 264563 | ERS3556858 |
| SB5514 | Klebsiella pneumoniae | Kp1 | 1087 | Madagascar | Rural | 2016 | 89 | 87  | 5332857 | 129436 | ERS3557033 |
| SB5515 | Klebsiella pneumoniae | Kp1 | 1296 | Madagascar | Rural | 2016 | 91 | 89  | 5407577 | 163767 | ERS3556852 |
| SB5516 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Rural | ND   | 90 | 119 | 5536122 | 107851 | ERS3556841 |
| SB5518 | Klebsiella pneumoniae | Kp1 | 500  | Madagascar | Rural | 2016 | 89 | 60  | 5320291 | 259971 | ERS3556802 |
| SB5521 | Klebsiella pneumoniae | Kp1 | 307  | Madagascar | Rural | 2016 | 91 | 52  | 5303204 | 214814 | ERS3556842 |
| SB5522 | Klebsiella pneumoniae | Kp1 | 2556 | Madagascar | Rural | 2016 | 84 | 141 | 5196671 | 99637  | ERS3557035 |
| SB5523 | Klebsiella pneumoniae | Kp1 | 2557 | Madagascar | Rural | 2016 | 84 | 80  | 5272792 | 186543 | ERS3557036 |
| SB5524 | Klebsiella pneumoniae | Kp1 | 491  | Madagascar | Rural | 2016 | 80 | 129 | 5257922 | 127044 | ERS3557037 |
| SB5525 | Klebsiella pneumoniae | Kp1 | 65   | Madagascar | Rural | 2016 | 87 | 58  | 5284478 | 180289 | ERS3556939 |
| SB5526 | Klebsiella pneumoniae | Kp1 | 43   | Madagascar | Rural | 2016 | 90 | 88  | 5378527 | 219565 | ERS3557038 |

|        |                       |     |      |            |       |      |          |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----------|----|-----|---------|--------|------------|
| SB5529 | Klebsiella pneumoniae | Kp1 | 1263 | Madagascar | Rural | 2016 |          | 80 | 60  | 5371814 | 263011 | ERS3556878 |
| SB5530 | Klebsiella pneumoniae | Kp1 | 2560 | Madagascar | Rural | 2016 |          | 91 | 568 | 5444783 | 28573  | ERS3556899 |
| SB5532 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Rural | 2016 | strain B | 90 | 38  | 5234949 | 329374 | ERS3557039 |
| SB5534 | Klebsiella pneumoniae | Kp1 | 36   | Madagascar | Rural | 2016 |          | 90 | 66  | 4986713 | 170149 | ERS3557040 |
| SB5535 | Klebsiella pneumoniae | Kp1 | 215  | Madagascar | Rural | 2016 |          | 82 | 60  | 5250714 | 197167 | ERS3556794 |
| SB5538 | Klebsiella pneumoniae | Kp1 | 441  | Madagascar | Rural | 2016 |          | 83 | 98  | 5440818 | 204880 | ERS3557042 |
| SB5540 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Rural | 2016 |          | 70 | 128 | 5370903 | 172812 | ERS3556795 |
| SB5546 | Klebsiella pneumoniae | Kp1 | 29   | Madagascar | Rural | 2016 |          | 70 | 64  | 5272828 | 303990 | ERS3556924 |
| SB5547 | Klebsiella pneumoniae | Kp1 | 12   | Madagascar | Rural | 2016 |          | 72 | 56  | 5189532 | 207116 | ERS3557046 |
| SB5548 | Klebsiella pneumoniae | Kp1 | 1213 | Madagascar | Rural | 2016 |          | 84 | 71  | 5142388 | 197728 | ERS3557047 |
| SB5549 | Klebsiella pneumoniae | Kp1 | 2570 | Madagascar | Rural | 2016 |          | 54 | 125 | 5153262 | 146750 | ERS3557048 |
| SB5550 | Klebsiella pneumoniae | Kp1 | 14   | Madagascar | Rural | 2016 |          | 62 | 94  | 5454149 | 195101 | ERS3557049 |
| SB5551 | Klebsiella pneumoniae | Kp1 | 36   | Madagascar | Rural | 2016 |          | 78 | 74  | 4967818 | 159973 | ERS3557050 |
| SB5555 | Klebsiella pneumoniae | Kp1 | 872  | Madagascar | Rural | 2016 |          | 90 | 93  | 5307910 | 121429 | ERS3556818 |
| SB5557 | Klebsiella pneumoniae | Kp1 | 76   | Madagascar | Rural | 2016 |          | 92 | 58  | 5318119 | 231978 | ERS3557053 |
| SB5558 | Klebsiella pneumoniae | Kp1 | 17   | Madagascar | Rural | 2016 |          | 88 | 70  | 5433470 | 369399 | ERS3556911 |
| SB5559 | Klebsiella pneumoniae | Kp1 | 1787 | Madagascar | Rural | 2016 |          | 90 | 78  | 5406219 | 287551 | ERS3556925 |
| SB5560 | Klebsiella pneumoniae | Kp1 | 101  | Madagascar | Rural | 2016 |          | 63 | 87  | 5391103 | 174864 | ERS3556890 |
| SB5562 | Klebsiella pneumoniae | Kp1 | 1787 | Madagascar | Rural | 2016 |          | 66 | 104 | 5389669 | 197174 | ERS3556926 |
| SB5563 | Klebsiella pneumoniae | Kp1 | 1787 | Madagascar | Rural | 2016 |          | 59 | 105 | 5376845 | 204126 | ERS3556927 |

|        |                       |     |      |            |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5564 | Klebsiella pneumoniae | Kp1 | 1447 | Madagascar | Rural | 2016 | 61 | 109 | 5344587 | 165070 | ERS3556819 |
| SB5565 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Rural | 2016 | 65 | 150 | 5262563 | 75592  | ERS3556820 |
| SB5568 | Klebsiella pneumoniae | Kp1 | 186  | Madagascar | Rural | 2016 | 53 | 76  | 5291083 | 172920 | ERS3557056 |
| SB5569 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Rural | 2016 | 60 | 136 | 5376455 | 129567 | ERS3556796 |
| SB5571 | Klebsiella pneumoniae | Kp1 | 60   | Madagascar | Rural | 2016 | 76 | 58  | 5141680 | 215109 | ERS3557057 |
| SB5572 | Klebsiella pneumoniae | Kp1 | 234  | Madagascar | Rural | 2016 | 67 | 76  | 5245852 | 234973 | ERS3556891 |
| SB5573 | Klebsiella pneumoniae | Kp1 | 2473 | Madagascar | Rural | 2016 | 74 | 157 | 5153367 | 89574  | ERS3557058 |
| SB5574 | Klebsiella pneumoniae | Kp1 | 1429 | Madagascar | Rural | 2016 | 74 | 93  | 5432728 | 150523 | ERS3557059 |
| SB5575 | Klebsiella pneumoniae | Kp1 | 2718 | Madagascar | Rural | 2016 | 78 | 89  | 5028703 | 190544 | ERS3557060 |
| SB5576 | Klebsiella pneumoniae | Kp1 | 2053 | Madagascar | Rural | 2016 | 88 | 140 | 5275264 | 112499 | ERS3556941 |
| SB5578 | Klebsiella pneumoniae | Kp1 | 375  | Madagascar | Rural | 2016 | 77 | 83  | 5302597 | 141350 | ERS3557061 |
| SB5579 | Klebsiella pneumoniae | Kp1 | 375  | Madagascar | Rural | 2016 | 52 | 101 | 5323233 | 112422 | ERS3557062 |
| SB5580 | Klebsiella pneumoniae | Kp1 | 375  | Madagascar | Rural | 2016 | 89 | 84  | 5427086 | 185754 | ERS3557063 |
| SB5581 | Klebsiella pneumoniae | Kp1 | 48   | Madagascar | Rural | 2016 | 86 | 63  | 5246421 | 218973 | ERS3556907 |
| SB5583 | Klebsiella pneumoniae | Kp1 | 22   | Madagascar | Rural | 2016 | 89 | 71  | 5470530 | 264543 | ERS3557064 |
| SB5584 | Klebsiella pneumoniae | Kp1 | 2484 | Madagascar | Rural | 2016 | 86 | 129 | 5144991 | 83161  | ERS3557065 |
| SB5585 | Klebsiella pneumoniae | Kp1 | 39   | Madagascar | Rural | 2016 | 91 | 69  | 5184265 | 222065 | ERS3556892 |
| SB5588 | Klebsiella pneumoniae | Kp1 | 2596 | Madagascar | Rural | 2016 | 71 | 106 | 5356391 | 193137 | ERS3557067 |
| SB5589 | Klebsiella pneumoniae | Kp1 | 904  | Madagascar | Rural | 2016 | 64 | 94  | 5317193 | 176576 | ERS3556862 |
| SB5590 | Klebsiella pneumoniae | Kp1 | 2612 | Madagascar | Rural | 2016 | 88 | 125 | 5293431 | 173362 | ERS3556779 |

|        |                       |     |      |            |       |      |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5592 | Klebsiella pneumoniae | Kp1 | 904  | Madagascar | Rural | ND   | 81 | 58  | 5315102 | 355285 | ERS3556864 |
| SB5593 | Klebsiella pneumoniae | Kp1 | 2505 | Madagascar | Rural | 2016 | 86 | 78  | 5462273 | 205884 | ERS3556789 |
| SB5594 | Klebsiella pneumoniae | Kp1 | 449  | Madagascar | Rural | 2016 | 91 | 55  | 5501182 | 240090 | ERS3556912 |
| SB5595 | Klebsiella pneumoniae | Kp1 | 449  | Madagascar | Rural | 2016 | 87 | 50  | 5495265 | 274270 | ERS3556913 |
| SB5596 | Klebsiella pneumoniae | Kp1 | 394  | Madagascar | Rural | 2016 | 84 | 64  | 5411353 | 271878 | ERS3556865 |
| SB5597 | Klebsiella pneumoniae | Kp1 | 39   | Madagascar | Rural | 2016 | 90 | 99  | 5365389 | 195528 | ERS3556928 |
| SB5599 | Klebsiella pneumoniae | Kp1 | 831  | Madagascar | Rural | 2016 | 76 | 64  | 5451335 | 348667 | ERS3556893 |
| SB5601 | Klebsiella pneumoniae | Kp1 | 394  | Madagascar | Rural | 2016 | 89 | 57  | 5454291 | 288135 | ERS3556866 |
| SB5602 | Klebsiella pneumoniae | Kp1 | 2595 | Madagascar | Rural | 2016 | 91 | 136 | 5311261 | 91065  | ERS3557070 |
| SB5603 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Rural | 2016 | 93 | 100 | 5386562 | 138910 | ERS3557071 |
| SB5604 | Klebsiella pneumoniae | Kp1 | 2596 | Madagascar | Rural | 2016 | 93 | 74  | 5343235 | 316377 | ERS3557072 |
| SB5605 | Klebsiella pneumoniae | Kp1 | 2597 | Madagascar | Rural | 2016 | 68 | 111 | 5199001 | 154482 | ERS3557073 |
| SB5606 | Klebsiella pneumoniae | Kp1 | 2598 | Madagascar | Rural | 2016 | 88 | 80  | 5270663 | 266403 | ERS3557074 |
| SB5608 | Klebsiella pneumoniae | Kp1 | 2599 | Madagascar | Rural | 2016 | 92 | 100 | 5442784 | 175920 | ERS3556880 |
| SB5609 | Klebsiella pneumoniae | Kp1 | 60   | Madagascar | Rural | 2016 | 90 | 52  | 5332478 | 240663 | ERS3557076 |
| SB5611 | Klebsiella pneumoniae | Kp1 | 2599 | Madagascar | Rural | 2016 | 89 | 97  | 5415046 | 242755 | ERS3556881 |
| SB5612 | Klebsiella pneumoniae | Kp1 | 2599 | Madagascar | Rural | 2016 | 90 | 100 | 5427357 | 176452 | ERS3556882 |
| SB5613 | Klebsiella pneumoniae | Kp1 | 380  | Madagascar | Rural | 2016 | 91 | 57  | 5424628 | 284246 | ERS3557077 |
| SB5614 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Rural | 2016 | 88 | 33  | 5236602 | 349311 | ERS3557078 |
| SB5616 | Klebsiella pneumoniae | Kp1 | 2459 | Madagascar | Rural | 2016 | 91 | 51  | 5218771 | 254988 | ERS3557080 |

|        |                       |     |      |            |       |      |          |    |     |         |        |            |
|--------|-----------------------|-----|------|------------|-------|------|----------|----|-----|---------|--------|------------|
| SB5618 | Klebsiella pneumoniae | Kp1 | 39   | Madagascar | Rural | 2016 |          | 73 | 77  | 5291111 | 225946 | ERS3556797 |
| SB5619 | Klebsiella pneumoniae | Kp1 | 2601 | Madagascar | Rural | 2016 |          | 85 | 88  | 5203937 | 293846 | ERS3557082 |
| SB5620 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Rural | 2016 |          | 94 | 51  | 5095900 | 204195 | ERS3556790 |
| SB5622 | Klebsiella pneumoniae | Kp1 | 2602 | Madagascar | Rural | 2016 |          | 88 | 57  | 5488160 | 286870 | ERS3557083 |
| SB5623 | Klebsiella pneumoniae | Kp1 | 1609 | Madagascar | Rural | 2016 |          | 90 | 57  | 5341447 | 204176 | ERS3557084 |
| SB5952 | Klebsiella pneumoniae | Kp1 | 2442 | Madagascar | Urban | 2016 |          | 92 | 47  | 5221079 | 263464 | ERS3557149 |
| SB5953 | Klebsiella pneumoniae | Kp1 | 337  | Madagascar | Urban | 2016 |          | 91 | 115 | 5316117 | 121259 | ERS3557150 |
| SB5954 | Klebsiella pneumoniae | Kp1 | 441  | Madagascar | Urban | 2016 |          | 90 | 67  | 5309293 | 227038 | ERS3557151 |
| SB5955 | Klebsiella pneumoniae | Kp1 | 1689 | Madagascar | Urban | 2016 |          | 83 | 60  | 5213238 | 263118 | ERS3556823 |
| SB5956 | Klebsiella pneumoniae | Kp1 | 76   | Madagascar | Urban | 2016 |          | 91 | 56  | 5248702 | 323844 | ERS3557152 |
| SB5958 | Klebsiella pneumoniae | Kp1 | 3102 | Madagascar | Urban | 2016 |          | 91 | 88  | 5199514 | 125901 | ERS3556806 |
| SB5959 | Klebsiella pneumoniae | Kp1 | 3102 | Madagascar | Rural | 2016 |          | 92 | 89  | 5215526 | 132250 | ERS3556807 |
| SB5961 | Klebsiella pneumoniae | Kp1 | 48   | Madagascar | Rural | 2016 | strain A | 91 | 79  | 5535868 | 184773 | ERS3556921 |
| SB5962 | Klebsiella pneumoniae | Kp1 | 48   | Madagascar | Rural | 2016 | strain A | 90 | 84  | 5505160 | 218871 | ERS3556922 |
| SB5963 | Klebsiella pneumoniae | Kp1 | 380  | Madagascar | Rural | 2016 |          | 92 | 85  | 5579823 | 283581 | ERS3556824 |
| SB5964 | Klebsiella pneumoniae | Kp1 | 22   | Madagascar | Rural | 2016 |          | 92 | 94  | 5411793 | 176235 | ERS3556808 |
| SB5965 | Klebsiella pneumoniae | Kp1 | 37   | Madagascar | Rural | 2016 |          | 92 | 48  | 5196099 | 255524 | ERS3557155 |
| SB5969 | Klebsiella pneumoniae | Kp1 | 1741 | Madagascar | Rural | 2016 |          | 90 | 70  | 5256203 | 172898 | ERS3556825 |
| SB5970 | Klebsiella pneumoniae | Kp1 | 45   | Madagascar | Rural | 2016 |          | 90 | 52  | 5368046 | 245611 | ERS3556826 |
| SB5973 | Klebsiella pneumoniae | Kp1 | 2445 | Madagascar | Rural | 2016 |          | 90 | 65  | 5380953 | 202741 | ERS3556871 |

|        |                                                   |     |      |            |       |      |    |     |         |        |            |
|--------|---------------------------------------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5974 | Klebsiella pneumoniae                             | Kp1 | 3106 | Madagascar | Rural | 2016 | 90 | 78  | 5247712 | 187053 | ERS3556791 |
| SB5487 | Klebsiella quasipneumoniae subsp. quasipneumoniae | Kp2 | 2507 | Madagascar | Rural | 2016 | 89 | 39  | 5161973 | 386493 | ERS3557014 |
| SB5517 | Klebsiella quasipneumoniae subsp. quasipneumoniae | Kp2 | 2507 | Madagascar | Rural | 2016 | 91 | 46  | 5164377 | 280690 | ERS3557034 |
| SB5542 | Klebsiella quasipneumoniae subsp. quasipneumoniae | Kp2 | 526  | Madagascar | Rural | 2016 | 50 | 77  | 5128810 | 218655 | ERS3557043 |
| SB5543 | Klebsiella quasipneumoniae subsp. quasipneumoniae | Kp2 | 526  | Madagascar | Rural | 2016 | 61 | 84  | 5125768 | 190557 | ERS3557044 |
| SB5545 | Klebsiella quasipneumoniae subsp. quasipneumoniae | Kp2 | 2604 | Madagascar | Rural | 2016 | 66 | 112 | 5146606 | 123181 | ERS3557045 |
| SB5617 | Klebsiella quasipneumoniae subsp. quasipneumoniae | Kp2 | 526  | Madagascar | Rural | 2016 | 90 | 82  | 5390302 | 184454 | ERS3557081 |
| SB5621 | Klebsiella quasipneumoniae subsp. quasipneumoniae | Kp2 | 526  | Madagascar | Rural | 2016 | 93 | 109 | 5417037 | 177026 | ERS3556845 |
| SB5373 | Klebsiella variicola subsp. variicola             | Kp3 | 2462 | Madagascar | Urban | 2015 | 89 | 47  | 5458458 | 365116 | ERS3556949 |
| SB5374 | Klebsiella variicola subsp. variicola             | Kp3 | 1615 | Madagascar | Urban | 2015 | 90 | 80  | 5657557 | 190187 | ERS3556798 |
| SB5375 | Klebsiella variicola subsp. variicola             | Kp3 | 2463 | Madagascar | Urban | 2015 | 76 | 58  | 5554336 | 270995 | ERS3556950 |
| SB5377 | Klebsiella variicola subsp. variicola             | Kp3 | 2464 | Madagascar | Urban | 2015 | 76 | 158 | 5644723 | 142194 | ERS3556952 |
| SB5379 | Klebsiella variicola subsp. variicola             | Kp3 | 906  | Madagascar | Urban | 2015 | 77 | 72  | 5767679 | 316215 | ERS3556874 |
| SB5381 | Klebsiella variicola subsp. variicola             | Kp3 | 925  | Madagascar | Urban | 2015 | 88 | 65  | 5639770 | 337248 | ERS3556955 |
| SB5386 | Klebsiella variicola subsp. variicola             | Kp3 | 2465 | Madagascar | Urban | 2015 | 88 | 55  | 5440752 | 341428 | ERS3556960 |
| SB5388 | Klebsiella variicola subsp. variicola             | Kp3 | 906  | Madagascar | Urban | 2015 | 90 | 68  | 5755776 | 262066 | ERS3556875 |
| SB5396 | Klebsiella variicola subsp. variicola             | Kp3 | 2468 | Madagascar | Urban | 2015 | 91 | 106 | 5685191 | 175804 | ERS3556965 |
| SB5400 | Klebsiella variicola subsp. variicola             | Kp3 | 2469 | Madagascar | Urban | 2015 | 77 | 40  | 5602483 | 404425 | ERS3556966 |
| SB5402 | Klebsiella variicola subsp. variicola             | Kp3 | 2470 | Madagascar | Urban | 2016 | 72 | 81  | 5487184 | 274238 | ERS3556968 |
| SB5405 | Klebsiella variicola subsp. variicola             | Kp3 | 2472 | Madagascar | Urban | 2016 | 56 | 135 | 5602972 | 151911 | ERS3556831 |

|        |                                           |      |            |       |      |    |     |         |        |            |
|--------|-------------------------------------------|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5409 | Klebsiella variicola subsp. variicola Kp3 | 2475 | Madagascar | Urban | 2016 | 68 | 102 | 5349267 | 106221 | ERS3556973 |
| SB5413 | Klebsiella variicola subsp. variicola Kp3 | 2477 | Madagascar | Urban | 2016 | 88 | 74  | 5357522 | 162652 | ERS3556975 |
| SB5416 | Klebsiella variicola subsp. variicola Kp3 | 2479 | Madagascar | Urban | 2016 | 75 | 113 | 5704285 | 286204 | ERS3556978 |
| SB5430 | Klebsiella variicola subsp. variicola Kp3 | 919  | Madagascar | Urban | 2016 | 56 | 192 | 5385660 | 93766  | ERS3556984 |
| SB5433 | Klebsiella variicola subsp. variicola Kp3 | 1562 | Madagascar | Urban | 2016 | 73 | 194 | 5952617 | 114346 | ERS3556835 |
| SB5437 | Klebsiella variicola subsp. variicola Kp3 | 919  | Madagascar | Urban | 2016 | 83 | 165 | 5346358 | 108291 | ERS3556988 |
| SB5444 | Klebsiella variicola subsp. variicola Kp3 | 2488 | Madagascar | Urban | 2016 | 92 | 44  | 5634621 | 373402 | ERS3556992 |
| SB5445 | Klebsiella variicola subsp. variicola Kp3 | 1843 | Madagascar | Urban | 2016 | 81 | 45  | 5365236 | 375469 | ERS3556993 |
| SB5449 | Klebsiella variicola subsp. variicola Kp3 | 2490 | Madagascar | Rural | 2015 | 75 | 52  | 5477885 | 212986 | ERS3556996 |
| SB5454 | Klebsiella variicola subsp. variicola Kp3 | 2491 | Madagascar | Rural | 2015 | 85 | 73  | 5442774 | 219515 | ERS3557001 |
| SB5465 | Klebsiella variicola subsp. variicola Kp3 | 906  | Madagascar | Rural | 2015 | 92 | 71  | 5763616 | 199460 | ERS3556877 |
| SB5485 | Klebsiella variicola subsp. variicola Kp3 | 2475 | Madagascar | Rural | 2016 | 67 | 60  | 5373490 | 222789 | ERS3557012 |
| SB5488 | Klebsiella variicola subsp. variicola Kp3 | 2508 | Madagascar | Rural | 2016 | 83 | 73  | 5425002 | 185824 | ERS3557015 |
| SB5493 | Klebsiella variicola subsp. variicola Kp3 | 2490 | Madagascar | Rural | 2016 | 66 | 68  | 5386564 | 216049 | ERS3557019 |
| SB5495 | Klebsiella variicola subsp. variicola Kp3 | 2551 | Madagascar | Rural | 2016 | 86 | 41  | 5486130 | 311060 | ERS3557020 |
| SB5496 | Klebsiella variicola subsp. variicola Kp3 | 2551 | Madagascar | Rural | 2016 | 87 | 37  | 5399987 | 394028 | ERS3557021 |
| SB5505 | Klebsiella variicola subsp. variicola Kp3 | 2553 | Madagascar | Rural | 2016 | 92 | 45  | 5575376 | 314335 | ERS3557026 |
| SB5519 | Klebsiella variicola subsp. variicola Kp3 | 697  | Madagascar | Rural | 2016 | 87 | 54  | 5529631 | 309412 | ERS3557157 |
| SB5520 | Klebsiella variicola subsp. variicola Kp3 | 919  | Madagascar | Rural | 2016 | 75 | 108 | 5514925 | 193275 | ERS3556859 |
| SB5552 | Klebsiella variicola subsp. variicola Kp3 | 2606 | Madagascar | Rural | 2016 | 61 | 77  | 5482237 | 209735 | ERS3557051 |

|        |                                                    |     |      |            |       |      |    |     |         |        |            |
|--------|----------------------------------------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5561 | Klebsiella variicola subsp. variicola              | Kp3 | 2609 | Madagascar | Rural | 2016 | 84 | 61  | 5402273 | 246602 | ERS3557054 |
| SB5566 | Klebsiella variicola subsp. variicola              | Kp3 | 357  | Madagascar | Rural | 2016 | 67 | 135 | 5613895 | 171203 | ERS3557055 |
| SB5582 | Klebsiella variicola subsp. variicola              | Kp3 | 2611 | Madagascar | Rural | 2016 | 90 | 75  | 5375242 | 213483 | ERS3556821 |
| SB5586 | Klebsiella variicola subsp. variicola              | Kp3 | 1984 | Madagascar | Rural | 2016 | 90 | 52  | 5332452 | 218499 | ERS3557066 |
| SB5600 | Klebsiella variicola subsp. variicola              | Kp3 | 2594 | Madagascar | Rural | 2016 | 92 | 48  | 5497328 | 281143 | ERS3557069 |
| SB5950 | Klebsiella variicola subsp. variicola              | Kp3 | 3101 | Madagascar | Urban | 2016 | 92 | 42  | 5355670 | 225689 | ERS3557147 |
| SB5951 | Klebsiella variicola subsp. variicola              | Kp3 | 3101 | Madagascar | Urban | 2016 | 92 | 37  | 5366750 | 246235 | ERS3557148 |
| SB5960 | Klebsiella variicola subsp. variicola              | Kp3 | 712  | Madagascar | Rural | 2016 | 92 | 78  | 5725724 | 258443 | ERS3557154 |
| SB5966 | Klebsiella variicola subsp. variicola              | Kp3 | 3103 | Madagascar | Rural | 2016 | 89 | 73  | 5700642 | 175109 | ERS3557156 |
| SB5967 | Klebsiella variicola subsp. variicola              | Kp3 | 3104 | Madagascar | Rural | 2016 | 89 | 75  | 5889124 | 220557 | ERS3556904 |
| SB5968 | Klebsiella variicola subsp. variicola              | Kp3 | 919  | Madagascar | Rural | 2016 | 90 | 90  | 5555178 | 172043 | ERS3556870 |
| SB5971 | Klebsiella variicola subsp. variicola              | Kp3 | 3104 | Madagascar | Rural | 2016 | 90 | 69  | 5886084 | 208898 | ERS3556905 |
| SB5972 | Klebsiella variicola subsp. variicola              | Kp3 | 3105 | Madagascar | Rural | 2016 | 90 | 58  | 5491854 | 188345 | ERS3556884 |
| SB5410 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2476 | Madagascar | Urban | 2016 | 88 | 60  | 5435462 | 296618 | ERS3556974 |
| SB5414 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2478 | Madagascar | Urban | 2016 | 92 | 58  | 5422494 | 295746 | ERS3556976 |
| SB5420 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 841  | Madagascar | Urban | 2016 | 88 | 69  | 5411541 | 297876 | ERS3556979 |
| SB5428 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2483 | Madagascar | Urban | 2016 | 83 | 32  | 5026681 | 379465 | ERS3556983 |
| SB5431 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2483 | Madagascar | Urban | 2016 | 57 | 44  | 5014161 | 371435 | ERS3556985 |
| SB5477 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2503 | Madagascar | Rural | 2015 | 62 | 97  | 5357585 | 228691 | ERS3556801 |
| SB5504 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2489 | Madagascar | Rural | 2016 | 80 | 127 | 5414623 | 153156 | ERS3556857 |

|        |                                                    |     |      |            |       |      |    |     |         |        |            |
|--------|----------------------------------------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5527 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2558 | Madagascar | Rural | 2016 | 85 | 101 | 5476532 | 212814 | ERS3556898 |
| SB5528 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2559 | Madagascar | Rural | 2016 | 80 | 94  | 5392586 | 218306 | ERS3556860 |
| SB5533 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2562 | Madagascar | Rural | 2016 | 89 | 143 | 5258337 | 105503 | ERS3556843 |
| SB5536 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2478 | Madagascar | Rural | 2016 | 92 | 64  | 5429262 | 295885 | ERS3557041 |
| SB5537 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1224 | Madagascar | Rural | 2016 | 64 | 135 | 5249042 | 152411 | ERS3556778 |
| SB5554 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2607 | Madagascar | Rural | 2016 | 90 | 35  | 5124474 | 294335 | ERS3557052 |
| SB5556 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2608 | Madagascar | Rural | 2016 | 90 | 123 | 5135771 | 100881 | ERS3556844 |
| SB5570 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2610 | Madagascar | Rural | 2016 | 59 | 197 | 5188694 | 72423  | ERS3556940 |
| SB5577 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 283  | Madagascar | Rural | 2016 | 78 | 92  | 5387217 | 198227 | ERS3556902 |
| SB5587 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2592 | Madagascar | Rural | 2016 | 60 | 64  | 5164824 | 207974 | ERS3556861 |
| SB5591 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2592 | Madagascar | Rural | ND   | 83 | 31  | 5116759 | 371534 | ERS3556863 |
| SB5598 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2593 | Madagascar | Rural | 2016 | 92 | 58  | 5222359 | 301855 | ERS3557068 |
| SB5607 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1584 | Madagascar | Rural | 2016 | 93 | 79  | 5202162 | 195586 | ERS3557075 |
| SB5615 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2059 | Madagascar | Rural | 2016 | 91 | 30  | 5113756 | 364932 | ERS3557079 |
| SB5957 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2721 | Madagascar | Urban | 2016 | 90 | 40  | 5188994 | 363415 | ERS3557153 |
| SB5387 | Klebsiella variicola subsp. tropicalensis          | Kp5 | 2466 | Madagascar | Urban | 2015 | 87 | 56  | 5698764 | 279038 | ERS2787526 |
| SB5439 | Klebsiella variicola subsp. tropicalensis          | Kp5 | 2486 | Madagascar | Urban | 2016 | 92 | 45  | 5613167 | 311118 | ERS2787527 |
| SB5455 | Klebsiella variicola subsp. tropicalensis          | Kp5 | 2630 | Madagascar | Rural | 2015 | 86 | 43  | 5338997 | 298293 | ERS3557002 |
| SB5531 | Klebsiella variicola subsp. tropicalensis          | Kp5 | 2561 | Madagascar | Rural | 2016 | 89 | 48  | 5442219 | 339248 | ERS2787528 |
| SB5544 | Klebsiella variicola subsp. tropicalensis          | Kp5 | 2605 | Madagascar | Rural | 2016 | 65 | 57  | 5488176 | 302938 | ERS2787529 |

|        |                                           |     |      |            |       |      |    |     |         |        |            |
|--------|-------------------------------------------|-----|------|------------|-------|------|----|-----|---------|--------|------------|
| SB5567 | Klebsiella variicola subsp. tropicalensis | Kp5 | 2717 | Madagascar | Rural | 2016 | 89 | 85  | 5730017 | 378142 | ERS3556879 |
| SB5610 | Klebsiella variicola subsp. tropicalensis | Kp5 | 2600 | Madagascar | Rural | 2016 | 82 | 64  | 5631209 | 376790 | ERS2787530 |
| SB5727 | Klebsiella pneumoniae                     | Kp1 | 1303 | Senegal    | Rural | 2016 | 89 | 180 | 5459317 | 93097  | ERS3556805 |
| SB5728 | Klebsiella pneumoniae                     | Kp1 | 1886 | Senegal    | Rural | 2016 | 89 | 93  | 5024483 | 105946 | ERS3556774 |
| SB5730 | Klebsiella pneumoniae                     | Kp1 | 3108 | Senegal    | Urban | 2016 | 91 | 144 | 5392688 | 126207 | ERS3557158 |
| SB5731 | Klebsiella pneumoniae                     | Kp1 | 3109 | Senegal    | Rural | 2016 | 89 | 85  | 5493388 | 203723 | ERS3557159 |
| SB5734 | Klebsiella pneumoniae                     | Kp1 | 281  | Senegal    | Rural | 2016 | 88 | 61  | 5248435 | 268545 | ERS3557161 |
| SB5735 | Klebsiella pneumoniae                     | Kp1 | 3111 | Senegal    | Rural | 2016 | 86 | 85  | 5259947 | 175089 | ERS3557162 |
| SB5736 | Klebsiella pneumoniae                     | Kp1 | 17   | Senegal    | Rural | 2016 | 81 | 72  | 5424208 | 297557 | ERS3557163 |
| SB5737 | Klebsiella pneumoniae                     | Kp1 | 1119 | Senegal    | Rural | 2016 | 91 | 146 | 5398689 | 164307 | ERS3556896 |
| SB5742 | Klebsiella pneumoniae                     | Kp1 | 528  | Senegal    | Rural | 2016 | 86 | 103 | 5558787 | 163712 | ERS3556872 |
| SB5743 | Klebsiella pneumoniae                     | Kp1 | 3113 | Senegal    | Rural | 2016 | 82 | 91  | 5282053 | 150704 | ERS3557164 |
| SB5749 | Klebsiella pneumoniae                     | Kp1 | 17   | Senegal    | Rural | 2016 | 89 | 92  | 5310170 | 171120 | ERS3557166 |
| SB5751 | Klebsiella pneumoniae                     | Kp1 | 3116 | Senegal    | Rural | 2016 | 90 | 75  | 5119808 | 174613 | ERS3557168 |
| SB5752 | Klebsiella pneumoniae                     | Kp1 | 36   | Senegal    | Rural | 2016 | 90 | 115 | 5306071 | 165411 | ERS3557169 |
| SB5753 | Klebsiella pneumoniae                     | Kp1 | 17   | Senegal    | Rural | 2016 | 84 | 48  | 5105070 | 310709 | ERS3557170 |
| SB5756 | Klebsiella pneumoniae                     | Kp1 | 3118 | Senegal    | Rural | 2016 | 92 | 67  | 5297658 | 211640 | ERS3557172 |
| SB5759 | Klebsiella pneumoniae                     | Kp1 | 111  | Senegal    | Rural | 2016 | 91 | 84  | 5325877 | 159857 | ERS3557174 |
| SB5762 | Klebsiella pneumoniae                     | Kp1 | 3258 | Senegal    | Rural | 2016 | 87 | 59  | 5350135 | 198994 | ERS3557176 |
| SB5765 | Klebsiella pneumoniae                     | Kp1 | 1962 | Senegal    | Rural | 2016 | 91 | 82  | 5419644 | 148060 | ERS3557178 |

|        |                       |     |      |         |       |      |        |    |     |         |        |            |
|--------|-----------------------|-----|------|---------|-------|------|--------|----|-----|---------|--------|------------|
| SB5768 | Klebsiella pneumoniae | Kp1 | 3123 | Senegal | Rural | 2016 |        | 88 | 59  | 5392118 | 194742 | ERS3557180 |
| SB5770 | Klebsiella pneumoniae | Kp1 | 152  | Senegal | Rural | 2016 |        | 89 | 85  | 5390648 | 176784 | ERS3557181 |
| SB5771 | Klebsiella pneumoniae | Kp1 | 3124 | Senegal | Rural | 2016 |        | 86 | 64  | 5208506 | 196298 | ERS3557182 |
| SB5773 | Klebsiella pneumoniae | Kp1 | 3260 | Senegal | Rural | 2016 |        | 91 | 86  | 5149261 | 130202 | ERS3557184 |
| SB5774 | Klebsiella pneumoniae | Kp1 | 3261 | Senegal | Rural | 2016 |        | 90 | 61  | 5206782 | 172719 | ERS3557185 |
| SB5775 | Klebsiella pneumoniae | Kp1 | 3262 | Senegal | Rural | 2016 |        | 88 | 54  | 5035131 | 205363 | ERS3557186 |
| SB5776 | Klebsiella pneumoniae | Kp1 | 337  | Senegal | Rural | 2016 |        | 90 | 94  | 5180324 | 154063 | ERS3557187 |
| SB5777 | Klebsiella pneumoniae | Kp1 | 397  | Senegal | Rural | 2016 |        | 90 | 127 | 5782991 | 151257 | ERS3557188 |
| SB5779 | Klebsiella pneumoniae | Kp1 | 629  | Senegal | Rural | 2016 |        | 90 | 58  | 5096179 | 210411 | ERS3557190 |
| SB5783 | Klebsiella pneumoniae | Kp1 | 3157 | Senegal | Rural | 2016 |        | 91 | 67  | 5170911 | 208422 | ERS3557194 |
| SB5786 | Klebsiella pneumoniae | Kp1 | 277  | Senegal | Rural | 2016 |        | 88 | 118 | 5275557 | 178172 | ERS3557195 |
| SB5787 | Klebsiella pneumoniae | Kp1 | 281  | Senegal | Rural | 2016 |        | 90 | 88  | 5178844 | 179779 | ERS3557196 |
| SB5788 | Klebsiella pneumoniae | Kp1 | 3267 | Senegal | Rural | 2016 |        | 87 | 81  | 5029390 | 163848 | ERS3557197 |
| SB5789 | Klebsiella pneumoniae | Kp1 | 3268 | Senegal | Rural | 2016 |        | 87 | 82  | 5156984 | 136129 | ERS3557198 |
| SB5790 | Klebsiella pneumoniae | Kp1 | 151  | Senegal | Rural | 2016 |        | 81 | 119 | 5431801 | 105949 | ERS3557199 |
| SB5791 | Klebsiella pneumoniae | Kp1 | 443  | Senegal | Rural | 2016 |        | 81 | 92  | 5391364 | 237290 | ERS3557200 |
| SB5792 | Klebsiella pneumoniae | Kp1 | 514  | Senegal | Rural | 2016 | Pair 1 | 90 | 113 | 5353868 | 153475 | ERS3557201 |
| SB5793 | Klebsiella pneumoniae | Kp1 | 36   | Senegal | Rural | 2016 | Pair 1 | 91 | 49  | 5091180 | 292649 | ERS3557202 |
| SB5794 | Klebsiella pneumoniae | Kp1 | 1087 | Senegal | Rural | 2016 |        | 89 | 90  | 5275598 | 130550 | ERS3556809 |
| SB5798 | Klebsiella pneumoniae | Kp1 | 3271 | Senegal | Rural | 2016 |        | 91 | 105 | 5407526 | 182184 | ERS3556785 |

|        |                       |     |      |         |       |      |        |    |     |         |        |            |
|--------|-----------------------|-----|------|---------|-------|------|--------|----|-----|---------|--------|------------|
| SB5800 | Klebsiella pneumoniae | Kp1 | 2141 | Senegal | Rural | 2016 |        | 88 | 114 | 5182647 | 145161 | ERS3557206 |
| SB5801 | Klebsiella pneumoniae | Kp1 | 225  | Senegal | Rural | 2016 |        | 78 | 146 | 5727134 | 167616 | ERS3557207 |
| SB5802 | Klebsiella pneumoniae | Kp1 | 3273 | Senegal | Rural | 2016 |        | 90 | 68  | 5410521 | 248210 | ERS3557208 |
| SB5805 | Klebsiella pneumoniae | Kp1 | 3275 | Senegal | Rural | 2016 |        | 80 | 94  | 5014660 | 150130 | ERS3557210 |
| SB5806 | Klebsiella pneumoniae | Kp1 | 3030 | Senegal | Rural | 2016 | Pair 3 | 89 | 116 | 5405559 | 169164 | ERS3557211 |
| SB5807 | Klebsiella pneumoniae | Kp1 | 1263 | Senegal | Rural | 2016 | Pair 3 | 84 | 147 | 5466219 | 139863 | ERS3556827 |
| SB5808 | Klebsiella pneumoniae | Kp1 | 1189 | Senegal | Rural | 2016 |        | 91 | 74  | 5427900 | 244942 | ERS3557212 |
| SB5810 | Klebsiella pneumoniae | Kp1 | 1189 | Senegal | Urban | 2016 |        | 91 | 80  | 5460049 | 244949 | ERS3557214 |
| SB5811 | Klebsiella pneumoniae | Kp1 | 3277 | Senegal | Urban | 2016 |        | 91 | 56  | 5239790 | 234654 | ERS3557215 |
| SB5812 | Klebsiella pneumoniae | Kp1 | 1661 | Senegal | Urban | 2016 |        | 83 | 52  | 5110071 | 235407 | ERS3557216 |
| SB5815 | Klebsiella pneumoniae | Kp1 | 37   | Senegal | Urban | 2016 | Pair 4 | 38 | 96  | 5513110 | 204218 | ERS3556851 |
| SB5816 | Klebsiella pneumoniae | Kp1 | 3278 | Senegal | Urban | 2016 | Pair 4 | 90 | 75  | 5306105 | 176071 | ERS3557217 |
| SB5817 | Klebsiella pneumoniae | Kp1 | 292  | Senegal | Urban | 2016 |        | 90 | 68  | 5408598 | 248952 | ERS3556811 |
| SB5818 | Klebsiella pneumoniae | Kp1 | 504  | Senegal | Urban | 2016 |        | 91 | 103 | 5414856 | 140340 | ERS3556828 |
| SB5819 | Klebsiella pneumoniae | Kp1 | 45   | Senegal | Urban | 2016 |        | 91 | 64  | 5416349 | 203813 | ERS3557218 |
| SB5822 | Klebsiella pneumoniae | Kp1 | 2108 | Senegal | Urban | 2016 | Pair 7 | 91 | 125 | 5288488 | 144314 | ERS3556812 |
| SB5823 | Klebsiella pneumoniae | Kp1 | 968  | Senegal | Urban | 2016 | Pair 7 | 90 | 114 | 5408865 | 135254 | ERS3556946 |
| SB5824 | Klebsiella pneumoniae | Kp1 | 2800 | Senegal | Urban | 2016 |        | 91 | 104 | 5286656 | 128390 | ERS3556947 |
| SB5825 | Klebsiella pneumoniae | Kp1 | 491  | Senegal | Urban | 2016 |        | 92 | 88  | 5301914 | 144465 | ERS3557219 |
| SB5826 | Klebsiella pneumoniae | Kp1 | 1401 | Senegal | Urban | 2016 |        | 92 | 106 | 5339690 | 158933 | ERS3557220 |

|        |                       |     |      |         |       |      |        |    |     |         |        |            |
|--------|-----------------------|-----|------|---------|-------|------|--------|----|-----|---------|--------|------------|
| SB5828 | Klebsiella pneumoniae | Kp1 | 1049 | Senegal | Urban | 2016 |        | 92 | 60  | 5346709 | 328795 | ERS3556873 |
| SB5829 | Klebsiella pneumoniae | Kp1 | 3280 | Senegal | Urban | 2016 |        | 92 | 67  | 5204522 | 215068 | ERS3557221 |
| SB5830 | Klebsiella pneumoniae | Kp1 | 3281 | Senegal | Urban | 2016 |        | 92 | 74  | 5165672 | 174742 | ERS3557222 |
| SB5831 | Klebsiella pneumoniae | Kp1 | 1401 | Senegal | Urban | 2016 |        | 89 | 118 | 5342371 | 195948 | ERS3557223 |
| SB5832 | Klebsiella pneumoniae | Kp1 | 2655 | Senegal | Urban | 2016 |        | 89 | 93  | 5356443 | 234283 | ERS3556906 |
| SB5833 | Klebsiella pneumoniae | Kp1 | 502  | Senegal | Urban | 2016 |        | 91 | 119 | 5261707 | 117110 | ERS3556775 |
| SB5834 | Klebsiella pneumoniae | Kp1 | 2715 | Senegal | Urban | 2016 |        | 91 | 57  | 5290405 | 220671 | ERS3556897 |
| SB5836 | Klebsiella pneumoniae | Kp1 | 1419 | Senegal | Urban | 2016 |        | 92 | 62  | 5123589 | 185639 | ERS3557224 |
| SB5837 | Klebsiella pneumoniae | Kp1 | 3282 | Senegal | Rural | 2016 |        | 92 | 57  | 5321722 | 199551 | ERS3556813 |
| SB5838 | Klebsiella pneumoniae | Kp1 | 3282 | Senegal | Urban | 2016 |        | 91 | 65  | 5316598 | 215700 | ERS3556814 |
| SB5839 | Klebsiella pneumoniae | Kp1 | 411  | Senegal | Urban | 2016 |        | 91 | 68  | 5315616 | 357999 | ERS3557225 |
| SB5841 | Klebsiella pneumoniae | Kp1 | 1666 | Senegal | Urban | 2016 |        | 92 | 126 | 5126909 | 121647 | ERS3556786 |
| SB5842 | Klebsiella pneumoniae | Kp1 | 3283 | Senegal | Rural | 2016 |        | 89 | 105 | 5528949 | 187384 | ERS3557226 |
| SB5847 | Klebsiella pneumoniae | Kp1 | 3286 | Senegal | Rural | 2016 |        | 71 | 108 | 5329435 | 140852 | ERS3557230 |
| SB5849 | Klebsiella pneumoniae | Kp1 | 29   | Senegal | Rural | 2016 | Pair 5 | 64 | 169 | 5672063 | 137034 | ERS3557232 |
| SB5850 | Klebsiella pneumoniae | Kp1 | 3287 | Senegal | Rural | 2016 | Pair 5 | 90 | 170 | 5494158 | 133356 | ERS3556792 |
| SB5851 | Klebsiella pneumoniae | Kp1 | 249  | Senegal | Rural | 2016 |        | 82 | 88  | 5342878 | 130037 | ERS3557233 |
| SB5854 | Klebsiella pneumoniae | Kp1 | 1419 | Senegal | Rural | 2016 |        | 79 | 79  | 5113371 | 156498 | ERS3557236 |
| SB5856 | Klebsiella pneumoniae | Kp1 | 3290 | Senegal | Urban | 2016 |        | 83 | 63  | 5140202 | 237840 | ERS3557237 |
| SB5858 | Klebsiella pneumoniae | Kp1 | 3286 | Senegal | Rural | 2016 |        | 87 | 72  | 5289021 | 162987 | ERS3557238 |

|        |                       |     |      |         |       |      |        |    |     |         |        |            |
|--------|-----------------------|-----|------|---------|-------|------|--------|----|-----|---------|--------|------------|
| SB5859 | Klebsiella pneumoniae | Kp1 | 1015 | Senegal | Rural | 2016 |        | 88 | 76  | 5206159 | 136951 | ERS3557239 |
| SB5862 | Klebsiella pneumoniae | Kp1 | 13   | Senegal | Rural | 2016 |        | 92 | 115 | 5520797 | 121910 | ERS3557241 |
| SB5863 | Klebsiella pneumoniae | Kp1 | 788  | Senegal | Rural | 2016 | Pair 6 | 82 | 49  | 5148220 | 218261 | ERS3557242 |
| SB5864 | Klebsiella pneumoniae | Kp1 | 3293 | Senegal | Rural | 2016 | Pair 6 | 92 | 50  | 5284856 | 299377 | ERS3557243 |
| SB5865 | Klebsiella pneumoniae | Kp1 | 3294 | Senegal | Urban | 2016 |        | 93 | 52  | 5212655 | 203902 | ERS3557244 |
| SB5866 | Klebsiella pneumoniae | Kp1 | 37   | Senegal | Urban | 2016 |        | 94 | 75  | 5288263 | 203773 | ERS3557245 |
| SB5867 | Klebsiella pneumoniae | Kp1 | 3295 | Senegal | Rural | 2016 |        | 93 | 115 | 5290947 | 133375 | ERS3557246 |
| SB5869 | Klebsiella pneumoniae | Kp1 | 3297 | Senegal | Rural | 2016 |        | 94 | 55  | 5227797 | 250406 | ERS3557248 |
| SB6073 | Klebsiella pneumoniae | Kp1 | 152  | Senegal | Rural | 2016 |        | 88 | 99  | 5408575 | 219852 | ERS3557249 |
| SB6074 | Klebsiella pneumoniae | Kp1 | 3337 | Senegal | Rural | 2016 |        | 86 | 62  | 5145564 | 286179 | ERS3557250 |
| SB6076 | Klebsiella pneumoniae | Kp1 | 3338 | Senegal | Rural | 2016 |        | 88 | 48  | 5264137 | 279955 | ERS3557252 |
| SB6077 | Klebsiella pneumoniae | Kp1 | 3339 | Senegal | Rural | 2016 |        | 87 | 33  | 5331978 | 389116 | ERS3557253 |
| SB6078 | Klebsiella pneumoniae | Kp1 | 661  | Senegal | Rural | 2016 |        | 87 | 39  | 5131974 | 366827 | ERS3557254 |
| SB6079 | Klebsiella pneumoniae | Kp1 | 867  | Senegal | Rural | 2016 |        | 89 | 80  | 5346442 | 180235 | ERS3556787 |
| SB6080 | Klebsiella pneumoniae | Kp1 | 1662 | Senegal | Rural | 2016 |        | 86 | 39  | 5120121 | 380942 | ERS3557255 |
| SB6081 | Klebsiella pneumoniae | Kp1 | 3340 | Senegal | Rural | 2016 |        | 87 | 39  | 5177609 | 303260 | ERS3557256 |
| SB6082 | Klebsiella pneumoniae | Kp1 | 3341 | Senegal | Rural | 2016 |        | 86 | 49  | 5091487 | 225350 | ERS3557257 |
| SB6085 | Klebsiella pneumoniae | Kp1 | 36   | Senegal | Rural | 2016 |        | 85 | 99  | 5307208 | 152475 | ERS3557258 |
| SB6086 | Klebsiella pneumoniae | Kp1 | 1962 | Senegal | Rural | 2016 |        | 84 | 60  | 5152762 | 310294 | ERS3557259 |
| SB6087 | Klebsiella pneumoniae | Kp1 | 397  | Senegal | Rural | 2016 |        | 60 | 120 | 5713771 | 240928 | ERS3557260 |

|        |                                                    |     |      |         |       |      |    |     |         |        |            |
|--------|----------------------------------------------------|-----|------|---------|-------|------|----|-----|---------|--------|------------|
| SB6088 | Klebsiella pneumoniae                              | Kp1 | 397  | Senegal | Rural | 2016 | 88 | 108 | 5706568 | 185261 | ERS3556829 |
| SB6089 | Klebsiella pneumoniae                              | Kp1 | 1658 | Senegal | Rural | 2016 | 92 | 53  | 5216422 | 283352 | ERS3557261 |
| SB6091 | Klebsiella pneumoniae                              | Kp1 | 39   | Senegal | Rural | 2016 | 88 | 61  | 5356306 | 213939 | ERS3556885 |
| SB5782 | Klebsiella quasipneumoniae subsp. quasipneumoniae  | Kp2 | 338  | Senegal | Rural | 2016 | 91 | 147 | 5417191 | 107073 | ERS3557193 |
| SB5799 | Klebsiella quasipneumoniae subsp. quasipneumoniae  | Kp2 | 3272 | Senegal | Rural | 2016 | 81 | 108 | 5268415 | 154994 | ERS3556810 |
| SB5803 | Klebsiella quasipneumoniae subsp. quasipneumoniae  | Kp2 | 3274 | Senegal | Rural | 2016 | 82 | 73  | 5193555 | 200431 | ERS3557209 |
| SB5745 | Klebsiella variicola subsp. variicola              | Kp3 | 3114 | Senegal | Rural | 2016 | 90 | 38  | 5324929 | 383397 | ERS3557165 |
| SB5754 | Klebsiella variicola subsp. variicola              | Kp3 | 3117 | Senegal | Rural | 2016 | 90 | 52  | 5618429 | 251495 | ERS3557171 |
| SB5767 | Klebsiella variicola subsp. variicola              | Kp3 | 3122 | Senegal | Rural | 2016 | 90 | 42  | 5443934 | 353056 | ERS3557179 |
| SB5868 | Klebsiella variicola subsp. variicola              | Kp3 | 3296 | Senegal | Rural | 2016 | 90 | 44  | 5411027 | 329853 | ERS3557247 |
| SB5726 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3107 | Senegal | Rural | 2016 | 88 | 78  | 5245485 | 185245 | ERS3556804 |
| SB5732 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3110 | Senegal | Rural | 2016 | 92 | 30  | 5017370 | 404872 | ERS3557160 |
| SB5739 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3112 | Senegal | Rural | 2016 | 81 | 171 | 5316922 | 119006 | ERS3556945 |
| SB5746 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3115 | Senegal | Rural | 2016 | 90 | 88  | 5402325 | 278229 | ERS3556783 |
| SB5750 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 1308 | Senegal | Rural | 2016 | 91 | 85  | 5173939 | 150803 | ERS3557167 |
| SB5758 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3120 | Senegal | Rural | 2016 | 86 | 48  | 5096656 | 259246 | ERS3557173 |
| SB5761 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3257 | Senegal | Rural | 2016 | 90 | 29  | 4912611 | 360617 | ERS3557175 |
| SB5764 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2685 | Senegal | Rural | 2016 | 92 | 22  | 5081748 | 375238 | ERS3557177 |
| SB5766 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3121 | Senegal | Rural | 2016 | 90 | 79  | 5331961 | 182382 | ERS3556784 |
| SB5772 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 138  | Senegal | Rural | 2016 | 87 | 67  | 5399904 | 223886 | ERS3557183 |

|        |                                                    |     |      |         |       |      |        |    |     |         |        |            |
|--------|----------------------------------------------------|-----|------|---------|-------|------|--------|----|-----|---------|--------|------------|
| SB5778 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3263 | Senegal | Rural | 2016 |        | 91 | 34  | 5027416 | 355298 | ERS3557189 |
| SB5780 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3264 | Senegal | Rural | 2016 |        | 91 | 85  | 5008109 | 111230 | ERS3557191 |
| SB5781 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3265 | Senegal | Rural | 2016 |        | 81 | 83  | 5012843 | 107757 | ERS3557192 |
| SB5784 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3266 | Senegal | Rural | 2016 |        | 90 | 100 | 5259734 | 124277 | ERS3556850 |
| SB5795 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3269 | Senegal | Rural | 2016 | Pair 2 | 90 | 56  | 5184927 | 295952 | ERS3557203 |
| SB5796 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3270 | Senegal | Rural | 2016 | Pair 2 | 85 | 73  | 5504592 | 260745 | ERS3557204 |
| SB5797 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 2559 | Senegal | Rural | 2016 |        | 90 | 47  | 5024763 | 253069 | ERS3557205 |
| SB5809 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3276 | Senegal | Rural | 2016 |        | 92 | 33  | 5110574 | 335878 | ERS3557213 |
| SB5843 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3284 | Senegal | Urban | 2016 |        | 92 | 44  | 5136034 | 265642 | ERS3557227 |
| SB5845 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 841  | Senegal | Rural | 2016 |        | 65 | 107 | 5277310 | 143617 | ERS3557228 |
| SB5846 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3285 | Senegal | Rural | 2016 |        | 71 | 86  | 5155621 | 119771 | ERS3557229 |
| SB5848 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 138  | Senegal | Rural | 2016 |        | 64 | 86  | 5159895 | 154440 | ERS3557231 |
| SB5852 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3288 | Senegal | Urban | 2016 |        | 59 | 159 | 5229734 | 108016 | ERS3557234 |
| SB5853 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3289 | Senegal | Urban | 2016 |        | 54 | 73  | 5065365 | 159405 | ERS3557235 |
| SB5860 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3292 | Senegal | Urban | 2016 |        | 55 | 87  | 5249338 | 193925 | ERS3557240 |
| SB6075 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | ND   | Senegal | Rural | 2016 |        | 82 | 38  | 4981468 | 282286 | ERS3557251 |
| SB6092 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3342 | Senegal | Rural | 2016 |        | 90 | 36  | 5010451 | 355888 | ERS3557262 |
| SB6093 | Klebsiella quasipneumoniae subsp. similipneumoniae | Kp4 | 3342 | Senegal | Rural | 2016 |        | 90 | 33  | 5001859 | 377541 | ERS3557263 |
| SB5857 | Klebsiella africanensis                            | Kp7 | 3291 | Senegal | Rural | 2016 |        | 87 | 50  | 5156720 | 195731 | ERS2787533 |

# defined as groups of isolates differing by less than 10 loci out of the 5,073 protein-coding genes of strain Kp616 (GenBank accession number GCA\_003076555.1).

**Table S2. High-risk clones**

| ST     | No. of assembled | Number of publication | Multidrug resistant (MDR) | Detected in this study |
|--------|------------------|-----------------------|---------------------------|------------------------|
| ST258  | 1095             | 209                   | MDR                       | No                     |
| ST11   | 744              | 236                   | MDR                       | No                     |
| ST15   | 329              | 109                   | MDR                       | Yes                    |
| ST512  | 308              | 41                    | MDR                       | No                     |
| ST101  | 206              | 68                    | MDR                       | Yes                    |
| ST307  | 197              | 34                    | MDR                       | Yes                    |
| ST147  | 176              | 86                    | MDR                       | No                     |
| ST37   | 137              | 31                    | MDR                       | Yes                    |
| ST45   | 134              | 9                     | MDR                       | Yes                    |
| ST14   | 133              | 32                    | MDR                       | Yes                    |
| ST17   | 97               | 19                    | MDR                       | Yes                    |
| ST16   | 97               | 24                    | MDR                       | No                     |
| ST23   | 87               | 66                    | HV                        | No                     |
| ST340  | 86               | 31                    | MDR                       | No                     |
| ST231  | 84               | 8                     | MDR                       | No                     |
| ST437  | 71               | 17                    | MDR                       | No                     |
| ST405  | 67               | 29                    | MDR                       | No                     |
| ST29   | 60               | 14                    | MDR                       | Yes                    |
| ST48   | 59               | 19                    | MDR                       | Yes                    |
| ST20   | 45               | 9                     | MDR                       | Yes                    |
| ST395  | 43               | 12                    | MDR                       | No                     |
| ST3128 | 41               | 1                     | MDR                       | No                     |
| ST25   | 39               | 9                     | HV                        | Yes                    |
| ST86   | 37               | 21                    | HV                        | No                     |
| ST152  | 35               | 6                     | MDR                       | Yes                    |
| ST268  | 32               | 3                     | MDR                       | No                     |
| ST36   | 32               | 5                     | MDR                       | Yes                    |
| ST336  | 32               | 8                     | MDR                       | No                     |
| ST3010 | 29               | 0                     | MDR                       | No                     |
| ST323  | 29               | 7                     | MDR                       | No                     |
| ST392  | 28               | 8                     | MDR                       | No                     |
| ST43   | 22               | 4                     | MDR                       | Yes                    |
| ST13   | 21               | 4                     | MDR                       | Yes                    |

|       |    |    |     |     |
|-------|----|----|-----|-----|
| ST39  | 20 | 7  | MDR | Yes |
| ST1   | 20 | 7  | MDR | Yes |
| ST65  | 15 | 20 | HV  | Yes |
| ST375 | 14 | 9  | HV  | Yes |
| ST274 | 12 | 5  | MDR | No  |
| ST383 | 12 | 12 | MDR | No  |
| ST76  | 11 | 5  | MDR | Yes |
| ST273 | 11 | 9  | MDR | No  |

<sup>1</sup>: As of April 2019: 6,258 available genomes

<sup>2</sup>: Klebsiella” + “pneumoniae” + “STxxx”, April 2019

**Table S3. Characteristics of the 496 Klebsiella isolates: antimicrobial susceptibility, capular types, resistance genes, virulence genes and plasmid replicons**

Please click on the link below

[https://drive.google.com/open?id=1o3XQfW\\_EGt6ffKdNp1O\\_ZyZeAmndWJAU](https://drive.google.com/open?id=1o3XQfW_EGt6ffKdNp1O_ZyZeAmndWJAU)

**Table S4.** Resistance phenotypes in urban and rural areas in Madagascar, Cambodia, Senegal and globally

| Resistance phenotype | Madagascar<br>Urban<br>(%)<br>N=84 | Rural<br>(%)<br>N=178 | p *                | Cambodia<br>Urban<br>(%)<br>N=40 | Rural<br>(%)<br>N=57 | p *                | Senegal<br>Urban<br>(%)<br>N=32 | Rural<br>(%)<br>N=105 | p *               | Total<br>(%)<br>N=496 | p **              |
|----------------------|------------------------------------|-----------------------|--------------------|----------------------------------|----------------------|--------------------|---------------------------------|-----------------------|-------------------|-----------------------|-------------------|
| AMC                  | 45.2                               | 42.7                  | 0.7                | 27.5                             | 5.3                  | 0.02               | 6.3                             | 13.3                  | 0.36 <sup>§</sup> | 29.0                  | <0.001            |
| TCC                  | 70.2                               | 50.6                  | 0.003              | 37.5                             | 10.5                 | 0.001 <sup>§</sup> | 6.3                             | 12.4                  | 0.52 <sup>§</sup> | 37.3                  | <0.001            |
| TZP                  | 25.0                               | 14.0                  | 0.03               | 25.0                             | 5.3                  | 0.005 <sup>§</sup> | 0.0                             | 1.0                   | 1 <sup>§</sup>    | 12.1                  | <0.001            |
| CEF                  | 4.8                                | 15.2                  | 0.02               | 30.0                             | 12.3                 | 0.03               | 6.3                             | 9.5                   | 0.73 <sup>§</sup> | 12.5                  | 0.04              |
| CTX                  | 3.6                                | 10.7                  | 0.05               | 27.5                             | 5.3                  | 0.002              | 0.0                             | 1.0                   | 1 <sup>§</sup>    | 7.5                   | <0.001            |
| CAZ                  | 3.6                                | 9.6                   | 0.09               | 27.5                             | 5.3                  | 0.002              | 3.1                             | 2.9                   | 1 <sup>§</sup>    | 7.7                   | 0.005             |
| FEP                  | 4.8                                | 10.7                  | 0.11               | 25.0                             | 5.3                  | 0.005              | 0.0                             | 1.0                   | 1 <sup>§</sup>    | 7.5                   | 0.001             |
| ATM                  | 3.6                                | 9.0                   | 0.11               | 27.5                             | 5.3                  | 0.002              | 0.0                             | 1.0                   | 1 <sup>§</sup>    | 6.9                   | <0.001            |
| IMI                  | 1.2                                | 1.1                   | 1 <sup>§</sup>     | 0.0                              | 0.0                  | -                  | 3.1                             | 0.0                   | 0.23 <sup>§</sup> | 0.8                   | 0.82 <sup>§</sup> |
| AMK                  | 1.2                                | 0.0                   | 0.32 <sup>§</sup>  | 0.0                              | 0.0                  | -                  | 0.0                             | 1.0                   | 1 <sup>§</sup>    | 0.4                   | 1 <sup>§</sup>    |
| GEN                  | 2.4                                | 5.1                   | 0.51 <sup>§</sup>  | 32.5                             | 7.0                  | 0.001              | 3.1                             | 1.0                   | 0.41 <sup>§</sup> | 6.0                   | <0.001            |
| TOB                  | 6.0                                | 9.0                   | 0.47 <sup>§</sup>  | 30.0                             | 7.0                  | 0.003              | 3.1                             | 2.9                   | 1 <sup>§</sup>    | 8.3                   | 0.001             |
| NAL                  | 1.2                                | 3.4                   | 0.44 <sup>§</sup>  | 22.5                             | 10.5                 | 0.1                | 0.0                             | 1.9                   | 1 <sup>§</sup>    | 4.8                   | <0.001            |
| CIP                  | 0.0                                | 7.9                   | 0.006 <sup>§</sup> | 15.0                             | 1.8                  | 0.02 <sup>§</sup>  | 3.1                             | 2.9                   | 1 <sup>§</sup>    | 5.0                   | 0.32              |
| TET                  | 36.9                               | 42.7                  | 0.37               | 45.0                             | 14.0                 | 0.001              | 18.8                            | 11.4                  | 0.37 <sup>§</sup> | 30.4                  | <0.001            |
| SXT                  | 27.4                               | 33.7                  | 0.3                | 40.0                             | 12.3                 | 0.002              | 50.0                            | 29.5                  | 0.03              | 30.8                  | 0.21              |
| DDST                 | 3.6                                | 10.7                  | 0.05               | 27.5                             | 5.3                  | 0.002              | 0.0                             | 1.0                   | 1 <sup>§</sup>    | 7.5                   | <0.001            |

\* Comparison between rural and urban sites; \*\* Comparison between countries

§ Fisher's exact test

**Table S5. Resistant genes rates among *K. pneumoniae* isolates, by country and site**

|               | Madagascar |       |       | Cambodia |       |         | Senegal |       |       | Total | p **   |
|---------------|------------|-------|-------|----------|-------|---------|---------|-------|-------|-------|--------|
|               | Urban      | Rural | p*    | Urban    | Rural | p*      | Urban   | Rural | p*    |       |        |
|               | (%)        | (%)   |       | (%)      | (%)   |         | (%)     | (%)   |       |       |        |
|               | N=84       | N=178 |       | N=40     | N=57  |         | N=32    | N=105 |       |       |        |
| CTX-M         | 3.6        | 8.4   | 0.15  | 25.0     | 3.5   | 0.003 § | 0.0     | 0.0   | -     | 6.0   | <0.001 |
| SHV-ESBL      | 0.0        | 2.2   | 0.31§ | 2.5      | 1.8   | 1       | 0.0     | 0.0   | -     | 1.2   | 0.24§  |
| pAmpC         | 1.2        | 1.1   | 1§    | 0.0      | 0.0   | -       | 0.0     | 0.0   | -     | 0.6   | 0.43§  |
| TEM-1         | 15.5       | 20.8  | 0.31  | 35.0     | 10.5  | 0.003   | 6.3     | 3.8   | 0.62§ | 15.3  | <0.001 |
| LAP-2         | 1.2        | 2.8   | 0.67§ | 7.5      | 3.5   | 0.4§    | 0.0     | 0.0   | -     | 2.2   | 0.02§  |
| OXA-1         | 2.4        | 2.8   | 1§    | 15.0     | 3.5   | 0.06§   | 0.0     | 0.0   | -     | 3.0   | 0.001§ |
| SCO-1         | 0.0        | 1.1   | 1§    | 0.0      | 0.0   | -       | 0.0     | 0.0   | -     | 0.4   | 0.71§  |
| AAC(3)-IIa/d  | 3.6        | 5.6   | 0.56§ | 32.5     | 7.0   | 0.001   | 3.1     | 0.0   | 0.23§ | 6.3   | <0.001 |
| AAC(6')-Ib-cr | 2.4        | 3.4   | 1§    | 20.0     | 3.5   | 0.01§   | 0.0     | 0.0   | -     | 3.6   | <0.001 |
| AAC(6')-Ib    | 0.0        | 0.0   | -     | 2.5      | 0.0   | 0.41§   | 0.0     | 0.0   | -     | 0.2   | 0.2§   |
| AadA          | 9.5        | 11.2  | 0.68  | 20.0     | 3.5   | 0.01§   | 15.6    | 7.6   | 0.18§ | 10.3  | 0.9    |
| AadB          | 0.0        | 0.0   | -     | 0.0      | 0.0   | -       | 0.0     | 1.0   | 1§    | 0.2   | 0.5§   |
| APH(6)-Id     | 14.3       | 21.9  | 0.15  | 32.5     | 8.8   | 0.003   | 15.6    | 5.7   | 0.13§ | 16.1  | 0.01   |
| APH(3'')-Ib   | 13.1       | 20.2  | 0.16  | 32.5     | 8.8   | 0.003   | 15.6    | 5.7   | 0.13§ | 15.3  | 0.02   |
| APH(3')-Ia    | 2.4        | 0.6   | 0.24§ | 0.0      | 0.0   | -       | 0.0     | 0.0   | -     | 0.6   | 0.43§  |
| QNDR          | 1.2        | 1.1   | 1§    | 7.5      | 1.8   | 0.3§    | 0.0     | 0.0   | -     | 1.4   | 0.03§  |
| QnrB          | 2.4        | 3.4   | 1§    | 15.0     | 1.8   | 0.02§   | 0.0     | 1.0   | 1     | 3.2   | 0.02§  |
| QnrS          | 1.2        | 3.9   | 0.44§ | 20.0     | 12.3  | 0.3     | 0.0     | 1.0   | 1§    | 4.8   | <0.001 |
| Sul1          | 20.2       | 23.0  | 0.61  | 17.5     | 3.5   | 0.03§   | 21.9    | 7.6   | 0.05§ | 16.5  | 0.002  |
| Sul2          | 17.9       | 28.1  | 0.07  | 32.5     | 8.8   | 0.003   | 12.5    | 4.8   | 0.21§ | 18.5  | <0.001 |
| Sul3          | 1.2        | 0.0   | 0.32§ | 5.0      | 3.5   | 1       | 0.0     | 0.0   | -     | 1.0   | 0.001§ |
| Dfr           | 27.4       | 35.4  | 0.2   | 37.5     | 14.0  | 0.007   | 25.0    | 9.5   | 0.04§ | 25.6  | <0.001 |
| TetA          | 15.5       | 21.3  | 0.26  | 40.0     | 12.3  | 0.002   | 9.4     | 6.7   | 0.7§  | 16.9  | 0.001  |
| TetB          | 0.0        | 0.6   | 1§    | 0.0      | 0.0   | -       | 0.0     | 0.0   | -     | 0.2   | 1§     |
| TetC          | 0.0        | 1.1   | 1§    | 0.0      | 1.8   | 1       | 0.0     | 0.0   | -     | 0.6   | 0.6§   |
| TetD          | 11.9       | 11.2  | 0.87  | 2.5      | 0.0   | 0.4§    | 9.4     | 1.9   | 0.08§ | 7.3   | 0.001  |
| MefB          | 0.0        | 0.0   | -     | 2.5      | 0.0   | 0.4§    | 0.0     | 0.0   | -     | 0.2   | 0.2§   |
| MphA          | 3.6        | 5.1   | 0.76§ | 7.5      | 1.8   | 0.3§    | 3.1     | 0.0   | 0.23§ | 3.4   | 0.1§   |
| CatB          | 2.4        | 2.8   | 1§    | 15.0     | 3.5   | 0.06§   | 0.0     | 0.0   | -     | 3.0   | 0.001§ |
| CatI          | 2.4        | 5.1   | 0.51§ | 2.5      | 0.0   | 0.41§   | 3.1     | 1.0   | 0.41§ | 2.8   | 0.23§  |
| CatII         | 0.0        | 0.6   | 1§    | 0.0      | 1.8   | 1§      | 0.0     | 0.0   | -     | 0.4   | 0.43§  |
| Cml           | 1.2        | 0.0   | 0.32§ | 2.5      | 1.8   | 1§      | 3.1     | 0.0   | 0.23§ | 0.8   | 0.24§  |
| FloR          | 0.0        | 1.1   | 1§    | 12.5     | 1.8   | 0.07§   | 0.0     | 0.0   | -     | 1.6   | 0.001§ |
| Arr           | 0.0        | 0.6   | 1§    | 5.0      | 0.0   | 0.17§   | 0.0     | 0.0   | -     | 0.6   | 0.17§  |
| QacEΔ1        | 19.0       | 24.7  | 0.31  | 17.5     | 5.3   | 0.09§   | 21.9    | 7.6   | 0.05  | 17.1  | 0.001  |
| QacL          | 0.0        | 0.0   | -     | 2.5      | 1.8   | 1       | 0.0     | 0.0   | -     | 0.4   | 0.04§  |

\*Comparison between rural and urban sites; \*\*Comparison between countries

§Fisher's exact test

**Table S6. Virulence genes rates by country and site**

| Virulence genes | Madagascar           |                    |                   | Cambodia             |                      |                   | Senegal              |                       |                   | Total<br>(%)<br>N=496 | p **                |
|-----------------|----------------------|--------------------|-------------------|----------------------|----------------------|-------------------|----------------------|-----------------------|-------------------|-----------------------|---------------------|
|                 | Urban<br>(%)<br>N=84 | Rural<br>(%) N=178 | p*                | Urban<br>(%)<br>N=40 | Rural<br>(%)<br>N=57 | p*                | Urban<br>(%)<br>N=32 | Rural<br>(%)<br>N=105 | p *               |                       |                     |
| Yersiniabactin  | 5.9                  | 11.8               | 0.14              | 22.5                 | 12.3                 | 0.18              | 6.2                  | 17.1                  | 0.16 <sup>§</sup> | 12.5                  | 0.17                |
| Colibactin      | 0                    | 0.6                | 1 <sup>§</sup>    | 0                    | 0                    | -                 | 0                    | 0.9                   | 1 <sup>§</sup>    | 0.4                   | 1 <sup>§</sup>      |
| Aerobactin      | 7.1                  | 4.5                | 0.4 <sup>§</sup>  | 0                    | 0                    | -                 | 0                    | 0                     | -                 | 2.8                   | 0.001 <sup>§</sup>  |
| Salmocheilin    | 2.4                  | 4.5                | 0.51 <sup>§</sup> | 2.5                  | 0                    | 0.41 <sup>§</sup> | 0                    | 0                     | -                 | 2.2                   | 0.03 <sup>§</sup>   |
| rmpA            | 1.2                  | 1.1                | 1 <sup>§</sup>    | 2.5                  | 0                    | 0.41 <sup>§</sup> | 0                    | 0                     | -                 | 0.8                   | 0.53 <sup>§</sup>   |
| rmpA2           | 1.2                  | 0.6                | 0.54 <sup>§</sup> | 0                    | 0                    | -                 | 0                    | 0                     | -                 | 0.4                   | 0.71 <sup>§</sup>   |
| Allantoinase    | 2.4                  | 3.4                | 1 <sup>§</sup>    | 10                   | 15.8                 | 0.41              | 9.4                  | 5.7                   | 0.44 <sup>§</sup> | 6.1                   | 0.001               |
| kfuABC          | 38.1                 | 38.2               | 1                 | 27.5                 | 61.4                 | 0.001             | 18.7                 | 40.9                  | 0.02              | 39.3                  | 0.17                |
| Microcin        | 0                    | 2.3                | 0.3 <sup>§</sup>  | 7.5                  | 8.8                  | 1 <sup>§</sup>    | 3.1                  | 3.8                   | 1 <sup>§</sup>    | 3.4                   | 0.01 <sup>§</sup>   |
| clpK            | 17.9                 | 24.2               | 0.25              | 5                    | 3.5                  | 1 <sup>§</sup>    | 25                   | 10.5                  | 0.07 <sup>§</sup> | 16.3                  | <0.001 <sup>§</sup> |

\*Comparison between rural and urban sites; \*\* Comparison between countries

§ Fisher's exact test

**Table S7.** Environmental exposures and colonization with Kp, MDR Kp and virulent Kp among pregnant women in Madagascar, Cambodia and Senegal 2015-2016

| Madagascar                                       | Kp (N=405)              |     |                       |                |      | MDR-Kp (N=262)      |     |                       |                |      | Virulent Kp (N=262) |      |                       |                |     |
|--------------------------------------------------|-------------------------|-----|-----------------------|----------------|------|---------------------|-----|-----------------------|----------------|------|---------------------|------|-----------------------|----------------|-----|
|                                                  | Univariate analysis     |     | Multivariate analysis |                |      | Univariate analysis |     | Multivariate analysis |                |      | Univariate analysis |      | Multivariate analysis |                |     |
|                                                  | Kp +<br>(n, %)<br>n=262 | OR  | p                     | aOR<br>[95%CI] | p    | MDR-Kp<br>n=104     | OR  | p                     | aOR<br>[95%CI] | p    | Virulent Kp<br>n=34 | OR   | p                     | aOR<br>[95%CI] | p   |
| <b>Rural site</b>                                | 178(67.9)               | 1.1 | 0.8                   | 1.1 [0.7-1.7]  | 0.8  | 72(69.2)            | 1.1 | 0.7                   | 1.2 [0.7-2.2]  | 0.5  | 27(79.4)            | 1.97 | 0.1                   | 1.7 [0.7-4.1]  | 0.3 |
| <b>Education</b>                                 |                         |     |                       |                |      |                     |     |                       |                |      |                     |      |                       |                |     |
| Absence/primary school                           | 56(21.4)                | Ref |                       |                |      | 14(13.5)            | ref |                       |                |      | 6(17.7)             | ref  |                       |                |     |
| Partial secondary school                         | 152(58.0)               | 1.2 | 0.6                   |                |      | 73(70.2)            | 2.8 | 0.003                 |                |      | 20(58.8)            | 0.9  | 0.7                   |                |     |
| Complete secondary or higher                     | 54(20.6)                | 1.1 | 0.8                   |                |      | 17(16.4)            | 1.4 | 0.5                   |                |      | 8(23.5)             | 0.9  | 0.8                   |                |     |
| <b>Primigravidae</b>                             | 180(68.7)               | 0.8 | 0.4                   |                |      | 71(68.3)            | 1.0 | 0.9                   |                |      | 24(70.6)            | 1.1  | 0.8                   |                |     |
| <b>No. of people living in household (&gt;3)</b> | 90(34.4)                | 1.3 | 0.3                   |                |      | 90(34.4)            | 0.7 | 0.2                   |                |      | 22(64.7)            | 1.6  | 0.2                   |                |     |
| <b>Hand washing after toilets use</b>            | 172(76.9)               | 0.5 | 0.03                  | 0.5 [0.3-0.9]  | 0.02 | 65(70.7)            | 0.6 | 0.07                  | 0.4 [0.2-0.9]  | 0.02 | 23(69.7)            | 0.6  | 0.3                   |                |     |
| <b>Antibiotic consumption during pregnancy</b>   | 42(16.0)                | 1.6 | 0.1                   | 2.1 [1.1-4.1]  | 0.03 | 15(14.4)            | 0.8 | 0.6                   |                |      | 9(26.5)             | 2.1  | 0.1                   |                |     |
| <b>Toilets outside the house</b>                 | 254(97.0)               | 2.1 | 0.1                   |                |      | 100(96.2)           | 0.6 | 0.55                  |                |      | 33(97.1)            | 1.0  | 0.9                   |                |     |
| <b>Use of fecal material as fertilizer</b>       | 23(8.8)                 | 0.8 | 0.6                   |                |      | 7(6.7)              | 0.6 | 0.3                   |                |      | 2(5.9)              | 0.6  | 0.6                   |                |     |
| <b>Domestic animal contact</b>                   |                         |     |                       |                |      |                     |     |                       |                |      |                     |      |                       |                |     |
| Dog                                              | 84(32.1)                | 1.1 | 0.7                   |                |      | 29(27.9)            | 0.7 | 0.2                   |                |      | 10(29.4)            | 0.9  | 0.7                   |                |     |
| Cat                                              | 84(32.1)                | 0.7 | 0.1                   |                |      | 30(28.9)            | 0.8 | 0.4                   |                |      | 11(32.4)            | 1.0  | 0.9                   |                |     |
| Pet bird                                         | 40(15.3)                | 0.7 | 0.2                   |                |      | 11(10.6)            | 0.5 | 0.1                   |                |      | 5(14.7)             | 0.9  | 0.9                   |                |     |
| <b>Poultry contact</b>                           |                         |     |                       |                |      |                     |     |                       |                |      |                     |      |                       |                |     |
| Chicken                                          | 133(50.8)               | 1.1 | 0.6                   |                |      | 45(43.3)            | 0.6 | 0.05                  |                |      | 11(32.4)            | 0.4  | 0.05                  |                |     |

|                                       |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
|---------------------------------------|-----------|-----|-------------|---------------|-------|-----------------------|-----|-------|-------------|------|-----------------------|------|-------|-------------|-------|
| Turkey                                | 48(18.3)  | 0.7 | 0.1         |               |       | 16(15.4)              | 0.7 | 0.3   |             |      | 3(8.8)                | 0.4  | 0.1   |             |       |
| Duck                                  | 52(19.9)  | 0.6 | 0.05        |               |       | 18(17.3)              | 0.8 | 0.4   |             |      | 4(11.8)               | 0.5  | 0.2   |             |       |
| <b>Livestock animals</b>              |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| <b>contact</b>                        |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| Beef                                  | 66(25.2)  | 0.8 | 0.3         |               |       | 21(20.2)              | 0.6 | 0.1   |             |      | 9(26.5)               | 1.1  | 0.9   |             |       |
| Pig                                   | 85(32.4)  | 0.8 | 0.4         |               |       | 28(26.9)              | 0.7 | 0.1   |             |      | 10(29.4)              | 0.9  | 0.7   |             |       |
| Horse                                 | 62(23.66) | 0.8 | 0.3         |               |       | 19(18.3)              | 0.6 | 0.1   |             |      | 7(20.6)               | 0.8  | 0.7   |             |       |
| Animals excrement manipulation        | 35(13.4)  | 0.6 | 0.03        | 0.5           | 0.008 | 11(10.6)              | 0.7 | 0.3   |             |      | 3(8.8)                | 0.6  | 0.4   |             |       |
| [0.2-0.8]                             |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| <b>Dietary habit</b>                  |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| Eats chicken ≥1/week                  | 100(38.5) | 0.8 | 0.4         |               |       | 39(37.5)              | 0.9 | 0.8   |             |      | 17(50)                | 1.7  | 0.1   |             |       |
| Eats beef ≥1/week                     | 160(61.5) | 0.8 | 0.4         |               |       | 62(59.6)              | 0.9 | 0.6   |             |      | 18(52.9)              | 0.67 | 0.3   |             |       |
| Eats pork ≥1/week                     | 122(46.9) | 0.7 | 0.2         |               |       | 48(46.2)              | 0.9 | 0.8   |             |      | 15(44.1)              | 0.9  | 0.7   |             |       |
| Eats rabbit                           | 22(8.5)   | 0.8 | 0.6         |               |       | 11(10.6)              | 1.6 | 0.3   |             |      | 7(20.6)               | 3.6  | 0.01  | 3.6         | 0.01  |
| [1.3-9.8]                             |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| Eats fish ≥1/week                     | 154(59.2) | 0.9 | 0.5         |               |       | 58(55.8)              | 0.8 | 0.4   |             |      | 22(64.7)              | 1.3  | 0.5   |             |       |
| Eats insects                          | 27(10.4)  | 0.8 | 0.5         |               |       | 15(14.4)              | 2.0 | 0.1   |             |      | 7(20.6)               | 2.7  | 0.04  |             |       |
| Eats raw meat                         | 32(12.4)  | 0.8 | 0.5         |               |       | 10(9.7)               | 0.7 | 0.3   |             |      | 6(17.7)               | 1.6  | 0.3   |             |       |
| Eats raw fish                         | 15(5.8)   | 0.9 | 0.8         |               |       | 4(3.9)                | 0.5 | 0.3   |             |      | 3(8.8)                | 1.7  | 0.4   |             |       |
| Eats dry meat                         | 44(17.3)  | 0.7 | 0.3         |               |       | 12(11.9)              | 0.5 | 0.07  | 0.4         | 0.01 | 7(20.6)               | 1.3  | 0.6   |             |       |
| [0.2-0.8]                             |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| Eats dry fish ≥1/week                 | 71(28.6)  | 0.7 | 0.1         |               |       | 34(35.8)              | 1.7 | 0.05  |             |      | 11(32.3)              | 1.2  | 0.6   |             |       |
| Eats raw vegetables ≥1/week           | 219(84.6) | 0.8 | 0.4         |               |       | 91(88.4)              | 1.7 | 0.2   | 2.4         | 0.03 | 28(82.4)              | 0.8  | 0.7   |             |       |
| [1.1-5.5]                             |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| Hospitalization of a household member | 20(7.7)   | 0.8 | 0.6         |               |       | 10(9.6)               | 1.6 | 0.3   |             |      | 7(20.6)               | 4.2  | 0.005 | 4.0         | 0.009 |
| [1.4-11.2]                            |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |
| Cambodia Kp (N=146)                   |           |     |             |               |       | MDR-KP (N=97)         |     |       |             |      | Virulent Kp (N=97)    |      |       |             |       |
| Univariate analysis                   |           |     |             |               |       | Univariate analysis   |     |       |             |      | Univariate analysis   |      |       |             |       |
| Multivariate analysis                 |           |     |             |               |       | Multivariate analysis |     |       |             |      | Multivariate analysis |      |       |             |       |
| Kp + (n, %) n=97                      | OR        | p   | aOR [95%CI] | p             |       | MDR-Kp n=25           | OR  | p     | aOR [95%CI] | p    | Virulent Kp n=29      | OR   | p     | aOR [95%CI] | p     |
| Rural site                            | 57(58.8)  | 1.5 | 0.26        | 1.4 [0.7-2.8] | 0.3   | 8(32.0)               | 0.2 | 0.002 |             |      | 16(55.2)              | 0.8  | 0.6   |             |       |
| Education                             |           |     |             |               |       |                       |     |       |             |      |                       |      |       |             |       |

|                                                  |          |     |      |          |     |     |          |      |     |
|--------------------------------------------------|----------|-----|------|----------|-----|-----|----------|------|-----|
| Absence/primary school                           | 52(53.6) | Ref |      | 13(52.0) | ref |     | 16(55.2) | Ref  |     |
| Partial secondary school                         | 29(29.9) | 0.7 | 0.3  | 8(32.0)  | 1.1 | 0.8 | 9(31.0)  | 1.0  | 0.9 |
| Complete secondary or higher                     | 16(16.5) | 4.0 | 0.1  | 4(16.0)  | 1.0 | 1.0 | 4(13.8)  | 0.75 | 0.7 |
| <b>Primigravidae</b>                             | 68(70.1) | 1.5 | 0.3  | 19(76.0) | 1.5 | 0.5 | 19(65.5) | 0.7  | 0.5 |
| <b>No. of people living in household (&gt;3)</b> | 68(70.1) | 1.4 | 0.4  | 68(70.1) | 1.1 | 0.8 | 23(79.3) | 2.0  | 0.2 |
| <b>Hand washing after toilets use</b>            | 81(83.5) | 2.2 | 0.05 | 20(80.0) | 0.7 | 0.6 | 24(82.8) | 0.9  | 0.9 |
| <b>Antibiotic consumption during pregnancy</b>   | 17(17.5) | 0.7 | 0.5  | 4(16.0)  | 0.9 | 0.8 | 4(13.8)  | 0.7  | 0.5 |
| <b>Toilets outside the house</b>                 | 45(46.4) | 0.8 | 0.6  | 6(24.0)  | 0.3 | 0.1 | 14(48.3) | 1.1  | 0.8 |
| <b>Use of fecal material as fertilizer</b>       | 3(3.1)   | 0.4 | 0.2  | 1(4)     | 0.3 | 0.3 | 2(6.9)   | 0.6  | 0.5 |
| <b>Domestic animal contact</b>                   |          |     |      |          |     |     |          |      |     |
| Dog                                              | 13(13.4) | 3.6 | 0.1  | 2(8.0)   | 0.5 | 0.4 | 5(17.2)  | 1.6  | 0.5 |
| Cat                                              | 8(8.3)   | 1.4 | 0.7  | 2(8.0)   | 0.9 | 0.9 | 3(10.3)  | 1.5  | 0.6 |
| Pet bird                                         | 1(1.0)   | -   | -    | 0(0)     | -   | -   | 0(0)     | -    | -   |
| <b>Poultry contact</b>                           |          |     |      |          |     |     |          |      |     |
| Chicken                                          | 11(11.3) | 2.0 | 0.3  | 4(16.0)  | 1.8 | 0.4 | 1(3.5)   | 0.2  | 0.1 |
| Turkey                                           | 0(0)     | -   | -    | 0(0)     | -   | -   | 0(0)     | -    | -   |
| Duck                                             | 0(0)     | -   | -    | 0(0)     | -   | -   | 0(0)     | -    | -   |
| <b>Livestock animals contact</b>                 |          |     |      |          |     |     |          |      |     |
| Beef                                             | 4(4.1)   | 0.7 | 0.6  | 0(0)     | -   | -   | 0(0)     | -    | -   |
| Pig                                              | 0(0)     | -   | -    | 0(0)     | -   | -   | 0(0)     | -    | -   |
| Horse                                            | 0(0)     | -   | -    | 0(0)     | -   | -   | 0(0)     | -    | -   |
| <b>Animals excrement manipulation</b>            | 7(7.2)   | 0.9 | 0.8  | 0(0)     | -   | -   | 1(3.5)   | 0.4  | 0.4 |
| <b>Dietary habit</b>                             |          |     |      |          |     |     |          |      |     |
| Eats chicken $\geq 1$ /week                      | 24(24.7) | 0.9 | 0.8  | 7(28.0)  | 1.3 | 0.7 | 9(31.0)  | 1.6  | 0.4 |

|                                           |                         |     |       |                |      |                                           |         |        |                |     |                                           |     |      |                |     |
|-------------------------------------------|-------------------------|-----|-------|----------------|------|-------------------------------------------|---------|--------|----------------|-----|-------------------------------------------|-----|------|----------------|-----|
| Eats beef ≥1/week                         | 29(29.9)                | 0.7 | 0.3   |                |      | 9(36.0)                                   | 1.5     | 0.4    |                |     | 7(24.1)                                   | 0.7 | 0.4  |                |     |
| Eats pork ≥1/week                         | 76(78.4)                | 0.5 | 0.2   |                |      | 19(76.0)                                  | 0.8     | 0.7    |                |     | 21(72.4)                                  | 0.6 | 0.4  |                |     |
| Eats rabbit                               | 1(1.0)                  | 0.5 | 0.6   |                |      | 1(4.0)                                    | -       | -      |                |     | 1(3.5)                                    | -   | -    |                |     |
| Eats fish ≥1/week                         | 83(85.6)                | 2.0 | 0.1   |                |      | 20(80.0)                                  | 0.6     | 0.4    |                |     | 23(79.3)                                  | 0.5 | 0.3  |                |     |
| Eats insects                              | 59(60.8)                | 1.5 | 0.3   |                |      | 14(56.0)                                  | 0.8     | 0.6    |                |     | 16(55.2)                                  | 0.7 | 0.5  |                |     |
| Eats raw meat                             | 19(19.6)                | 1.1 | 0.9   |                |      | 4(16.0)                                   | 0.7     | 0.6    |                |     | 4(13.8)                                   | 0.6 | 0.4  |                |     |
| Eats raw fish                             | 7(6)                    | 1.8 | 0.5   |                |      | 2(8.0)                                    | 1.2     | 0.9    |                |     | 0(0)                                      | -   | -    |                |     |
| Eats dry meat                             | 56(57.7)                | 0.9 | 0.7   |                |      | 14(56.0)                                  | 0.9     | 0.8    |                |     | 18(62.1)                                  | 1.3 | 0.6  |                |     |
| Eats dry fish ≥1/week                     | 51(52.6)                | 2.1 | 0.04  | 2.0            | 0.05 | 12(48.0)                                  | 0.8     | 0.6    |                |     | 15(51.7)                                  | 0.9 | 0.9  |                |     |
|                                           |                         |     |       | [1.0-4.1]      |      |                                           |         |        |                |     |                                           |     |      |                |     |
| Eats raw vegetables<br>≥1/week            | 51(52.6)                | 1.4 | 0.4   |                |      | 13(52.0)                                  | 1.0     | 0.9    |                |     | 13(44.8)                                  | 0.6 | 0.3  |                |     |
| Hospitalization of a<br>household member  | 9(9.3)                  | 0.9 | 0.9   |                |      | 3(12.0)                                   | 1.5     | 0.6    |                |     | 2(6.9)                                    | 0.6 | 0.6  |                |     |
|                                           |                         |     |       |                |      |                                           |         |        |                |     |                                           |     |      |                |     |
| Senegal Kp (N=330)                        |                         |     |       |                |      | MDR-KP (N=137)                            |         |        |                |     | Virulent Kp (N=137)                       |     |      |                |     |
|                                           |                         |     |       |                |      |                                           |         |        |                |     |                                           |     |      |                |     |
|                                           |                         |     |       |                |      |                                           |         |        |                |     |                                           |     |      |                |     |
| Univariate analysis Multivariate analysis |                         |     |       |                |      | Univariate analysis Multivariate analysis |         |        |                |     | Univariate analysis Multivariate analysis |     |      |                |     |
|                                           | Kp +<br>(n, %)<br>n=137 | OR  | p     | aOR<br>[95%CI] | p    | MDR-Kp<br>n=16                            | OR      | p      | aOR<br>[95%CI] | p   | Virulent Kp<br>n=29                       | OR  | p    | aOR<br>[95%CI] | p   |
| Rural site                                | 104(75.9)               | 2.0 | 0.007 | 1.5 [0.9-2.6]  | 0.2  | 12(75.0)                                  | 1.9     | 0.9    | 1.8 [0.5-6.8]  | 0.4 | 24(82.8)                                  | 1.7 | 0.3  | 1.2 [0.4-3.5]  | 0.8 |
| Education                                 |                         |     |       |                |      |                                           |         |        |                |     |                                           |     |      |                |     |
| Absence/primary<br>school                 | 92(67.2)                | ref |       |                |      | 11(68.8)                                  | ref     |        |                |     | 17(58.6)                                  | ref |      |                |     |
| Partial secondary<br>school               | 31(22.6)                | 1.4 | 0.3   |                |      | 3(18.8)                                   | 0.8     | 0.7    |                |     | 9(31.0)                                   | 1.8 | 0.2  |                |     |
| Complete secondary<br>or higher           | 14(10.2)                | 1.0 | 1     |                |      | 2(12.5)                                   | 1.2     | 0.8    |                |     | 3(10.3)                                   | 1.2 | 0.8  |                |     |
| Primigravidae                             | 110(80.3)               | 1.2 | 0.6   |                |      | 13(81.3)                                  | 1.1     | 0.9    |                |     | 24(82.8)                                  | 1.2 | 0.7  |                |     |
| No. of people living<br>in household (>3) | 67(48.9)                | 1.1 | 0.5   |                |      | 67(48.9)                                  | 0.8     | 0.7    |                |     | 13(44.8)                                  | 0.8 | 0.6  |                |     |
| Hand washing after<br>toilets use         | 132(96.4)               | 1.3 | 0.6   |                |      | 16(100)                                   | -.<br>- | -<br>- |                |     | 26(89.7)                                  | 0.2 | 0.05 |                |     |

|                                         |           |     |       |           |       |  |          |     |      |            |      |  |          |     |      |            |      |  |
|-----------------------------------------|-----------|-----|-------|-----------|-------|--|----------|-----|------|------------|------|--|----------|-----|------|------------|------|--|
| Antibiotic consumption during pregnancy | 11(8.0)   | 0.7 | 0.4   |           |       |  | 1(6.3)   | 0.7 | 0.8  |            |      |  | 2(6.9)   | 0.8 | 0.8  |            |      |  |
| Toilets outside the house               | 129(94.2) | 1.2 | 0.7   |           |       |  | 14(87.5) | 0.4 | 0.2  |            |      |  | 29(100)  | -   | -    |            |      |  |
| Use of fecal material as fertilizer     | 19(13.9)  | 1.7 | 0.1   |           |       |  | 2(12.5)  | 0.9 | 0.9  |            |      |  | 8(27.6)  | 3.4 | 0.02 | 3.7        | 0.02 |  |
| Domestic animal contact                 |           |     |       |           |       |  |          |     |      |            |      |  |          |     |      | [1.7-11.0] |      |  |
| Dog                                     | 10(7.3)   | 1.2 | 0.7   |           |       |  | 2(12.5)  | 2.0 | 0.4  |            |      |  | 2(2.1)   | 0.9 | 0.9  |            |      |  |
| Cat                                     | 41(29.9)  | 1.0 | 0.9   |           |       |  | 1(6.3)   | 0.1 | 0.06 |            |      |  | 13(44.8) | 2.3 | 0.05 |            |      |  |
| Pet bird                                | 1(0.7)    | 1.4 | 0.8   |           |       |  | 0(0)     | -   | -    |            |      |  | 0(0)     | -   | -    |            |      |  |
| Poultry contact                         |           |     |       |           |       |  |          |     |      |            |      |  |          |     |      |            |      |  |
| Chicken                                 | 82(59.9)  | 2.2 | 0.001 | 1.9       | 0.008 |  | 5(31.3)  | 0.3 | 0.02 | 0.2        | 0.01 |  | 23(79.3) | 3.2 | 0.05 |            |      |  |
|                                         |           |     |       | [1.2-3.1] |       |  |          |     |      | [0.06-0.7] |      |  |          |     |      |            |      |  |
| Turkey                                  | 1(0.7)    | 0.3 | 0.2   |           |       |  | 0(0)     | -   | -    |            |      |  | 0(0)     | -   | -    |            |      |  |
| Duck                                    | 13(9.5)   | 1.2 | 0.7   |           |       |  | 1(6.3)   | 0.6 | 0.6  |            |      |  | 2(6.9)   | 0.7 | 0.6  |            |      |  |
| Livestock animals contact               |           |     |       |           |       |  |          |     |      |            |      |  |          |     |      |            |      |  |
| Beef                                    | 2(1.5)    | 2.8 | 0.4   |           |       |  | 1(6.3)   | 8.0 | 0.2  |            |      |  | 0(0)     | -   | -    |            |      |  |
| Pig                                     | 2(1.5)    | 1.4 | 0.7   |           |       |  | 0(0)     | -   | -    |            |      |  | 0(0)     | -   | -    |            |      |  |
| Horse                                   | 4(2.9)    | 5.8 | 0.1   |           |       |  | 1(6.3)   | 2.6 | 0.4  |            |      |  | 1(3.5)   | 1.3 | 0.8  |            |      |  |
| Animals excrement manipulation          | 49(35.8)  | 2.0 | 0.1   |           |       |  | 2(12.5)  | 0.2 | 0.06 |            |      |  | 13(44.8) | 1.6 | 0.3  |            |      |  |
| Dietary habit                           |           |     |       |           |       |  |          |     |      |            |      |  |          |     |      |            |      |  |
| Eats chicken ≥1/week                    | 55(40.2)  | 1.7 | 0.04  |           |       |  | 4(25.0)  | 0.5 | 0.2  |            |      |  | 11(37.9) | 0.9 | 0.8  |            |      |  |
| Eats beef ≥1/week                       | 44(32.1)  | 1.5 | 0.1   |           |       |  | 4(25.0)  | 0.7 | 0.5  |            |      |  | 6(20.7)  | 0.5 | 0.1  |            |      |  |
| Eats pork ≥1/week                       | 0(0)      | -   | -     |           |       |  | 0(0)     | -   | -    |            |      |  | 0(0)     | -   | -    |            |      |  |
| Eats rabbit                             | 7(5.1)    | 1.2 | 0.7   |           |       |  | 0(0)     | -   | -    |            |      |  | 3(10.3)  | 3.0 | 0.2  |            |      |  |
| Eats fish ≥1/week                       | 124(90.5) | 1.6 | 0.2   |           |       |  | 15(93.8) | 1.7 | 0.6  |            |      |  | 23(79.3) | 0.3 | 0.03 | 0.2        | 0.02 |  |
|                                         |           |     |       |           |       |  |          |     |      |            |      |  |          |     |      | [0.1-0.8]  |      |  |
| Eats insects                            | 3(2.2)    | 4.3 | 0.2   |           |       |  | 0(0)     | -   | -    |            |      |  | 1(3.5)   | 1.9 | 0.6  |            |      |  |
| Eats raw meat                           | 0(0)      | -   | -     |           |       |  | 0(0)     | -   | -    |            |      |  | 0(0)     | -   | -    |            |      |  |
| Eats raw fish                           | 1(0.7)    | -   | -     |           |       |  | 0(0)     | -   | -    |            |      |  | 0(0)     | -   | -    |            |      |  |
| Eats dry meat                           | 37(27.0)  | 1.0 | 0.9   |           |       |  | 4(25.0)  | 0.9 | 0.8  |            |      |  | 11(37.9) | 1.9 | 0.1  |            |      |  |
| Eats dry fish ≥1/week                   | 135(98.5) | 2.7 | 0.2   |           |       |  | 16(100)  | -   | -    |            |      |  | 29(100)  | -   | -    |            |      |  |
| Eats raw vegetables ≥1/week             | 104(75.9) | 1.0 | 0.9   |           |       |  | 13(81.3) | 1.4 | 0.6  |            |      |  | 23(79.3) | 1.3 | 0.6  |            |      |  |

|                                       |          |     |     |  |        |     |     |  |         |     |      |
|---------------------------------------|----------|-----|-----|--|--------|-----|-----|--|---------|-----|------|
| Hospitalization of a household member | 21(15.3) | 1.8 | 0.1 |  | 1(6.3) | 0.3 | 0.3 |  | 8(27.6) | 2.8 | 0.05 |
|---------------------------------------|----------|-----|-----|--|--------|-----|-----|--|---------|-----|------|

Figure S1: study sites




Figure S2. Flow diagram of inclusions






Figure S3. MLST sequence type (ST) diversity



☐ ST associated with multidrug resistance

ST associated with hypervirulence

Diversity of the beta-lactamases-encoding genes (*bla*<sub>SHV</sub>, *bla*<sub>OKP</sub>, *bla*<sub>LEN</sub>)Diversity of the beta-lactamases-encoding genes (*bla*<sub>SHV</sub>, *bla*<sub>OKP</sub>, *bla*<sub>LEN</sub>)

ND, not detected

**Figure S5.** Rate of virulence clusters or genes, by country of isolation





Figure S6. Details of presence per isolate of antimicrobial resistance phenotype, and of genes coding for antimicrobial resistance, virulence or plasmid replicons and metal tolerance (MT).



Isolates are ordered vertically according to the same phylogeny (left) as in the main phylogeny Figure. Note that cluster kfuABC was conserved in non-Kp1 isolates but not within Kp1.