



Highly Resistant Cholera Outbreak Strain in Zimbabwe

Tapfumanei Mashe, Daryl Domman, Andrew Tarupiwa, Portia Manangazira, Isaac Phiri, Masunda Kudzai, Prosper Chonzi, Elisabeth Njamkepo, Masindi Ramudzulu, Sekesai Mtapuri-Zinyowera, et al.

► To cite this version:

Tapfumanei Mashe, Daryl Domman, Andrew Tarupiwa, Portia Manangazira, Isaac Phiri, et al.. Highly Resistant Cholera Outbreak Strain in Zimbabwe. *New England Journal of Medicine*, 2020, 383 (7), pp.687-689. 10.1056/NEJMc2004773 . pasteur-02928196

HAL Id: pasteur-02928196

<https://pasteur.hal.science/pasteur-02928196>

Submitted on 2 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Highly drug-resistant cholera outbreak strain in Zimbabwe, 2018 - 2019

Mashe Tapfumanei*, MSc

Ministry of Health and Child Care, Zimbabwe

Daryl Domman*, Ph.D

University of New Mexico Health Sciences Center, Albuquerque, New Mexico, USA

Tarupiwa Andrew, MSc

Ministry of Health and Child Care, Zimbabwe

Manangazira Portia, M.D., M.P.H

Phiri Isaac, M.D., M.P.H

Ministry of Health and Child Care, Zimbabwe

Masunda Kudzai, M.D., M.P.H

Chonzi Prosper, M.D., M.P.H

Beatrice Road Infectious Diseases Hospital, Harare, Zimbabwe

Elisabeth Njamkepo, Ph.D

Institut Pasteur, Paris, France

Masindi Ramudzulu, B.Sc (Hons)

National Institute for Communicable Diseases, South Africa

Mtapuri-Zinyowera Sekesai[‡], Ph.D

Ministry of Health and Child Care, Zimbabwe

Anthony Marius Smith[‡], Ph.D,

National Institute for Communicable Diseases, South Africa

François-Xavier Weill[‡], M.D., Ph.D

Institut Pasteur, Paris, France

Corresponding author:

F-X.W., Unité des Bactéries Pathogènes Entériques, Institut Pasteur, 28 rue du Dr Roux, 75015 Paris, France, francois-xavier.weill@pasteur.fr, Tel +33 1 45688345

*These authors contributed equally; [‡]these authors jointly supervised this work

Key words: Cholera, antimicrobial resistance, whole genome sequencing, Zimbabwe

Word count: 507.

45 TO THE EDITOR: From 6 September 2018 to 26 March 2019, Zimbabwe experienced a
46 large cholera outbreak with 10,730 suspected cholera cases and 68 deaths (Fig. S1).
47 Antimicrobial susceptibility data available on 65% (241/371) of the confirmed cases
48 showed that the *Vibrio cholerae* O1 serotype Ogawa isolates were multidrug resistant,
49 with unexpectedly high-level resistance to ciprofloxacin (96.7%, 233/241) and
50 ceftriaxone (99.6%, 240/241). Details of the methods are provided in the Supplementary
51 Appendix, available with the full text of this letter at NEJM.org. We sequenced the
52 whole-genomes of 11 *V. cholerae* O1 isolates – including ten collected during the
53 cholera outbreak in Zimbabwe, and one originating from a South African patient with
54 travel history to Zimbabwe (Fig. S1, Table S1) – to investigate the determinants of
55 antimicrobial resistance as well as their phylogenetic relationships to >1,200 global
56 seventh cholera pandemic *V. cholerae* El Tor (7PET) genomes.^{1,2} Phylogenomic
57 analyses show that the 11 studied genomes belonged to sublineage T13 of the 7PET
58 lineage (Figs 1A and 1B, Table S2). Sublineage T13 was recently introduced from
59 South Asia into East Africa and from there to Yemen (Figs 1B and 1C).² The T13
60 isolates were previously shown to have a narrow antimicrobial resistance (AMR)
61 pattern, mostly due to a ~10-kb deletion in the SXT/R391 genomic island (called
62 ICE *Vchl*Ind5), resulting in the loss of four AMR genes.² However, the 2018 Zimbabwean
63 isolates differed from previous T13 isolates by having 14 additional antimicrobial
64 resistance (AMR) genes carried on a ~160 kb IncA/C2 plasmid, leading to a broader
65 resistance profile (Fig. S2, Tables S3-S4). In particular, the isolates were intermediately
66 resistant or resistant to tetracycline (presence of the *tet(A)* gene), intermediately
67 resistant or resistant to ciprofloxacin (mutations of the *gyrA* and *parC* genes and
68 presence of the *aac6'-Ib-cr* gene) and produced a CTX-M-15 extended spectrum beta-


lactamase. Antibiotics such as tetracyclines, macrolides, and fluoroquinolones are commonly used in the treatment of moderate-to-severe cholera cases – as an adjunct to rehydration therapy – to shorten the duration and volume of diarrhea, thereby limiting bacterial transmission.³ However, beta-lactams are not. Resistance to extended-spectrum cephalosporins (ESCs) is uncommon in 7PET isolates, and only ESC-resistant (ESC^R) 7PET isolates from sporadic or small outbreak cases were observed in Africa before 2018.¹ It is noteworthy that a large typhoid outbreak (> 3000 suspected cases) caused by a ciprofloxacin-resistant strain – requiring the use of ESCs – occurred in Harare between October 2017 – February 2018.^{4,5} This might have contributed to the ESC^R patterns seen among the cholera outbreak strain several months later.

The 2018 Zimbabwean isolates were susceptible to azithromycin, which was consequently used to treat severe cholera cases during the outbreak. However, azithromycin resistance genes, such as *mph(A)* or *mph(E)* were sporadically identified in CTX-M-15-producing 7PET isolates collected in Zimbabwe (Table S4), the Democratic Republic of the Congo¹, and Kenya² from 2010 to 2015. The acquisition of an IncA/C2 plasmid carrying both *mph* and *tet* genes by this T13 7PET strain would jeopardize the oral antibiotic treatment of cholera.

We emphasize the need for cross-border collaboration and continued laboratory surveillance to stemming this highly-drug resistant T13 cholera strain in Africa.

References

1. Weill FX, Domman D, Njamkepo E, et al. Genomic history of the seventh pandemic of cholera in Africa. *Science* 2017;358:785-9.
2. Weill FX, Domman D, Njamkepo E, et al. Genomic insights into the 2016–2017 cholera epidemic in Yemen. *Nature* 2019;565:230-3.
3. Leibovici-Weissman Y, Neuberger A, Bitterman R, et al. Antimicrobial drugs for treating cholera. *Cochrane Database Syst Rev* 2014;19;CD008625.
4. N'cho HS, Masunda KPE, Mukeredzi I, et al. Notes from the field: typhoid fever outbreak—Harare, Zimbabwe, October 2017–February 2018. *MMWR Morb Mortal Wkly Rep* 2019;68:44-5.
5. Mashe T, Gudza-Mugabe M, Tarupiwa A, et al. Laboratory characterisation of *Salmonella enterica* serotype Typhi isolates from Zimbabwe, 2009-2017. *BMC Infect Dis* 2019;19:487.



Legend to Figure

Phylogenetic relatedness of the *V. cholerae* O1 El Tor isolates from the 2018 outbreak in Zimbabwe. **a**, Maximum likelihood phylogeny of 1,213 genomic sequences. A6 was used as an outgroup. The scale bar denotes substitutions per variable site (SNVs). Branches are colored according to geographic location, inferred by stochastic mapping of the geographic origin of each isolate onto the tree. The inferred introduction events into Africa are indicated by the letter 'T'. One branch was artificially shortened and as denoted by the hash mark. **b**, Maximum clade credibility tree produced with BEAST for a subset of 92 representative isolates of the distal part of genomic wave 3 (i.e., those with the *ctxB7* allele). Geographic location of the isolates is indicated in the same colors as in panel **a**. Selected nodes supported by bootstrap values ≥ 0.95 are shown. Acquisition of the multidrug-resistant plasmid is shown **c**, The geographical distribution

121 of selected 7PET sublineages T10 - T13. The date ranges shown for introductions are
122 the 95% credible interval estimate of the MRCA in years. Dashed lines do not represent
123 precise routes of transmission.

124

125

126