



Celebrating the career and legacy of Professor Pascale Cossart

Carmen Buchrieser

► To cite this version:

Carmen Buchrieser. Celebrating the career and legacy of Professor Pascale Cossart. *Molecular Microbiology*, 2020, 113 (3), pp.535-537. 10.1111/mmi.14478 . pasteur-03153599

HAL Id: pasteur-03153599

<https://pasteur.hal.science/pasteur-03153599>

Submitted on 26 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

**EDITORIAL: CELEBRATING THE CAREER AND LEGACY OF PROFESSOR
PASCALE COSSART**

Carmen Buchrieser

*Institut Pasteur, Biology of Intracellular Bacteria and CNRS UMR 3525, 28, Rue du Docteur
Roux, 75724 Paris, France*

It is an enormous pleasure and honour for me to contribute this editorial to introduce the special issue that we have assembled to honour the numerous, unique and outstanding scientific achievements of Pascale Cossart and her group. Pascale Cossart, a distinguished Professor at the Institut Pasteur in Paris, France, has been a worldwide leader in the study of fundamental processes in molecular and cellular microbiology and host pathogen interactions for more than 30 years. Using mainly *Listeria monocytogenes*, but also *Rickettsia conorii*, as models she has made several seminal findings in the field of infection biology, cell biology and epigenetics. Her work on fundamental RNA biology and bacterial regulation, and more recently the discovery of a new antibiotic resistance mechanism that is called ribosome splitting, are examples of her elegant contributions to the field.

In this Special issue of *Molecular Microbiology*, colleagues, former students, postdoctoral fellows and friends of Pascale Cossart have gathered together to contribute reviews and articles in the area of her major research interests. The special issue starts with a personal tribute to Pascale Cossart, contributed by *Werner Goebel*, a longstanding colleague and competitor who traces many of the ground-breaking findings made by Pascale Cossart and her group.

Listeria monocytogenes is reported in all textbooks as an intracellular pathogen that replicates in the cytosol of human cells after its escape from the vacuole. However, recently this paradigm was challenged by observations that *L. monocytogenes* can also adopt a vacuolar lifestyle when persisting within cells (Kortebi *et al.*, 2017). As another example, *Mycobacterium tuberculosis*, which was thought for a long time to be a strictly intra-vacuolar pathogen, was recently shown to rupture the phagosome and to contact the cytosol of the phagocytes it infects (Russell, 2016, Simeone *et al.*, 2009). This discovery was very important to better understand former unsolved steps of the disease process, as it is associated with altered intracellular signalling. In particular, this knowledge is important for the development

of vaccines, a priority for tuberculosis research (Groschel *et al.*, 2017). Thus a strict distinction between vacuolar and cytosolic intracellular pathogens is not evident. Similarly, the strict distinction between extracellular and intracellular pathogens does not hold true anymore. In a very interesting opinion article *Arturo Casadevall and Ferric Fang* show that new findings blur the boundaries between extracellular and intracellular pathogens and they discuss the benefits and caveats arising from the intracellular pathogen concept.

Today, multidrug resistant bacteria are a worldwide public health threat with increasing mortality and tremendous costs. Thus multidisciplinary research efforts are of utmost importance to better understand resistance emergence, persistence, dissemination and to find new, effective drugs. In an illustration of how the knowledge of ancient civilisations might help to solve this modern problem, *Julian Davis* and *Shekooh Behroozian* provide insights on how to use native environmental resources for extraction of natural products, so called medicinals. In particular, they present a personal view on the antimicrobial properties of clay minerals and propose that these underexplored natural resources may provide an alternative means to fight pathogens resistant to commonly used drugs.

Another fascinating approach to fight antibiotic resistance is proposed by *Jörg Vogel* who provides new ideas on how species-specific programmable RNA antibiotics could be developed. Indeed, broad-spectrum antibiotics are killing many bacteria present in and on our body, including both beneficial as well as pathogenic ones. Thus, the possibility of selective killing of pathogenic and/or multidrug resistant bacteria is very appealing, and might lead to microbiome editing for health. While proposing different ways to use short antisense oligonucleotides as antimicrobials, he also highlights the caveats including specificity, delivery, resistance and persistence.

Antibiotics are mainly seen as drugs to treat bacterial infections by inhibiting bacterial growth or killing them. However, as most antibiotics are naturally produced by soil organisms, their primary ecological function in the environment may be different. *Pishchany Gleb* and *Roberto Kolter* consider possible ecological roles of antibiotics and discuss their alternative functions at sub-inhibitory concentrations, such as the modulation of gene expression or biofilm formation, their influence on the stability of RNA molecules, and effects on bacterial evolution by increasing mutagenesis rates and horizontal gene transfer through conjugation. They argue that it is important to start studying the importance of these molecules in the

context of multispecies communities and in environmental conditions to understand how antibiotics might affect microbial communities and to expand our knowledge on these other, understudied effects of antibiotics.

Fernando Baquero, Val Fernandez-Lanza, Melodie Duval and Teresa Coque consider another idea that puts the environment into the focus of antibiotic resistance, namely the impact of gene – environment interactions on the acquisition of antibiotic resistance determinants, the ecogenetics of antibiotic resistance. As a specific example of how the ecology of an organism may shape the acquisition of antibiotic resistance, they examine *L. monocytogenes*, as it is intriguing that these bacteria have retained susceptibility to the key antimicrobials used to treat infections. This is even more surprising as *L. monocytogenes* is found ubiquitous in the environment and in a large spectrum of animals and plants. The authors discuss the impacts of global ecology, population size, genome structure, environmental stresses, and suggest that the many biocide and heavy metal resistance genes that are present in the *L. monocytogenes* genome may also play a role.

Cameron Parson, Sangmi Lee and Sophia Kathariou discuss these heavy metal resistance genes in *L. monocytogenes* and other Gram-positive bacteria further. They give a comprehensive update on the plasmid- and chromosomal-borne cadmium and arsenic resistance genes, and discuss their importance for environmental persistence. They suggest that the food-processing environment exerts a major selection pressure for cadmium resistance in *L. monocytogenes*, however many natural *Listeria* isolates are also highly resistant to several toxic metals. Thus, further studies will help to understand the ecological and evolutionary factors that drive acquisition, spread and diversification of the genomic elements conferring heavy metal resistance.

Two important new concepts in RNA biology have been discovered by the group of Pascale Cossart that are highlighted in the two following articles. *Pierre Mandin and Jörgen Johansson* give a short overview on how bacteria sense temperature changes and the impact this might have on virulence. For many years it was known that bacteria can sense temperature changes, and that in pathogenic bacteria virulence gene expression is often correlated with the transition from the environment (low temperature) to the human host (37°C). An example of temperature-regulated virulence gene expression the PrfA thermosensor of *L. monocytogenes* is described and compared to other bacterial

thermosensors. *Alejandro Toledo-Arana* and *Iñigo Lasa* discuss additional regulatory concepts by giving a masterly account of the advances made and new insights gained through the implementation of high-throughput methods for RNA profiling. Through genome-wide transcriptional profiling massive amount of antisense transcription, as well as conditional transcriptional termination and non-contiguous operons, have been discovered. Furthermore, the authors highlight riboswitch-dependent regulation of antisense RNA and overlapping transcription between neighbouring genes and present the related excludon concept, a new paradigm of regulation based on overlapping transcription.

Another rapidly expanding area of gene regulation studies was initiated by the discovery and characterization of small noncoding regulatory RNAs. Many of these control bacterial pathogenicity and the adaptation of bacteria to environmental stresses and to their hosts (Oliva *et al.*, 2015). *Jens Georg, David Lalaouna, Shengwei Hou, Steffen Lott, Isabelle Caldelari, Stefano Marzi, Wolfgang Hess* and *Pascale Romby* present a comprehensive and exciting overview on how targets of small RNAs can be identified, a still very challenging task. They show how by combining powerful computational approaches with experimental approaches the full extent of small RNA-dependent regulatory networks can be elucidated.

Peptidoglycan is an essential molecule for bacteria that maintains their shape, cell integrity and provides a protective function against changing environmental conditions. Furthermore, peptidoglycan can be altered within the host to impair responses triggered by pattern recognition receptors (Pucciarelli & Garcia-del Portillo, 2018). Peptidoglycan is also studied as a target to control bacterial infections. However, these studies are rarely undertaken in natural environments but under laboratory conditions. Until now our knowledge on peptidoglycan stems mainly from few model bacteria, and intracellular bacteria have only been studied very recently. *Francisco Garcia-del-Portillo* provides an illuminating account of recent advances on probing peptidoglycan synthesis and assembly within eukaryotic host cells, with a particular focus on the enzymes involved in peptidoglycan metabolism in the context of symbiotic growth and persistent infection. He gives a detailed description of the enzymatic machinery identified mainly in *Escherichia coli* and of peptidoglycan recycling utilized by bacterial pathogens to evade immune defences. Furthermore, the current techniques that are used for analysis and isolation of cell wall muropeptides are described and examples of intracellular bacteria that have been analysed with these methods are summarised.

139

140 Peptidoglycan is also one of the three major constituents of the *L. monocytogenes* cell wall,
141 along with anionic teichoic acid polymers and wall-associated and wall-anchored proteins.
142 Polymeric teichoic acid is divided into wall teichoic-acid, which is directly conjugated to the
143 peptidoglycan, and lipoteichoic acid, which is tethered to the plasma membrane. The *Listeria*
144 teichoic acid polymers have important functions such as regulating bacterial growth, ion-
145 homeostasis, biofilm formation, interaction with bacteriophages, host cell invasion and
146 virulence. The *L. monocytogenes* serotyping scheme classifies strains into different serotypes.
147 It recognizes structural and compositional changes on the cell wall surface and led to the
148 discovery of strains belonging to specific serovars, which cause most human infections. In
149 their MicroReview *Eric Sumrall, Anja Keller, Yang Shen and Martin Loessner* provide a
150 comprehensive update on the synthesis, structural variability and function of teichoic acids in
151 the genus *Listeria*. They discuss the impacts of teichoic acid decorations on serotype-specific
152 designations, on phage resistance and virulence *via* wall-associated virulence proteins. They
153 propose that the chemical synthesis of teichoic acids could be used for vaccine development.
154 These engineered, teichoic acid-based vaccines could fully exploit their immunogenic
155 potential and might represent a new strategy to fight infections with Gram-positive bacteria.

156

157 In the following article, *Eric Sumrall, Christopher Schefer, Jeanine Rismonod, Angelika*
158 *Gründling, Martin Loessner and Yang Shen* present new research results on the function of
159 teichoic acids in virulence of *L. monocytogenes*. They show that two different
160 glycosyltransferases are required for the galactosylation of lipoteichoic acid (LTA) and wall
161 teichoic acid (WTA), respectively. Thus *L. monocytogenes* encodes two surface-acting
162 galactosyltransferases with distinct substrate specificities, which have an impact on the cell
163 surface modification and InlB (an important virulence factor for invasion of eukaryotic cells)
164 retention and function. The demonstration that modification of WTA alone is critical for InlB
165 localization is interesting in that LTA and WTA are often referred to in the literature with
166 minimal functional distinction. This study clearly indicates that these surface moieties have
167 important functional differences, and leads one to wonder what other functions might be
168 solely attributed to WTA *versus* LTA, or *vice versa*. Another important surface structure is
169 pili. They have a pivotal role in the colonization of specific host tissues in many pathogenic
170 bacteria. *Frederico Iovino, Priyanka Nannapaneni, Brigitta Henriques-Normark and Staffan*
171 *Normark* provide a state-of-the art overview of the recent findings of type 1 pilus and its
172 adhesive tip protein RrgA from *Streptococcus pneumoniae*. The type 1 pilus is an important

yet less studied surface associated virulence determinant of pneumococcal surface proteins, and thus the presentation of biochemical, structural and host pathogen interaction studies showing the impact of type 1 pilus in colonization and pathogenicity are very timely. Pneumococcal type 1 pili seem to confer a competitive advantage for nasopharyngeal colonization of humans, and the RrgA tip protein might be a good addition to the group of proteins used for the pneumococcal vaccine.

How did these different bacterial cell envelopes evolve? In particular, how did monoderm and diderm bacteria evolve? In the penultimate article of this issue *Daniela Megrian, Najwa Taib, Jerzy Witwinowski, Christophe Beloin and Simonetta Gribaldo* address the important question of cell envelope diversity across the bacterial domain. Based on the existence of diderm bacteria among the generally monoderm Firmicutes, the division of Gram-positive (monoderm) and Gram-negative (diderm bacteria) is discussed. Using an in depth and comprehensive phylogenomic analyses, together with results from a new experimental model, *Veillonella parvula*, a diderm Firmicute, the authors propose that the diderm envelope is ancestral within the Firmicutes and the monoderm envelope evolved multiple times through loss of the outer membrane. This provocative hypothesis will stimulate discussions and further studies to understand bacterial cell envelope evolution. Finally, *Megan de Ste Croix, Jonathan Holmes, Joseph Wanford, Richard Moxon, Marco Oggioni, and Christopher Bayliss* discuss the role of bottlenecks in the evolution of diversity-generating mechanisms of microbes during bacterial transmission and infection. They provide an authoritative summary of progress in the field and challenges for future studies.

This collection of articles reflects many of the areas of research to which Pascale Cossart has contributed, highlighting and building upon her outstanding findings over the last decades. We are confident that these contributions and ideas will inspire young researchers to embrace the upcoming challenges of the New Microbiology (Cossart, 2018).

References

- Cossart, P., (2018) *The New Microbiology: From Microbiomes to CRISPR*. American Society of Microbiology.
- Groschel, M.I., F. Sayes, S.J. Shin, W. Frigui, A. Pawlik, M. Orgeur, R. Canetti, N. Honore, R. Simeone, T.S. van der Werf, W. Bitter, S.N. Cho, L. Majlessi & R. Brosch, (2017) Recombinant BCG Expressing ESX-1 of *Mycobacterium marinum* Combines Low Virulence with Cytosolic Immune Signaling and Improved TB Protection. *Cell Rep* **18**: 2752-2765.

- 209 Kortebe, M., E. Milohanic, G. Mitchell, C. Pechoux, M.C. Prevost, P. Cossart & H. Bierne,
210 (2017) *Listeria monocytogenes* switches from dissemination to persistence by
211 adopting a vacuolar lifestyle in epithelial cells. *PLoS Pathog* **13**: e1006734.
- 212 Oliva, G., T. Sahr & C. Buchrieser, (2015) Small RNAs, 5' UTR elements and RNA-binding
213 proteins in intracellular bacteria: impact on metabolism and virulence. *FEMS*
214 *Microbiol Rev* **39**: 331-349.
- 215 Pucciarelli, M.G. & F. Garcia-del Portillo, (2018) Within-Host Envelope Remodelling and its
216 Impact in Bacterial Pathogen Recognition. *Curr Issues Mol Biol* **25**: 43-60.
- 217 Russell, D.G., (2016) The ins and outs of the *Mycobacterium tuberculosis*-containing
218 vacuole. *Cell Microbiol* **18**: 1065-1069.
- 219 Simeone, R., D. Bottai & R. Brosch, (2009) ESX/type VII secretion systems and their role in
220 host-pathogen interaction. *Curr Opin Microbiol* **12**: 4-10.
- 221