



HAL
open science

Closed and high quality bacterial genomes of the Oligo Mouse Microbiota community

Quentin Lamy-Besnier, Romain Koszul, Laurent Debarbieux, Martial Marbouty

► To cite this version:

Quentin Lamy-Besnier, Romain Koszul, Laurent Debarbieux, Martial Marbouty. Closed and high quality bacterial genomes of the Oligo Mouse Microbiota community. *Microbiology Resource Announcements*, 2021, 10 (17), pp.e01396-20. 10.1128/MRA.01396-20 . pasteur-03204821

HAL Id: pasteur-03204821

<https://pasteur.hal.science/pasteur-03204821>

Submitted on 21 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.





L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Closed and High-Quality Bacterial Genome Sequences of the Oligo-Mouse-Microbiota Community

 Quentin Lamy-Besnier,^{a,b,c}  Romain Koszul,^b  Laurent Debarbieux,^a  Martial Marbouty^b

^aBacteriophage, Bacterium, Host Laboratory, Department of Microbiology, Institut Pasteur, Paris, France

^bUnité Régulation Spatiale des Génomes, Institut Pasteur, CNRS, UMR 3525, Paris, France

^cUniversité Paris Descartes, Paris, France

ABSTRACT The Oligo-Mouse-Microbiota (OMM¹²) gnotobiotic murine model is an increasingly popular model in microbiota studies. However, following Illumina and PacBio sequencing, the genomes of the 12 strains could not be closed. Here, we used genomic chromosome conformation capture (Hi-C) data to reorganize, close, and improve the quality of these 12 genomes.

The Oligo-Mouse-Microbiota (OMM¹²) is a murine bacterial synthetic community (12 strains) introduced in axenic mice. The resulting gnotobiotic murine model has been increasingly used for diverse gut microbiota studies across several animal facilities (1–4). This bacterial consortium comprises members from the 5 major phyla naturally present in mice microbiota and reconstitutes its main functionalities, such as metabolism and colonization resistance against pathogens (5).

Despite two rounds of sequencing using PacBio and Illumina technologies (6, 7), one-half of the genomes currently accessible remain made of 2 to 20 contigs (Table 1). In particular, the genome of *Bacteroides caecimuris*, the most abundant bacterium of the OMM¹² consortium, consists of 19 contigs (5). Improving scaffolding would facilitate a number of downstream analyses, such as prophage prediction.

We used genomic chromosome conformation capture (Hi-C), a technique that uses chemical fixation (formaldehyde) to *in vivo* cross-link nearby DNA sequences, digestion with restriction enzymes, DNA extremities refilling with biotinylated nucleotides, and finally, proximity ligation, DNA extraction, and deep sequencing. The relative ligation frequencies between nonadjacent DNA segments reflect the average three-dimensional (3D) organization of the genome of interest (8) and also, because of the polymer nature of DNA, the relative distance separating them along the chromosome. The latter property has therefore been exploited to bridge scaffolding gaps resulting, for instance, from the presence of repeated elements (9). Hi-C scaffolding is now commonly used in sequencing projects of large eukaryotic genomes but applies to bacteria as well (10, 11).

A Hi-C metagenomic protocol for mammalian gut samples developed in our laboratory (12) was applied to fecal pellets from the OMM¹² mice bred at Institut Pasteur (protocol 20.173 approved by the veterinary staff and under authorization APAFIS 26874 by the national ethics committee). Libraries were prepared using streptavidin beads to capture biotinylated ligation junction as described previously (13) and sequenced with an Illumina NextSeq 550 system to generate a total of 101,182,905 reads of 35 bp. The quality of the reads was first verified with FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc>). Then, for each bacterium, the python library hicstuff (<https://github.com/koszullab/hicstuff>) was used to generate Hi-C contact maps using the most recent OMM¹² genomes (6). Contigs were reorganized (i.e., scaffolded) based on their relative contact frequencies (14). When necessary, preexisting contigs were split and rearranged to better fit the expected 3D structure, based on the contact data. In order for users to detect junctions

Citation Lamy-Besnier Q, Koszul R, Debarbieux L, Marbouty M. 2021. Closed and high-quality bacterial genome sequences of the Oligo-Mouse-Microbiota community. *Microbiol Resour Announc* 10:e01396-20. <https://doi.org/10.1128/MRA.01396-20>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2021 Lamy-Besnier et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Laurent Debarbieux, laurent.debarbieux@pasteur.fr, or Martial Marbouty, martial.marbouty@pasteur.fr.

Received 22 January 2021

Accepted 6 April 2021

Published 29 April 2021

TABLE 1 Description and accession numbers of the OMM¹² genomes^a

Strain	Genome size (bp)	Previous no. of contigs ^b	No. of genes	DSM no.	Accession no.	Reads (%) ^c
<i>Akkermansia muciniphila</i> YL44	2,737,357	1	2,717	26127	CP065322	2.8
<i>Acutalibacter muris</i> KB18	3,802,913	1	4,271	26090	CP065321	0.002
<i>Bifidobacterium animalis</i> YL2	2,021,936	2	1,740	26074	CP065311	0.0007
<i>Bacteroides caecimuris</i> I48	4,800,606	19	4,322	26085	CP065319	55.8
<i>Blautia coccoides</i> YL58	5,128,582	1	5,351	26115	CP065312	20.5
<i>Enterocloster clostridioformis</i> YL32	7,157,610	16	7,955	26114	CP065314	4.9
<i>Clostridium innocuum</i> I46	4,452,146	1	4,966	26113	CP065320	5.6
<i>Enterococcus faecalis</i> KB1	3,025,655	1	2,919	32036	CP065317	0.1
<i>Flavonifractor plautii</i> YL31	3,813,715	5	4,049	26117	CP065315	0.38
<i>Limosilactobacillus reuteri</i> I49	2,063,624	3	1,958	32035	CP065318	0.99
<i>Muribaculum intestinale</i> YL27	3,307,069	1	2,957	28989	CP065316	4.5
<i>Turicimonas muris</i> YL45	2,887,949	20	2,753	26109	CP065313	0.089

^aRaw data can be accessed at [SRX9907181](https://www.ncbi.nlm.nih.gov/sra/SRX9907181).

^bGarzetti et al. (6).

^cUnmapped reads = 4.34 %.

between contigs, contigs were separated by $10 \times N$ in the final fasta file. We successfully closed the genomes of the 12 strains except for *Flavonifractor plautii*, where the position of a 70,760-bp region could not be assigned into the main scaffold. The genome of *F. plautii* therefore remains split in 2 scaffolds. An automatic annotation was then performed using RAST (15). The access to a single scaffold of high quality will greatly improve the accuracy of genomic analyses performed by the community of users of the OMM¹² gnotobiotic mice.

Data availability. The 12 reassembled and closed genome sequences of the OMM¹² bacteria as well as the FastQ reads have been deposited under the accession numbers provided in Table 1 under the BioProject number [PRJNA680355](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA680355).

ACKNOWLEDGMENTS

We thank Cyril Matthey Doret and Agnès Thierry for assistance during computational and experimental work, respectively.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication. This research was supported by funding to L.D. and M.M. from ANR-20-CE92-0048 and to R.K. from the European Research Council under the Horizon 2020 Program (ERC grant agreement 771813) and from JPI-EC-AMR STARCS ANR-16-JPEC-0003-05. Q.L.-B. is funded by École Doctorale FIRE-Program Bettencourt.

REFERENCES

- Bolsega S, Basic M, Smoczek A, Buettner M, Eberl C, Ahrens D, Odum KA, Stecher B, Bleich A. 2019. Composition of the intestinal microbiota determines the outcome of virus-triggered colitis in mice. *Front Immunol* 10:1708. <https://doi.org/10.3389/fimmu.2019.01708>.
- Eberl C, Ring D, Münch PC, Beutler M, Basic M, Slack EC, Schwarzer M, Srutkova D, Lange A, Frick JS, Bleich A, Stecher B. 2019. Reproducible colonization of germ-free mice with the Oligo-Mouse-Microbiota in different animal facilities. *Front Microbiol* 10:2999. <https://doi.org/10.3389/fmicb.2019.02999>.
- Lourenço M, Chaffringeon L, Lamy-Besnier Q, Pédrón T, Campagne P, Eberl C, Bérard M, Stecher B, Debarbieux L, De Sordi L. 2020. The spatial heterogeneity of the gut limits predation and fosters coexistence of bacteria and bacteriophages. *Cell Host Microbe* 28:390–401.e395. <https://doi.org/10.1016/j.chom.2020.06.002>.
- Studer N, Desharnais L, Beutler M, Brugiroux S, Terrazos MA, Menin L, Schürch CM, McCoy KD, Kuehne SA, Minton NP, Stecher B, Bernier-Latmani R, Hapfelmeier S. 2016. Functional intestinal bile acid 7 α -dehydroxylation by *Clostridium scindens* associated with protection from *Clostridium difficile* infection in a gnotobiotic mouse model. *Front Cell Infect Microbiol* 6:191. <https://doi.org/10.3389/fcimb.2016.00191>.
- Brugiroux S, Beutler M, Pfann C, Garzetti D, Ruscheweyh HJ, Ring D, Diehl M, Herp S, Löttscher Y, Hussain S, Bunk B, Pukall R, Huson DH, Münch PC, McHardy AC, McCoy KD, Macpherson AJ, Loy A, Clavel T, Berry D, Stecher B. 2016. Genome-guided design of a defined mouse microbiota that confers colonization resistance against *Salmonella enterica* serovar Typhimurium. *Nat Microbiol* 2:16215. <https://doi.org/10.1038/nmicrobiol.2016.215>.
- Garzetti D, Brugiroux S, Bunk B, Pukall R, McCoy KD, Macpherson AJ, Stecher B. 2017. High-quality whole-genome sequences of the Oligo-Mouse-Microbiota bacterial community. *Genome Announc* 5:e00758-17. <https://doi.org/10.1128/genomeA.00758-17>.
- Uchimura Y, Wyss M, Brugiroux S, Limenitakis JP, Stecher B, McCoy KD, Macpherson AJ. 2016. Complete genome sequences of 12 species of stable defined moderately diverse mouse microbiota 2. *Genome Announc* 4:e00951-16. <https://doi.org/10.1128/genomeA.00951-16>.
- Dekker J, Rippe K, Dekker M, Kleckner N. 2002. Capturing chromosome conformation. *Science* 295:1306–1311. <https://doi.org/10.1126/science.1067799>.
- Flot JF, Marie-Nelly H, Koszul R. 2015. Contact genomics: scaffolding and phasing (meta)genomes using chromosome 3D physical signatures. *FEBS Lett* 589:2966–2974. <https://doi.org/10.1016/j.febslet.2015.04.034>.
- Baudry L, Guiguelmoni N, Marie-Nelly H, Cormier A, Marbouty M, Avia K, Mie YL, Godfroy O, Sterck L, Cock JM, Zimmer C, Coelho SM, Koszul R. 2020. instaGRAAL: chromosome-level quality scaffolding of genomes

- using a proximity ligation-based scaffold. *Genome Biol* 21:148. <https://doi.org/10.1186/s13059-020-02041-z>.
11. Marbouty M, Baudry L, Cournac A, Koszul R. 2017. Scaffolding bacterial genomes and probing host-virus interactions in gut microbiome by proximity ligation (chromosome capture) assay. *Sci Adv* 3:e1602105. <https://doi.org/10.1126/sciadv.1602105>.
 12. Marbouty M, Thierry A, Millot GA, Koszul R. 2021. MetaHiC phage-bacteria infection network reveals active cycling phages of the healthy human gut. *Elife* 10:e60608. <https://doi.org/10.7554/eLife.60608>.
 13. Moreau P, Cournac A, Palumbo GA, Marbouty M, Mortaza S, Thierry A, Cairo S, Lavigne M, Koszul R, Neuveut C. 2018. Tridimensional infiltration of DNA viruses into the host genome shows preferential contact with active chromatin. *Nat Commun* 9:4268. <https://doi.org/10.1038/s41467-018-06739-4>.
 14. Marie-Nelly H, Marbouty M, Cournac A, Flot JF, Liti G, Parodi DP, Syan S, Guillén N, Margeot A, Zimmer C, Koszul R. 2014. High-quality genome (re)assembly using chromosomal contact data. *Nat Commun* 5:5695. <https://doi.org/10.1038/ncomms6695>.
 15. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.