

# Heterogeneous expression of Pil3 pilus is critical for Streptococcus gallolyticus translocation across polarized colonic epithelial monolayers

Mariana Martins, Laurence Du Merle, Patrick Trieu-Cuot, Shaynoor Dramsi

## ▶ To cite this version:

Mariana Martins, Laurence Du Merle, Patrick Trieu-Cuot, Shaynoor Dramsi. Heterogeneous expression of Pil3 pilus is critical for Streptococcus gallolyticus translocation across polarized colonic epithelial monolayers. Microbes and Infection, 2020, 22 (1), pp.55-59. 10.1016/j.micinf.2019.12.001. pasteur-03262258

## HAL Id: pasteur-03262258 https://pasteur.hal.science/pasteur-03262258

Submitted on 16 Jun2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License

Microbes and Infection 22 (2020) 55-59

Contents lists available at ScienceDirect

## Microbes and Infection

journal homepage: www.elsevier.com/locate/micinf

Short communication

# Heterogeneous expression of Pil3 pilus is critical for *Streptococcus* gallolyticus translocation across polarized colonic epithelial monolayers



Microbes an Infection

### Mariana Martins<sup>a, b, 1</sup>, Laurence du Merle<sup>a, b</sup>, Patrick Trieu-Cuot<sup>a, b</sup>, Shaynoor Dramsi<sup>a, b, \*</sup>

<sup>a</sup> Department of Microbiology, Biology of Gram-positive Pathogens Unit, Institut Pasteur, Paris, France
<sup>b</sup> Centre National de la Recherche Scientifique (CNRS) ERL, 6002, Paris, France

#### ARTICLE INFO

Article history: Received 8 July 2019 Accepted 2 December 2019 Available online 16 December 2019

Keywords: Pil3 pilus Streptococcus gallolyticus Colon Translocation S. bovis

#### ABSTRACT

Streptococcus gallolyticus is an opportunistic pathogen responsible for septicemia and endocarditis. We report that *S. gallolyticus* UCN34 adheres and crosses epithelial monolayers in a Pil3 dependent manner. Confocal images revealed a paracellular passage. Both the  $\Delta pil3$  mutant and the Pil3+ overexpressing variant were unable to cross Caco-2 and T84 barriers. However, combining live  $\Delta pil3$  mutant with fixed Pil3+ variant in a 9:1 ratio allowed efficient translocation of the  $\Delta pil3$  mutant. These results demonstrate that heterogeneous expression of Pil3 plays a key role for UCN34 translocation across the intestinal barrier. Through this skilful strategy, *S. gallolyticus* probably evade host immune responses.

© 2019 The Author(s). Published by Elsevier Masson SAS on behalf of Institut Pasteur. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Streptococcus gallolyticus subsp. gallolyticus (Sgg), formerly known as Streptococcus bovis biotype I, is an emerging opportunistic pathogen responsible for septicemia and infective endocarditis in the elderly [1,2]. This Gram-positive coccus is one of the few intestinal bacteria that have been consistently linked to colorectal cancer (CRC) [3,4]. Recent experimental data indicate that Sgg can act both as a driver and a passenger of CRC [5–10].

The first complete *S. gallolyticus* genome of strain UCN34, isolated from a patient suffering from endocarditis and later diagnosed for colon cancer, provided important insights on the adaptation and virulence strategies developed by this bacterium [11]. We previously showed that the Pil1 pilus is important for binding to collagen as well as colonization of heart valves in a rat model of experimental endocarditis [12]. We also showed that the Pil3 pilus is important for colonization of the host colon via attachment to the mucus covering colonic cells [13,14].

Pil1 and Pil3 are both expressed heterogeneously in the UCN34 population, with a majority of cells weakly piliated and a minority highly piliated, through a mechanism combining phase variation and attenuation [15].

In this report we investigated how Sgg adheres to and translocates across tight epithelial barriers in the absence of a secreted mucus layer. We demonstrate a key role of the Pil3 pilus in this process and visualized Sgg passage through a paracellular pathway. Our results indicate that in the UCN34 WT population, the highly piliated bacteria activate opening of the tight junctions to allow paracellular crossing of Pil3<sub>low</sub> expressing bacteria. This demonstrates the functional relevance of Pil3 heterogeneity.

#### 1. Materials and methods

#### 1.1. Cell cultures and bacterial strains

Caco-2 and T84 cells were routinely grown in Dulbecco's Modified Eagle Medium supplemented with 10% heat-inactivated fetal bovine serum (FBS). To obtain polarized Caco-2 and T84 monolayers, cells were seeded on inverted 12 mm polycarbonate, 3  $\mu$ m-pore, tissue culture inserts (Corning) at a density of 10<sup>6</sup> cells/ cm<sup>2</sup>. After 6 h at 37 °C, transwell inserts were placed back into the wells and supplemented with fresh media every two days for 14 days. Trans Epithelial Resistance (TER) was measured with an Ohmmeter (Millicell-ERS, Millipore) and paracellular permeability was measured using the nonionic macromolecular tracer FITC-Dextran 4000 Da (Sigma). The cell medium in both compartments was removed. The lower compartment (corresponding to

<sup>\*</sup> Corresponding author. 28, rue du Dr Roux, 75015, Paris, France.

E-mail address: shaynoor.dramsi@pasteur.fr (S. Dramsi).

<sup>&</sup>lt;sup>1</sup> Current address: Volumina Medical SA, Route de la Corniche 5, Epalinges 1066, Switzerland.

https://doi.org/10.1016/j.micinf.2019.12.001

<sup>1286-4579/© 2019</sup> The Author(s). Published by Elsevier Masson SAS on behalf of Institut Pasteur. This is an open access article under the CC BY-NC-ND license (http:// creativecommons.org/licenses/by-nc-nd/4.0/).

the apical surface of the epithelium) was replaced with RPMI without phenol red (Invitrogen) supplemented with FITC-dextran 4000 (5 mg/ml) and the upper compartment (i.e. basolateral side) with RPMI without phenol red. After incubation for 1 h, the upper compartment was sampled and fluorescence at 490 nm measured. S. *gallolyticus* strains were grown at 37 °C in Todd-Hewitt (TH) broth in standing filled flasks.

#### 1.2. Adherence/invasion assays

Caco-2 and T84 cells were seeded at  $3 \times 10^5$  cells ml<sup>-1</sup> in 24well plates and incubated at 37 °C in 5% CO2 until 100% confluence. Overnight cultures of S. gallolyticus strains were washed once in PBS and resuspended in DMEM low glucose medium prior to infecting cells at a MOI of 10 bacteria per cell. Bacteria added to confluent monolayers were centrifuged at 500 RPM  $\approx$  90g to synchronize infections. After 2, 4, 6 h of incubation at 37 °C under 5% CO<sub>2</sub> atmosphere, monolayers were washed 4 times to remove non-adherent bacteria, cells were then lysed in cold water and plated on Todd-Hewitt agar to count cell-associated bacteria. The percentage of adherence was calculated as follows: (CFU on plate count/CFU in inoculum) X100. Assays were performed in triplicate and were repeated in at least 3 independent experiments. For invasion assays, following bacterial adhesion, cells were washed three times with DMEM and then incubated for 60 min with DMEM containing gentamicin 100 µg/mL (Sigma) to kill extracellular bacteria. Cells were then lysed and enumerated as indicated above to assess the number of viable intracellular bacteria.

#### 1.3. Bacterial translocation assays

Overnight cultures of S. gallolyticus were washed in PBS and resuspended at  $1 \times 10^8$  CFU/ml in pre-warmed DMEM. Cell monolayers were washed with DMEM and inverted in 6-well plates. 50 µL of bacterial inoculum was then added to the apical side of the cells and incubated for 1 h at 37 °C in 5% CO<sub>2</sub> atmosphere. The transwell inserts were placed back into the corresponding wells and fresh culture media was added to both compartments. At each time point of infection, the medium from the upper compartment was recovered for CFU determination and replaced by fresh media to prevent bacterial planktonic growth. For the preparation of the artificial mixture, Sgg Pil3+ cells were washed in PBS and fixed with 4% PFA for 20 min. Following fixation, bacteria were washed 4 times in DMEM. We verified that no CFU could be recovered after this treatment. Live  $\Delta pil3$  mutant cells were then added to the PFA killed Pil3+ variant, a ratio of 9:1. For the blocking experiments with antibodies directed against Pil3, a combination of antibodies directed against the C- and N-terminal domains of Pil3A were added to the fixed Pil3+ variant and preincubated for 30 min at room temperature. This mixture was then added to the live  $\Delta pil3$  mutant for cell monolayer infection as indicated above.

#### 1.4. Confocal microscopy

At 6 h post-infection, cell monolayers were washed once with PBS and then fixed with PFA 4% for 10 min at room temperature. Monolayers were then washed three times with PBS and subsequently quenched with Glycine 0.1 M. For confocal microscopy, cells were permeabilized in PBS-Triton-X100 0.2% for 10 min at 4 °C. Cell monolayers were then incubated with anti-UCN34 polyclonal antibody at a 1:200 dilution, to specifically label *S. gallolyticus*, followed by incubation with secondary DyLight-488 conjugated goat anti-rabbit antibody (1:200). In addition, E-cadherin was stained using an anti-E-cadherin HECD-1 monoclonal

antibody (Invitrogen) at 1:100 with subsequent incubation with the secondary DyLight594 conjugated anti-mouse antibody (1:100). Finally, Hoecht 33342 (1:2000) was added to visualize cell nuclei and Alexa Fluor 647 phalloidin (1:50) to detect the actin cytoskeleton. Samples were mounted using ProLong Gold Antifade reagent and Z-stacks of 300 nm step size were acquired using a Leica TCS SP5 confocal microscope with a 63x oil objective. Immunofluorescence images were analyzed using the Fiji software.

#### 2. Results

#### 2.1. The Sgg Pil3 pilus enhances adherence to human colonic cells

Streptococcal pili have been implicated both in adherence to eukaryotic cells as well as in bacterial translocation across host epithelial barriers [16,17]. We previously showed that the Sgg Pil3 pilus strongly contributes to bacterial attachment to human mucus-producing cells HT29-MTX [14]. Therefore, we wondered whether Pil3 could also play a role in bacterial translocation. However, HT29-MTX cells were not able to form very tight epithelial barriers on Transwell filters. We therefore tested two other well-studied human colonic cell lines Caco-2 and T84, which are both able to form tightly polarized epithelial monolayers *in vitro* [18].

We first compared adherence of Sgg UCN34 (WT), a highly Pil3 piliated variant (Pil3+) and a deletion mutant ( $\Delta pil3$ ) to Caco-2 cells at different time points. These assays were carried out as previously described in HT29 and HT29-MTX cells [14]. As shown in Fig. 1, higher Pil3 expression levels led to increased bacterial adhesion to Caco-2 cells as compared to the WT while bacterial adherence in the absence of Pil3 was decreased at 6 h post-infection. Very similar results were observed in T84 cells (Fig. S1A). Of note, Sgg strain UCN34, which displays intermediate adherence, is composed of a heterogeneous population with approximately 10–20% highly expressing Pil3 pilus (Pil3<sup>high</sup>) and 80–90% weakly piliated (Pil3<sup>low</sup>) [14]. Together, these results indicate that the Pil3 pilus contributes to Sgg adhesion to human colonic epithelial cells. We next investigated Sgg translocation.

#### 2.2. S. gallolyticus translocates across colonic epithelial barriers

In order to study translocation of Sgg, we first established an *in vitro* model of polarized cells using human colonic Caco-2 and T84 cell lines (Figs. S1 and S2). Both cell lines were cultured on Transwell inserts for 3–21 days which allowed their polarization and



**Fig. 1. Role of Pil3 pilus in the adherence of Sgg UCN34 to human colorectal cancer cells Caco-2.** Adherence is presented as percentage of bacterial inoculum after 4 h and 6 h at 37 °C at a multiplicity of infection of 10 bacteria per cell. Planktonic growth of the three Sgg variants in this cell medium was monitored and did not change during the infection period.

differentiation [18]. On these filters, fully differentiated Caco-2 and T84 cells expressed well-organized cell-to-cell junctions, forming a cell monolayer mimicking the intestinal epithelial barrier. Integrity of the epithelial barrier was monitored over time using two complementary methods: transepithelial electrical resistance (TER) of the monolayers and permeability to 4 kDa FITC-Dextran molecular ruler (Figs. S1CD and S2). After 11 days in culture, the Caco-2 and T84 monolavers were already impermeable to the 4 kDa dextran molecules and reached a TER of approximately 300 and 2500  $\Omega$  cm<sup>2</sup>, respectively (Figs. S1C and S2B). The capacity of Sgg UCN34 to translocate across the impermeable Caco-2 and T84 barriers was assessed at days 7, 14, and 21 and although translocation of UCN34 WT increased with cell differentiation (Fig. S3), day 14 was chosen as the time point giving the most consistent results. In order to demonstrate that bacterial translocation is an active process, cell monolayers were apically infected with Sgg UCN34 by inverting the transwell (lower chamber of the transwell insert) and translocated bacteria were recovered in the upper chamber at 2, 4, and 6 h post-infection (Fig. 2A). The rate of translocation increased over time with a maximum translocation of about 10% in Caco-2 (Fig. 2B) and 12% in T84 monolayers (Fig. S1B) at 6 h post-infection. The effect of Sgg on epithelial barrier function was also assessed by measuring the TER following infection. TER values remained stable upon infection and were comparable to non-infected control monolayers, indicating no major disruption of the epithelial barrier by Sgg UCN34 (Figs. S2C and S1D). In order to visualize bacteria during the translocation process, confocal imaging was carried out on infected Caco-2 cells grown on filters at 6 h post-infection. Caco-2 cells were stained with E-cadherin which localizes at epithelial junctions, actin was stained with phalloidin to visualize the cytoskeleton and bacteria were labeled using a specific polyclonal antibody directed against strain UCN34. During translocation Sgg UCN34 was found primarily close to E-cadherin, at epithelial junctions between adjacent cells (Fig. 2C). These images suggest that UCN34 uses a paracellular route with a transient opening of cell junctions, as suggested by a previous study (3). Cell monolayers were considered well polarized, as demonstrated by the specific accumulation of actin close to the apical membrane of the cells. No intracellular Sgg UCN34 could be detected inside Caco-2 cells (Fig. S4). It is important to mention that the adhesion/invasion experiments shown in Fig. S4 were performed in low glucose conditions to mimic the conditions of Transwell experiments. Altogether, these results strongly suggest that Sgg UCN34 is able to translocate epithelial



**Fig. 2. Translocation of Sgg UCN34 across Caco-2 cells.** A- Schematic representation of the infection process using inverted transwell system. The transwell inserts were washed, inverted and incubated for about 1 h to allow bacterial attachment to the cells. Then the inserts were placed back into a new 6 well plate. At each time point of infection, the medium for the upper compartment was completely recovered for CFU quantification and replaced by fresh medium. B- Translocation of *Sgg* UCN34 across Caco-2 monolayer. Cells were infected for 2, 4 and 6 h with a multiplicity of infection of 10 bacteria per cell. Translocation values are relative to the inoculum and represent 5 independent experiments performed in duplicate. C- Visualization of *Sgg* UCN34 translocation using confocal microscopy. After 6 h of infection, Caco-2 monolayer were fixed, permeabilized and stained with a monoclonal antibody against E-cadherin (in red), A547 conjugated-phalloidin to visualize actin (purple), and Hoescht 33342 to reveal cells nuclei (cyan). Bacteria were detected with a specific polyclonal antibody raised against the whole bacterium (green). The upper panels show an YX view of the filter, whereas the lower panels show an YZ view. The arrows are pointing to UCN34. In the upper panels the scale bar represents 10 µm and in the yz planes 5 µm. Right panel: representative image from another independent experiment. The scale bar corresponds to 10 µm.

barriers of human intestinal cells through a process involving a transient and subtle opening of tight junctions.

# 2.3. Pil3 pilus heterogeneity in Sgg is required for efficient bacterial translocation

To gain further mechanistic insights about Sgg translocation across epithelial barriers, we analyzed the possible contribution of the Pil3 pilus in this process. Caco-2 (Fig. 3A) and also T84 (Fig. S1B) monolayers of cells were infected with Sgg UCN34 (WT), a Pil3+ variant and the  $\Delta pil3$  mutant for 2, 4, and 6 h. WT Sgg was able to translocate these barriers at a rate of 5–10%. In contrast, the otherwise isogenic Pil3+ variant and  $\Delta pil3$  mutant were significantly impaired by about 5-fold in this process as compared to WT UCN34. These results demonstrate that: i) the Pil3 pilus is essential for Sgg translocation across intestinal barriers as the  $\Delta pil3$  mutant is unable to translocate and more surprisingly ii) that Pil3 pilus heterogeneity is also functionally important since a highly expressing Pil3+ variant cannot translocate these cell monolayers. We hypothesize that increased adherence of the Pil3+ variant at the apical surface of the cells impaired bacterial translocation. Hence, it appears that a delicate balance in Pil3 expression is important for efficient Sgg translocation. To test this hypothesis, we artificially mimicked the expression levels of the Pil3 pilus in the natural Sgg UCN34 population (around 90% of Pil3low and 10% of Pil3<sub>high</sub>). To achieve this, we mixed the  $\Delta pil3$  mutant with PFA killed Pil3+ variant in a proportion of 9:1, respectively. We then infected Caco-2 monolayers with this mixture and observed the capacity of the live  $\Delta pil3$  mutant to translocate (Fig. 3B). Interestingly, the presence of fixed Pil3+ bacteria allowed the  $\Delta pil3$  mutant to cross the Caco-2 monolayer with the same efficiency as Sgg UCN34. These data demonstrate that heterogeneous expression of Pil3 in Sgg is crucial for efficient translocation across intestinal barriers. Furthermore, blocking experiments were carried out in this experimental setting using specific antibodies against Pil3 (Fig. 3C). First, we showed that the addition of antibodies against both Pil3 subunits impaired translocation of the  $\Delta pil3$  mutant in the presence of PFA killed Pil3+ variant cells (Fig. 3C). The Pil3 pilus is composed of a putative tip-located Pil3A adhesin and of a major Pil3B pilin subunit constituting the backbone of the Pil3 filamentous structure. In order to demonstrate the specific role of the Pil3A adhesin in Sgg translocation, antibodies against Pil3A were tested in the same experimental setting and were shown to be sufficient to block translocation of the  $\Delta pil3$  mutant with PFA killed Pil3+ cells (Fig. 3D). As a control, we used a similar type of antibody raised against Pil1 that did not prevent translocation of live  $\Delta pil3$  mutant cells in the presence of PFA killed Pil3+ bacteria. Altogether, these results demonstrate that heterogeneous expression of the Pil3 pilus in the Sgg UCN34 is critical for its ability to translocate efficiently across tight epithelial barriers, and that the paracellular opening process depends on Pil3A adhesin interaction with one or several as yet unknown host cell receptor(s).

#### 3. Discussion

*S. gallolyticus subsp. gallolyticus* (*Sgg*) belongs to the *S. bovis/S. equinus* complex, a diverse group of streptococci that are commensals of the gut, opportunistic pathogens and used in dairy product fermentation. *Sgg* is described as a weak colonizer of the gastrointestinal tract with a fecal carriage of about 2.5–15%. It is



Fig. 3. Bacterial translocation across Caco-2 monolayer is dependent on Pil3 pilus heterogeneous expression. A- UCN34, Pil3+ and  $\Delta pil3$  translocation across Caco-2 monolayer after 2 h, 4 h and 6 h of infection. B- translocation of a mix of 9:1  $\Delta pil3$  and fixed killed Pil3+ as compared to WT UCN34 and  $\Delta pil3$  alone. C- inhibition of translocation with antibodies against Pil3. D-inhibition of translocation with specific antibodies directed against Pil3A adhesin but not with control antibodies directed against Pil1. Results are means  $\pm$  SD from 5 independent experiments performed in duplicate. Asterisks represent statistical differences relative to WT strain UCN34 with \* p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001 using two-way ANOVA with Bonferroni's post-test in GraphPad Prism version 5.

believed that under certain specific physiological conditions such as development of colon malignancies, *Sgg* is able to overgrow by benefiting from specific tumoral nutrients [8] and outcompeting closely related microbiota gut commensals [9]. This increase in *Sgg* load and the changes in the gut intestinal barrier resulting from tumor development are suspected to favor *Sgg* translocation across the tight intestinal barrier, which in turn can lead to invasive infections such as septicemia and infective endocarditis [19,20].

In this work, we investigated the ability of Sgg strain UCN34 to translocate across intestinal barriers using Caco-2 and T84, two widely used model cell lines derived from human colon adenocarcinoma. Our results are in perfect agreement with a previous report by Boleij et al. showing that Sgg UCN34 could efficiently translocate across polarized Caco-2 cells while the closely related non-pathogenic *S. gallolyticus subsp. macedonicus* (Sgm) was not able to do so [21]. Since Sgm does not possess any pili, we hypothesized that the Pil3 pilus could be involved in translocation. Pili have long been considered important players in bacterial attachment to host tissues and their role in translocation across intestinal epithelia was elucidated for GBS [17].

Here, we demonstrate that translocation of Sgg UCN34 across polarized intestinal cells is a Pil3-dependent process. Indeed, the  $\Delta pil3$  mutant was unable to translocate across Caco-2 and T84 monolayers. Expression of Pil3 in WT UCN34 is known to be heterogeneous at the population level, with a majority of cells weakly piliated (90%) and a minority highly piliated (10%). Interestingly we found that a Pil3+ variant homogeneously expressing high levels of Pil3 pilus is unable to translocate intestinal barriers. This result suggested that heterogeneous expression of Pil3 plays a key role in the translocation process. We were able to mimic this heterogeneity *in vitro* by mixing live  $\Delta pil3$  mutant cells with the PFA-killed Pil3+ variant. Both the  $\Delta pil3$  and Pil3+ variants were unable to translocate the intestinal barrier alone. Strikingly, when combined in a proportion similar to that found in the UCN34 WT population, about 9  $\Delta pil3$  for 1 Pil3+, we found that the presence of Pil3+ variant cells allowed translocation of the  $\Delta pil3$  mutant. Based on these results we propose the following model to explain crossing of epithelial junctions by Sgg. Highly piliated Pil3 bacteria in the UCN34 population interact with an unknown cell surface receptor, or a component of the tight junctions, likely through the Pil3A adhesin, activating signaling pathway(s) involved in regulation of epithelial cell junctions. This then leads to opening of cell junctions allowing loosely bound bacteria with low Pil3 levels to pass in between adjacent cells. This translocation occurs without major disruption of the epithelial junctions. As shown for Pil1 [15], this heterogeneity of Pil3 pilus expression may also mitigate host immune responses allowing Sgg to more efficiently evade host intestinal immune defenses in the lamina propria and later in the blood. Future studies will be aimed at investigating the identity of the Pil3A receptor on polarized colonic cells.

#### **Declaration of Competing Interest**

We herein declare that all authors have seen and approved the content of this manuscript and contributed significantly to this work. None of the authors have a financial, personal, or other relationships with other people or organizations within three years of beginning the submitted work that could inappropriately influence, or be perceived to influence, their work.

#### Acknowledgments

We sincerely thank Tarek Msadek for the critical reading of the manuscript. This work was supported by the French Government's "Investissement d'Avenir" program Laboratoire d'Excellence "Integrative Biology of Emerging Infectious Diseases" Grant ANR-10-LABX-62-IBEID. Mariana Martins was funded by the PPU program and the ARC foundation.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.micinf.2019.12.001.

#### References

- Vogkou CT, Vlachogiannis NI, Palaiodimos L, Kousoulis AA. The causative agents in infective endocarditis: a systematic review comprising 33,214 cases. Eur J Clin Microbiol Infect Dis 2016;35:1227–45.
- [2] Hoen B, Chirouze C, Cabell CH, Selton-Suty C, Duchene F, Olaison L, et al. Emergence of endocarditis due to group D streptococci: findings derived from the merged database of the International Collaboration on Endocarditis. Eur J Clin Microbiol Infect Dis 2005;24:12–6.
- [3] Boleij A, van Gelder MM, Swinkels DW, Tjalsma H. Clinical Importance of Streptococcus gallolyticus infection among colorectal cancer patients: systematic review and meta-analysis. Clin Infect Dis 2011;53:870–8.
- [4] Kwong TNY, Wang X, Nakatsu G, Chow TC, Tipoe T, Dai RZW, et al. Association between bacteremia from specific microbes and subsequent diagnosis of colorectal cancer. Gastroenterology 2018;155:383–90. e8.
- [5] Zhang Y, Weng Y, Gan H, Zhao X, Zhi F. Streptococcus gallolyticus conspires myeloid cells to promote tumorigenesis of inflammatory bowel disease. Biochem Biophys Res Commun 2018;506:907–11.
- [6] Kumar R, Herold JL, Taylor J, Xu J, Xu Y. Variations among *Streptococcus gal-lolyticus* subsp. *gallolyticus* strains in connection with colorectal cancer. Sci Rep 2018;8:1514.
- [7] Kumar R, Herold JL, Schady D, Davis J, Kopetz S, Martinez-Moczygemba M, et al. *Streptococcus gallolyticus* subsp. *gallolyticus* promotes colorectal tumor development. PLoS Pathog 2017;13:e1006440.
- [8] Boleij A, Dutilh BE, Kortman GA, Roelofs R, Laarakkers CM, Engelke UF, et al. Bacterial responses to a simulated colon tumor microenvironment. Mol Cell Proteom 2012;11:851–62.
- [9] Aymeric L, Donnadieu F, Mulet C, du Merle L, Nigro G, Saffarian A, et al. Colorectal cancer specific conditions promote *Streptococcus gallolyticus* gut colonization. Proc Natl Acad Sci U S A 2018;115:E283–91.
- [10] Abdulamir AS, Hafidh RR, Bakar FA. Molecular detection, quantification, and isolation of *Streptococcus gallolyticus* bacteria colonizing colorectal tumors: inflammation-driven potential of carcinogenesis via IL-1, COX-2, and IL-8. Mol Cancer 2010;9:249.
- [11] Rusniok C, Couve E, Da Cunha V, El Gana R, Zidane N, Bouchier C, et al. Genome sequence of *Streptococcus gallolyticus*: insights into its adaptation to the bovine rumen and its ability to cause endocarditis. J Bacteriol 2010;192: 2266–76.
- [12] Danne C, Entenza JM, Mallet A, Briandet R, Debarbouille M, Nato F, et al. Molecular characterization of a *Streptococcus gallolyticus* genomic island encoding a pilus involved in endocarditis. J Infect Dis 2011;204:1960–70.
- [13] Martins M, Porrini C, du Merle L, Danne C, Robbe-Masselot C, Trieu-Cuot P, et al. The Pil3 pilus of *Streptococcus gallolyticus* binds to intestinal mucins and to fibrinogen. Gut Microb 2016;7:526–32.
- [14] Martins M, Aymeric L, du Merle L, Danne C, Robbe-Masselot C, Trieu-Cuot P, et al. *Streptococcus gallolyticus* Pil3 pilus is required for adhesion to colonic mucus and for colonization of mouse distal colon. J Infect Dis 2015;212: 1646–55.
- [15] Danne C, Dubrac S, Trieu-Cuot P, Dramsi S. Single cell stochastic regulation of pilus phase variation by an attenuation-like mechanism. PLoS Pathog 2014;10:e1003860.
- [16] Soriani M, Santi I, Taddei A, Rappuoli R, Grandi G, Telford JL. Group B Streptococcus crosses human epithelial cells by a paracellular route. J Infect Dis 2006;193:241–50.
- [17] Pezzicoli A, Santi I, Lauer P, Rosini R, Rinaudo D, Grandi G, et al. Pilus backbone contributes to group B *Streptococcus* paracellular translocation through epithelial cells. J Infect Dis 2008;198:890–8.
- [18] Dharmsathaphorn K, Madara JL. Established intestinal cell lines as model systems for electrolyte transport studies. Methods Enzymol 1990;192: 354–89.
- [19] Pasquereau-Kotula E, Martins M, Aymeric L, Dramsi S. Significance of *Strep-tococcus gallolyticus* subsp. gallolyticus association with colorectal cancer. Front Microbiol 2018;9:614.
- [20] Boleij A, Tjalsma H. The itinerary of *Streptococcus gallolyticus* infection in patients with colonic malignant disease. Lancet Infect Dis 2013;13:719–24.
- [21] Boleij A, Muytjens CM, Bukhari SI, Cayet N, Glaser P, Hermans PW, et al. Novel clues on the specific association of *Streptococcus gallolyticus* subsp gallolyticus with colorectal cancer. J Infect Dis 2011;203:1101–9.