

Adnaviria: a New Realm for Archaeal Filamentous Viruses with Linear A-Form Double-Stranded DNA Genomes

Mart Krupovic, Jens H Kuhn, Fengbin Wang, Diana P Baquero, Valerian V Dolja, Edward H Egelman, David Prangishvili, Eugene V Koonin

▶ To cite this version:

Mart Krupovic, Jens H Kuhn, Fengbin Wang, Diana P Baquero, Valerian V Dolja, et al.. Adnaviria: a New Realm for Archaeal Filamentous Viruses with Linear A-Form Double-Stranded DNA Genomes. Journal of Virology, 2021, 95 (15), pp.e00673-21. 10.1128/JVI.00673-21. pasteur-03284792

HAL Id: pasteur-03284792 https://pasteur.hal.science/pasteur-03284792

Submitted on 12 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

JVI Accepted Manuscript Posted Online 19 May 2021 J Virol doi:10.1128/JVI.00673-21 Copyright © 2021 American Society for Microbiology. All Rights Reserved.

1 Adnaviria: a new realm for archaeal filamentous viruses with linear A-form double-

2 stranded DNA genomes

3

- 4 Mart Krupovic^a*, Jens H. Kuhn^b, Fengbin Wang^c, Diana P. Baquero^a, Valerian V. Dolja^d,
- 5 Edward H. Egelman^c, David Prangishvili^{a,e}, Eugene V. Koonin^f

6

- 7 ^aArchaeal Virology Unit, Institut Pasteur, Paris, France
- ^bNIH/NIAID/DCR/Integrated Research Facility at Fort Detrick (IRF-Frederick), Fort Detrick,
 Frederick, MD, USA
- ^cDepartment of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville,
 VA, USA
- ^dDepartment of Botany and Plant Pathology, Oregon State University, Corvallis, OR, USA
- 13 ^eIvane Javakhishvili Tbilisi State University, Tbilisi, Georgia
- ¹⁴ ^fNational Center for Biotechnology Information, National Library of Medicine, National
- 15 Institutes of Health, Bethesda, MD, USA

16

17 *Correspondence to Mart Krupovic, mart.krupovic@pasteur.fr

- 19 Keywords: A-form DNA, Ligamenvirales, Tokiviricetes, Rudiviridae, Lipothrixviridae,
- 20 Tristromaviridae, hyperthermophilic archaea, virus evolution, virus classification, virus
- 21 taxonomy, International Committee on Taxonomy of Viruses (ICTV), major capsid protein, virus
- 22 structure and assembly
- 23

The International Committee on Taxonomy of Viruses (ICTV) has recently adopted a comprehensive, hierarchical system of virus taxa. The highest rank in this hierarchy are realms, each of which is considered monophyletic but apparently originated independently of other realms. Here we announce the creation of a new realm, *Adnaviria*, which unifies archaeal filamentous viruses with linear A-form double-stranded DNA genomes and characteristic major capsid proteins unrelated to those encoded by other known viruses.

31

32 Text

In 2018, the International Committee on Taxonomy of Viruses (ICTV) expanded the number of 33 taxonomic ranks available for virus classification from 5 to 15 (1, 2). This development enabled 34 35 formal recognition of the evolutionary kinships among distantly related viruses and led to creation of four realms-virus taxa that are roughly equivalent to the domain rank in cellular 36 taxonomy (3): The realm Riboviria encompasses all RNA viruses and reverse-transcribing 37 viruses that encode homologous RNA-directed RNA polymerases and reverse transcriptases, 38 39 respectively; Monodnaviria comprises viruses with predominantly single-stranded (ss) DNA genomes and encoding rolling-circle replication initiation endonucleases of the HUH 40 superfamily; Duplodnaviria unifies viruses with double-stranded DNA genomes that produce 41 virions with icosahedral capsids formed by major capsid proteins (MCPs) with the HK97 fold; 42 and Varidnaviria comprises dsDNA viruses with icosahedral capsids built from MCPs with 43 44 double-jelly roll fold (4). However, many virus families remained unassigned to higher taxa due to the lack of demonstrable evolutionary relationships to viruses assigned to the four realms. 45

lournal of Virology

announce the creation, and recent official acceptance by the ICTV 46 Here we 47 (https://talk.ictvonline.org/taxonomy/), of a new realm, Adnaviria, which encompasses structurally related archaeal filamentous viruses with dsDNA genomes that adopt the A-form 48 conformation within their virions. 49

Filamentous viruses infect hosts of all cellular domains but, despite overall similar morphology, 50 virions of eukaryotic, bacterial, and archaeal filamentous viruses are built from capsid proteins 51 with different structural folds (5). Furthermore, the types of the viral genomes that are protected 52 53 by the corresponding capsid proteins are radically different among viruses infecting hosts from the different domains of life (6-11): All known eukaryotic filamentous viruses have linear ssRNA 54 genomes and are classified in realm Riboviria (4); all known bacterial filamentous viruses have 55 56 circular ssDNA genomes and belong to realm Monodnaviria (4, 12); and all known archaeal 57 filamentous viruses have dsDNA genomes and, until recently, have remained unassigned to taxa ranked higher than order. 58

59 Archaeal viruses that form filamentous virions are currently classified into four families: 60 Clavaviridae, Lipothrixviridae, Rudiviridae, and Tristromaviridae (13). Based on the shared gene content, structural similarity of their virions, and homology of the MCPs, families 61 62 Lipothrixviridae and Rudiviridae were included in order Ligamenvirales (14). In virions of both families, the nucleoprotein helix is composed of asymmetric units containing two MCP 63 molecules, a homodimer in the case of rudivirids and a heterodimer of paralogous MCPs in the 64 case of lipothrixvirids (Figure 1) (11, 15). The MCPs of ligamenviral particles have a unique α -65 helical fold first found in the MCP of rudivirid Sulfolobus islandicus rod-shaped virus 2 (SIRV2) 66 67 (16). Lipothrixvirids and rudivirids share a characteristic feature in that the interaction between

Σ

the MCP dimer and the linear dsDNA genome maintains the DNA in the A form. Consequently,
the entire genome adopts the A-form in virions (<u>15-18</u>).

70 Clavavirids (19, 20) and tristromavirids (21) do not encode proteins with recognizable sequence 71 similarity to proteins of ligamenvirals (22). Hence, the two families were not included in Ligamenvirales. Recently, the structure of particles produced by tristromavirid Pyrobaculum 72 73 filamentous virus 2 (PFV2) has been characterized by cryogenic electron microscopy (cryo-EM) 74 (23). Unexpectedly, the two nucleocapsid proteins of tristromavirids were found to be structurally related to those of ligamenvirals, and the virion organizations were discovered to be 75 remarkably similar, including the A-form conformation of the genomic dsDNA (23) (Figure 1). 76 77 Similar to lipothrixvirids, virions of tristromavirids are enveloped by a lipid membrane (24). 78 Based on the structural similarity between members of *Ligamenvirales* and *Tristromaviridae*, we inferred that they share a common origin (23). To formalize this evolutionary relationship, 79 Tristromaviridae was assigned to a new order, Primavirales, which, along with Ligamenvirales, 80 81 was included in a new class, Tokiviricetes.

82 Members of the *Tokiviricetes* represent one of the most commonly detected virus groups in terrestrial hot springs and are globally distributed (25-27). Rudivirids have been extensively 83 studied in particular, and many aspects of their life cycles are understood in considerable detail 84 (28). Notably, the mechanisms of genome replication do not appear to be conserved among 85 86 tokiviricetes. It has been proposed that rudivirus SIRV2 uses a combination of stranddisplacement, rolling-circle, and strand-coupled genome replication mechanisms, which generate 87 88 multimeric, highly branched, "brush-like" intermediates reaching lengths of >1,200 kb (~34 genome units) (29). The latter are then processed into unit-length, linear genomes with 89 covalently linked hairpin ends. The genome replication of the lipothixvirid Acidianus 90

91 filamentous virus 1 has been suggested to start by the formation of a D loop and to progress by 92 the strand displacement replication mechanism, whereas termination relies on recombination 93 events through the formation of terminal loop-like structures, though the genes involved in this 94 unique mechanism of replication remain unknown (<u>30</u>). Similar to rudivirid genomes, 95 tristromavirid genomes have terminal inverted repeats (<u>21</u>), but the replication mechanism of 96 these viruses has not been investigated.

97 High-resolution structures are now available for virions of six distinct tokiviricetes (11, 15). Like many structurally related viruses in the two other realms of dsDNA viruses (Duplodnaviria and 98 Varidnaviria), there is no detectable sequence similarity among the capsid proteins of viruses 99 100 from different tokiviricete families, suggesting a vast undescribed diversity of viruses in this part 101 of the virosphere. Indeed, it has been suggested that tokiviricetes were present in the last archaeal 102 common ancestor, and possibly even in the last universal cellular ancestor (LUCA) (31). Regardless, the available data unequivocally show that archaeal filamentous viruses do not fall 103 104 into any of the four established realms. Thus, given the lack of detectable relationship with other 105 viruses and officially acknowledging the uniqueness of the three families of archaeal filamentous 106 viruses that contain A-form DNA in their virions, a new taxon of the highest rank, realm 107 Adnaviria, was created for their classification. To bridge the gap between the class and realm 108 taxa, intermediate kingdom and phylum taxa, named Zilligvirae and Taleaviricota, respectively, 109 were established.

In contrast, structural characterization of clavavirid Aeropyrum pernix bacilliform virus 1
(APBV1) particles by cryo-EM confirmed that the fold of its MCP and its overall virion
organization are unrelated to that of ligamenviral, tristromavirid, or other characterized virus

113 particles (<u>32</u>). Therefore, notwithstanding their filamentous virions, clavavirids are not included

114 in realm *Adnaviria* and likely represent another realm to be established in the future.

115

116 Etymology of new taxa recently accepted by the ICTV:

- *Adnaviria*; from <u>A</u>-form <u>DNA</u> characteristic of viruses in this realm and suffix *-viria* for
 realm taxa.
- *Zilligvirae*; after Wolfram Zillig (1925–2005), a pioneer of research on
- 120 hyperthermophilic archaeal viruses, and the suffix *-virae* for kingdom taxa.
- *Taleaviricota*; from Latin *talea*, meaning "rod" (referring to the virion morphology), and
 suffix *-viricota* for phylum taxa.
- *Tokiviricetes*; from Georgian *orego* (*toki*), meaning "thread", and suffix -*viricetes* for
 class taxa.
- *Primavirales*; from Latin *prima*, meaning "first", referring to the fact that Thermoproteus
 tenax virus 1, classified in this order, was the first hyperthermophilic archaeal virus to be
 isolated in 1983 (33).

128

129 ACKNOWLEDGMENTS

- 130 We are grateful to Anya Crane (National Institutes of Health [NIH] National Institute of Allergy
- and Infectious Diseases [NIAID] Division of Clinical Research [DCR] Integrated Research
- 132 Facility at Fort Detrick [IRF-Frederick], Frederick, MD, USA) for editing the manuscript.

133	M.K. was supported by l'Agence Nationale de la Recherche (grant ANR-20-CE20-0009-02) and
134	Emergence(s) project MEMREMA from Ville de Paris. E.V.K. is supported by the Intramural
135	Research Program of the National Institutes of Health of the USA (National Library of
136	Medicine). E.H.E. was supported by NIH R35GM122510. This work was supported in part
137	through Laulima Government Solutions, LLC prime contract with the US National Institute of
138	Allergy and Infectious Diseases (NIAID) under Contract No. HHSN272201800013C. J.H.K.
139	performed this work as an employee of Tunnell Government Services (TGS), a subcontractor of
140	Laulima Government Solutions, LLC under Contract No. HHSN272201800013C.
141	The views and conclusions contained in this document are those of the authors and should not be
142	interpreted as necessarily representing the official policies, either expressed or implied, of the US
143	Department of Health and Human Services or of the institutions and companies affiliated with

the authors.

146 Figure legend

147	Figure 1. Taxonomic organization of realm Adnaviria. A. Taxonomic ranks from family to
148	realm are rendered with different colors. B-D. Cryogenic electron microscopy shows structures
149	of the major capsid protein (MCP) homodimer (characteristic of rudivirids) and heterodimer
150	(characteristic of lipothrixvirids and tristromavirids) (B), MCPs bound to viral double-stranded
151	(ds) DNA genome in A-form conformation (C), and side views of the viral nucleocapsid are
152	shown for each constituent family (D). Rudivirdae, Lipothrixviridae, and Tristromaviridae are
153	represented by structures of Sulfolobus islandicus rod-shaped virus 2 (PDB entry: 3J9X),
154	Acidianus filamentous virus 1 (PDB entry: 5WTG), and Pyrobaculum filamentous virus 2 (PDB

155 entry: 6VTB), respectively. The figure is modified from ref $(\underline{11})$.

156	56 References		
157	1.	Gorbalenya AE, Krupovic M, Mushegian A, Kropinski AM, Siddell SG, Varsani A,	
158		Adams MJ, Davison AJ, Dutilh BE, Harrach B, Harrison RL, Junglen S, King AMQ,	
159		Knowles NJ, Lefkowitz EJ, Nibert ML, Rubino L, Sabanadzovic S, Sanfaçon H,	
160		Simmonds P, Walker PJ, Zerbini FM, Kuhn JH. 2020. The new scope of virus taxonomy:	
161		partitioning the virosphere into 15 hierarchical ranks. Nat Microbiol 5:668-674.	
162	2.	Siddell SG, Walker PJ, Lefkowitz EJ, Mushegian AR, Adams MJ, Dutilh BE,	
163		Gorbalenya AE, Harrach Bz, Harrison RL, Junglen S, Knowles NJ, Kropinski AM,	
164		Krupovic M, Kuhn JH, Nibert M, Rubino L, Sabanadzovic S, Sanfaçon H, Simmonds P,	
165		Varsani A, Zerbini FM, Davison AJ. 2019. Additional changes to taxonomy ratified in a	
166		special vote by the International Committee on Taxonomy of Viruses (October 2018).	
167		Arch Virol 164:943-946.	
168	3.	Walker PJ, Siddell SG, Lefkowitz EJ, Mushegian AR, Dempsey DM, Dutilh BE, Harrach	
169		B, Harrison RL, Hendrickson RC, Junglen S, Knowles NJ, Kropinski AM, Krupovic M,	
170		Kuhn JH, Nibert M, Rubino L, Sabanadzovic S, Simmonds P, Varsani A, Zerbini FM,	
171		Davison AJ. 2019. Changes to virus taxonomy and the International Code of Virus	
172		Classification and Nomenclature ratified by the International Committee on Taxonomy of	
173		Viruses (2019). Arch Virol 164:2417-2429.	
174	4.	Koonin EV, Dolja VV, Krupovic M, Varsani A, Wolf YI, Yutin N, Zerbini FM, Kuhn	
175		JH. 2020. Global organization and proposed megataxonomy of the virus world. Microbiol	
176		Mol Biol Rev 84:e00061-19.	
177	5.	Krupovic M, Koonin EV. 2017. Multiple origins of viral capsid proteins from cellular	
178		ancestors. Proc Natl Acad Sci U S A 114:E2401-E2410.	

Σ

179	6.	DiMaio F, Chen C-C, Yu X, Frenz B, Hsu Y-H, Lin N-S, Egelman EH. 2015. The
180		molecular basis for flexibility in the flexible filamentous plant viruses. Nat Struct Mol
181		Biol 22:642-4.
182	7.	Klug A. 1999. The tobacco mosaic virus particle: structure and assembly. Philos Trans R
183		Soc Lond B Biol Sci 354:531-5.
184	8.	Stubbs G, Kendall A. 2012. Helical viruses. Adv Exp Med Biol 726:631-58.
185	9.	Tarafder AK, von Kügelgen A, Mellul AJ, Schulze U, Aarts DGAL, Bharat TAM. 2020.
186		Phage liquid crystalline droplets form occlusive sheaths that encapsulate and protect
187		infectious rod-shaped bacteria. Proc Natl Acad Sci U S A 117:4724-4731.
188	10.	Zamora M, Méndez-López E, Agirrezabala X, Cuesta R, Lavín JL, Sánchez-Pina MA,
189		Aranda MA, Valle M. 2017. Potyvirus virion structure shows conserved protein fold and
190		RNA binding site in ssRNA viruses. Sci Adv 3:eaao2182.
191	11.	Baquero DP, Liu Y, Wang F, Egelman EH, Prangishvili D, Krupovic M. 2020. Structure
192		and assembly of archaeal viruses. Adv Virus Res 108:127-164.
193	12.	Roux S, Krupovic M, Daly RA, Borges AL, Nayfach S, Schulz F, Sharrar A, Matheus
194		Carnevali PB, Cheng J-F, Ivanova NN, Bondy-Denomy J, Wrighton KC, Woyke T, Visel
195		A, Kyrpides NC, Eloe-Fadrosh EA. 2019. Cryptic inoviruses revealed as pervasive in
196		bacteria and archaea across Earth's biomes. Nat Microbiol 4:1895-1906.
197	13.	Prangishvili D, Bamford DH, Forterre P, Iranzo J, Koonin EV, Krupovic M. 2017. The
198		enigmatic archaeal virosphere. Nat Rev Microbiol 15:724-739.
199	14.	Prangishvili D, Krupovic M. 2012. A new proposed taxon for double-stranded DNA
200		viruses, the order "Ligamenvirales". Arch Virol 157:791-5.

10

201	15.	Wang F, Baquero DP, Beltran LC, Su Z, Osinski T, Zheng W, Prangishvili D, Krupovic
202		M, Egelman EH. 2020. Structures of filamentous viruses infecting hyperthermophilic
203		archaea explain DNA stabilization in extreme environments. Proc Natl Acad Sci U S A
204		117:19643-19652.
205	16.	DiMaio F, Yu X, Rensen E, Krupovic M, Prangishvili D, Egelman EH. 2015. A virus that
206		infects a hyperthermophile encapsidates A-form DNA. Science 348:914-7.
207	17.	Kasson P, DiMaio F, Yu X, Lucas-Staat S, Krupovic M, Schouten S, Prangishvili D,
208		Egelman EH. 2017. Model for a novel membrane envelope in a filamentous
209		hyperthermophilic virus. Elife 6:e26268.
210	18.	Liu Y, Osinski T, Wang F, Krupovic M, Schouten S, Kasson P, Prangishvili D, Egelman
211		EH. 2018. Structural conservation in a membrane-enveloped filamentous virus infecting a
212		hyperthermophilic acidophile. Nat Commun 9:3360.
213	19.	Prangishvili D, Mochizuki T, Liu Y, Krupovic M, ICTV Report Consortium. 2019. ICTV
214		Virus Taxonomy Profile: Clavaviridae. J Gen Virol 100:1267-1268.
215	20.	Mochizuki T, Yoshida T, Tanaka R, Forterre P, Sako Y, Prangishvili D. 2010. Diversity
216		of viruses of the hyperthermophilic archaeal genus Aeropyrum, and isolation of the
217		Aeropyrum pernix bacilliform virus 1, APBV1, the first representative of the family
218		Clavaviridae. Virology 402:347-54.
219	21.	Prangishvili D, Rensen E, Mochizuki T, Krupovic M, ICTV Report Consortium. 2019.
220		ICTV Virus Taxonomy Profile: Tristromaviridae. J Gen Virol 100:135-136.
221	22.	Iranzo J, Koonin EV, Prangishvili D, Krupovic M. 2016. Bipartite network analysis of
222		the archaeal virosphere: evolutionary connections between viruses and capsidless mobile
223		elements. J Virol 90:11043-11055.

Z

23.

225		Structure of a filamentous virus uncovers familial ties within the archaeal virosphere.
226		Virus Evol 6:veaa023.
227	24.	Rensen EI, Mochizuki T, Quemin E, Schouten S, Krupovic M, Prangishvili D. 2016. A
228		virus of hyperthermophilic archaea with a unique architecture among DNA viruses. Proc
229		Natl Acad Sci U S A 113:2478-83.
230	25.	Baquero DP, Contursi P, Piochi M, Bartolucci S, Liu Y, Cvirkaite-Krupovic V,
231		Prangishvili D, Krupovic M. 2020. New virus isolates from Italian hydrothermal
232		environments underscore the biogeographic pattern in archaeal virus communities. ISME
233		J 14:1821-1833.
234	26.	Bautista MA, Black JA, Youngblut ND, Whitaker RJ. 2017. Differentiation and structure
235		in Sulfolobus islandicus rod-shaped virus populations. Viruses 9:120.
236	27.	Prangishvili D, Arnold HP, Gotz D, Ziese U, Holz I, Kristjansson JK, Zillig W. 1999. A
237		novel virus family, the Rudiviridae: Structure, virus-host interactions and genome
238		variability of the sulfolobus viruses SIRV1 and SIRV2. Genetics 152:1387-96.
239	28.	Prangishvili D, Koonin EV, Krupovic M. 2013. Genomics and biology of rudiviruses, a
240		model for the study of virus-host interactions in Archaea. Biochem Soc Trans 41:443-50.
241	29.	Martínez-Alvarez L, Bell SD, Peng X. 2016. Multiple consecutive initiation of
242		replication producing novel brush-like intermediates at the termini of linear viral dsDNA
243		genomes with hairpin ends. Nucleic Acids Res 44:8799-8809.
244	30.	Pina M, Basta T, Quax TE, Joubert A, Baconnais S, Cortez D, Lambert S, Le Cam E,
245		Bell SD, Forterre P, Prangishvili D. 2014. Unique genome replication mechanism of the
246		archaeal virus AFV1. Mol Microbiol 92:1313-25.

Wang F, Baquero DP, Su Z, Osinski T, Prangishvili D, Egelman EH, Krupovic M. 2020.

12

 \sum

247	31.	Krupovic M, Dolja VV, Koonin EV. 2020. The LUCA and its complex virome. Nat Rev
248		Microbiol 18:661-670.
249	32.	Ptchelkine D, Gillum A, Mochizuki T, Lucas-Staat S, Liu Y, Krupovic M, Phillips SEV,
250		Prangishvili D, Huiskonen JT. 2017. Unique architecture of thermophilic archaeal virus
251		APBV1 and its genome packaging. Nat Commun 8:1436.
252	33.	Janekovic D, Wunderl S, Holz I, Zillig W, Gierl A, Neumann H. 1983. TTV1, TTV2 and
253		TTV3, a family of viruses of the extremely thermophilic, anaerobic, sulfur reducing
254		archaebacterium Thermoproteus tenax. Mol Gen Genet 192:39-45.

255

 \sum

Journal of Virology

