



Lockdown impact on COVID-19 epidemics in regions across metropolitan France

Simon Cauchemez, Cécile Tran Kiem, Juliette Paireau, Patrick Rolland, Arnaud Fontanet

► To cite this version:

Simon Cauchemez, Cécile Tran Kiem, Juliette Paireau, Patrick Rolland, Arnaud Fontanet. Lockdown impact on COVID-19 epidemics in regions across metropolitan France. *The Lancet*, 2020, 396 (10257), pp.1068-1069. 10.1016/S0140-6736(20)32034-1 . pasteur-03326010

HAL Id: pasteur-03326010

<https://pasteur.hal.science/pasteur-03326010>

Submitted on 27 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lockdown impact on COVID-19 epidemics in regions across metropolitan France

Simon Cauchemez, PhD¹, Cécile Tran Kiem, MSc^{1,2}, Juliette Paireau, PhD^{1,3}, Patrick Rolland, MSc³, Arnaud Fontanet, Prof^{4,5}

1. Mathematical Modelling of Infectious Diseases Unit, Institut Pasteur, UMR2000, CNRS, Paris, France.
2. Collège Doctoral, Sorbonne Université, Paris, France.
3. Santé Publique France, French National Public Health Agency, Saint-Maurice, France.
4. Epidemiology of Emerging Diseases Unit, Institut Pasteur, Paris, France.
5. PACRI unit, Conservatoire National des Arts et Métiers, Paris, France

Correspondence:

Simon Cauchemez

Mathematical Modelling of Infectious Diseases Unit

Institut Pasteur

28 rue du Dr Roux

75015 Paris

France

Word count: 390

Lockdowns have been used by most European countries in response to the COVID-19 pandemic. In France, a national lockdown was implemented on 17 March 2020. Some have questioned the need for a nationwide implementation given that most hospitalizations were concentrated in two of thirteen regions; others have even questioned the impact of the lockdown on SARS-CoV-2 spread, arguing that the natural epidemic peak was about to be reached.

On 17 March, daily hospital admissions were indeed highest in Grand-Est (5.3 per 100,000 inhabitants) and Île-de-France (3.6 per 100,000 inhabitants) regions. Yet, a surge in COVID-19 hospital admissions was occurring at that time across all regions of metropolitan France (Figure 1). The COVID-19 epidemic spread from the Eastern to the Western parts of France, crossing the daily hospitalization threshold of 1 per 100,000 inhabitants between 10 (Grand-Est) and 23 March (Bretagne and Nouvelle-Aquitaine). Île-de-France (Paris region) experienced the highest rate of hospitalization per day (10.0 per 100,000 inhabitants), and Bretagne the lowest (1.3 per 100,000 inhabitants). Regardless of the time the epidemic started in the region, and its scale, 12 out of 13 regions experienced a peak in daily hospitalizations 11 +/- 3 days after the lockdown was implemented. This figure corresponds to the mean duration between infection and hospitalization for the patients experiencing severe forms of disease¹. Since the different regions were at different stages of the pandemic at the time the lockdown was implemented, the synchrony in regional peaks strongly suggests that the lockdown, rather than the natural course of the epidemic, explains the peak in hospitalizations. Moreover, most regions were experiencing exponential growth in hospitalizations (Figure 1), such that saturation of local intensive care units might have occurred in those regions in the absence of any lockdown.

Lockdown therefore appears to have been successful not only in alleviating the burden on the intensive care units of the two most severely affected regions of France, but also in preventing uncontrolled epidemics in other regions. These simple observations support results from other studies which have estimated the impact of lockdown on SARS-CoV-2 spread to be strong²⁻⁵. Enforcement of public health and social measures in combination with important testing, tracing, and isolating capacities will be critical in case of an epidemic rebound to avoid re-introducing a lockdown – a situation for which the economic cost and broader impact on society are considerable.

Data sharing statement: The data are further described in Salje et al³ and are available at the following link <https://doi.org/10.5281/zenodo.3889894>.

Author contributions and conflicts of interest: The funders had no role in the writing of the letter. SC had access to all the data and final responsibility for the decision to submit for publication. SC, CTK, JP, PR, AF wrote the letter. SC and AF are members of the French COVID19 Scientific Council of the French Ministry of Health. Other authors declare no conflict of interest.

Acknowledgements: We acknowledge financial support from the Investissement d’Avenir program, the Laboratoire d’Excellence Integrative Biology of Emerging Infectious Diseases program (grant ANR-10-LABX-62-IBEID), the INCEPTION project (PIA/ANR-16-CONV-0005), and the European Union’s Horizon 2020 research and innovation program under grant 101003589 (RECOVER).

References:

1. Bi Q, Wu Y, Mei S, et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis* 2020.
2. Flaxman S, Mishra S, Gandy A, et al. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature* 2020.
3. Salje H, Tran Kiem C, Lefrancq N, et al. Estimating the burden of SARS-CoV-2 in France. *Science* 2020.
4. Dehning J, Zierenberg J, Spitzner FP, et al. Inferring change points in the spread of COVID-19 reveals the effectiveness of interventions. *Science* 2020.
5. Okell LC, Verity R, Watson OJ, et al. Have deaths from COVID-19 in Europe plateaued due to herd immunity? *Lancet* 2020; **395**(10241): e110-e1.

Figure Legend:

Figure 1: Daily hospital admissions of COVID-19 patients per 100,000 inhabitants in the 13 regions of metropolitan France (SI-VIC database). All COVID-19 patients are either biologically confirmed or present with a computed tomographic image highly suggestive of SARS-CoV-2 infection. The grey lines correspond to raw data. The purple lines correspond to the 7-day moving average. The horizontal black dotted lines indicate a level of 1 daily hospital admissions per 100,000 inhabitants. The vertical orange lines indicate the day the nationwide lockdown was enforced. The vertical blue lines indicate the national peak in hospital admissions. The vertical red dotted lines indicate the regional peaks in hospital admissions. Abbreviations for the names of the regions: ARA – Auvergne-Rhône-Alpes ; BFC – Bourgogne-Franche-Comté ; BRE – Bretagne ; CVL – Centre-Val de Loire ; COR – Corse ; GES – Grand Est ; HDF – Hauts-de-France ; IDF – Île-de-France ; NAQ – Nouvelle-Aquitaine ; NOR – Normandie ; OCC – Occitanie ; PAC – Provence-Alpes-Côte d’Azur ; PDL – Pays de la Loire.

