



Identification of bipotent progenitors that give rise to myogenic and connective tissues in mouse

Alexandre Grimaldi, G. Comai, Sébastien Mella, Shahragim Tajbakhsh

► To cite this version:

Alexandre Grimaldi, G. Comai, Sébastien Mella, Shahragim Tajbakhsh. Identification of bipotent progenitors that give rise to myogenic and connective tissues in mouse. 2021. pasteur-03386399v1

HAL Id: pasteur-03386399

<https://pasteur.hal.science/pasteur-03386399v1>

Preprint submitted on 19 Oct 2021 (v1), last revised 12 Mar 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Identification of bipotent progenitors that give rise to myogenic and connective tissues in mouse

Alexandre Grimaldi^{1,2}, Glenda Comai^{1,2}, Sébastien Mella^{1,2}, Shahragim Tajbakhsh^{1,2,*}

¹Stem Cells & Development Unit, 25 rue du Dr. Roux, Institut Pasteur, 75015 Paris, France

²UMR CNRS 3738, Institut Pasteur, Paris, France

*corresponding author

ABSTRACT

How distinct cell fates are manifested by direct lineage ancestry from bipotent progenitors, or by specification of individual cell types within a field of cells is a key question for understanding the emergence of tissues. The interplay between skeletal muscle progenitors and associated connective tissue cells provides a model for examining how muscle functional units are established. Most craniofacial structures originate from the vertebrate-specific neural crest cells except in the dorsal portion of the head, where they arise from cranial mesoderm. Here, using multiple lineage-traced single cell RNAseq, advanced computational methods and in situ analyses, we identify Myf5⁺ bipotent progenitors that give rise to both muscle and juxtaposed connective tissue. Following this bifurcation, muscle and connective tissue cells retain complementary signalling features and maintain spatial proximity. Interruption of upstream myogenic identity shifts muscle progenitors to a connective tissue fate. Interestingly, Myf5-derived connective tissue cells, which adopt a novel regulatory signature, were not observed in ventral craniofacial structures that are colonised by neural crest cells. Therefore, we propose that an ancestral program gives rise to bifated muscle and connective tissue cells in skeletal muscles that are deprived of neural crest.

INTRODUCTION

Stromal cells that are associated with skeletal muscles play critical roles in providing structural support and molecular cues (Biferali et al., 2019; Kardon et al., 2003; Sefton and Kardon, 2019). The majority of muscle-associated connective tissues in the head is derived from cranial neural crest cells (NCCs), an embryonic cell population that contributes to most of the structural components of the "new head", a vertebrate innovation (Douarin and Kalcheim, 1999; Gans and Northcutt, 1983; Grenier et al., 2009; Heude et al., 2018; Noden and Trainor, 2005). Recently, the extent of this contribution was redefined in muscles derived from cranial mesoderm, including extraocular (EOM), laryngeal and pharyngeal muscles (Comai et al., 2020; Grimaldi et al., 2015; Heude et al., 2018; Noden and Epstein, 2010). Interestingly, these muscles contain mesenchyme that is mesoderm-derived in their dorso-medial component, whereas the remaining muscle mass is embedded in mesenchyme that is neural crest-derived. It is unclear how the coordinated emergence of myogenic and connective tissues takes place during development, and how they establish long lasting paracrine communication.

Along the trunk axis, paraxial somitic mesoderm gives rise to skeletal muscles and associated connective tissues (Burke and Nowicki, 2003). Upon signals emanating from the neural tube, notochord, ectoderm and lateral plate mesoderm, the dermomyotome (dorsal portion of the somite) undergoes an epithelial-to-mesenchymal transition and gives rise to several cell types including all skeletal muscles of the body, vasculature, tendons and bones (Ben-Yair and Kalcheim, 2008; Christ et al., 2007). Similarly, cranial mesodermal progenitors give rise to these diverse cell types yet, the unsegmented nature of head mesoderm raises the question of how spatiotemporal control of these cellular identities is established. Moreover, cardiopharyngeal mesoderm, which constitutes the major portion of cranial mesoderm, has cardiovascular potential, which manifests in the embryo as regions of clonally related cardiac and craniofacial skeletal muscles (Diogo et al., 2015; Swedlund and Lescroart, 2019). This skeletal muscle/cardiac branchpoint has been the subject of intense investigation in several model organisms including ascidians, avians, and mouse (Wang et al., 2019). However, the issue of connective tissue divergence from this or another lineage has not been addressed.

Recently, advanced pipelines integrating scRNAseq data and modern algorithms have been instrumental for identifying new lineage relationships during development (Cao et al., 2019; He et al., 2020). Here, we employed unbiased and lineage-restricted single-cell transcriptomics using multiple transgenic mouse lines combined with new computational methods, in situ labelling and loss of function experiments, and show that bipotent progenitors expressing the muscle determination gene *Myf5* give rise to skeletal muscle and anatomically associated connective tissues. Notably, this property was restricted to muscles with partial contribution from NCCs, suggesting that in the absence of NCCs, cranial mesoderm acts as a source of connective tissue.

RESULTS

Myogenic and non-myogenic mesodermal populations coexist within distinct head lineages

Somitic (*Pax3*-dependent) and cranial (*Tbx1/Pitx2*-dependent) mesoderm give rise to diverse cell types including those of the musculoskeletal system (Figure 1A). We first set out to explore the emergence of skeletal muscles and their associated mesodermal tissue within these programs. To that end, we employed a broad anterior mesoderm lineage-tracing strategy using *Mesp1*^{Cre/+}; *R26*^{mTmG/+} mouse embryos at E10.5 when craniofacial skeletal muscles are being established (Heude et al., 2018). The upper third (anterior to forelimb) of the embryos was dissected, the live GFP+ cells were isolated by FACS and processed for scRNAseq analysis (Figure S1A, C-D). After removal of doublets and lower quality cells (see Methods), a large portion of the cells obtained corresponded to adipogenic, chondrogenic, sclerotomal, endothelial, and cardiovascular cells (Figure 1B, Figure S2A-B). *Pax3*, *Pitx2*, *Tbx1*, *Myf5* and *Myod* expression distinguished clusters containing the cranial myogenic progenitors (Figure 1C, Figure S2A).

After subsetting these clusters, a few subclusters clearly separated based on their origin and anatomical location (Figure 1D-E, Figure S2C). Surprisingly, about half of the cells exhibited a connective tissue signature, including a strong bias towards *Prrx1*, a marker for lateral plate mesoderm (Durland et al., 2008), *Col1a1*, a major extracellular matrix component of connective cells (Micheli et al., 2020), and *Twist1*, previously reported to confer mesenchymal properties to cranial mesoderm (Bildsoe et al., 2016) (Figure 1F). Furthermore, the expression of *Pdgfra*, a well-defined marker of stromal cells (Farahani and Xaymardan, 2015), was robustly anticorrelated with the expression of its ligand *Pdgfa*, and associated with non-myogenic genes. Conversely, *Pdgfa* expression correlated with a myogenic cell state (Figure 1F, Figure S2D). Myogenic *Pdgfa* expression was shown to promote adjacent sclerotomal cells to adopt a rib cartilage fate (Tallquist et al., 2000). Therefore, this analysis identified anatomically distinct muscle and closely-associated connective tissue progenitors and highlights a potential PDGFR-mediated crosstalk between these 2 cells types.

Transcriptional trajectories reveal a myogenic to non-myogenic cell state transition

To understand the lineage relationship between myogenic and non-myogenic cells, we exploited the unspliced and spliced variants of our scRNAseq data, and computed the RNA velocity in each cell, using a recently described tool (Bergen et al., 2020) (Figure 2, Figure S3). RNA velocity interrogates the relative abundance of unspliced and spliced gene variants, which depends on the rates of transcription, degradation, and splicing to infer directional trajectories (Bergen et al., 2020; Manno et al., 2018). The cell cycle status constitutes a potential bias in scRNAseq data, especially when heterogeneous populations undergo cellular expansion, commitment and differentiation (McDavid et al., 2016). To eliminate this potential bias, cell cycle genes were consistently regressed out during preprocessing and directional trajectories were overlaid with cell cycle phase visualization for comparisons (Figure S3A, Methods). Notably, RNA velocity-inferred trajectories indicated that *Myf5*+ cells from the myogenic compartment contributed to non-myogenic cells (Figure 2A). These calculations were based on gene-

and cluster-specific dynamics, which yield higher accuracy than the initially described RNA velocity method, while providing quantitative metrics for quality control (Figure S3B and Methods).

Another powerful feature of this method is the ability to infer “driver genes” that are responsible for most of the calculated RNA velocity, hence actively transcribed, or repressed (Bergen et al., 2020). Therefore, these genes can identify transitory states underlying cell fate decisions. We used this approach to uncover the driver genes that were responsible for the velocity found in anterior somites, as these cells displayed the most consistent directionality, and appeared to be independent of cell cycle (Figure 2B, Figure S3A-B, Table1). Top transcribed driver genes included *Foxp1* (Shao and Wei, 2018), *Meox2* (Noizet et al., 2016), *Meis1* (López-Delgado et al., 2020), *Twist2* (Franco et al., 2009), *Fap* (Puré and Blomberg, 2018), *Pdgfra* (Tallquist et al., 2000), *Prrx1* (Leavitt et al., 2020) and *Pcolce* (Bildsoe et al., 2016), that are associated with fibrosis and connective tissue development (Figure 2C). Interestingly, we noted that *Pdgfra* appeared as a driver gene and was activated along this inferred trajectory, whereas *Pdgfa* expression decreased rapidly (Figure 2D). Taken together, RNA velocity analysis for anterior somite mesodermal progenitors showed that *Myf5*⁺/*Pdgfa*⁺ cells shifted towards a non-myogenic fate, by downregulating these 2 markers and activating *Pdgfra* expression (Figure 2E).

***Myf5*-derived lineage contributes to connective tissue cells in the absence of neural crest**

Given that the number of cells examined in the EOM and pharyngeal arch mesodermal clusters from the E10.5 dataset was lower than for anterior somites, we decided to validate the relevance of *Myf5*-derived non-myogenic cells in these cranial regions directly in vivo. We thus examined the EOM, larynx and upper back muscles in the early fetus at E14.5 using a *Myf5*-lineage reporter mouse (*Myf5*^{Cre/+}; *R26*^{TdTomato/+}) combined with a contemporary reporter for *Pdgfra*⁺ non-myogenic cells (*Pdgfra*^{H2BGFP/+}) (Figure 3). Notably, we observed tdTom/H2BGFP double-positive cells in regions of EOM, laryngeal and upper back muscles that were partially or fully deprived of neural crest (Adachi et al., 2020; Comai et al., 2020; Heude et al., 2018) (Figure 3A-C'). Conversely, no double-positive cells were detected in muscles that are fully embedded in neural crest derived connective tissues such as mandibular and tongue muscles (Heude et al., 2018) (Figure 3D-E').

Mesenchymal tissue that is associated with the EOM arises from mesoderm in its most dorso-medial portion and from neural crest in its ventro-lateral portion (Comai et al., 2020). This dual origin makes it a prime candidate to explore the relative contribution of *Myf5*-derived cells to the associated connective tissues within a single functional unit. Using *Wnt1*^{Cre/+}; *R26*^{mTmG/+}; *Myf5*^{nlacZ/+} (NCC tracing) and *Mesp1*^{Cre/+}; *R26*^{mTmG/+}; *Myf5*^{nlacZ/+} (mesoderm tracing) at E13.5, we found that *Myf5*-expressing cells (assessed by β-gal expression) were exclusively in *Mesp1*-derived domains and absent from the *Wnt1* lineage (Figure 3F-I'). By E14.5, we observed a medio-lateral gradient of *Myf5*-lineage contribution to EOM associated connective tissues, and this was anticorrelated with the local contribution of neural crest cells to connective tissues (Figure 3J-K).

In agreement with our scRNAseq velocity analysis, these observations suggest that the mesodermal *Myf5*-lineage compensates for the lack of muscle-associated connective tissue in domains that are deprived of neural crest.

***Myf5*-derived cells can maintain a molecular crosstalk following bifurcation in cell fate**

To investigate in more detail potential paracrine cell-cell communication between myogenic and non-myogenic cells, we examined their signalling complementarity together with their anatomical proximity. We first generated an E11.5 *Myf5*-lineage traced (*Myf5*^{Cre/+}; *R26*^{mTmG/+}) sc-RNAseq dataset (Figure S1A, E-F, Figure S3A, C) and focused on the EOM, which was clearly distinguished as an independent cluster based on the co-expression of several markers including *Pitx2* and *Alx4* (Bothe and Dietrich, 2006) (Figure 4A). Here, RNA velocity revealed strong myogenic/non-myogenic bi-directional cell-fate transitions (Figure 4B). In agreement with the E10.5 mesodermal sc-RNAseq dataset, EOM progenitors presented a strong dichotomy in *Pdgfa* and *Pdgfra* expression between myogenic and non-myogenic cells, respectively. To confirm if the potential crosstalk between these clusters is maintained at later stages of EOM development, we interrogated the anatomical proximity of these cells by performing single molecule fluorescent in situ hybridization (RNAscope) for *Pdgfa* and *Pdgfra* on E14.5 lineage-traced *Myf5*^{Cre/+}; *R26*^{mTmG/+} fetuses (Figure 4C-D'). In accordance with the scRNAseq analysis, we observed a complementary pattern of expression of *Pdgfa* (membrane GFP+) and *Pdgfra* (membrane GFP-) transcripts (Figure 4C-D').

Gene set enrichment analysis of EOM myogenic and non-myogenic driver genes revealed that transmembrane receptor protein kinase and SMAD activity were shared terms between the 2 clusters, indicating that specific complementary signalling networks could be actively maintained between these two populations (Figure S4A). Therefore, we examined the dynamic induction of tyrosine kinase ligands and receptors in the EOM. As observed in the scRNAseq dataset of anterior somites (Figure 2) and by RNAscope on tissue sections at the EOM level (Figure 4C-D'), *Pdgfra* expression was actively induced in non-myogenic *Myf5*-derived cells while *Pdgfa* expression was found in myogenic cells (Figure 4E-F). Notably, two additional tyrosine kinase receptors, namely *Bmpr1b* and *Ephb1*, were found to be among the top 100 driver genes of the myogenic EOM compartment, indicating that myogenic commitment is associated with upregulation of these receptors in the EOM (Figure 4F, Table1). Strikingly, two of their respective ligands *Bmp4* and *Efnb1*, were found to be specifically expressed in non-myogenic cells (Figure 4E). These results favor a model where complementary paracrine signalling networks operates between myogenic and non-myogenic *Myf5*-derived cells (Figure 4G), while their cellular juxtaposition is maintained through fetal stages.

Obstructing myogenesis expands connective tissue formation from bipotent cells

The directional trajectories identified by RNA velocity in the EOM at E11.5 showed a strong bipolarity in fate with a higher velocity confidence index at each end of the myogenic and non-myogenic domains, and lower at their interface (Figure S4B). This suggests that the anticipated cell fate is bipotential before cell fate bifurcation. Conversely, cells that were located on either side of this central region were

identified with greater confidence as committed either to a myogenic or a non-myogenic fate (Figure S4B). To identify the regulatory factors underlying this bipotency, we used SCENIC, a regulatory network inference algorithm (Aibar et al., 2017). This tool allows regrouping of sets of correlated genes into regulons (i.e. a transcription factor and its targets) based on binding motifs and co-expression. Use of this pipeline significantly reduced the number of variables from thousands of genes to a few hundred regulatory modules, while preserving the general aspect of the data, particularly the bipartite distribution of myogenic and non-myogenic cells (Figure 5A). The top regulons of this analysis revealed active transcription factors underlying myogenic and non-myogenic cell fates in the EOM at E11.5. Notably, *Myf5*, *Pitx1*, *Mef2a* and *Six1*, transcription factors known to be implicated in myogenic development (Buckingham and Rigby, 2014), appeared among the top regulons in myogenic cells whereas *Fli1*, *Ebf1*, *Ets1*, *Foxc1*, *Meis1* and *Six2*, genes known for their involvement in adipogenic, vascular, mesenchymal and tendon development (Jimenez et al., 2006; López-Delgado et al., 2020; Noizet et al., 2016; Truong and Ben-David, 2000; Whitesell et al., 2019; Yamamoto-Shiraishi and Kuroiwa, 2013), constituted some of the highly active non-myogenic transcription factors (Figure 5B). Given that *Myf5* appeared as a top regulatory factor of the myogenic program, we interrogated the fate of *Myf5*-expressing progenitors in a *Myf5^{nlacZ/nlacZ}* null genetic background. Interestingly, some β -gal⁺ cells were found in the cartilage primordium (Sox9⁺) of the EOM in the heterozygous control at E12.5 indicating that cells with recent *Myf5* activity diverged to a non-myogenic fate (Figure 5C-C'). As previously reported, the extraocular muscles are absent in this mutant (Figure 5D, asterisk) (Sambasivan et al., 2009). Notably, disruption of *Myf5* activity led to a 3-fold increase in the proportion of non-myogenic *Myf5*-derived cells in this region (Figure 5E). In contrast, no double-positive cells were found in the masseter, a muscle fully embedded in neural crest, even in the absence of *Myf5* (Figure 5E). Robust *Myf5* expression is thus necessary to maintain a balance between myogenic and non-myogenic cell fates of *Myf5*⁺ bipotent progenitors only in neural crest-depleted regions. Conversely, virtually no *Pdgfra*⁺ cells were found to be derived from *Myod* expressing cells in most muscles of *Myod^{lCre};R26^{TdTomato/+};Pdgfra^{H2BGFP/+}* fetuses at E14.5, and only rare double positive cells were found in the EOM (Figure S5). This observation indicates that bipotency is associated with *Myf5*, and that subsequent activation of *Myod* within this lineage locks cell fate into the myogenic program thereby suppressing their connective tissue potential (Figure 5F).

***Myf5*-derived contribution to connective tissues is sustained through muscle initiation**

Although we identified *Myf5*-derived non-myogenic cells in various regions of the embryo, it was not clear if this population was self-sustaining, or continuously generated throughout development. To address this issue, we performed 2 more scRNAseq experiments at E12.5 and E14.5, using contemporary *Myf5* labelling (*Myf5^{GFP-P/+}*; Figures 6, S1B, G-J, S3A, D-E). In accordance with the earlier datasets, cells that appeared to belong to muscle anlagen of EOM, somites and caudal arches progressed towards a non-myogenic state (Figure 6A-C'). To assess the identity of these cells, we performed a gene set enrichment network analysis combining the differentially expressed genes of non-myogenic clusters of all stages. We found that all stages contributed equally to "GO Molecular Function" and "Reactome pathways" terms in spite of their relatively diverse gene expression signatures (Figure

S6). This finding suggests that these non-myogenic cells are relatively homogeneous in gene signatures throughout cranial muscles when they emerge from common bipotent progenitors. Highly significant terms hinted at a myogenic-supporting role, providing muscle progenitors with extracellular matrix components, and contributing to neuronal guidance (Figure 6E). Among these terms, presence of Pdgf signalling and receptor kinase activity indicated, once again, that the interactions found in the EOM could occur also at later stages in various craniofacial muscles that are deprived of neural crest derived connective tissue.

A novel regulatory network underlies the non-myogenic cell fate

Myf5+ bipotent progenitors were observed at multiple stages and anatomical locations, and they yielded a relatively homogeneous population expressing common markers associated with extracellular matrix components, cell adhesion molecules, and tyrosine kinase signalling. To assess whether the regulatory mechanisms guiding this transition are distinct in different locations in the head, we set out to explore the common molecular switches underlying cell fate decisions. To do so, we developed a pipeline where we combined the list of driver genes at the start of the non-myogenic trajectory (Table 1) with the most active regulons in the non-myogenic region (Methods, code in open access). This resulted in a network consisting of the most active transcription factors and the most transcriptionally dynamic genes found at the non-myogenic branchpoint. We performed this operation for each dataset independently and displayed them as individual networks (Figure 7A-D). Finally, we overlapped the list of these “driver regulators” to identify the common transcription factors guiding the non-myogenic cell fate decision (Figure 7E). Notably, *Foxp2*, *Hmga2*, *Meis1*, *Meox2* and *Tcf7l2* were identified in all 4 datasets as key driver regulators, and thus are likely to play significant role in the non-myogenic transition (Figure 7E, Table 2).

Interestingly, Tcfs and Lef1 were among these top common regulators and they form a complex effector for the canonical Wnt pathway. Previous work showed that during cranial myogenesis, neural crest cells release inhibitors of the Wnt pathway to promote myogenesis (Tzahor et al., 2003). It is thus tempting to speculate that in the absence of neural crest, mesoderm-derived bipotent progenitors can give rise to connective tissue by maintaining canonical Wnt activity. To assess this hypothesis, we examined the expression of Axin2, a common readout for Wnt/ β -cat activity (Babb et al., 2017; Moosdijk et al., 2020). Interestingly, Axin2 levels were elevated in the non-myogenic portion of all the different datasets (Figure 7F-I). Additionally, Dkk2, which has been described as an activator of Wnt/ β -cat pathway in the neural crest (Devotta et al., 2018), was also found to be elevated, indicative of a putative positive-feedback loop mechanism supporting the maintenance of this population.

DISCUSSION

Distinct fates can emerge through the specification of individual cell types, or through direct lineage ancestry from bipotent or multipotent cells. Here we addressed this issue in the context of the emergence of myogenic and associated connective tissue cells during the formation of craniofacial muscles. By combining state-of-the-art analytical methods, we identified the transcriptional dynamics, the intercellular communication networks, and the regulators controlling the balance between two complementary cell fates. Specifically, our work provides evidence for a novel mesoderm-derived bipotent cell population that gives rise to muscle and associated connective tissue cells spatiotemporally, and only in regions deprived of neural crest cells (Figure S7).

Brown adipocytes, neurons, pericytes and rib cartilage have been reported to express *Myf5* in ancestral cells (Daubas et al., 2000; Haldar et al., 2008; Sebo et al., 2018; Stuelsatz et al., 2014). Interestingly, when *Myf5* expression is disrupted, cells can acquire non-myogenic fates and contribute to connective tissue (this study), cartilage, and dermis (Tajbakhsh et al., 1996). These studies suggest that *Myf5*-expression alone is not sufficient to promote robust myogenic fate in multiple regions of developing embryos. Consistent with these observations, *Myod*⁺ cells do not contribute to rib cartilage (Wood et al., 2020) and only give rise to very few connective tissue cells in the periocular region (this study). These findings are also consistent with the higher chromatin-remodelling capacity of *Myod* compared to *Myf5*, and its role as defining the committed myogenic cell state (Conerly et al., 2016; Tapscott, 2005). In contrast to a previous study (Stuelsatz et al., 2014), we found no neural-crest derived cells expressing *Myf5* during EOM tissueogenesis at E13.5 (using *Wnt1*^{Cre/+}; *R26*^{mTmG/+}; *Myf5*^{nlacZ/+}). We note that *Myf5*-expressing cells contribute to non-myogenic cells from early embryonic stages (E10.5) and continue to do so in the fetus, indicating that these bipotent cells persist well after muscles are established.

Here, we also identified a core set of transcription factors specifically active in the non-myogenic cells across all datasets. We propose that these genes guide bipotent cells to a non-myogenic fate and thus confer mesenchymal properties to non-committed progenitors. Recent studies have identified anatomically distinct fibroblastic populations using single-cell transcriptomics, yet unique markers could not be identified (Muhl et al., 2020; Sacchetti et al., 2016), making characterisation of cell subtypes challenging. *Tcf4/Tcf7l2* was identified as a master regulator of fibroblastic fate during muscle-associated connective tissue development although it is also expressed in myogenic progenitors at lower levels (Kardon et al., 2003; Mathew et al., 2011; Sefton and Kardon, 2019). We also report that this gene is one of the main regulators of connective tissue fate. Other transcription factors have been linked to skin fibroblast fates including *Tcf4*, *Six2*, *Meox2*, *Egr2* and *Foxs1*, and their repression favours a myofibroblastic potential (Noizet et al., 2016). *Six2* and *Meox2* were also identified in our analysis, which raises the question of the shared genetic programs between myofibroblastic cells and fibroblastic cells derived from progenitors primed for myogenesis during development.

Interestingly, *Prrx1*, a marker for lateral plate mesoderm (Durland et al., 2008), was differentially expressed in the connective tissue population at various stages. Although lateral plate mesoderm is identifiable in the trunk, its anterior boundaries in the head are unclear (Prummel et al., 2020). More detailed analyses of *Prrx1*, *Isl1* and *Myf5* lineages need to be carried out to delineate the specific boundaries of each progenitor contribution to cranial connective tissue.

Tyrosine kinase receptors have been implicated in a number of developmental programs for both muscle and associated connective tissues (Arnold et al., 2020; Knight and Kothary, 2011; Olson and Soriano, 2009; Tallquist et al., 2000; Tzahor et al., 2003; Vinagre et al., 2010). For example, the differentiation of fetal myoblasts is inhibited by growth factors Tgf β and Bmp4 (Cossu et al., 2000). Epha7 signalling is expressed in embryonic and adult myocytes and promotes their differentiation (Arnold et al., 2020). Significantly, we noticed a striking and lasting complementary expression of *Pdgfa* and *Pdgfra* throughout embryonic stages, in the myogenic and non-myogenic progenitors respectively. Pdgf ligands emanating from hypaxial myogenic cells under the control of *Myf5* were shown to be necessary for rib cartilage development (Tallquist et al., 2000; Vinagre et al., 2010). Additionally, *Pdgfra* promotes expansion of fibroblasts during fibrosis (Olson and Soriano, 2009). Interestingly, we found that *Pdgfa* expression was reduced in cells expressing high levels of *Myog* at the fetal stage (data not shown). Therefore, *Myf5*-derived myogenic progenitor cells might guide non-myogenic *Myf5*-derived expansion, which in turn provides ligands and extracellular matrix components to favour myogenic development and patterning. Moreover, unlike trunk myogenesis, cranial muscle development relies on the expression of Wnt and Bmp inhibitors from surrounding tissues (Tzahor et al., 2003). Interestingly, we showed that the *Myf5*-derived non-myogenic cells express Bmp4, Dkk2, and Axin2. Additionally, we showed that the Wnt effector complex Tcf/Lef is active in these cells. It is thus likely that these cells maintain their non-myogenic fate by promoting Bmp production and Wnt activity cell-autonomously. Further studies could provide further insights into the evolutionary ancestry of this bipotency by studying other model organisms devoid of neural crest.

MATERIALS & METHODS

scRNAseq data generation

For E10.5 to E12.5 embryos, the cranial region above the forelimb was dissected in ice-cold 3% FBS in PBS and mechanically dissociated with forceps and pipetting. The same procedure was applied at E14.5 but the dissection was refined to the pharyngeal and laryngeal regions. Tissues were then digested in TrypLE (ThermoFisher Cat #: 12604013) during 3 rounds of 5-min incubation (37°C, 1400 RPM), interspersed with gentle pipetting to further dissociate the tissue. Cells were resuspended in FBS 3%, filtered, and incubated with Calcein Blue (eBioscience, Cat #: 65-0855-39) and Propidium Iodide (ThermoFisher Cat #: P1304MP) to check for viability. Viable cells were sorted on BD FACSAria™ III and manually counted using a hemocytometer. RNA integrity was assessed with Agilent Bioanalyzer 2100 to validate the isolation protocol prior to scRNAseq (RIN>8 was considered acceptable). 4000 to 13000 cells were loaded onto 10X Genomics Chromium microfluidic chip and cDNA libraries were

generated following manufacturer's protocol. Concentrations and fragment sizes were measured using Agilent Bioanalyzer and Invitrogen Qubit. cDNA libraries were sequenced using NextSeq 500 and High Output v2.5 (75 cycles) kits. Genome mapping and count matrix generation were done following 10X Genomics Cell Ranger pipeline.

RNA velocity and driver genes

RNA velocity analyses were performed using scvelo (Bergen et al., 2020) in python. This tool allows inferring velocity flow and driver genes using scRNAseq data, with major improvements from previous methods (Manno et al., 2018). First, unspliced and spliced transcript matrices were generated using velocity (Manno et al., 2018) command line function, which outputs unspliced, spliced, and ambiguous matrices as a single loom file. These files were combined with filtered Seurat objects to yield objects with unspliced and spliced matrices, as well as Seurat-generated annotations and cell-embeddings (UMAP, tSNE, PCA). These datasets were then processed following scvelo online guide and documentation. Velocity was calculated based on the dynamical model (using `scv.tl.recover_dynamics(adata)`, and `scv.tl.velocity(adata, mode='dynamical')`) and when outliers were detected, differential kinetics based on top driver genes were calculated and added to the model (using `scv.tl.velocity(adata, diff_kinetics=True)`). Specific driver genes were identified by determining the top likelihood genes in the selected cluster. The lists of top 100 drivers for each stage are given in Table1.

Seurat preprocessing

scRNAseq datasets were preprocessed using Seurat in R (<https://satijalab.org/seurat/>) (Butler et al., 2018). Cells with more than 20% of mitochondrial gene fraction were discarded. The number of genes expressed averaged to 4000 in all 4 datasets. Dimension reduction and UMAP generation were performed following Seurat workflow. Doublets were inferred using DoubletFinder v3 (McGinnis et al., 2019). Cell cycle genes, mitochondrial fraction, number of genes, number of UMI were regressed in all datasets following Seurat dedicated vignette. We noticed that cell cycle regression, although clarifying anatomical diversity, seemed to induce low and high UMI clustering (Figure S1E-F). For the E10.5 and E11.5 datasets, 2 replicates were generated from littermates and merged after confirming their similitude. For subsequent datasets (E12.5 and E14.5), no replicates were used. Annotation and subsetting were also performed in Seurat. "Myogenic" and "Non-myogenic" annotations were based on *Pdgfa* and *Pdgfra* expression and myogenic genes *Myf5*, *Myod*, and *Myog*. Cells not expressing *Pdgfa* were annotated as "non-myogenic" unless they express myogenic genes. Cells expressing *Pdgfa* were annotated as "myogenic". We noticed that at later stages, *Pdgfa* expression decreases in Myog+ cells. Thus, driver genes of connective tissue at E12.5 and E14.5 were determined using cluster annotations obtained from Leiden-based clustering.

Gene regulatory network inference

Gene regulatory networks were inferred using SCENIC (R implementation) (Aibar et al., 2017) and pySCENIC (python implementation) (Sande et al., 2020). This algorithm allows regrouping of sets of correlated genes into regulons (i.e. a transcription factor and its targets) based on motif binding and co-

expression. UMAP and heatmap were generated using regulon AUC matrix (Area Under Curve) which refers to the activity level of each regulon in each cell.

Driver regulons

Results from SCENIC and scvelo were combined to identify regulons that could be responsible for the transcriptomic induction of driver genes. Similar to the steps mentioned above, SCENIC lists of regulons were used to infer connections between transcription factors and driver gene. Networks were generated as explained above, and annotated with “Active regulon” or “driver gene”. The lists of individual driver regulons of each dataset were then combined and the most recurring driver regulons were identified. The code is available at this address: <https://github.com/TajbakhshLab/DriverRegulators>

Gene set enrichment analysis

Gene set enrichment analyses were performed on either the top markers (obtained from Seurat function FindAllMarkers) or from driver genes (obtained from scvelo), using Cluego (Bindea et al., 2009). “GO Molecular Pathway”, “GO Biological Process” and “Reactome pathways” were used independently to identify common and unique pathways involved in each dataset. In all analyses, an enrichment/depletion two-sided hypergeometric test was performed and p-values were corrected using the Bonferroni step down method.

Mouse strains

Animals were handled according to European Community guidelines and the ethics committee of the Institut Pasteur (CETEA) approved protocols. The following strains were previously described: *Myf5^{Cre}* (Haldar et al., 2008), *Mesp1^{Cre}* (Saga et al., 1999), *Tg:Wnt1Cre* (Danielian et al., 1998), *R26^{TdTom}* (Ai9;(Madisen et al., 2009)), *R26^{mTmG}* (Muzumdar et al., 2007), *Myf5^{nlacZ}* (Tajbakhsh et al., 1996), *Pdgfra^{H2BGFP}* (Hamilton et al., 2003) and *Myf5^{GFP-P}* (Kassar-Duchossoy et al., 2004). To generate *Myf5^{Cre/+};R26^{TdTomato/+};Pdgfra^{H2BGFP/+}* embryos, *Myf5^{Cre/+}* females were crossed with *Pdgfra^{H2BGFP/+};R26^{TdTomato/TdTomato}* males. Mice were kept on a mixed genetic background C57BL/6JRj and DBA/2JRj (B6D2F1, Janvier Labs). Mouse embryos and fetuses were collected between embryonic day (E) E10.5 and E14.5, with noon on the day of the vaginal plug considered as E0.5.

Immunofluorescence

Collected embryonic and adult tissues were fixed 2.5h in 4% paraformaldehyde (Electron Microscopy Sciences, Cat #:15710) in PBS with 0,2-0,5% Triton X-100 (according to their stage) at 4°C and washed overnight at 4°C in PBS. In preparation for cryosectioning, embryos were equilibrated in 30% sucrose in PBS overnight at 4°C and embedded in OCT. Cryosections (16-20µm) were left to dry at RT for 30 min and washed in PBS. The primary antibodies used in this study are chicken polyclonal anti-β-gal (Abcam, Cat #: ab9361, dilution 1:1000), mouse monoclonal IgG1, mouse monoclonal IgG1 anti-Myod (BD Biosciences, Cat# 554130, dilution 1:100), mouse monoclonal IgG1 anti-Pax7 (DSHB, Cat. #: AB_528428, dilution 1:20), rabbit anti-mouse Sox9 (Millipore, Cat. #: AB5535, dilution 1/2000), rabbit polyclonal anti-Tomato (Clontech Cat. #: 632496, dilution 1:400) and chicken polyclonal anti-GFP

(Abcam Cat. #: 13970, dilution 1:1000). Images were acquired using Zeiss LSM780 or LSM700 confocal microscopes and processed using ZEN software (Carl Zeiss).

RNAscope in situ hybridization

Embryos for in situ hybridization were fixed overnight in 4% PFA. Embryos were equilibrated in 30% sucrose in PBS and sectioned as described for immunofluorescence. RNAscope probes Mm-Pdgfa (411361) and Mm-Pdgfra (480661-C2) were purchased from Advanced Cell Diagnostics, Inc. In situ hybridization was performed using the RNAscope Multiplex Fluorescent Reagent Kit V2 as described previously (Comai et al., 2019).

Data Availability

The data that support the findings of this study are available from the corresponding author, S.T, upon request. The code that was used to generate the driver regulators is available at this address: <https://github.com/TajbakhshLab/DriverRegulators>

Acknowledgements

We acknowledge funding support from the Institut Pasteur, Association Française contre le Myopathies, Agence Nationale de la Recherche (Laboratoire d'Excellence Revive, Investissement d'Avenir; ANR-10-LABX-73) and MyoHead, Association Française contre les Myopathies (Grant #20510), Fondation pour la Recherche Médicale (Grant # FDT201904008277), and the Centre National de la Recherche Scientifique. We gratefully acknowledge the UtechS Photonic BioImaging, C2RT, Institut Pasteur, supported by the French National Research Agency (France BioImaging; ANR-10-INSB-04; Investments for the Future).

Competing interests

The authors declare no competing interests.

REFERENCES:

Adachi, N., Bilio, M., Baldini, A. and Kelly, R. G. (2020). Cardiopharyngeal mesoderm origins of musculoskeletal and connective tissues in the mammalian pharynx. *Development* **147**, dev185256.

Aibar, S., González-Blas, C. B., Moerman, T., Huynh-Thu, V. A. A., Imrichova, H., Hulselmans, G., Rambow, F., Marine, J.-C. C., Geurts, P., Aerts, J., et al. (2017). SCENIC: single-cell regulatory network inference and clustering. *Nature methods* **14**, 1083–1086.

Arnold, L. L., Cecchini, A., Stark, D. A., Ihnat, J., Craigg, R. N., Carter, A., Zino, S. and Cornelison, D. (2020). EphA7 promotes myogenic differentiation via cell-cell contact. *Elife* **9**, e53689.

Babb, R., Chandrasekaran, D., Neves, V. C. M. and Sharpe, P. T. (2017). Axin2-expressing cells differentiate into reparative odontoblasts via autocrine Wnt/ β -catenin signaling in response to tooth damage. *Sci Rep-uk* **7**, 3102.

- Ben-Yair, R. and Kalcheim, C. (2008). Notch and bone morphogenetic protein differentially act on dermomyotome cells to generate endothelium, smooth, and striated muscle. *J Cell Biol* **180**, 607–618.
- Bergen, V., Lange, M., Peidli, S., Wolf, F. A. and Theis, F. J. (2020). Generalizing RNA velocity to transient cell states through dynamical modeling. *Nat Biotechnol* 1–7.
- Biferali, B., Proietti, D., Mozzetta, C. and Madaro, L. (2019). Fibro–Adipogenic Progenitors Cross-Talk in Skeletal Muscle: The Social Network. *Front Physiol* **10**, 1074.
- Bildsoe, H., Fan, X., Wilkie, E. E., Ashoti, A., Jones, V. J., Power, M., Qin, J., Wang, J., Tam, P. P. L. and Loebel, D. A. F. (2016). Transcriptional targets of TWIST1 in the cranial mesoderm regulate cell-matrix interactions and mesenchyme maintenance. *Dev Biol* **418**, 189–203.
- Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W.-H., Pagès, F., Trajanoski, Z. and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* **25**, 1091–1093.
- Bothe, I. and Dietrich, S. (2006). The molecular setup of the avian head mesoderm and its implication for craniofacial myogenesis. *Developmental Dynamics* **235**, 2845–2860.
- Buckingham, M. and Rigby, P. W. J. (2014). Gene Regulatory Networks and Transcriptional Mechanisms that Control Myogenesis. *Dev Cell* **28**, 225–238.
- Burke, A. C. and Nowicki, J. L. (2003). A New View of Patterning Domains in the Vertebrate Mesoderm. *Dev Cell* **4**, 159–165.
- Butler, A., Hoffman, P., Smibert, P., Papalexi, E. and Satija, R. (2018). Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411–420.
- Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D. M., Hill, A. J., Zhang, F., Mundlos, S., Christiansen, L., Steemers, F. J., et al. (2019). The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**, 496–502.
- Christ, B., Huang, R. and Scaal, M. (2007). Amniote somite derivatives. *Dev Dynam* **236**, 2382–2396.
- Comai, G., Heude, E., Mella, S., Paisant, S., Pala, F., Gallardo, M., Langa, F., Kardon, G., Gopalakrishnan, S. and Tajbakhsh, S. (2019). A distinct cardiopharyngeal mesoderm genetic hierarchy establishes antero-posterior patterning of esophagus striated muscle. *Elife* **8**, e47460.
- Comai, G. E., Tesařová, M., Dupé, V., Rhinn, M., Vallecillo-García, P., Silva, F. da, Feret, B., Exelby, K., Dollé, P., Carlsson, L., et al. (2020). Local retinoic acid signaling directs emergence of the extraocular muscle functional unit. *Plos Biol* **18**, e3000902.
- Conerly, M. L., Yao, Z., Zhong, J. W., Groudine, M. and Tapscott, S. J. (2016). Distinct Activities of Myf5 and MyoD Indicate Separate Roles in Skeletal Muscle Lineage Specification and Differentiation. *Dev Cell* **36**, 375–85.
- Danielian, P. S., Muccino, D., Rowitch, D. H., Michael, S. K. and McMahon, A. P. (1998). Modification of gene activity in mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. *Curr Biol* **8**, 1323–S2.
- Daubas, P., Tajbakhsh, S., Hadchouel, J., Primig, M. and Buckingham, M. (2000). Myf5 is a novel early axonal marker in the mouse brain and is subjected to post-transcriptional regulation in neurons. *Dev Camb Engl* **127**, 319–31.
- Devotta, A., Hong, C.-S. and Saint-Jeannet, J.-P. (2018). Dkk2 promotes neural crest specification by activating Wnt/β-catenin signaling in a GSK3β independent manner. *Elife* **7**, e34404.

- 506 **Diogo, R., Kelly, R. G., Christiaen, L., Levine, M., Ziermann, J. M., Molnar, J. L., Noden, D. M. and**
507 **Tzahor, E.** (2015). A new heart for a new head in vertebrate cardiopharyngeal evolution. *Nature* **520**, 466–
508 73.
- 509 **Douarin, N. L. and Kalcheim, C.** (1999). *The Neural Crest*. 2nd ed., Developmental and Cell Biology Series.
510 Cambridge University Press.
- 511 **Durland, J. L., Sferlazzo, M., Logan, M. and Burke, A. C.** (2008). Visualizing the lateral somitic frontier in
512 the Prx1Cre transgenic mouse. *J Anat* **212**, 590–602.
- 513 **Farahani, R. M. and Xaymardan, M.** (2015). Platelet-Derived Growth Factor Receptor Alpha as a Marker of
514 Mesenchymal Stem Cells in Development and Stem Cell Biology. *Stem Cells Int* **2015**, 362753.
- 515 **Franco, H., Casasnovas, J. J. and Cadilla, C. L.** (2009). Periostin gene expression is regulated by the bHLH
516 transcription factor TWIST2 in human skin fibroblasts. *Faseb J* **23**, 660.11-660.11.
- 517 **Gans, C. and Northcutt, R. G.** (1983). Neural Crest and the Origin of Vertebrates: A New Head. *Science* **220**,
518 268–273.
- 519 **Grenier, J., Teillet, M.-A., Grifone, R., Kelly, R. G. and Duprez, D.** (2009). Relationship between neural
520 crest cells and cranial mesoderm during head muscle development. *Plos One* **4**, e4381.
- 521 **Grimaldi, A., Parada, C. and Chai, Y.** (2015). A Comprehensive Study of Soft Palate Development in Mice.
522 *PloS one* **10**, e0145018.
- 523 **Haldar, M., Karan, G., Tvrdik, P. and Capecchi, M. R.** (2008). Two cell lineages, myf5 and myf5-
524 independent, participate in mouse skeletal myogenesis. *Dev Cell* **14**, 437–45.
- 525 **Hamilton, T. G., Klinghoffer, R. A., Corrin, P. D. and Soriano, P.** (2003). Evolutionary Divergence of
526 Platelet-Derived Growth Factor Alpha Receptor Signaling Mechanisms. *Mol Cell Biol* **23**, 4013–4025.
- 527 **He, P., Williams, B. A., Trout, D., Marinov, G. K., Amrhein, H., Berghella, L., Goh, S.-T., Plajzer-Frick,**
528 **I., Afzal, V., Pennacchio, L. A., et al.** (2020). The changing mouse embryo transcriptome at whole tissue
529 and single-cell resolution. *Nature* **583**, 760–767.
- 530 **Heude, E., Tesarova, M., Sefton, E. M., Jullian, E., Adachi, N., Grimaldi, A., Zikmund, T., Kaiser, J.,**
531 **Kardon, G., Kelly, R. G., et al.** (2018). Unique morphogenetic signatures define mammalian neck muscles
532 and associated connective tissues. *eLife* **7**,.
- 533 **Jimenez, M. A., Åkerblad, P., Sigvardsson, M. and Rosen, E. D.** (2006). Critical Role for Ebf1 and Ebf2 in
534 the Adipogenic Transcriptional Cascade ▽ †. *Mol Cell Biol* **27**, 743–757.
- 535 **Kardon, G., Harfe, B. D. and Tabin, C. J.** (2003). A Tcf4-Positive Mesodermal Population Provides a
536 Prepatter for Vertebrate Limb Muscle Patterning. *Dev Cell* **5**, 937–944.
- 537 **Kassar-Duchossoy, L., Gayraud-Morel, B., Gomès, D., Rocancourt, D., Buckingham, M., Shinin, V. and**
538 **Tajbakhsh, S.** (2004). Mrf4 determines skeletal muscle identity in Myf5:Myod double-mutant mice. *Nature*
539 **431**, 466–471.
- 540 **Knight, J. D. and Kothary, R.** (2011). The myogenic kinome: protein kinases critical to mammalian skeletal
541 myogenesis. *Skelet Muscle* **1**, 29.
- 542 **Leavitt, T., Hu, M. S., Borrelli, M. R., Januszyk, M., Garcia, J. T., Ransom, R. C., Mascharak, S.,**
543 **desJardins-Park, H. E., Litzenburger, U. M., Walmsley, G. G., et al.** (2020). Prrx1 Fibroblasts Represent
544 a Pro-fibrotic Lineage in the Mouse Ventral Dermis. *Cell Reports* **33**, 108356.

- 545 **López-Delgado, A. C., Delgado, I., Cadenas, V., Sánchez-Cabo, F. and Torres, M.** (2020). Axial skeleton
546 anterior-posterior patterning is regulated through feedback regulation between Meis transcription factors and
547 retinoic acid. *Biorxiv* 2020.03.09.983106.
- 548 **Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter, R.**
549 **D., Hawrylycz, M. J., Jones, A. R., et al.** (2009). A robust and high-throughput Cre reporting and
550 characterization system for the whole mouse brain. *Nat Neurosci* **13**, 133–40.
- 551 **Manno, G. L., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber, K., Kastri,**
552 **M. E., Lönnerberg, P., Furlan, A., et al.** (2018). RNA velocity of single cells. *Nature* **560**, 494–498.
- 553 **Mathew, S. J., Hansen, J. M., Merrell, A. J., Murphy, M. M., Lawson, J. A., Hutcheson, D. A., Hansen, M.**
554 **S., Angus-Hill, M. and Kardon, G.** (2011). Connective tissue fibroblasts and Tcf4 regulate myogenesis.
555 *Dev Camb Engl* **138**, 371–84.
- 556 **McDavid, A., Finak, G. and Gottardo, R.** (2016). The contribution of cell cycle to heterogeneity in single-cell
557 RNA-seq data. *Nat Biotechnol* **34**, 591–593.
- 558 **McGinnis, C. S., Murrow, L. M. and Gartner, Z. J.** (2019). DoubletFinder: Doublet Detection in Single-Cell
559 RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337.e4.
- 560 **Micheli, A. J. D., Swanson, J. B., Disser, N. P., Martinez, L. M., Walker, N. R., Oliver, D. J., Cosgrove, B.**
561 **D. and Mendias, C. L.** (2020). Single-cell transcriptomic analysis identifies extensive heterogeneity in the
562 cellular composition of mouse Achilles tendons. *Am J Physiol-cell Ph* **319**, C885–C894.
- 563 **Moosdijk, A. A. A., Grift, Y. B. C., Man, S. M. A., Zeeman, A. L. and Amerongen, R.** (2020). A novel
564 Axin2 knock-in mouse model for visualization and lineage tracing of WNT/CTNNB1 responsive cells.
565 *Genesis* **58**, e23387.
- 566 **Muhl, L., Genové, G., Leptidis, S., Liu, J., He, L., Mocci, G., Sun, Y., Gustafsson, S., Buyandelger, B.,**
567 **Chivukula, I. V., et al.** (2020). Single-cell analysis uncovers fibroblast heterogeneity and criteria for
568 fibroblast and mural cell identification and discrimination. *Nat Commun* **11**, 3953.
- 569 **Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. and Luo, L.** (2007). A global double-fluorescent Cre
570 reporter mouse. *Genesis (New York, N.Y. : 2000)* **45**, 593–605.
- 571 **Noden, D. M. and Epstein, M. L.** (2010). Embryonic origins of avian and mammalian laryngeal
572 musculoskeletal structures. *Faseb J* **24**, 172.1-172.1.
- 573 **Noden, D. M. and Trainor, P. A.** (2005). Relations and interactions between cranial mesoderm and neural crest
574 populations. *Journal of anatomy* **207**, 575–601.
- 575 **Noizet, M., Lagoutte, E., Gratigny, M., Bouschbacher, M., Lazareth, I., Crollius, H. R., Darzacq, X. and**
576 **Dugast-Darzacq, C.** (2016). Master regulators in primary skin fibroblast fate reprogramming in a human ex
577 vivo model of chronic wounds. *Wound Repair Regen* **24**, 247–262.
- 578 **Olson, L. E. and Soriano, P.** (2009). Increased PDGFRalpha activation disrupts connective tissue development
579 and drives systemic fibrosis. *Dev Cell* **16**, 303–13.
- 580 **Prummel, K. D., Nieuwenhuize, S. and Mosimann, C.** (2020). The lateral plate mesoderm. *Dev Camb Engl*
581 **147**, dev175059.
- 582 **Puré, E. and Blomberg, R.** (2018). Pro-tumorigenic roles of fibroblast activation protein in cancer: back to the
583 basics. *Oncogene* **37**, 4343–4357.
- 584 **Sacchetti, B., Funari, A., Remoli, C., Giannicola, G., Kogler, G., Liedtke, S., Cossu, G., Serafini, M.,**
585 **Sampaioles, M., Tagliafico, E., et al.** (2016). No Identical “Mesenchymal Stem Cells” at Different Times

586 and Sites: Human Committed Progenitors of Distinct Origin and Differentiation Potential Are Incorporated
587 as Adventitial Cells in Microvessels. *Stem Cell Rep* **6**, 897–913.

588 **Saga, Y., Miyagawa-Tomita, S., Takagi, A., Kitajima, S., Miyazaki, J. i and Inoue, T.** (1999). MesP1 is
589 expressed in the heart precursor cells and required for the formation of a single heart tube. *Dev Camb Engl*
590 **126**, 3437–47.

591 **Sambasivan, R., Gayraud-Morel, B., Dumas, G., Cimper, C., Paisant, S., Kelly, R. G., Kelly, R. and**
592 **Tajbakhsh, S.** (2009). Distinct regulatory cascades govern extraocular and pharyngeal arch muscle
593 progenitor cell fates. *Developmental cell* **16**, 810–21.

594 **Sande, B. V. de, Flerin, C., Davie, K., Waegeneer, M. D., Hulselmans, G., Aibar, S., Seurinck, R., Saelens,**
595 **W., Cannoodt, R., Rouchon, Q., et al.** (2020). A scalable SCENIC workflow for single-cell gene regulatory
596 network analysis. *Nat Protoc* 1–30.

597 **Sebo, Z. L., Jeffery, E., Holtrup, B. and Rodeheffer, M. S.** (2018). A mesodermal fate map for adipose tissue.
598 *Development* **145**, dev166801.

599 **Sefton, E. M. and Kardon, G.** (2019). Connecting muscle development, birth defects, and evolution: An
600 essential role for muscle connective tissue. *Curr Top Dev Biol* **132**, 137–176.

601 **Shao, X. and Wei, X.** (2018). FOXP1 enhances fibrosis via activating Wnt/ β -catenin signaling pathway in
602 endometriosis. *Am J Transl Res* **10**, 3610–3618.

603 **Stuelsatz, P., Shearer, A. and Yablonka-Reuveni, Z.** (2014). Ancestral Myf5 gene activity in periocular
604 connective tissue identifies a subset of fibro/adipogenic progenitors but does not connote a myogenic origin.
605 *Developmental biology* **385**, 366–79.

606 **Swedlund, B. and Lescroart, F.** (2019). Cardiopharyngeal Progenitor Specification: Multiple Roads to the
607 Heart and Head Muscles. *Csh Perspect Biol* a036731.

608 **Tajbakhsh, S., Rocancourt, D. and Buckingham, M.** (1996). Muscle progenitor cells failing to respond to
609 positional cues adopt non-myogenic fates in myf-5 null mice. *Nature* **384**, 266–270.

610 **Tallquist, M., Weismann, K. and Development, H.-M.** (2000). Early myotome specification regulates PDGFA
611 expression and axial skeleton development.

612 **Tapscott, S. J.** (2005). The circuitry of a master switch: Myod and the regulation of skeletal muscle gene
613 transcription. *Development* **132**, 2685–2695.

614 **Truong, A. H. and Ben-David, Y.** (2000). The role of Fli-1 in normal cell function and malignant
615 transformation. *Oncogene* **19**, 6482–6489.

616 **Tzahor, E., Kempf, H., Mootoosamy, R. C., Poon, A. C., Abzhanov, A., Tabin, C. J., Dietrich, S. and**
617 **Lassar, A. B.** (2003). Antagonists of Wnt and BMP signaling promote the formation of vertebrate head
618 muscle. *Gene Dev* **17**, 3087–3099.

619 **Vinagre, T., Moncaut, N., Carapuço, M., Nóvoa, A., Bom, J. and Mallo, M.** (2010). Evidence for a
620 Myotomal Hox/Myf Cascade Governing Nonautonomous Control of Rib Specification within Global
621 Vertebral Domains. *Dev Cell* **18**, 655–661.

622 **Wang, W., Niu, X., Stuart, T., Jullian, E., Mauck, W. M., Kelly, R. G., Satija, R. and Christiaen, L.** (2019).
623 A single-cell transcriptional roadmap for cardiopharyngeal fate diversification. *Nature Cell Biology* **21**, 674–
624 686.

625 **Whitesell, T. R., Chrystal, P. W., Ryu, J.-R. R., Munsie, N., Grosse, A., French, C. R., Workentine, M. L.,**
626 **Li, R., Zhu, L. J., Waskiewicz, A., et al. (2019).** foxc1 is required for embryonic head vascular smooth
627 muscle differentiation in zebrafish. *Developmental biology* **453**, 34–47.

628 **Wood, W. M., Otis, C., Etemad, S. and Goldhamer, D. J. (2020).** Development and patterning of rib
629 primordia are dependent on associated musculature. *Dev Biol*.

630 **Yamamoto-Shiraishi, Y. and Kuroiwa, A. (2013).** Wnt and BMP signaling cooperate with Hox in the control
631 of Six2 expression in limb tendon precursor. *Dev Biol* **377**, 363–374.

632

633

Figure 1




Figure 1. scRNAseq reveals non-myogenic populations of cranial mesoderm lineages.

- A) Scheme of connective tissue origin in the head and known mesodermal upstream regulators. E: Eye, 1-4: Pharyngeal arches 1-4.
- B) UMAP of *Mesp1*^{Cre/+}; *R26*^{mTmG/+} E10.5 scRNAseq with main cell types highlighted.
- C) UMAP expression plots of *Pitx2* (EOM), *Tbx1* (cranial mesoderm except EOM) and *Pax3* (somitic mesoderm), indicating the clusters of progenitors.
- D) UMAP of progenitor subset annotated as myogenic and non-myogenic based on expression patterns found in E and F.
- E) UMAP expression plots *Pitx2*, *Tbx1* and *Pax3*.
- F) Heatmap of top 20 markers of myogenic versus non-myogenic clusters. *Pdgfra*/*Pdgfa* genes are highlighted.

Figure 2

RNA velocity & Driver genes
(*scvelo*)




Figure 2. Transcriptomic dynamics reveal a myogenic to non-myogenic transition in anterior somite progenitors.

A) Velocity UMAP plots displaying myogenic and non-myogenic clusters. Arrows represent the lineage progression based on RNA velocity (relative abundance of unspliced and spliced transcripts).

B) Heatmap of driver genes accounting for anterior somite velocity, highlighting *Pdgfra*.

C) Phase portraits of few selected driver genes in the anterior somites: *Foxp1*, *Meox2*, *Meis1*, *Twist2*, *Fap*, *Pdgfra*, *Prrx1* and *Pcolce*. Y-axis represents the amount of unspliced transcript per cell; X-axis represents the number of spliced transcripts per cell. A high fraction of unspliced variants indicates an active transcription of the locus, while the inverse indicates inactive/repressed transcription. Dynamics of transcription were inferred at a gene- and cluster-specific level (see Methods).

D) Phase portraits, velocity and expression plots of *Pdgfa* and *Pdgfra* showing splicing dynamics of these 2 genes.

E) Working model of myogenic and non-myogenic fate decision from a common bipotent progenitor in anterior somites.

Figure 3




Figure 3. Myf5-derived mesodermal connective tissue partially compensates for the lack of neural crest.

A-E') Transverse sections of an E14.5 *Myf5*^{Cre/+}; *R26*^{TdTomato/+}; *Pdgfra*^{H2BGFP/+} embryo immunostained for Myod/Myog. White arrowheads indicated cells double-positive GFP/Tomato and negative for Myod/Myog (n=3 embryos).

F-I') Transverse cryosections of the EOM at E13.5 of *Wnt1*^{Cre/+}; *R26*^{mTmG/+}; *Myf5*^{nlacZ/+} (G,I) and *Mesp1*^{Cre/+}; *R26*^{mTmG/+}; *Myf5*^{nlacZ/+} (F,H) immunostained for β-gal, at the level of the medial attachment (F,G) and lateral muscle masses (H,I). Yellow arrowhead indicates *Myf5*-expressing cells in the context of mesodermal and neural crest lineages. (n=2 embryos with 5 tissue sections analyzed per embryo)

J) Quantifications of the proportion of double positive cells in E14.5 *Myf5*^{Cre/+}; *R26*^{TdTomato/+}; *Pdgfra*^{H2BGFP/+} embryo in various regions throughout the EOM (n=3 embryos, with 5 tissue sections analyzed per embryo).

K) Scheme highlighting the quantified regions in (J), and summarising the contribution of each population to connective tissue.

Figure 4

Extraocular muscles




Figure 4. Maintenance of signalling cues between Myf5-derived myogenic and non-myogenic cells in EOM.

- A) UMAPs of *Myf5*^{Cre/+}; *R26*^{mTmG/+} E11.5 EOM subset and the respective RNA velocity trajectories (top). Expression plots of *Pitx2* and *Alx4*, marking the EOM cluster (bottom).
- B) Velocities within myogenic and non-myogenic clusters and expression plots of *Pdgfra* and *Pdgfra*.
- C-D') RNAscope on *Myf5*^{Cre/+}; *R26*^{mTmG/+} E14.5 tissue sections with *Pdgfra* (cyan) and *Pdgfra* (red) probes. *Myf5*-derived cells are labelled by membrane GFP staining.
- D') High magnification of D. Yellow arrowheads indicate *Myf5*-derived *Pdgfra*-expressing cells (non-myogenic). Red arrowheads indicate *Myf5*-derived *Pdgfra*-expressing cells (myogenic).
- E) Expression pattern of ligands *Pdgfra*, *Bmp4* and *Efnb1*.
- F) Phase portrait, gene velocity and expression plots of receptors *Pdgfra*, *Bmpr1b* and *Ephb1*.
- G) Model of myogenic and non-myogenic cell communication following bifurcation

Figure 5




Figure 5. Disruption of *Myf5* increases the connective tissue output from bipotent cells

A) UMAP of *Myf5*^{Cre/+}; *R26*^{mTmG/+} E11.5 EOM based on SCENIC Regulon activity (Area Under Curve score).

B) Heatmap of top regulons (transcription factor and associated targets). The suffix “_extended” indicates that the regulon includes motifs that have been linked to the TF by lower confidence annotations, for instance, inferred by motif similarity. Number in brackets indicates number of genes comprising the regulon.

C-D) Transverse sections of *Myf5*^{nlacZ/+} (C-C'), and *Myf5*^{nlacZ/nlacZ} (D-D') in the EOM region at E12.5 immunostained for β-gal (green), Sox9 (red) and Myod/Myog/Pax7 (grey). Yellow arrowheads indicate β-gal/Sox9 double positive cells and show an expansion these cells in the mutant. Asterisk highlights the lack of myogenic progenitors in the EOM region of the mutant embryo, indicated by the absence of Myod/Myog/Pax7 staining.

E) Quantification of proportion of β-gal+;Sox9+ double positive cells in the total Sox9+ population of the EOM and Masseter muscles. Each dot is a different sample, the center line of the boxplot is the median value. (n=3 embryos, p-values were calculated using a two-sided Mann-Whitney U test).

F) Model of lineage progression from bipotent cells in a *Myf5* null background.

Figure 6




Figure 6. Myf5-derived non-myogenic cells are continuously generated up to fetal stages

A-C') RNA velocity plots of *Myf5^{Cre/+}; R26^{mTmG/+}* E11.5, *Myf5^{GFP-P/+}* E12.5 and *Myf5^{GFP-P/+}* E14.5 datasets displaying cell-type annotation (A-C) and myogenic and non-myogenic clustering (A'-C').

D-E) Gene ontology network of GO Molecular Function and Reactome pathway performed on combined top 100 markers.

(D'-E') Relative contribution of each stage to term node.

Figure 7




Figure 7. A shared program including Wnt/ β -cat activity supports non-myogenic fate transition at various stages and anatomical locations

A-D) Velocity UMAP highlighting the non-myogenic cluster at each stage (dotted box), from which the underlying network was inferred. Driver genes and regulatory networks (regulons) were produced for each stage independently, and a stage-specific network of active transcription factor and associated driver gene targets was built. Size of nodes corresponds to the number of edges (connections) they have, i.e. the number of driver genes the transcription factor regulates.

E) Histogram displaying frequency of appearance of most predominant transcription factors as driver regulators (4= present in all 4 datasets as driver regulon, 1= present in a single dataset). Red arrowheads highlight members of the Wnt/ β -cat pathway Tcf4, Tcf7l2 and Lef1.

F-I) Dotplot of the expression levels and percent of Axin2 and Dkk2 in the myogenic and the non-myogenic portions of all 4 datasets.

SUPPLEMENTAL INFORMATION

Identification of bipotent progenitors that give rise to myogenic and connective tissues in mouse


Alexandre Grimaldi^{1,2}, Glenda Comai^{1,2}, Sébastien Mella^{1,2}, Shahrarim Tajbakhsh^{1,2,*}

¹Stem Cells & Development Unit, 25 rue du Dr. Roux, Institut Pasteur, 75015 Paris, France

²UMR CNRS 3738, Institut Pasteur, Paris, France

*corresponding author

Figure S1




Supplemental Figure S1. Seurat library preprocessing metrics.

A-B) Gating strategy used to isolate by FACS *Myf5*^{Cre/+}; *R26*^{mTomG/+} (A) and *Myf5*^{GFP-P/+} cells (B). To isolate *Mesp1*^{Cre/+}; *R26*^{mTomG/+} cells, the same strategy as in (A) was used. The Alexa Fluor 488 and FITC channels were used interchangeably to identify GFP+ cells. The BV421 and Alexa Fluor 405 were used interchangeably to identify the Calcein Blue+ live cells. The PE-Texas Red channel was used to discard mTomato+ cells and Propidium Iodide + cells. The percentage of cells captured by each gate is displayed on each plot.

C, E, G, I) Violin plots of gene count, UMI count and mitochondrial fraction for each dataset.

D, F, H, J) Gene count and UMI count per cell type for each dataset. A “High count” cluster was found in the E11.5 dataset, and was removed in downstream analysis.


Figure S2



Supplemental Figure S2. Myogenic and non-myogenic markers define anterior mesodermal tissues.

- A) *Mesp1*^{Cre/+};*R26*^{mTmG/+} E10.5 UMAP expression plots of markers of various mesodermal lineages.
- B) Heatmap of top 5 markers of each cluster of *Mesp1*^{Cre/+};*R26*^{mTmG/+} E10.5.
- C) UMAP expression plot of the *Mesp1*^{Cre/+};*R26*^{mTmG/+} E10.5 subset. *En2*: marker of pharyngeal arch 1 (Knight et al., 2008), *En1*: marker of epaxial somitic progenitors (Cheng et al., 2004), *Lbx1*: marker for tongue progenitors (Gross et al., 2000), *Isl1*: marker of cardiopharyngeal mesoderm of pharyngeal arch 2-6 (Comai et al., 2019), *Shox2*: marker of caudal cardiopharyngeal mesoderm (Wang et al., 2020), *Pitx2*: marker of the extraocular region (Zacharias et al., 2010).
- D) Pearson correlation plot of myogenic (*Pdgfra*, *Myf5*, *Myod1*, *Myog*, *Acta2*) and non-myogenic (*Pdgfra*, *Prrx1*, *Meis1*, *Twist1*, *Osr1*, *Col1a1*) genes. The size of the dots is inversely proportional to their p-value. A cross indicates a p-value higher than 0.05. The color of the dots indicates the strength of the a positive (blue) or negative (red) correlation.

Figure S3




Supplemental Figure S3. Cell cycle phases and scvelo metrics.

A) UMAP of each dataset with overlaid velocity and cell cycle phase.


B-E) Quality control metrics of scvelo, including velocity length, velocity confidence and spliced/unspliced abundance per dataset and cell type.

Figure S4

A



B





Supplemental Figure S4. EOM non-myogenic cells arise from a myogenic compartment and crosstalk with myogenic cells.

A) GO Molecular Function network, including relative contribution of each cluster to the term and significance levels. Insert show the significance of each term.

B) UMAP of *Myf5*^{Cre/+}; *R26*^{mTmG/+} E11.5 EOM illustrating velocity confidence and velocity length. Higher confidence is found on both ends of the EOM cluster.

Figure S5



EOM: lateral → medial

Mandible


Supplemental Figure S5. Myod⁺ cells are restricted to a myogenic fate.

A-E) Transverse sections of *Myod*^{Cre/+}; *R26*^{TdTomato/+}; *Pdgfra*^{H2BGFP/+} embryos at E14.5 immunostained for Myod/Myog (committed and differentiating myoblasts) in the extraocular (A), mandibular (B), laryngeal (C), tongue (D) and upper neck (E) regions. White arrowhead indicates rare double positive cells (GFP+/Tom+).

Supplemental Figure S6. Non-myogenic Myf5-derived cells display a similar gene ontology.

Gene ontology analysis for Reactome pathways, including genes underlying each term, and their representation in each dataset. Specific genes of each stage appear related.

Figure S7



Supplemental Figure S7. Model of Myf5+ bipotent progenitors giving rise to muscle and associated connective tissues.

Model for bipotent Myf5+/Pdgfa+ progenitors giving rise to myogenic and non-myogenic cells and discrete parts of the head, deprived of neural crest. Upon activation of a set of transcription factors including Prrx1/2, Foxp2, Hmga2, Meis1, Meox2, Fli1, Twist1, Ets1, Tcf7l2 and Tcf4, a fibrogenic fate is acquired. A molecular dialogue is initiated at the branchpoint including extracellular matrix components and tyrosine kinase signalling such as Pdgf, Ephrins and Bmps. The non-myogenic fate is maintained cell-autonomously by a canonical Wnt positive feedback loop.

E10.5 Anterior somites	E11.5 EOM Myogenic	E11.5 EOM Non-myogenic	E12.5 Non-myogenic	E14.5 Non-myogenic
Tshz2	Ccdc141	Zfp62	Mgat4c	Dnm1
Eya1	Mcm6	Plxna4	Cenpv	Pid1
C1qtnf3	Dync1i1	Col23a1	C130073E24Rik	Nrp2
Meis2	Tpm2	Edil3	Tbx3os1	Ntrk3
Limch1	Celf2	Map2	E330013P04Rik	Tmem132c
Moxd1	Sox6	Rora	Stk26	Egflam
Epha4	Tnc	Sema5a	Edil3	Gpr153
Pitx2	Magi3	Colec12	Fdft1	Efemp1
Parm1	Sh3glb1	Smoc1	Lima1	Adamts2
Hpse2	Parm1	Ptptr	Trim59	Brinp1
Lrm1	Ephb1	Ror1	Meg3	Vegfc
Dmrt2	Bmpr1b	Dock5	Gins3	Twist2
Myl3	Hells	Map1b	Tpm2	Itgb5
Fap	Pdgfc	Fn1	Cdh6	Gria1
Hs6st2	Ptprd	Limch1	Csmd3	Sned1
Ddr2	Cnr1	Tenm4	Tceal5	Sorcs3
Cald1	Sema3d	Rbms3	Pclaf	Ebf2
Pmx1	Clcn5	Srgap3	Tspan9	Fam19a1
Magi3	Chd7	Tmem132c	Eps8	Trabd2b
Ntn1	Col25a1	Sdc2	Lmna	Plxdc2
Zfx3	Reep1	Add3	Dmrt2	Sh3gl3
Meis1	Cttna1	Pdgfra	Cpeb4	Luzp2
Tnni1	Tpm1	Gmcs	Hpgd	Pdzd2
Crym	Zim1	St6galnac3	Rcsd1	Sema3e
Ebf1	Lmx1a	Epb41l3	Pdgfra	Rims1
Nr2f1	Neb	Pde3a	Plac1	Epha3
Ntng1	Atad2	Tox	Palmd	Cyp7b1
Pgm5	Dapk2	Smarca2	Gucy1a1	Gem
Cdh6	Prox1	Ctdspl	Wif1	Ldb2
Foxp1	Lsomp	Magi2	Naalad2	Scube1
Celf2	Ttn	Dpysl3	Smoc2	Pdgfra
Tbx1	Pls3	Fgfr2	Rassf4	Pde1a
Bdnf	Slf2	Ldb2	Pttg1	Nde1
Colec12	Vat1l	Igf1	Josd2	Enpp2
Eya4	E2f1	Elk3	Plxna4	Fam107b
Sobp	Epb41l2	Zmiz1	Eya2	Sbxp6
Peg3	Gm28653	Dlc1	Nrsn1	Rerg
Pdgfra	Lmn1	Nhs	Fign	Prex2
Nrk	Mef2c	Cdkn1c	Inpp1	Man1a
Ptn	St8sia2	Plpp3	Rnf152	Tmem45a
Daam1	Tshz1	Ebf1	Lasp1	Sh3bp4
Dlk1	Wee1	Sorbs2	Mrln	Mcc
Unc5c	Slc24a3	Baz1a	Cdt1	Ncald
Lpar1	Ncoa1	Fat4	Notch3	Kdelr2
Syne2	Dek	Golgb1	Pax3	Pcdh19
Nkd2	Kdm5b	Hpse2	Egfr	Gas7
Brinp1	Unc13c	Samd4	Dbf4	Cpt1c
Zfx4	Ddr1	Itga9	Bcr	Adam22
Nnat	Pip4k2a	Magi1	Milt3	Itgb8
Gxylt2	Fndc3c1	Pcdh9	Nectin1	Dchs2
Cimp	Rbm24	Tgfr2	Grin3a	Cep350
Ror2	Rreb1	Ntf3	Cbfa2t3	Oat
Nfia	Rragd	Col11a1	Cdh2	Rab30
Ebf2	Acsl3	Runx1t1	Anln	Aff2
Ednra	Acvr2a	Tnrc18	Ccdc6	Gna14
Fli1	Zeb1	Crym	Mcu	Slc29a1
Tspan12	Rgma	Fap	Fnip2	Pls3
Ttc28	Arpp21	Ppp1r1a	Kcnk13	Traf3ip1
Nfib	Lef1	Tes	Sned1	Rcsd1
Ccdc88c	Nr2f2	Bicc1	Nde1	Lgr4
Col13a1	Foxo1	Il1rap1	Hipk3	Zfp9
2700069118Rik	Pdzm4	Alcam	Arhgap11a	Hs3st5
Pcolce	Hmga2	2700069118Rik	Fam8a1	Aspn

E10.5 Anterior somites	E11.5 EOM Myogenic	E11.5 EOM Non-myogenic	E12.5 Non-myogenic	E14.5 Non-myogenic
Scn3a	Lurap1l	Dab2	Klf21a	Nrxn1
Acvr2a	Pkig	Cntln	Mtss1	Rrm1
Auts2	Ncl	Clmn	Abcd2	Igf1bp7
Col3a1	CT025619.1	Rbms1	Irx5	Slc35f3
Gap43	Erb4	Tmem2	Pacs2	Klf15
Mrln	Cdk14	Cdh6	Nab1	Slc1a3
Pax3	Klf21a	Lypd6	Cond2	Bmp6
Sim1	Zfp704	Mmp2	Bok	Dkk2
Epb41l2	Nasp	Klf5c	Dok5	Tspan9
Ppp3ca	Plekha5	Cadm2	Ncapg	Ets1
Tnfrsf6	Cap2	Prkg2	Rfx8	Gria3
Tmem132c	Snca	Cped1	Fhod3	Sox8
Tmem2	Epha4	Dtl	Tk1	Melk
Epb41l3	Atad5	Ror2	Asf1b	Ntm
Crybg3	Cntn3	Utrn	Tek	Synpo2l
Nrxn1	Cacna2d1	Foxp1	Arfgef3	Hlf
Farp1	Pak3	L3mbtl3	Rnf182	Adamts5
Sulf1	Megf10	Cdh23	Klf14	Picb4
Tmtc2	Tnnt1	Negr1	1810041L15Rik	Cdc25b
Pde4dip	Acta2	Hmgn1	Rrm2	Mgat4a
Phldb2	Barx2	Col26a1	Fgf5	Mdfic
Plpp3	Mrln	Fbn2	Barx2	Trpc5
Ybx3	Pgm5	Ankrd12	Fli1	Klf4
Ppm1l	Fmr1	Lhfp	Jph2	Pice1
Twist2	Smc4	Hs3st3b1	Dtx4	Il17rd
Nuak1	Cimp	Adgrl3	Ncald	Mmp16
Tgfb2	Alpk2	Svil	Zic4	Hhip
Sfrp1	Kctd1	Mob3b	Dlc1	Tpx2
Sncaip	Meg3	Trabd2b	Cdc45	Ndc80
Tenm3	Samd5	Rmst	Gatm	Bub1b
Cdh2	Nrk	Prx1	Ssc5d	Hmmr
Iqgap2	Piezo2	5330434G04Rik	Phactr2	Kank4
App	Robo1	Zfx3	Ppp1r14c	Tmeff2
Pgam2	Col1a2	Foxp2	Agl	Nr4a1
Rspo3	Cntrl	Mpp6	Tox3	Aurkb
Cdon	Mllt3	Crispld1	Aurka	Lrrtm3
Ebf3	Peg3	Eya1	Cdh15	Cenpq

Table 1: Driver genes underlying cell fate decision in each dataset.

	E10.5	E11.5	E12.5	E14.5
Foxp2	(+)	(+)	(+)	(+)
Hmga2	(+)	(+)	(+)	(+)
Meis1	(+)	(+)	(+)	(+)
Meox2	(+)	(+)	(+)	(+)
Tcf7l2	(+)	(+)	(+)	(+)
Fli1	(+)	(+)	(+)	(-)
Lef1	(-)	(+)	(+)	(+)
Prrx1	(+)	(+)	(-)	(+)
Prrx2	(-)	(+)	(+)	(+)
Six2	(+)	(+)	(+)	(-)
Creb3l1	(-)	(+)	(-)	(+)
Ebf1	(+)	(-)	(+)	(-)
Ets1	(-)	(+)	(-)	(+)
Foxp4	(+)	(+)	(-)	(-)
Hoxb3	(-)	(+)	(+)	(-)
Klf6	(-)	(+)	(-)	(+)
Nfatc4	(-)	(+)	(-)	(+)
Nfib	(-)	(+)	(+)	(-)
Pax7	(-)	(-)	(+)	(+)
Pbx1	(-)	(+)	(-)	(+)
Rreb1	(-)	(-)	(+)	(+)
Tbx15	(+)	(+)	(-)	(-)
Tcf4	(+)	(-)	(+)	(-)
Twist1	(+)	(+)	(-)	(-)
Zic4	(+)	(-)	(+)	(-)
Zmiz1	(-)	(+)	(+)	(-)
Ar	(-)	(-)	(-)	(+)
Arid5b	(-)	(-)	(+)	(-)
Atf3	(-)	(-)	(-)	(+)
Chd2	(+)	(-)	(-)	(-)

Table 2: Driver regulators of non-myogenic fate in each dataset.

(+): Present, (-): Absent.