

Srand specific PCR amplification of low copy number DNA

Andreas Meyerhans, Jean Pierre Vartanian, Simon Wain-Hobson

▶ To cite this version:

Andreas Meyerhans, Jean Pierre Vartanian, Simon Wain-Hobson. Srand specific PCR amplification of low copy number DNA. Nucleic Acids Research, 1992, 20 (3), pp.521-523. 10.1093/nar/20.3.521 . pasteur-03520147

HAL Id: pasteur-03520147 https://pasteur.hal.science/pasteur-03520147

Submitted on 10 Jan2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Strand specific PCR amplification of low copy number DNA

Andreas Meyerhans⁺, Jean-Pierre Vartanian and Simon Wain-Hobson^{*} Laboratoire de Rétrovirologie Moléculaire, Institut Pasteur, 28 Rue de Dr. Roux, 75724 Paris Cedex 15, France

Received October 3, 1991; Revised and Accepted January 7, 1992

ABSTRACT

A method for the amplification of a single DNA strand at low copy number is described. It is a wholly PCR based approach which involves an initial linear amplification of the target using a tagged strand specific primer. This is followed by classical PCR amplification of the progeny using a pair of primers, one specific for the sequence tagged onto the 5' end of the first round primer, the second specific for the target sequence. Given the protocol used the ratio of the two strands in the final amplification product was 50:1.

INTRODUCTION

The replication of viruses frequently results in the unequal production of one of the two DNA or RNA strands. Replication of single stranded phage DNA by a rolling circle model being a case in point. Retroviral replication is also highly asymmetric involving RNA-dependent and DNA-dependent DNA polymerization by the viral reverse transcriptase and DNA-dependent RNA polymer-ization by the cellular RNA polymerase II. Recently we have described the phenomenon of $G \rightarrow A$ hypermutation of the human immunodeficiency genome whereby up to 40% of all G residues become substituted by A (1). It is not known during which step of retroviral replication this phenomenon occurs.

In order to analyse better asymmetric viral replication processes we have developed a means of amplifying a low copy number single DNA strand in a two step PCR procedure involving both linear and exponential amplification cycles.

MATERIALS AND METHODS

Target sequences and amplification primers

The two single stranded M13 recombinant sequences used (L602R and L610) encoded 430bp complementary sequences derived from the second coding exon of the human immunodeficiecy virus type 1 (HIV-1) *rev* gene. These sequences map to 7880-8306 on the HIV-1 Lai sequence (2, 3). They were

identified as part of a longitudinal study of HIV-1 rev coding sequences in vivo (4). Within the region analysed in this study four mismatches were apparent between these complementary sequences (Fig. 1).

The sequences of the amplification primers were: Sp1A 5' CACCGCATGGATAGGATTGAGGACAGGCCCGACG-GAAC; Sp1B 5' GGCAAGCTTCACCGCATGGATAGGA-TTGAG; and Sp2 5' CCCGAATTCCAGAAGTTCCACAAT-CC. The sequences specific for the HIV-1 rev sequences are shown underligned, while a randomly chosen sequence tagged onto the 5' end of the Sp1A primer is shown in boldface. Restriction sites and clamps are shown in normal type. The second round primer Sp1B encoded only the random sequence along with a HindIII restriction site doubly (underligned) at the 5' extremity so as to facilitate cloning. Finally the HIV-1 specific Sp2 primer encoded an EcoRI site (doubly underligned) for the same reasons. All oligonucleotides were synthesized using phosphoramidate chemistry. After detritylation and alkaline deprotection, crude material was used for all applications without further purification.

PCR amplification

Approximately 3.10⁵ molecules of both single stranded recombinant M13 DNA (i.e. L602R and L610) were mixed together. Nine reactions were carried out involving three different Sp1A primer concentrations (100fmol, 1fmol and 0.1fmol) with respectively 10, 50 and 100 linear amplification cycles. One fourth of the initial linear amplification reaction was taken and amplified for a further 35 cycles in an exponential amplification reaction using the Sp2 and Sp1B primer pair at a concentration of 100pmol/reaction. Amplification conditions were: 2.5mM MgCl₂, 50mM KCl, 10mM Tris-HCl pH 8.3, 0.2mM dNTP, 2.5U Tag polymerase in a final volume of 100µl overlaid with 50μ l of heavy white mineral oil (Sigma). All amplifications were performed in a Perkin Elmer-Cetus thermal cycler. For linear amplifications the samples were first heated at 90°C for 5 min. The denaturation, annealing and extension temperatures and times were 95°C (30s), 50°C (1 min) and 72°C (1 min) respectively. For exponential amplification the parameters for denaturation,

^{*} To whom correspondence should be addressed

⁺ Present address: Department of Virology, Institut for Medical Microbiology and Hygiene, University of Freiburg, D-7800 Freiburg, FRG

annealing and extension were 95°C (30s), 50°C (30s) and 72°C (30s) respectively followed by a final elongation step of 10 min at 72°C. As a control the target sequences were amplified using the Sp2 and Sp1A primers at 100pmol each for 35 cycles using the same conditions as described above. The final product size was 212bp.

Cloning and sequencing

After amplification the reaction products were separated on a 0.8% low melting point agarose gel. The purified DNA was digested with EcoRI and HindIII and subsequently ligated into similarly cleaved M13mp18 RF DNA. The control DNA amplified by the Sp1A-Sp2 pair only was kinased and blunt end cloned into alkaline phosphatased treated SmaI cleaved M13mp18 RF DNA. After transformation of E. coli strain TG1, recombinant M13 plaques were transferred to nitrocellulose filters and screened by hybridization with strand specific probes P1 and P2 which span the four base mismatches shown in Fig. 1. Their sequences are P1 5' AGACCTGCGGAGCCTGTT and P2 5' CGACCTACGGAGCCCGTG. Filters were prehybridized at 50°C for 1hr in 4×SSC, 0.5% SDS, 1×Denhardt's solution and 5mg/ml denatured salmon sperm DNA. They were then transferred to a fresh solution (as above) containing 5' ³²P endlabeled and heat-denatured probe mix (106cpm/ml solution) and hybridized for 1 hour at 50°C. Filters were washed three times with 0.1% SDS, 2×SSC at 50°C. After autoradiography the ratio of P1/P2 hybridizing plaques was established. To confirm the



hybridization results 20 P1 positive recombinant M13 clones were grown up and sequenced by the standard dideoxy method using $[\alpha$ -³⁵S]dATP (600Ci/mmol) and resolved on buffer gradient gels.

RESULTS

The experimental plan is outlined in Figure 1. The four sites at which base mismatches occur between DNA strands are indicated. There are three distinct steps, A, B and C. Step A involved 100 cycles using the Sp1A primer of which the 17 most 3' bases are complementary to the HIV-1 rev target sequence. This resulted in a linear amplification of one of the two strands yielding a relative ratio of approximately 100:1. Part of this reaction was taken and reamplified using the Sp2/Sp1B primer pair resulting in a classical PCR reaction. During the first cycle of exponential, or classical PCR, the Sp2 primer copies both the plus strands preserving the ratio of 100:1. In the second cycle Sp1B, by virtue of its sequence identity to the 5' most 21 bases of Sp1A will ensure amplification of sequences tagged by the Sp1A primer. It must be noted that there will still be some remaining Sp1A primer in the reaction (step C). During the second cycle of classical PCR the remaining Sp1A primer may anneal to the untagged complementary sequence and now copy this strand. In all subsequent cycles both strands will be amplified by the Sp2/Sp1B primer pair. However it is to be expected that the relative ratio of the strands will be > 100:1 since priming by Sp1B, being in molar excess over Sp1A, predominates. This assumes, of course, that the 100 single strand cycles are completely efficient.

A 25μ l aliquot of products from the 10, 50 and 100 linear amplifications was followed by 35 classical cycles of PCR. Two 10 μ l samples from each of the final reactions were electrophoresed on a 1.5% agarose gel and transferred to nitrocellulose. One filter was hybridized with the radiolabelled P1 probe, the other with P2. The autoradiograms are shown in Figure 2. The band intensities have been normalized to the control which contained an equimolar proportion of the two sequences as amplified by the Sp1A and Sp2 primer pair alone. As can be seen the signal intensities were much stronger using the P2 probe as opposed to the P1 probe. In order to quantitate precisely the proportion of the two strands the products from 3A, 3B and 3C (100 linear cycles) were cloned into the *Eco*RI and *Hind*III sites of M13mp18 RF DNA. The recombinant plaques were screened



Figure 1. Schema representing strand specific amplification. The initial double stranded heterologous molecule is shown on top with the four mismatched base pairs. Using the Sp1A primer one of the two strands is amplified linearly $100 \times$ (A). Thereafter the Sp1B (specific for Sp1A) and Sp2 (specific for rev) primer pair are added and exponential amplification is now possible. During the first such cycle only Sp2 may serve as a primer. During the second Sp1B will prime polymerization from sequences tagged by the initial Sp1A primer (B). However there remains a little Sp1A primer, albeit at a non optimal concentration, which ccan now prime from the complementary copy to the undesired strand (C). Subsequent rounds results in the co-amplification of both strands. It is to be expected that the molar ratio of the two products will be of the order of 100:1 as shown.

Figure 2. Analysis of strand specific DNA in the final PCR product. Probes P1 and P2 are specific for the desired and undesired DNA strands. Samples 1, 2 and 3 represent 10, 50 and 100 linear amplification cycles while A, B and C refer to initial Sp1A primer concentrations of 100, 1 and 0.1fmol/reaction respectively. The control lane (T) represents a classical amplification of the target sequences using the Sp1A/Sp2 primer pair.

with 5' labelled 32P end-labelled P2 and P1 probes. The proportion of P2:P1 positive plaques was 60:1, 40:1 and 50:1 for samples 3A, 3B and 3C respectively indicating that background did not vary in proportion to the initial Sp1A primer concentration. From other work with these PCR amplified rev sequences we know that there is no preferential bias in the blunt-

end cloning of this fragment into M13. To strengthen the results of the hybridization experiments 20 P2 positive recombinant M13 clones from reaction 3C were grown up and sequenced by the dideoxy chain terminator procedure. All 20 yielded the expected sequence shown in Fig. 1.

DISCUSSION

The protocol described allowed amplification of a specific DNA strand. The proportion of clones corresponding to the desired:undesired DNA strand was approximately 50:1 as identified upon direct screening of the cloned 3A, 3B and 3C products. That the complementary strand was detected may be attributed to the fact that during the second cycle of classical PCR the Sp1A primer, which still remained in the reaction, could anneal to the untagged strand. The ratio of the two products was less than the 100:1 ratio predicted were amplification to proceed with 100% efficiencies probably because the concentration of Sp1A primer was far from optimal, 100-0.1 fmol as opposed to 100pmol in a normal PCR reaction. Presumably at such concentrations not all target molecules were saturated by primer within the 1 minute annealing time employed in the protocol. The concentration of primer was kept low so that any amplification between Sp1A and Sp2 in the round of PCR would be inefficient with respect to Sp1B/Sp2 amplification. However the relative amplification factor of 50 is sufficient to allow the analysis of a number of biological processes.

An advantage of the method is that it is simple involving only two steps. As the sequence of the Sp2 primer was defined randomly it can be used to amplify any sequence so long as the corresponding sequence is included at the 5' end of the initial linear amplification primer. Furthermore as there is little manipulation between the two amplification steps the risk of contamination is reduced.

Strand specific amplification of DNA has been exploited for a number of purposes. The most noticeable example being in the sequencing of single stranded DNA derived from a plasmid or phage (5). Other applications include strategies for chromosomal walking (6). The degree of amplification depends on the number of linear cycles of PCR. This works well when the imput DNA concentration is sufficiently high. A normal PCR amplification of a sequence from mammalian DNA employs 1µg, or approximately 150,000 cell genome equivalents. The 3.105 copies used in the protocol thus correspond to 1 copy/haploid DNA equivalent indicating that the procedure is capable of amplifying single mammalian gene sequences. Subsequent experiments with DNA dilutions showed that as little as 10 copies of single stranded M13 DNA could be detected (data not shown).

This simple method could be used for the amplification of one strand over another where the differences anticipated are of the order of < 50. The study of DNA repair in vivo of artificially mismatched DNA sequences is an example. In these instances the use of viral vectors and plaque purification enabled analysis of progeny after transfection of cells by heteroduplex molecules. Restriction enzyme and sequence analysis showed that upon migration to the nucleus certain heteroduplexes were very

efficiently repaired while others were much less so (7-9). However these techniques rely on high copy numbers and a viral replicon and are therefore not readily applicable.

The method could be adapted to the analysis and quantitation of replication intermediates of RNA viruses if preceeded by an initial reverse transcriptase step. Finally it should prove useful in determining during which stage of the HIV-1 replication cycle $G \rightarrow A$ hypermutation occurs (1).

ACKNOWLEDGEMENTS

This work was supported by grants from the Institut Pasteur and l'Agence National de Recherches sur le SIDA (ANRS). A.M. was a post-doctoral research fellow of the Centre International des Etudiants et Stagiaires and J.P.V. was supported by the Ministère de la Recherche et de la Technologie.

REFERENCES

- 1. Vartanian, J.P., Meyerhans, A., Asjö, B. and Wain-Hobson, S. (1991) J. Virol., 65. 1779-1788.
- 2. Wain-Hobson, S., Sonigo, P., Danos, O., Cole, S. and Alizon, M. (1985) Cell, 40.9-17.
- 3. Wain-Hobson, S., Vartanian, J.P., Henry, M., Chenciner, N., Cheynier, R., Delassus, S., Pedroza Martins, L., Sala, M., Nugeyere, M.T., Guétard, D., Klatzmann, D., Gluckman, J.C., Rozenbaum, W., Barré-Sinoussi, F. and Montagnier, L. (1991) Science 252, 961-965.
- Pedroza Martins, L., Chenciner, N., Åsjö, B., Meyerhans, A. and Wain-Hobson, S. (1991) J. Virol., 65, 4502-4507.
- 5. Gyllensten, U.B. and Erlich, H.A. (1988) Proc. Natl. Acad. Sci. USA, 85, 7652 - 7656.
- 6. Rosenthal, A. and Jones, D.S.C. (1990) Nuc. Acids Res., 18, 3095-3096.
- 7. Brown, T.C. and Jiricny, J. (1987) Cell, 50, 945-950.
- 8. Brown, T.C., and Jiricny, J. (1988) Cell, 54, 705-711.
- 9. Wiebauer, K., and Jiricny, J. (1989) Nature, 339, 234-236.