



HAL
open science

Linking microbiota composition with antimalarial antibody response

Ottavia Romoli, Liliana Mancio-Silva, Mathilde Gendrin

► **To cite this version:**

Ottavia Romoli, Liliana Mancio-Silva, Mathilde Gendrin. Linking microbiota composition with antimalarial antibody response. *Trends in Parasitology*, 2021, 37 (10), pp.853-855. 10.1016/j.pt.2021.07.007 . pasteur-03672692

HAL Id: pasteur-03672692

<https://pasteur.hal.science/pasteur-03672692v1>

Submitted on 16 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Linking microbiota composition with antimalarial antibody response**

2

3 Ottavia Romoli¹, Liliana Mancio-Silva^{2, 3, 4}, Mathilde Gendrin^{1,4,*}

4

5 ¹ Microbiota of insect vectors group, Institut Pasteur de la Guyane, Cayenne, French Guiana,
6 France.

7 ² Institut national de la santé et de la recherche médicale, Unité 1201, Paris, France.

8 ³ Biology of host-parasite interactions unit, Institut Pasteur, Paris, France.

9 ⁴ Parasites and insect vectors department, Institut Pasteur, Paris, France.

10 * Correspondence: mathilde.gendrin@pasteur.fr (M. Gendrin)

11 ORCID IDs:

12 OR: 0000-0002-7608-2650

13 LMS: 0000-0002-5585-018X

14 MG: 0000-0001-6405-6458

15

16 Keywords: microbiota, malaria, severity, *Plasmodium*

17

18 **Abstract**

19 Microbiota composition recently arose as a factor correlating with malaria infection. Mandal *et al.*
20 showed via cecal transplant and antibacterial treatment that the mouse microbiota modulates
21 parasitemia by affecting spleen germinal centers where B-cells are matured. They further
22 identified correlations between microbiota composition and malaria severity in Ugandan children.

23

24 In mammals, the gut microbiota shapes both mucosal and systemic immunity, impacting the
25 outcome of infections, even those that take place distally from the gut. Malaria infections
26 represent an interesting working model on this regard, as the causing *Plasmodium* parasites
27 primarily infect the liver and blood of the mammalian host, without colonizing the gastrointestinal
28 lumen. *Plasmodium* parasites are transmitted by *Anopheles* mosquitoes. After being introduced
29 into the skin during a bite, parasites travel to the liver where they multiply inside hepatocytes. The
30 new progeny released in the bloodstream will then cyclically infect and proliferate in red blood
31 cells. Antibodies against *Plasmodium* parasites, which are produced by germinal centers in the
32 spleen, are critical to control infection.

33 *Plasmodium* infection has recently been found to correlate with microbiota composition in
34 mice [1,2]. These correlations were linked to the presence of specific bacteria [1] or to a gut
35 microbiota-dependent production of *Plasmodium*-specific antibodies in germinal center immune
36 cells [3]. To further investigate these associations, Mandal and collaborators took advantage of a
37 mouse model (strain C57BL/6) purchased from two different vendors (Tac, for *Taconic Biosciences*,
38 and CR, for *Charles River Laboratories*; [4]). These mice represent a unique system to study
39 associations between gut microbiota composition and malaria infection, as they are characterized
40 by a distinct response to *Plasmodium yoelii* 17XNL, with Tac mice efficiently restricting parasitemia
41 and CR being more permissive [2]. Using cecal transplants, Mandal *et al.* showed that the CR
42 enterotype (i.e. microbiota composition) “dominates” over the Tac one: after cecal transplants
43 from CR to Tac, mice shift from a Tac to a CR enterotype, but after a transplant from Tac to CR,
44 mice keep a CR-enterotype. Accordingly, Tac mice receiving CR caecum content exhibit higher
45 parasitemia, but not vice versa (**Figure 1A**).

46 Mandal and colleagues then perturbed the microbiota by providing mice with five
47 alternative antibiotic treatments. They observed a strong decrease in parasite burden when

48 treating CR susceptible mice with these antibiotics, which links the bacterial component of the gut
49 microbiota to *P. yoelii* infection. Using vancomycin, an antibiotic that does not cross the gut
50 barrier, at different time points prior to or after *Plasmodium* infection, they linked microbiota
51 perturbation with efficiency of parasite clearance. First, they found that antibiotic treatment
52 several weeks before infection reduced *P. yoelii* parasitemia in CR mice and protected animals
53 against experimental cerebral malaria upon subsequent *Plasmodium berghei* ANKA infection.
54 Second, treatment 7 days after infection was still able to significantly reduce *P. yoelii* infection
55 burden, indicating that microbiota perturbation can rapidly influence defense against infection.
56 Finally, to investigate connections between microbiota and immunity, the authors examined
57 germinal center reactions in CR mice treated with vancomycin. They found that vancomycin
58 treatment induced the production of CD4+ follicular helper T cells and germinal center B cells.
59 Disruption of germinal center functions mitigated the effect of antibiotic-treatment on immune
60 cells and on parasitemia (**Figure 1B**).

61 The authors also assessed correlations between malaria infection and microbiota
62 interactions in a small cohort of children in Uganda. They found different bacteria compositions in
63 the asymptomatic *Plasmodium falciparum*-infected patients, the infected with severe malarial
64 anemia and the non-infected children. They did not identify any bacteria that appear protective
65 against severity, but rather found that the frequency of specific bacteria, such as *Flavonifractor*
66 *plautii*, *Escherichia coli* and *Faecalibacterium prausnitzii* positively correlated with severe malaria.
67 This profiling allowed them to build a supervised machine learning model that predicted *P.*
68 *falciparum* status with 74% accuracy.

69 This work sheds new light into the interplay between microbiota and immune system. It
70 demonstrates that bacteria, rather than fungal, eukaryotic, or viral components of the microbiota,
71 are responsible for the observed effects on parasitemia in mice, as antibacterial treatment fully

72 recapitulates the impact of cecal transplant. By providing an intriguing example of antibiotic
73 treatment strengthening the immune system, this study may have important implications in
74 humans, especially in malaria-endemic regions where antibiotics are often prescribed to treat
75 non-specific febrile symptoms or used as self-medication [5,6]. An association between malaria
76 infection and microbiota composition in humans has been described in few independent studies
77 [1,7]. Such correlations may be explained both by an influence of specific bacteria on host
78 immunity affecting malaria infection [1], as highlighted in Mandal *et al.*'s study, and by the
79 alteration of host physiology by *Plasmodium*, promoting colonization by specific bacteria [8,9].

80 On an experimental perspective, this work further highlights that besides host and parasite
81 genetics, the host microbiota can strongly influence the outcome of malaria infections in the
82 laboratory. Microbiota, which is directly related to host diet and nutrition, should be monitored as
83 an additional experimental variable to achieve reproducible and solid results across replicates and
84 laboratories [10]. Reproducibility could be improved by using experimental models with a known
85 microbiota composition, using for instance germ-free mice recolonized with specific microbiota
86 communities.

87 Our understanding of interactions between gut microbiota and malaria severity is still
88 preliminary and further studies are needed to understand the mechanisms behind these results,
89 to identify bacterial species responsible for these observations and to assess whether these results
90 could be recapitulated in other mouse models or with other parasites. While we are far from
91 discovering the “magic pill” that will make us resistant to infectious diseases, such advances in the
92 study of microbiota-immune system crosstalk may eventually lead to the development of
93 probiotics as living therapeutics for improving human health, even when dealing with pathogens
94 that never infect the gut. This also opens perspectives to personalized medicine, which considers
95 each patient’s identity in terms of genome, proteome and metabolism, and which may also

96 include the microbiota composition as an important parameter to monitor and to link to specific
97 medical conditions.

98

99 **Acknowledgements**

100 This work is funded by the French Government's Investissement d'Avenir program, Laboratoire
101 d'Excellence "Integrative Biology of Emerging Infectious Diseases" (grant no. ANR-10-LABX-62-
102 IBEID) and by ANR JCJC MosMi to MG (grant no. ANR-18-CE15-0007).

103

104 **Declaration of Interests**

105 The authors declare no competing interests.

106

107 **References**

108

109 1 Yilmaz, B. *et al.* (2014) Gut Microbiota Elicits a Protective Immune Response against Malaria
110 Transmission. *Cell* 159, 1277–1289

111 2 Villarino, N.F. *et al.* (2016) Composition of the gut microbiota modulates the severity of malaria.
112 *Proc National Acad Sci* 113, 2235–2240

113 3 Waide, M.L. *et al.* (2020) Gut Microbiota Composition Modulates the Magnitude and Quality of
114 Germinal Centers during Plasmodium Infections. *Cell Reports* 33, 108503

115 4 Mandal, R.K. *et al.* (2021) Dynamic modulation of spleen germinal center reactions by gut
116 bacteria during Plasmodium infection. *Cell Reports* 35, 109094

117 5 Griensven, J. van *et al.* (2020) Point of care biomarkers to guide antibiotic prescription for acute
118 febrile illness in sub-Saharan Africa: promises and caveats. *Open Forum Infect Dis* 7, ofaa260-

119 6 Yeika, E.V. *et al.* (2021) Comparative assessment of the prevalence, practices and factors
120 associated with self-medication with antibiotics in Africa. *Trop Med Int Health* DOI:
121 10.1111/tmi.13600

122 7 Yooseph, S. *et al.* (2015) Stool microbiota composition is associated with the prospective risk of
123 Plasmodium falciparum infection. *Bmc Genomics* 16, 631

124 8 Walker, G.T. *et al.* (2021) Malaria parasite infection compromises colonization resistance to an
125 enteric pathogen by reducing gastric acidity. *Sci Adv* 7, eabd6232

126 9 Mukherjee, D. *et al.* (2020) Microbiota, a Third Player in the Host–Plasmodium Affair. *Trends*
127 *Parasitol* 36, 11–18

128 10 Mandal, R.K. *et al.* (2020) Temporospatial shifts within commercial laboratory mouse gut
129 microbiota impact experimental reproducibility. *Bmc Biol* 18, 83

130

131 **Figure 1. The gut bacterial microbiota modulates spleen germinal center reactions during**

132 ***Plasmodium yoelii* infections. (A)** Gut microbiota composition differs in mice purchased from two

133 distinct vendors (CR, Charles Rivers Laboratories; Tac, Taconic Biosciences) and correlates with

134 different ability to control parasitemia during *P. yoelii* infection (CR: susceptible to

135 hyperparasitemia; Tac: resistant to hyperparasitemia). Fecal transplants can induce the CR

136 phenotype in Tac mice but not vice versa. **(B)** An antibacterial treatment reduces parasite burden


137 and induces the production of germinal center B cells and CD4+ follicular helper T cells. Disruption

138 of germinal center functions mitigates the effect of antibiotic treatment on immune cells and on

139 parasitemia.

140

A



B

