



## Draft Genome Sequences of Six Isolates of the *Bacillus cereus* Group Isolated from Pet Reptiles

Milena Skóra, Magdalena Zajac, Renata Kwit, Magdalena Skarżyńska, Paulina Pasim, Emilia Mikos-Wojewoda, Arkadiusz Bomba, Aleksandra Giza, Olivier Chesneau, Rene S Hendriksen, et al.

### ► To cite this version:

Milena Skóra, Magdalena Zajac, Renata Kwit, Magdalena Skarżyńska, Paulina Pasim, et al.. Draft Genome Sequences of Six Isolates of the *Bacillus cereus* Group Isolated from Pet Reptiles. Microbiology Resource Announcements, 2022, 11 (9), pp.e0038522. 10.1128/mra.00385-22 . pasteur-03791460

**HAL Id: pasteur-03791460**

**<https://pasteur.hal.science/pasteur-03791460>**

Submitted on 29 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Draft Genome Sequences of Six Isolates of the *Bacillus cereus* Group Isolated from Pet Reptiles

Milena Skóra,<sup>a</sup>  Magdalena Zając,<sup>a</sup> Renata Kwit,<sup>a</sup>  Magdalena Skarżyńska,<sup>a</sup> Paulina Pasim,<sup>a</sup> Emilia Mikos-Wojewoda,<sup>a</sup> Arkadiusz Bomba,<sup>b</sup> Aleksandra Giza,<sup>b</sup> Olivier Chesneau,<sup>c</sup> Rene S. Hendriksen,<sup>d</sup>  Dariusz Wasyl<sup>a,b</sup>

<sup>a</sup>Department of Microbiology, National Veterinary Research Institute, Pulawy, Poland

<sup>b</sup>Department of Omic Analyses, National Veterinary Research Institute, Pulawy, Poland

<sup>c</sup>Collection de l'Institut Pasteur (CIP), Department of Microbiology, Institut Pasteur, Paris, France

<sup>d</sup>Research Group for Genomic Epidemiology, WHO Collaborating Centre for Antimicrobial Resistance in Foodborne Pathogens, Food and Agriculture Organization Reference Laboratory for Antimicrobial Resistance, European Union Reference Laboratory for Antimicrobial Resistance, National Food Institute, Technical University of Denmark, Kgs. Lyngby, Denmark

**ABSTRACT** Bacteria of the *Bacillus cereus* group are Gram-positive rods and are widespread in nature, but little information is currently available about their presence in reptiles. Here, we report draft genome sequences of six *Bacillus* isolates belonging to three species, namely, *Bacillus cereus*, *Bacillus paranthracis*, and *Bacillus toyonensis*, isolated from pet reptiles in Poland.

The *Bacillus cereus* group consists of Gram-positive, rod-shaped, aerobic, or facultatively anaerobic bacteria that belong to 21 closely related species (1, 2). One of the most relevant characteristics of the *Bacillus* genus is the ability of the bacteria to form spores which allow them to survive under harsh environmental conditions. The bacilli are widely distributed in the environment, being present in the soil, water, and other ecological niches (3). The presence of *Bacillus* species was also confirmed in the feces of cows as well as in the gastrointestinal tracts of poultry (4). In this study, we present draft genome sequences of six fecal isolates belonging to three species, namely, *Bacillus cereus*, *Bacillus paranthracis*, and *Bacillus toyonensis*, and compare these sequences with reference genomes deposited in GenBank (5).

Fecal samples were incubated in buffered peptone water (BWP) for 18 h at 37°C, and then 10  $\mu$ L of each culture was streaked onto polymyxin pyruvate-egg yolk-mannitol-bromothymol blue agar (PEMBA) (homemade) and incubated for 24 h at 37°C (6). Single colonies were passaged onto the nutrition agar and incubated overnight at 37°C (7). The pure culture was identified by matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI-TOF MS) using the extraction method following the manufacturers' guidelines (Bruker Daltonik GmbH). After the incubation onto the nutrition agar at 37°C for 24 h, genomic DNA was extracted with the Maxwell Rapid Sample Concentrator (RSC) cultured cells DNA kit (Promega) according to the manufacturer's instructions. The concentration and quality parameters of the DNA were defined with a spectrophotometer (NanoDrop One) and a Qubit 3 fluorometer. The DNA libraries were prepared using the Nextera XT sample preparation kit following the manufacturer's recommendations. Sequencing was performed using the MiSeq reagent kit on the MiSeq platform (Illumina) with the 2  $\times$  300-bp paired-end protocol, to 100 $\times$  depth of sequencing. All bioinformatic tools were used with default parameters, except where otherwise noted. Raw paired-end reads were quality filtered using FastQC v0.11.5 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc>) (8), and trimming and adapter sequence removal were executed using Trimmomatic 0.36 (ILLUMINACLIP: 2:30:10, LEADING: 3, TRAILING: 3, SLIDINGWINDOW: 4:15, MINLEN: 36) (9). Trimmed reads were then merged using BBMerge from the bbtools software suite (<https://jgi.doe.gov/data-and-tools/software-tools/bbtools/bb-tools-user-guide/bbmerge-guide/>) and assembled using SPAdes v3.9.0 with the “careful” flag (10). The draft genome sizes varied from 5,135,335 to 5,937,563 bp, and

**Editor** Steven R. Gill, University of Rochester School of Medicine and Dentistry

**Copyright** © 2022 Skóra et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Magdalena Zając, [magdalena.zajac@piwet.pulawy.pl](mailto:magdalena.zajac@piwet.pulawy.pl).

The authors declare no conflict of interest.

**Received** 9 May 2022

**Accepted** 30 July 2022

**Published** 15 August 2022

**TABLE 1** Genome properties of the *Bacillus* strains isolated from reptiles in Poland

| Isolate | Species                      | Host                       | Genome size (bp) | No. of contigs | Total no. of reads | Coverage | GC content (%) | <i>N</i> <sub>50</sub> (bp) | MLST type                               | SRA accession no. | Genome accession no. |
|---------|------------------------------|----------------------------|------------------|----------------|--------------------|----------|----------------|-----------------------------|-----------------------------------------|-------------------|----------------------|
| PIW23   | <i>Bacillus paranthracis</i> | <i>Eumeces schneideri</i>  | 5,135,275        | 35             | 3,949,460          | 205×     | 35.3           | 1,174,839                   | Unknown<br>(nearest STs: 869, 182, 761) | SRR18728991       | JAKJQA0000000000     |
| PIW25   | <i>Bacillus toyonensis</i>   | <i>Testudo hermanni</i>    | 5,937,563        | 58             | 4,533,936          | 208×     | 35.0           | 648,267                     | ST972                                   | SRR18728986       | JAKJPV0000000000     |
| PIW26   | <i>Bacillus cereus</i>       | <i>Testudo hermanni</i>    | 5,866,545        | 103            | 2,083,574          | 89×      | 34.9           | 334,014                     | Unknown<br>(nearest STs: 1197, 193)     | SRR18728987       | JAKJPW0000000000     |
| PIW27   | <i>Bacillus toyonensis</i>   | <i>Testudo horsfieldii</i> | 5,646,249        | 63             | 3,602,000          | 192×     | 35.0           | 556,185                     | ST718                                   | SRR18728988       | JAKJPX0000000000     |
| PIW28   | <i>Bacillus cereus</i>       | <i>Testudo horsfieldii</i> | 5,534,170        | 93             | 3,694,768          | 200×     | 35.0           | 328,652                     | ST197                                   | SRR18728989       | JAKJPY0000000000     |
| PIW162  | <i>Bacillus cereus</i>       | <i>Testudo horsfieldii</i> | 5,442,449        | 29             | 4,761,076          | 263×     | 35.1           | 1,736,072                   | ST511                                   | SRR18728990       | JAKJPZ0000000000     |

the GC content was approximately 35.1%. Genomic features (number of contigs,  $N_{50}$  value, GC content, and total size) were defined with QUAST (11) and included in Table 1. Average nucleotide identity (ANI) was analyzed using the JSpeciesWS online service (12) which revealed that the PIW23 genome sequence had the highest ANI based on MUMer (ANIm) value to *B. paranthracis* (GenBank accession number GCF\_001883995.1; 99.23%); PIW25 and PIW27 to *B. toyonensis* (GCA\_000496285.1; >99.58%); and PIW26, PIW28, and PIW162 to *B. cereus* (GCA\_000007825.1; >96.05%). The multilocus sequence type (MLST) was validated by using MLST 2.0 (<https://cge.cbs.dtu.dk/services/MLST/>) (13). The MLST scheme for *B. cereus* was applied to all isolates. The MLST type was found in PIW25, PIW27, PIW28, and PIW162, and other isolates (PIW23, PIW26) were identified as an unknown sequence type (ST).

**Data availability.** The draft genome sequences reported here were deposited in GenBank under the BioProject accession number [PRJNA799608](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA799608). The raw sequence read and genome assembly accession numbers are listed in Table 1.

## ACKNOWLEDGMENTS

This work was supported by funding from the European Union Horizon 2020 Research and Innovation Program under grant agreement number 773830, One Health European Joint Program, and from Ministry of Science funds for science in 2018 to 2022 allocated to the implementation of an international cofinanced project, agreement number 3932/H2020/2018/2.

## REFERENCES

1. Liu Y, Du J, Lai Q, Zeng R, Ye D, Xu J, Shao Z. 2017. Proposal of nine novel species of the *Bacillus cereus* group. *Int J Syst Evol Microbiol* 67:2499–2508. <https://doi.org/10.1099/ijsem.0.001821>.
2. Bukharin OV, Perunova NB, Andryuschenko SV, Ivanova EV, Bondarenko TA, Chainikova IN. 2019. Genome sequence announcement of *Bacillus paranthracis* strain ICIS-279, isolated from human intestine. *Microbiol Resour Announc* 8:e00662-19. <https://doi.org/10.1128/MRA.00662-19>.
3. Fira D, Dimkić I, Berić T, Lozo J, Stanković S. 2018. Biological control of plant pathogens by *Bacillus* species. *J Biotechnol* 285:44–55. <https://doi.org/10.1016/j.jbiotec.2018.07.044>.
4. Nfor NN, Lapin CN, Richard McLaughlin W. 2015. Isolation of *Bacillus cereus* group from the fecal material of endangered wood turtles. *Curr Microbiol* 71:524–527. <https://doi.org/10.1007/s00284-015-0875-x>.
5. Karen Clark G, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2016. Genbank. *Nucleic Acids Res* 44:D67–D72. <https://doi.org/10.1093/nar/gkv1276>.
6. International Organization for Standardization. 2006. ISO 21871:2006. Microbiology of food and feed stuffs. Horizontal method for the determination of low numbers of presumptive *Bacillus cereus*—most probable number technique and detection method. International Organization for Standardization, Geneva, Switzerland.
7. International Organization for Standardization. 2014. ISO/TR 6579-3:2014 (E): microbiology of the food chain: horizontal method for the detection, enumeration and serotyping of *Salmonella*. Part 3. Guidelines for serotyping of *Salmonella* spp. International Organization for Standardization, Geneva, Switzerland.
8. Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
9. Bolger AM, Lohse M, Usadel B. 2014. Genome analysis Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
11. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. Genome analysis QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29: 1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
12. Richter M, Rossell O-M, Ora R, Glö Ckner FO, Rg Peplies J. 2016. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931. <https://doi.org/10.1093/bioinformatics/btv681>.
13. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>.