



# Regulation of neurogenesis in the adult and aging brain

Lida Katsimpardi, Pierre-Marie Lledo

## ► To cite this version:

Lida Katsimpardi, Pierre-Marie Lledo. Regulation of neurogenesis in the adult and aging brain. Current Opinion in Neurobiology, 2018, 53, pp.131-138. 10.1016/j.conb.2018.07.006 . pasteur-03887176

**HAL Id: pasteur-03887176**

**<https://pasteur.hal.science/pasteur-03887176>**

Submitted on 6 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



# Regulation of neurogenesis in the adult and aging brain

Lida Katsimpardi<sup>1,2</sup> and Pierre-Marie Lledo<sup>1,2</sup>

Neural stem cells (NSCs) represent a remarkable developmental unit, necessary for the proper functioning of neurogenesis, by retaining their plasticity to self-renew and give rise to progeny throughout life in specific regions of the adult brain. Although NSCs were thought to merely represent a stem cell type in the brain, recent advances have demonstrated the incredible complexity of NSC identity and functions. Ranging between quiescence, activation and intermediary subtypes, NSCs choose their fate through their developmental inheritance, regional positioning within the niche, as well as dynamic transcriptional and metabolic states. The plasticity of their developmental program is reflected in the tremendous changes they undergo upon external environmental cues and extrinsic manipulations, and harnessing these potentials can open new avenues to fight against brain injury, neurodegenerative and age-related diseases.

## Addresses

<sup>1</sup> Perception and Memory Lab, Neuroscience Department, Institut Pasteur, Paris, France

<sup>2</sup> Centre National de la Recherche Scientifique, Unité Mixte de Recherche 3571, F-75015 Paris, France

Corresponding authors: Katsimpardi, Lida ([lida.katsimpardi@pasteur.fr](mailto:lida.katsimpardi@pasteur.fr)), Lledo, Pierre-Marie ([pmllledo@pasteur.fr](mailto:pmllledo@pasteur.fr))

**Current Opinion in Neurobiology** 2018, **53**:131–138

This review comes from a themed issue on **Developmental neuroscience**

Edited by **Alex Kolodkin** and **Guillermina López-Bendito**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 2nd August 2018

<https://doi.org/10.1016/j.conb.2018.07.006>

0959-4388/© 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The process of adult mammalian neurogenesis, the birth of new neurons in the brain, was discovered over 50 years ago [1], and during these last decades enormous progress has been made in deciphering the mechanistic aspects of adult neurogenesis, regulation of intrinsic NSC machinery, as well as systemic regulation of the niche in animal models; yet this field of research still remains fruitful. Identification of pathways involved in adult neurogenesis and exploration of the mechanistic regulation of NSCs can increase our understanding of this process and give us further insight into the relationship of adult neurogenesis and neurological and psychiatric disorders, as well as brain injury. For

this, open questions about the biology of NSCs, and the way their activity could be regulated, need to be addressed. Recently, in addition to the study of the overall process of neurogenesis, much effort has focused on deciphering the intrinsic regulation of stem cells in the brain, both in the hippocampus as well as the subventricular zone (SVZ) niche. In this review we discuss recent advances and new insights into *in vivo* NSC heterogeneity, the balance between quiescence and activation, the role of cell metabolism, as well as transcriptomic and metabolomic analyses that have pushed the field forward into the exploration of the multiple facets of the adult NSC as a cell type that can dynamically transit between different states upon external cues.

## Human adult neurogenesis

The main reason behind the continuing interest in understanding the process of mammalian adult neurogenesis is the notion that similar processes might be involved in the human brain. Whether neurogenesis in humans exists has been investigated using several and distinct approaches that brought compelling evidence about the presence of adult hippocampal neurogenesis in human brains. Because of the difficulty in accessing human adult tissue and measuring the incorporation of thymidine analogs to label proliferating cells in humans *in vivo*, several studies were based on the analysis of postmortem human brains, which had been labeled for different purposes. For example, the remarkable study by Eriksson *et al.* provided strong evidence for the birth of newborn neurons and incorporation of BrdU in cycling progenitor cells in brains of cancer patients [2], whereas Spalding *et al.* took advantage of the concentration of nuclear bomb test-derived <sup>14</sup>C in genomic DNA to demonstrate the existence and to calculate the turnover rate of newborn neurons throughout adult life in humans [3]. Interestingly, two very recent -but opposing- publications brought back the debate concerning the existence of human adult neurogenesis. Sorrells *et al.*, using postmortem and fresh tissue, reported that there was no evidence of neurogenesis in humans after adolescence whatsoever [4<sup>•</sup>], while the study by Boldrini *et al.* demonstrated the exact opposite by showing that adult neurogenesis persists during life in humans, albeit with a small decrease with aging, while the volume of the dentate gyrus remains the same [5<sup>•</sup>]. Further exploration of this complex question is necessary in order to conclude on the processes underlying the timeline and the mechanisms of neurogenesis in humans.

## Neural stem cell heterogeneity: not all NSCs are born equal

In the adult mammalian brain, NSCs reside mainly in two areas of the adult brain, the SVZ and the dentate gyrus of

the hippocampus [6,7], and they represent a pool of self-renewing cells that can differentiate into neurons upon different stimuli [8]. Despite the preconceived notion that adult NSCs are merely stem cells residing in the brain, increasing evidence suggests that NSCs constitute an extremely diverse population of cells. NSCs exhibit different characteristics and functions depending on their proliferative state, as well as their regional identity. In the SVZ, NSCs have restricted positional information depending on the specific location where they reside, which will determine the neuronal type into which they will terminally differentiate and mature in the olfactory bulb [9]. Depending on the respective microdomain in the SVZ niche, patterned by specific transcription factors such as *Nkx6.2*, *Zic* [10], *Gsx2* [11], *Nkx2.1* [12] or *Pax6* [13,14], NSCs generate several different subtypes of interneurons that regulate the olfactory bulb [15], revealing the complexity and inter-regulation between cell types in the neurogenic niche [16,17]. In addition, adult NSCs and their embryonic counterparts generate functionally distinct subpopulations of dopaminergic neurons [18], while exposure to reward-associated odors specifically increases the activity of adult-born neurons but not preexisting neurons [19]. The remarkable plasticity of NSCs is also demonstrated by the capacity of SVZ NSCs to convert to reactive astrocytes and contribute to the astrocyte scar following brain injury, and these SVZ-derived reactive astrocytes can also be converted to neurons by *Mash1* [20]. Interestingly, it was recently reported that a subset of CD133+ ependymal cells throughout the central nervous system (CNS) can be reactivated into neuronal differentiation upon specific cues, such as VEGF and bFGF, suggesting that these cells are dormant ependymal NSCs [21], lending more credence to the stem cell identity of ependymal cells [16,22–24].


The above findings show that regional and developmental identity plays a pivotal role for NSC lineage progression.

### The dynamic state of a neural stem cell

Although adult stem cells in brain niches are broadly referred to as NSCs, we can distinguish different subtypes of NSCs, mainly based on their state of quiescence or activation. NSCs in the adult SVZ niche originate from a subpopulation of embryonic radial glia cells, which become specified during development and maintain their quiescent state until reactivation in adulthood [25]. Specifically, most adult SVZ NSCs originate from a distinct population of slowly dividing neural progenitors in the ganglionic eminence of the embryonic brain [26]. Single-cell transcriptomic analyses confirmed that adult NSCs share a core transcriptional phenotype with their radial glial progenitors and that the transition to the adult NSC state occurs during late neurogenesis [27]. Interestingly, the number of the embryonic stem cells that will become adult NSCs is regulated by the type of cell

division during embryonic development [28<sup>••</sup>]. Modification of the embryonic program, such as deletion of VCAM1, a molecule necessary for the maintenance of adult quiescence [29], results in a reduction of the adult quiescent NSC (qNSC) pool, showing that maintenance of NSC properties is a continuous developmental process, operating in a temporal-dependent and region-dependent mechanism [30]. Chronic live imaging of the hippocampus showed that NSCs divide within a limited time window and that their division patterns are associated with each NSC's cell division history [31]. Lineage tracing techniques in the SVZ showed that a small subset of adult dividing NSCs go through symmetric self-renewing divisions, whereas the majority, around 75%, undergoes lineage progression to generate neural progenitors, which rapidly differentiate into neuroblasts at the expense of NSCs [32<sup>•</sup>]. While some activated NSCs (aNSCs) rapidly cease their neurogenic activity, other NSCs are re-activated from their quiescent state to take over lineage progression and thus safeguard the continuation of neurogenesis [33]. Simultaneously, a small fraction of NSCs can revert back to a transient quiescent state through degradation of proactivation factor *Ascl1* in order to maintain life-long hippocampal neurogenesis and avoid stem cell pool exhaustion [34]. However, the majority of NSCs, once activated they divide until they become exhausted, which could explain NSC depletion with aging. The advance of single-cell transcriptomics has provided extremely useful information about the different states of a NSC, from quiescence to activation, suggesting a high degree of transcriptional dynamics throughout these states. Purification of acutely isolated SVZ NSCs revealed four types: dormant NSCs, qNSCs, aNSCs and progenitor cells (Figure 1a). NSCs present a heterogeneous molecular profile and multiple states of activation in the adult SVZ niche [35]. Most NSCs are ciliated, quiescent, express GFAP and CD133, and they give rise to cycling, activated EGFR+ NSCs, which in turn differentiate into progenitors and finally neuroblasts [36,37]. Activated NSCs retain the ability to form spheres *in vitro*, unlike qNSCs [38]. However, additional NSC subpopulations in intermediate states have recently been discovered. Pseudotemporal ordering of single-cell transcriptomic analyses revealed three subpopulations of aNSCs (early, mid and late activation states), which exhibit variations in cell cycle timing and progression, together with differential expression of specific genes, placing these subpopulations in a continuum between quiescence and activation [39<sup>•</sup>]. In these cells, activation is associated with protein synthesis and differentiation priming, while dormancy is coupled to high glycolytic and lipid metabolism [35]. Interestingly, single-cell RNA-Seq in the dentate gyrus revealed that hippocampal NSCs also exhibit a molecular heterogeneity and take part in a progressive continuum of transcriptional dynamics from quiescence to neuronal differentiation [40,41<sup>•</sup>].

Figure 1



Schematic representation of the subventricular zone neurogenic niche. **(a)** Cell subtypes involved in the progression from dormant niche astrocytes to quiescent NSCs to activated NSCs to neural progenitors to neuroblasts. **(b)** Cytoarchitecture of the SVZ: the ventricular area (V) is composed of ependymal cells (dark and light brown) and the apical cilium of qNSCs, while the SVZ contains qNSCs (dark blue with cilium), aNSCs (light blue without cilium), transit-amplifying neural progenitors (green) and chains of migrating neuroblasts (pink). NSCs make contact with the cerebrospinal fluid on the apical side and blood vessels (red) through a basal process. **(c)** Sagittal section of the brain depicting the SVZ area (small square).

Most adult NSCs depend on Notch signaling for their maintenance and self-renewal [42]; however, different Notch family members have distinct roles in adult NSC regulation. Notch1 maintains actively proliferating NSCs, whereas Notch2 maintains NSC quiescence

[43]. Similarly, Notch3 plays a crucial role in the maintenance of qNSCs residing in the lateral and ventral walls, but not in the medial and dorsal walls, of the SVZ [44]. Within the Notch-dependent NSC pool, *Hes5* +BLBP + cells were characterized as a long-term



neurogenic population that can transition between quiescence and activation and are greatly affected by aging [45]. In addition, non-canonical Wnt signaling can induce activation of Cdc42 in qNSCs which in turn modulates the expression of Notch1, Id1 and N-Cadherin, resulting in anchorage of NSCs to the niche, as well as maintenance of their self-renewal capacity [46].

Stem cell fate is tightly related to metabolic changes in the surrounding environment, regulated by energy-sensing pathways in NSCs [47]. For example, insulin stimulates proliferation, but not self-renewal, of adult NSCs through IRS2-mediated regulation of Cdk4 activity [48]. Recently, metabolomic analyses of NSCs showed high levels of taurine and glucose in these cells compared to other stem cell types [49]. Interestingly, it was also demonstrated that qNSCs in the dentate gyrus require high levels of fatty acid oxidation to maintain quiescence, while a shift in lipid metabolism can induce a change of NSC state [50\*\*] and impairment of lipogenesis in adult NSCs results in a sharp decline of neurogenesis [51]. Along those lines, deregulation of lipid metabolism in the neurogenic niche is sufficient to induce a disease phenotype, such as Alzheimer's [52], demonstrating the importance of proper metabolic regulation within the cell.

### Extrinsic regulation of NSC state

The neurogenic niche is an extensive microenvironment supporting and nurturing NSC through structural scaffolding, secretion of local factors, nutrients and oxygen necessary for their maintenance. It is composed of different types of neural and non-neural cells that interact with each other: ependymal cells (in the SVZ), astrocytes, pericytes, microglia and blood vessels, as well as the NSC progeny (transit-amplifying neural progenitors and neuroblasts) [53–55] (Figure 1b,c). Local stimuli from the niche, as well as circulating blood factors secreted from remote organs, can positively or negatively affect the NSC state and differentiation potential, thereby modulating neurogenesis in the adult brain [56–58]. Astrocytes and microglia can affect neurogenesis and induce cognitive dysfunction by secretion of pro-inflammatory cytokines in the niche, such as IL-1, [59]. A main non-neural component of the niche is the vasculature, which is intertwined within the niche [60–63]. Strikingly, not only the neurogenic niche vasculature, but also blood vessels from non-neurogenic, cortical areas can secrete diffusible signals, such as PlGF-2, to affect NSC proliferation [64\*,65]. In a reciprocal fashion, adult NSCs secrete factors, such as VEGF [66], or the neurovascular protein EGFL7, which is secreted by both NSCs and vascular cells and modulates the quiescence state of NSCs [67]. Additionally, NSCs can decrease the inflammatory metabolite succinate and thus push niche microglia towards an anti-inflammatory phenotype [68]. In addition to the local microenvironment, the cerebrospinal fluid, which is produced by the choroid plexus and circulates in

the ventricles, brings secreted factors that regulate NSC fate and, consequently, neurogenesis. Some of these factors include IGF2 (which regulates proliferation of NSCs), Sonic Hedgehog, Wnts, retinoic acid, NT-3 and bone morphogenetic proteins [69–73]. Regulation of the NSC state can also take place through modulation by neuronal activity, which is a hallmark of adult neurogenesis [74]. For example, long-range GABAergic projection neurons depolarize GABA signaling onto local parvalbumin interneurons which, in turn, keep hippocampal NSC quiescent [75\*]. Remarkably, it was recently shown that adult neurogenesis could be modulated depending on hunger or satiety states via hypothalamic control. In this case, proopiomelanocortin neurons selectively innervate the anterior SVZ and promote proliferation of Nkx2.1+ NSCs and the generation of deep granule neurons [76\*\*]. Additionally, a mouse model of kainic acid-induced epilepsy demonstrated that neuronal hyper-excitation accelerates depletion of NSCs by inducing their activation *en masse*, converting them into reactive astrocytes thereby exhausting adult hippocampal neurogenesis [77].

### Aging and rejuvenation of NSCs

Aging negatively affects neurogenesis by inducing a sharp and continuous decrease in cell production in both the SVZ and hippocampal neurogenic niches of the brain [7,78–80]. With aging, aNSCs lose their proliferative potential and become quiescent [81], but, remarkably, they can be reactivated to a certain extent upon stimulation, such as exercise or even seizure [79], indicating that NSC plasticity is preserved to a certain extent in the aged organism. For example, it was recently demonstrated that high mobility group B family 2 (HMGB2) is associated with the transition of NSCs from quiescence to proliferation and that aging negatively impacted these cell populations, while running exercise stimulated the proliferation of HMGB2+ cells [82]. Furthermore, global SVZ transcriptome analyses of multiple time points during the aging process (2, 6, 18 and 22 months of age) showed that SVZ transcriptome is not modified in a linear manner with aging, since processes such as proliferation of Mash1+ progenitors decrease until 18 months of age, but then sharply increase at 22 months [83\*]. A recent analysis of the molecular profiles between 2-month-old and 6-month-old mice showed sharp molecular modifications in the aNSC program, and a characteristic lengthening of their cell cycle [84\*]. Age-related changes in the cell cycle are likely due to an accumulation of damaged proteins resulting in a reduction in the NSC proliferation rate [85]. An impairment of diffusion barriers has been shown to cause symmetric inheritance of ubiquitinated or damaged proteins, leaving all NSC progeny with excess cellular damage [86]. Deficient proteostasis was shown to be due to defective lysosomes in qNSCs, but enhancement of the lysosomal pathway, via transient expression of the active form of TFEB in aged qNSCs or systemic

treatment with rapamycin, resulted in reversal of the quiescent to an active state [87<sup>••</sup>]. Taken together, the above findings suggest that an inherent developmental program dictates the NSC transcriptional dynamic state throughout life.

In addition to the intrinsic aging program, environmental influences from neighboring niche cells, or even remote organs, can affect NSC state and fate [88]. Transcriptional dynamics observations in the hippocampal niche reveal age-associated changes in the numbers and molecular profiles of NSCs, progenitors and microglia [41<sup>•</sup>]. With aging, microglia become progressively activated in the SVZ niche and secrete proinflammatory cytokines resulting in a hostile environment for NSCs and, subsequently, reduce neurogenesis [89]. In the vicinity of the SVZ, the choroid plexus was shown to tightly regulate age-related behavior. Blocking IFN- $\gamma$  signaling at the choroid plexus attenuated chronic neuroinflammation and restored cognitive functions and hippocampal neurogenesis [90]. Moreover, the choroid plexus transcriptome and secretome, the proteins that circulate in the cerebrospinal fluid, were recently shown to be differentially expressed at different ages, and those changes directly affected NSC behavior and fate [91<sup>••</sup>].

Because of the enormous consequences of aging on NSCs, a lot of effort has focused on identifying mechanisms that could potentially reset the aging clock. Systemic manipulations such as exercise, calorie restriction and heterochronic blood transfer have demonstrated that it is possible to reactivate the intrinsic program in order to rejuvenate NSCs and, consequently, the brain [92,93]. We, and others, have shown that it is possible to rejuvenate aged NSCs by infusing young blood, through heterochronic parabiosis [94] or young plasma injections [95]. Systemic factors found in the blood are pro-aging (CCL11,  $\beta$ 2-microglobulin [96,97]) or rejuvenating (GDF11, TIMP2 [94,98<sup>••</sup>,99,100]), raising the exciting possibility that blocking or inducing these factors, respectively, could help fight the age-associated declines in neurogenesis and cognitive function. Recently, virally mediated overexpression of ten-eleven translocation-2 (Tet2) methylcytosine dioxygenase in the dentate gyrus was able to rescue hippocampal neurogenesis and enhance cognition through an increase in the production of 5-hydroxymethylcytosine, involved in DNA methylation, proposing an epigenetic-mediated rejuvenation of NSC [101]. Furthermore, treatment of 6-month-old mice with resveratrol, a molecule linked to activation of Sirtuins and longevity, resulted in the enhancement of hippocampal neurogenesis, brain plasticity and cognition [102].

## Outlook

Research on NSC biology over the past few years has greatly increased our understanding of the molecular

mechanisms governing NSC behavior within physiological and pathological contexts. The delicate balance between NSC quiescence and activation is easily shifted depending on the different stimuli and could be used to better manipulate NSC fate *in vitro* and *in vivo*. Moreover, recent findings point to the conclusion that aging is not necessarily a permanent state, but could be malleable, and that finding ways to interfere in the cell intrinsic machinery in order to slow down or even reverse this process will be the challenge for years to come.

## Conflict of interest

Nothing declared.

## Acknowledgements

We apologize to colleagues whose work could not be cited due to space limitations. We thank Nicolas Kuperwasser for proofreading the manuscript. L.K. received a fellowship from the Laboratory for Excellence Programme "Revive". This work was supported by Institut Pasteur, Centre National pour la Recherche Scientifique, the life insurance company "AG2R-La-Mondiale", Agence Nationale de la Recherche (ANR-15-CE37-0004-01), NIH US-French Research Proposal Grants #1R01DC015137-01 and ANR-15-NEUC-0004 (Circuit-OPL), the Laboratory for Excellence Programme "Revive" (Investissement d'Avenir, ANR-10-LABX-73) and "Biopsy" (Investissement d'Avenir, ANR-11-IDEX-0004-02).

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Altman J: **Autoradiographic study of degenerative and regenerative proliferation of neuroglia cells with tritiated thymidine.** *Exp Neurol* 1962, **5**:302-318.  
This study shows that human adult neurogenesis declines rapidly during the first years of life and that the generation of hippocampal new neurons is non-existent or rare after adolescence.
2. Eriksson PS *et al.*: **Neurogenesis in the adult human hippocampus.** *Nat Med* 1998, **4**:1313-1317.
3. Spalding KL *et al.*: **Dynamics of hippocampal neurogenesis in adult humans.** *Cell* 2013, **153**:1219-1227.
4. Sorrells SF *et al.*: **Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults.** *Nature* 2018, **555**:377-381.  
•
5. Boldrini M *et al.*: **Human hippocampal neurogenesis persists throughout aging.** *Cell Stem Cell* 2018, **22**:589-599 e5.
6. Doetsch F, Garcia-Verdugo JM, Alvarez-Buylla A: **Cellular composition and three-dimensional organization of the subventricular germinal zone in the adult mammalian brain.** *J Neurosci* 1997, **17**:5046-5061.  
This study identifies the persistence of proliferating neural progenitors, immature and mature granule neurons, and glia throughout human life span.
7. Kuhn HG, Dickinson-Anson H, Gage FH: **Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation.** *J Neurosci* 1996, **16**:2027-2033.
8. Bond AM, Ming GL, Song H: **Adult mammalian neural stem cells and neurogenesis: five decades later.** *Cell Stem Cell* 2015, **17**:385-395.
9. Merkle FT, Mirzadeh Z, Alvarez-Buylla A: **Mosaic organization of neural stem cells in the adult brain.** *Science* 2007, **317**:381-384.

10. Merkle FT *et al.*: **Adult neural stem cells in distinct microdomains generate previously unknown interneuron types.** *Nat Neurosci* 2014, **17**:207-214.
11. Lopez-Juarez A *et al.*: **Gsx2 controls region-specific activation of neural stem cells and injury-induced neurogenesis in the adult subventricular zone.** *Genes Dev* 2013, **27**:1272-1287.
12. Delgado RN, Lim DA: **Embryonic Nkx2.1-expressing neural precursor cells contribute to the regional heterogeneity of adult V-SVZ neural stem cells.** *Dev Biol* 2015, **407**:265-274.
13. Kohwi M *et al.*: **Pax6 is required for making specific subpopulations of granule and periglomerular neurons in the olfactory bulb.** *J Neurosci* 2005, **25**:6997-7003.
14. Hack MA *et al.*: **Neuronal fate determinants of adult olfactory bulb neurogenesis.** *Nat Neurosci* 2005, **8**:865-872.
15. Young KM *et al.*: **Subventricular zone stem cells are heterogeneous with respect to their embryonic origins and neurogenic fates in the adult olfactory bulb.** *J Neurosci* 2007, **27**:8286-8296.
16. Chaker Z, Codega P, Doetsch F: **A mosaic world: puzzles revealed by adult neural stem cell heterogeneity.** *Wiley Interdiscip Rev Dev Biol* 2016, **5**:640-658.
17. Lledo PM, Merkle FT, Alvarez-Buylla A: **Origin and function of olfactory bulb interneuron diversity.** *Trends Neurosci* 2008, **31**:392-400.
18. Galliano E *et al.*: **Embryonic and postnatal neurogenesis produce functionally distinct subclasses of dopaminergic neuron.** *Elife* 2018, **7**.
19. Grelat A *et al.*: **Adult-born neurons boost odor-reward association.** *Proc Natl Acad Sci U S A* 2018, **115**:2514-2519.
20. Faiz M *et al.*: **Adult neural stem cells from the subventricular zone give rise to reactive astrocytes in the cortex after Stroke.** *Cell Stem Cell* 2015, **17**:624-634.
21. Luo Y *et al.*: **Single-cell transcriptome analyses reveal signals to activate dormant neural stem cells.** *Cell* 2015, **161**:1175-1186.
22. Coskun V *et al.*: **CD133+ neural stem cells in the ependyma of mammalian postnatal forebrain.** *Proc Natl Acad Sci U S A* 2008, **105**:1026-1031.
23. Chaker Z *et al.*: **Suppression of IGF-I signals in neural stem cells enhances neurogenesis and olfactory function during aging.** *Aging Cell* 2015, **14**:847-856.
24. Sachewsky N *et al.*: **Primitive neural stem cells in the adult mammalian brain give rise to GFAP-expressing neural stem cells.** *Stem Cell Rep* 2014, **2**:810-824.
25. Fuentealba LC *et al.*: **Embryonic origin of postnatal neural stem cells.** *Cell* 2015, **161**:1644-1655.
26. Furutachi S *et al.*: **Slowly dividing neural progenitors are an embryonic origin of adult neural stem cells.** *Nat Neurosci* 2015, **18**:657-665.
27. Yuzwa SA *et al.*: **Developmental emergence of adult neural stem cells as revealed by single-cell transcriptional profiling.** *Cell Rep* 2017, **21**:3970-3986.
28. Falk S *et al.*: **Time-specific effects of spindle positioning on embryonic progenitor pool composition and adult neural stem cell seeding.** *Neuron* 2017, **93**:777-791 e3.  
 This study shows that spindle orientation in murine embryonic radial glia cells regulates the division type (symmetric or asymmetric) and, in consequence, the numbers of neural stem cells that will remain in the adult age.
29. Kokovay E *et al.*: **VCAM1 is essential to maintain the structure of the SVZ niche and acts as an environmental sensor to regulate SVZ lineage progression.** *Cell Stem Cell* 2012, **11**:220-230.
30. Hu XL *et al.*: **Persistent expression of VCAM1 in radial glial cells is required for the embryonic origin of postnatal neural stem cells.** *Neuron* 2017, **95**:309-325 e6.
31. Pilz GA *et al.*: **Live imaging of neurogenesis in the adult mouse hippocampus.** *Science* 2018, **359**:658-662.
32. Obernier K *et al.*: **Adult neurogenesis is sustained by symmetric self-renewal and differentiation.** *Cell Stem Cell* 2018, **22**:221-234 e8.  
 This study demonstrates that lifelong SVZ/olfactory neurogenesis is sustained mainly through symmetric divisions, decoupling proliferation from differentiation.
33. Calzolari F *et al.*: **Fast clonal expansion and limited neural stem cell self-renewal in the adult subependymal zone.** *Nat Neurosci* 2015, **18**:490-492.
34. Urban N *et al.*: **Return to quiescence of mouse neural stem cells by degradation of a proactivation protein.** *Science* 2016, **353**:292-295.
35. Llorens-Bobadilla E *et al.*: **Single-cell transcriptomics reveals a population of dormant neural stem cells that become activated upon brain injury.** *Cell Stem Cell* 2015, **17**:329-340.
36. Codega P *et al.*: **Prospective identification and purification of quiescent adult neural stem cells from their in vivo niche.** *Neuron* 2014, **82**:545-559.
37. Khatri P *et al.*: **Proliferation and cilia dynamics in neural stem cells prospectively isolated from the SEZ.** *Sci Rep* 2014, **4**:3803.
38. Mich JK *et al.*: **Prospective identification of functionally distinct stem cells and neurosphere-initiating cells in adult mouse forebrain.** *Elife* 2014, **3**:e02669.
39. Dulken BW *et al.*: **Single-cell transcriptomic analysis defines heterogeneity and transcriptional dynamics in the adult neural stem cell lineage.** *Cell Rep* 2017, **18**:777-790.  
 This single-cell transcriptomic analysis identifies intermediate types between quiescent and activated SVZ neural stem cells that have distinct molecular profiles.
40. Shin J *et al.*: **Single-cell RNA-Seq with waterfall reveals molecular cascades underlying adult neurogenesis.** *Cell Stem Cell* 2015, **17**:360-372.
41. Artegiani B *et al.*: **A single-cell RNA sequencing study reveals cellular and molecular dynamics of the hippocampal neurogenic niche.** *Cell Rep* 2017, **21**:3271-3284.  
 Comprehensive resource paper based on single-cell transcriptomic analysis of the hippocampal neurogenic niche uncovering the molecular signature and dynamics of niche cells in the adult hippocampus.
42. Hitoshi S *et al.*: **Notch pathway molecules are essential for the maintenance, but not the generation, of mammalian neural stem cells.** *Genes Dev* 2002, **16**:846-858.
43. Engler A *et al.*: **Notch2 signaling maintains NSC quiescence in the murine ventricular-subventricular zone.** *Cell Rep* 2018, **22**:992-1002.
44. Kawai H *et al.*: **Area-specific regulation of quiescent neural stem cells by Notch3 in the adult mouse subependymal zone.** *J Neurosci* 2017, **37**:11867-11880.
45. Giachino C *et al.*: **Molecular diversity subdivides the adult forebrain neural stem cell population.** *Stem Cells* 2014, **32**:70-84.
46. Chavali M *et al.*: **Non-canonical Wnt signaling regulates neural stem cell quiescence during homeostasis and after demyelination.** *Nat Commun* 2018, **9**:36.
47. Rafalski VA, Mancini E, Brunet A: **Energy metabolism and energy-sensing pathways in mammalian embryonic and adult stem cell fate.** *J Cell Sci* 2012, **125**:5597-5608.
48. Chirivella L *et al.*: **Cyclin-dependent kinase 4 regulates adult neural stem cell proliferation and differentiation in response to insulin.** *Stem Cells* 2017, **35**:2403-2416.
49. Castiglione F *et al.*: **NMR metabolomics for stem cell type discrimination.** *Sci Rep* 2017, **7**:15808.
50. Knobloch M *et al.*: **A fatty acid oxidation-dependent metabolic shift regulates adult neural stem cell activity.** *Cell Rep* 2017, **20**:2144-2155.



Elegant research paper demonstrating the crucial role of fatty acid metabolism in the switch between quiescence and activation of hippocampal neural stem cells.

51. Knobloch M *et al.*: **Metabolic control of adult neural stem cell activity by Fasn-dependent lipogenesis.** *Nature* 2013, **493**:226-230.
  52. Hamilton LK *et al.*: **Aberrant lipid metabolism in the forebrain niche suppresses adult neural stem cell proliferation in an animal model of Alzheimer's disease.** *Cell Stem Cell* 2015, **17**:397-411.
  53. Doetsch F *et al.*: **Subventricular zone astrocytes are neural stem cells in the adult mammalian brain.** *Cell* 1999, **97**:703-716.
  54. Bell RD *et al.*: **Pericytes control key neurovascular functions and neuronal phenotype in the adult brain and during brain aging.** *Neuron* 2010, **68**:409-427.
  55. Culver JC, Vadakkan TJ, Dickinson ME: **A specialized microvascular domain in the mouse neural stem cell niche.** *PLoS ONE* 2013, **8**:e53546.
  56. Lim DA, Alvarez-Buylla A: **The adult ventricular-subventricular zone (V-SVZ) and olfactory bulb (OB) neurogenesis.** *Cold Spring Harb Perspect Biol* 2016, **8**.
  57. Ramirez-Castillejo C *et al.*: **Pigment epithelium-derived factor is a niche signal for neural stem cell renewal.** *Nat Neurosci* 2006, **9**:331-339.
  58. Bjornsson CS *et al.*: **It takes a village: constructing the neurogenic niche.** *Dev Cell* 2015, **32**:435-446.
  59. Garber C *et al.*: **Astrocytes decrease adult neurogenesis during virus-induced memory dysfunction via IL-1.** *Nat Immunol* 2018, **19**:151-161.
  60. Tavazoie M *et al.*: **A specialized vascular niche for adult neural stem cells.** *Cell Stem Cell* 2008, **3**:279-288.
  61. Shen Q *et al.*: **Endothelial cells stimulate self-renewal and expand neurogenesis of neural stem cells.** *Science* 2004, **304**:1338-1340.
  62. Shen Q *et al.*: **Adult SVZ stem cells lie in a vascular niche: a quantitative analysis of niche cell-cell interactions.** *Cell Stem Cell* 2008, **3**:289-300.
  63. Mirzadeh Z *et al.*: **Neural stem cells confer unique pinwheel architecture to the ventricular surface in neurogenic regions of the adult brain.** *Cell Stem Cell* 2008, **3**:265-278.
  64. Crouch EE *et al.*: **Regional and stage-specific effects of prospectively purified vascular cells on the adult V-SVZ neural stem cell lineage.** *J Neurosci* 2015, **35**:4528-4539.
- Rapid purification strategy for endothelial cells and pericytes, demonstrating the importance of both neurogenic and non-neurogenic vasculature in secreting diffusible signals that affect neural stem cell behavior.
65. Crouch EE, Doetsch F: **FACS isolation of endothelial cells and pericytes from mouse brain microregions.** *Nat Protoc* 2018, **13**:738-751.
  66. Kirby ED *et al.*: **Adult hippocampal neural stem and progenitor cells regulate the neurogenic niche by secreting VEGF.** *Proc Natl Acad Sci U S A* 2015, **112**:4128-4133.
  67. Bicker F *et al.*: **Neurovascular EGFL7 regulates adult neurogenesis in the subventricular zone and thereby affects olfactory perception.** *Nat Commun* 2017, **8**:15922.
  68. Peruzzotti-Jametti L *et al.*: **Macrophage-derived extracellular succinate licenses neural stem cells to suppress chronic neuroinflammation.** *Cell Stem Cell* 2018, **22**:355-368 e13.
  69. Lehtinen MK *et al.*: **The cerebrospinal fluid provides a proliferative niche for neural progenitor cells.** *Neuron* 2011, **69**:893-905.
  70. Lehtinen MK, Walsh CA: **Neurogenesis at the brain-cerebrospinal fluid interface.** *Annu Rev Cell Dev Biol* 2011, **27**:653-679.
  71. Huang X *et al.*: **Transventricular delivery of sonic hedgehog is essential to cerebellar ventricular zone development.** *Proc Natl Acad Sci U S A* 2010, **107**:8422-8427.
  72. Daynac M *et al.*: **Hedgehog controls quiescence and activation of neural stem cells in the adult ventricular-subventricular zone.** *Stem Cell Rep* 2016, **7**:735-748.
  73. Delgado AC *et al.*: **Endothelial NT-3 delivered by vasculature and CSF promotes quiescence of subependymal neural stem cells through nitric oxide induction.** *Neuron* 2014, **83**:572-585.
  74. Lepousez G, Nissant A, Lledo PM: **Adult neurogenesis and the future of the rejuvenating brain circuits.** *Neuron* 2015, **86**:387-401.
  75. Bao H *et al.*: **Long-range GABAergic inputs regulate neural stem cell quiescence and control adult hippocampal neurogenesis.** *Cell Stem Cell* 2017, **21**:604-617 e5.
- This study demonstrates the importance of long-range GABAergic signaling onto local parvalbumin neurons to control neural stem cell quiescence.
76. Paul A, Chaker Z, Doetsch F: **Hypothalamic regulation of regionally distinct adult neural stem cells and neurogenesis.** *Science* 2017, **356**:1383-1386.
- This fascinating study reveals that adult neurogenesis can be modulated upon hunger states, via hypothalamic control and mosaic innervation of the SVZ niche.
77. Sierra A *et al.*: **Neuronal hyperactivity accelerates depletion of neural stem cells and impairs hippocampal neurogenesis.** *Cell Stem Cell* 2015, **16**:488-503.
  78. Ben Abdallah NM *et al.*: **Early age-related changes in adult hippocampal neurogenesis in C57 mice.** *Neurobiol Aging* 2010, **31**:151-161.
  79. Lugert S *et al.*: **Quiescent and active hippocampal neural stem cells with distinct morphologies respond selectively to physiological and pathological stimuli and aging.** *Cell Stem Cell* 2010, **6**:445-456.
  80. Encinas JM *et al.*: **Division-coupled astrocytic differentiation and age-related depletion of neural stem cells in the adult hippocampus.** *Cell Stem Cell* 2011, **8**:566-579.
  81. Bouab M *et al.*: **Aging of the subventricular zone neural stem cell niche: evidence for quiescence-associated changes between early and mid-adulthood.** *Neuroscience* 2011, **173**:135-149.
  82. Kimura A *et al.*: **HMGB2 expression is associated with transition from a quiescent to an activated state of adult neural stem cells.** *Dev Dyn* 2018, **247**:229-238.
  83. Apostolopoulou M *et al.*: **Non-monotonic changes in progenitor cell behavior and gene expression during aging of the adult V-SVZ neural stem cell niche.** *Stem Cell Rep* 2017, **9**:1931-1947.
- This study demonstrates that SVZ transcriptome is not modified in a linear manner with aging.
84. Daynac M *et al.*: **Age-related neurogenesis decline in the subventricular zone is associated with specific cell cycle regulation changes in activated neural stem cells.** *Sci Rep* 2016, **6**:21505.
- This study shows cell cycle alterations during aging of SVZ neural stem cells together with distinct molecular profiles.
85. Vilchez D, Saez I, Dillin A: **The role of protein clearance mechanisms in organismal ageing and age-related diseases.** *Nat Commun* 2014, **5**:5659.
  86. Moore DL *et al.*: **A mechanism for the segregation of age in mammalian neural stem cells.** *Science* 2015, **349**:1334-1338.
  87. Leeman DS *et al.*: **Lysosome activation clears aggregates and enhances quiescent neural stem cell activation during aging.** *Science* 2018, **359**:1277-1283.
- Fascinating study demonstrating differences in proteostasis between quiescent and activated neural stem cells, as well as the importance of lysosomal activity for activation and in aging.
88. DeCarolis NA *et al.*: **The role of the microenvironmental niche in declining stem-cell functions associated with biological aging.** *Cold Spring Harb Perspect Med* 2015, **5**.



89. Solano Fonseca R *et al.*: **Neurogenic niche microglia undergo positional remodeling and progressive activation contributing to age-associated reductions in neurogenesis.** *Stem Cells Dev* 2016, **25**:542-555.
  90. Baruch K *et al.*: **Aging-induced type I interferon response at the choroid plexus negatively affects brain function.** *Science* 2014, **346**:89-93.
  91. Silva-Vargas V *et al.*: **Age-dependent niche signals from the choroid plexus regulate adult neural stem cells.** *Cell Stem Cell* 2016.
- This elegant study establishes the crucial role of the choroid plexus and its secretome on SVZ neural stem cell behavior during adult life and aging.
92. Katsimpardi L, Rubin LL: **Young systemic factors as a medicine for age-related neurodegenerative diseases.** *Neurogenesis (Austin)* 2015, **2**:e1004971.
  93. Bouchard J, Villeda SA: **Aging and brain rejuvenation as systemic events.** *J Neurochem* 2015, **132**:5-19.
  94. Katsimpardi L *et al.*: **Vascular and neurogenic rejuvenation of the aging mouse brain by young systemic factors.** *Science* 2014, **344**:630-634.
  95. Villeda SA *et al.*: **Young blood reverses age-related impairments in cognitive function and synaptic plasticity in mice.** *Nat Med* 2014, **20**:659-663.
  96. Villeda SA *et al.*: **The ageing systemic milieu negatively regulates neurogenesis and cognitive function.** *Nature* 2011, **477**:90-94.
  97. Smith LK *et al.*: **Beta2-microglobulin is a systemic pro-aging factor that impairs cognitive function and neurogenesis.** *Nat Med* 2015, **21**:932-937.
  98. Castellano JM *et al.*: **Human umbilical cord plasma proteins revitalize hippocampal function in aged mice.** *Nature* 2017, **544**:488-492.
- This compelling study demonstrates that human umbilical cord plasma revitalizes hippocampal and cognitive function in mice, and reveals TIMP2 as a novel rejuvenating factor for the aged brain.
99. Loffredo FS *et al.*: **Growth differentiation factor 11 is a circulating factor that reverses age-related cardiac hypertrophy.** *Cell* 2013, **153**:828-839.
  100. Sinha M *et al.*: **Restoring systemic GDF11 levels reverses age-related dysfunction in mouse skeletal muscle.** *Science* 2014, **344**:649-652.
  101. Gontier G *et al.*: **Tet2 rescues age-related regenerative decline and enhances cognitive function in the adult mouse brain.** *Cell Rep* 2018, **22**:1974-1981.
  102. Torres-Perez M *et al.*: **Resveratrol enhances neuroplastic changes, including hippocampal neurogenesis, and memory in Balb/C mice at six months of age.** *PLoS ONE* 2015, **10**: e0145687.