



HAL
open science

Genomic Surveillance of *Listeria monocytogenes* in Taiwan, 2014 to 2019

Yu-Huan Tsai, Alexandra Moura, Zi-Qi Gu, Jui-Hsien Chang, Ying-Shu Liao,
Ru-Hsiou Teng, Kuo-Yao Tseng, Dai-Ling Chang, Wei-Ren Liu, Yu-Tsung
Huang, et al.

► **To cite this version:**

Yu-Huan Tsai, Alexandra Moura, Zi-Qi Gu, Jui-Hsien Chang, Ying-Shu Liao, et al.. Genomic Surveillance of *Listeria monocytogenes* in Taiwan, 2014 to 2019. *Microbiology Spectrum*, 2022, 10 (6), 10.1128/spectrum.01825-22 . pasteur-03922803v1

HAL Id: pasteur-03922803

<https://pasteur.hal.science/pasteur-03922803v1>

Submitted on 4 Jan 2023 (v1), last revised 6 Jan 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Genomic surveillance of *Listeria monocytogenes* in Taiwan, 2014-2019

Yu-Huan Tsai,^{a,*} Alexandra Moura,^{b,c,*} Zi-Qi Gu,^a Jui-Hsien Chang,^d Ying-Shu Liao,^d Ru-Hsiou Teng,^d Kuo-Yao Tseng,^a Dai-Ling Chang,^a Wei-Ren Liu,^a Yu-Tsung Huang,^{e,f} Alexandre Leclercq,^{b,c} Hsiu-Jung Lo,^{g,h,i} Marc Lecuit,^{b,c,j#}, Chien-Shun Chiou^{d#}

^aLaboratory of Host-Microbe Interactions and Cell Dynamics, Institute of Microbiology and Immunology, National Yang Ming Chiao Tung University, Taipei, Taiwan

^bInstitut Pasteur, Université Paris Cité, Inserm U1117, Biology of Infection Unit, Paris, France

^cInstitut Pasteur, National Reference Centre and WHO Collaborating Centre *Listeria*, Paris, France

^dCenter for Diagnostics and Vaccine Development, Centers for Disease Control, Taichung, Taiwan

^eDepartments of Laboratory Medicine and Internal Medicine, National Taiwan University Hospital, National Taiwan University College of Medicine, Taipei, Taiwan

^fGraduate Institute of Clinical Laboratory Sciences and Medical Biotechnology, National Taiwan University, Taipei, Taiwan

^gNational Institute of Infectious Disease and Vaccinology, National Health Research Institutes, Miaoli, Taiwan

^hSchool of Dentistry, China Medical University, Taichung, Taiwan

ⁱDepartment of Biological Science and Technology, National Yang Ming Chiao Tung University, Hsinchu, Taiwan

^jNecker-Enfants Malades University Hospital, Division of Infectious Diseases and Tropical Medicine, Institut Imagine, APHP, Paris, France

Running Head: *L. monocytogenes* in Taiwan

Address correspondence to Chien-Shun Chiou, nipmesc@cdc.gov.tw

* These authors contributed equally. Author order was determined by mutual agreement.

These authors share senior authorship. Author order was determined by mutual agreement.

1 **Abstract**

2 *Listeria monocytogenes* is a life-threatening foodborne pathogen. Here we report the genomic
3 characterization of a nationwide dataset of 411 clinical and 82 food isolates collected in Taiwan
4 between 2014 and 2019. The observed incidence of listeriosis increased from 0.83 to 7 cases
5 per million population upon implementation of mandatory notification in 2018. Pregnancy-
6 associated cases accounted for 2.8% of human listeriosis and all-cause 7-day mortality was of
7 11.9% in non-maternal-neonatal listeriosis. *L. monocytogenes* was isolated from 90% of raw
8 pork and 34% of chicken products collected in the supermarkets. Sublineages SL87, SL5 and
9 SL378 accounted for the majority (65%) of clinical cases. SL87 and SL378 were also
10 predominant (57%) in food products. Five cgMLST clusters accounted for 57% clinical cases,
11 suggesting unnoticed outbreaks spanning up to 6 years. Mandatory notification allowed
12 identifying the magnitude of listeriosis in Taiwan. Continuous real-time genomic surveillance
13 will allow reducing contaminating sources and disease burden.

14
15 **Importance**

16 Understanding the phylogenetic relationship between clinical and food isolates is important to
17 identify the transmission routes of foodborne diseases. Here we performed a nationwide study
18 between 2014 and 2019 of both clinical and food *Listeria monocytogenes* isolates and
19 sequenced their genomes. We show a 9-fold increase in listeriosis reporting upon
20 implementation of mandatory notification. We found that sublineages SL87 and SL378
21 predominated among both clinical (50%) and food (57%) isolates, and identified 5 cgMLST
22 clusters accounting for 57% of clinical cases, suggestive of potential protracted sources of
23 contamination over up to 6 years in Taiwan. These findings highlight that mandatory
24 declaration is critical in identifying the burden of listeriosis, and the importance of genome
25 sequencing for a detailed characterization of the pathogenic genotypes circulating in Asia.

26 **Introduction**

27 Listeriosis is a foodborne infection caused by the Gram-positive bacterial pathogen
28 *Listeria monocytogenes*. Human listeriosis is a severe infection characterized by bacteremia,
29 meningoencephalitis and maternal-neonatal infection (1, 2). While most human foodborne
30 infections are associated with a high incidence but by a relative low morbidity and mortality,
31 listeriosis has a low incidence varying between 0.1 and 12 per million people, but a high
32 mortality rate (15-21%), even when active antimicrobial therapy is prescribed (3-5).

33 The population structure of *L. monocytogenes* is highly clonal, composed of four
34 phylogenetic lineages, 15 serotypes that can be grouped into six PCR serogroups (6, 7), and
35 hundreds of clonal complexes (CCs) and sublineages (SLs) as defined by multilocus sequence
36 typing (MLST) and core genome MLST (cgMLST), respectively (8). Multiple listeriosis
37 outbreaks associated with contaminated food products have been reported worldwide (9-13),
38 mainly associated to serogroups IVb (lineage I) and IIa (lineage II). The virulence of *L.*
39 *monocytogenes* is heterogeneous: some CCs are hypervirulent and more efficient in gut
40 colonization (e.g., CC1, CC2, CC4 and CC6 from serogroup IVb), resulting in high clinical
41 frequency; others (e.g., CC121 and CC9, from serogroups IIa and IIc) are hypovirulent, express
42 a truncated form of internalin (InlA) (14, 15), a major virulence factor of *L. monocytogenes*
43 (16, 17). The hypovirulent clones exhibit higher tolerance to surface disinfectants and are more
44 prevalent in food isolates, infecting mainly highly immunocompromised individuals (15, 18).

45 In Asia, the incidence of listeriosis was thought to be low before the early 2000s. However,
46 an increase of listeriosis incidence comparable to that of Western countries has been observed
47 in the last decade (19-23), likely as a consequence of enhanced surveillance. In Taiwan, patient
48 information and isolates of notifiable diseases are collected by the Taiwan Centers for Disease
49 Control. No listeriosis outbreak had been reported prior to mandatory notification and only few
50 cases have been sporadically identified in medical centers, mostly associated with CC87, CC19

51 and CC155 (20, 24). While listeriosis has been a notifiable disease in many Western countries
52 since the 1990s, it was only implemented in Taiwan in January 2018, which led to a significant
53 increase in reported listeriosis cases. Here we provide a comprehensive nationwide overview
54 of listeriosis incidence prior and after mandatory declaration and a detailed genome-based
55 characterization of *L. monocytogenes* genotypes circulating in Taiwan.

56 **Methods**

57 **Sampling**

58 We collected 411 clinical *L. monocytogenes* isolates from 2014 to 2019, of which 333 were
59 obtained in the context of the nationwide mandatory notification between Jan. 1, 2018 and
60 December 31, 2019 by the Taiwan Centers for Disease Control (CDC). The use of mandatory
61 notification data was approved by the ethics committee of Institutional Review Board of
62 Centers for Disease Control, Ministry of Health and Welfare, Taiwan. To assist epidemiological
63 investigations, 280 food samples (113 raw chicken, 117 raw ground pork, and 50 ready-to-eat
64 vegetable mixed salad products; Table S1) were also collected in 2016 and 2019 from
65 supermarkets and open-air markets in the cities where the incidence of human listeriosis was
66 the highest (Taipei, New Taipei and Taichung) and in Hsinchu (Table S1). Each food item of
67 individual markets was sampled once for vegetable products from supermarkets and meat
68 products from open-air markets. Raw meat commodities from supermarkets were sampled
69 twice 2 weeks apart. The raw meat products were sampled from 46 markets of 5 districts in
70 Taichung, 16 markets of 4 districts in Taipei, and 16 markets of 3 districts in New Taipei cities.
71 *L. monocytogenes* isolation from food samples was performed according to the guidelines of
72 the Taiwan food and drug administration (<http://www.fda.gov.tw>). Briefly, 25 g of each food
73 sample was pre-enriched in 100 ml of *Listeria* enrichment broth UVM (CMP, Taiwan) and
74 incubated at 37 °C, 200 rpm for 18 to 24 h. Secondary enrichment was performed by
75 transferring 100 µL of the overnight culture to 10 ml of *Listeria* enrichment Fraser broth. After
76 overnight incubation at 37 °C with 200 rpm agitation, 10 µL of the secondary enrichment broth
77 was seeded onto Chromogenic Listeria Agar (CMP, Taiwan) and incubated at 37 °C for up to
78 48 h. The suspected colonies were purified in Brain Heart Infusion Agar (CMP, Taiwan)
79 cultured at 37 °C for 18 to 24 h. Species identification of the colonies were confirmed by
80 MALDI-TOF mass-spectrometry as previously described (25). To reduce redundancy, isolates

81 obtained from the same sample and belonging to the same CT were considered as duplicates,
82 and only one representative was considered in epidemiological analyses.

83

84 **Whole genome sequencing and sequence analysis**

85 DNA libraries were prepared with Illumina DNA Prep and TruSeq® DNA PCR-Free Low
86 Throughput Library Preparation Kits (Illumina Inc. USA). Whole genome sequencing was
87 performed using Illumina MiSeq sequencing platform (Illumina Inc. USA) with MiSeq
88 Reagent Kit v3 (2X 300 bp). Reads were trimmed with AlienTrimer (26) and assembled with
89 SPAdes 3.11. Reads were trimmed from adapter sequences and non-confident bases using
90 AlienTrimmer v.0.455 (minimum read length of 30 bases and minimum quality Phred score 20,
91 i.e. 99% base call accuracy) and corrected with Musket v.1.156, implemented in fqCleaner
92 v.3.0 (Alexis Criscuolo, Institut Pasteur). High-accuracy assemblies were obtained from using
93 SPAdes 3.11 with the automatic k-mer, --only-assembler and --careful options, to reduce the
94 number of mismatches and short indels and used in downstream analyses (27). Molecular
95 typing based on genoserogrouping, MLST and cgMLST was performed with BIGSdb v.1.30
96 (<https://bigsdbs.pasteur.fr/listeria/>) (8, 28). The presence of plasmids, intact prophages and
97 *Listeria* genomic regions was inferred from the assemblies using MOB-suite v.2.0.1 (29),
98 PHASTER (30) and BIGSdb-*Lm* (<http://bigsdbs.pasteur.fr/listeria/>), respectively. Antibiotic
99 resistance genes were detected using BIGSdb-*Lm* and ABRicate v.1.0.1
100 (<https://github.com/tseemann/abricate>). Minimum spanning trees and single linkage
101 dendrograms were built from cgMLST profiles using BioNumerics v.7.6 and visualized in iTOL.

102

103 **Statistical analysis**

104 The significance of associations of clones with each disease type or region was analyzed with
105 χ^2 tests as described by creating a contingency table for each clone (15).

106

107 **Data availability**

108 All sequence data are available on NCBI Sequence Read Archive under the BioProject

109 accession number PRJNA493675.

110 **Results**

111 **Epidemiology of listeriosis in Taiwan**

112 We included 411 cases of human listeriosis in this study. Before nationwide mandatory
113 notification, only 78 listeriosis cases were reported from 2014 to 2017 in 10 cities of Taiwan,
114 and an incidence of 75 cases per million people was estimated based on total patient admissions
115 to a medical center in Taipei (20). Following mandatory notification, 333 human listeriosis
116 cases were reported in Taiwan in 2018-2019 and disease information was available from 327
117 of them. These comprised 278 (85%) bacteremia cases, 32 (9.8%) CNS infections, nine (2.8%)
118 maternal-neonatal infections, and eight of other infections (peritonitis, $n=5$, cholecystitis, $n=2$,
119 pleurisy, $n=1$) (Fig. 1A and Table 1). The annual incidence was of 7.16 and 6.95 listeriosis cases
120 per million people in 2018 and 2019, respectively (Fig. 1A). Cases were mainly in the main
121 island of Taiwan, with very few cases in the offshore islands (2 cases in Penghu and 1 case in
122 Kinmen) (Fig. 1B).

123 The median age of the patients was 68.5 years (range, 0-96 years) for the bacteremia cases,
124 65.5 years (range, 20-93 years) for the CNS infections, 60 years (range 41-87 years) for the
125 peritonitis and cholecystitis, and 30 years (range, 26-38 years) for the maternal infections
126 (Table 1). Diarrhea was noted in 5.71% of the patients and only in bacteremia patients (6.47%)
127 (Table 1). Among the 318 non-maternal-neonatal listeriosis cases, 166 (52.2%) were female
128 (Table 1). All-cause 7-day mortality was 11.9% in non-perinatal listeriosis (Table 1).

129

130 **Prevalence of *L. monocytogenes* in food samples**

131 *L. monocytogenes* was isolated from raw chicken products (25 of 113, 22.12%) and raw ground
132 pork products (69 of 117, 58.97%) (Table S1). Notably, 90.41% (66/73) of raw ground pork
133 products in the supermarkets were positive for *L. monocytogenes*, whereas only 6.8% (3/44) of
134 those sampled in the open-air markets were positive for *L. monocytogenes*. In the case of

135 chicken products, 33.8% (25/74) of those sold in supermarkets were positive for *L.*
136 *monocytogenes*, and all the chicken meat sampled from open-air markets were negative for *L.*
137 *monocytogenes*. By contrast, all the vegetable products sampled in 2019 ($n=50$) were negative
138 for *L. monocytogenes*.

139

140 **Molecular diversity of *L. monocytogenes* clinical and food isolates**

141 A total of 493 isolates were sequenced (Table S2). Clinical isolates ($n=411$) were distributed
142 across lineages I ($n=267$, 65%) and II ($n=144$, 35%) and 22 different cgMLST sublineages (of
143 20 MLST CCs). SL87 ($n=147$, 36%), SL5 ($n=62$, 15%) and SL378 ($n=58$, 14%) were the most
144 prevalent sublineages, accounting for 65% of clinical cases (Fig. S1B). SL5 was associated to
145 CNS infection (odds ratio 2.495, $p=0.0373$, Fig. S1A).

146 Food isolates ($n=82$) were distributed across lineages I ($n=31$, 38%) and II ($n=51$, 62%) and
147 eight different cgMLST sublineages (of 7 CCs). The three most prevalent sublineages were
148 SL87 ($n=26$, 32%), SL378 ($n=21$, 26%) and SL9 ($n=15$, 18%), which accounted for 76% of
149 the food isolates. SL87, SL378, SL155 and SL9 predominated the raw meat products ($n=72$,
150 87.8%, Fig. S1B and S1C). SL155 was only isolated from raw ground pork products ($n=10$,
151 20.4% of the pork products) and SL8 was only isolated from raw chicken products ($n=4$, 12.1%
152 of the chicken products) (Fig. S1C). Although frequently isolated from listeriosis patients,
153 neither SL5 nor SL1 was isolated from the food products (Fig. S1B).

154 A total of 117 cgMLST types (CT) were identified: 96 ($n=291$ isolates) found exclusively in
155 clinical cases, 14 exclusively in food ($n=18$) and seven found in both human and food isolates
156 ($n=184$). All isolates collected from the same food sample belonged to the same CT, except in
157 two raw ground pork products sampled in Taichung, for which two different strains were
158 observed in each (cgMLST types L1-SL87-ST87-CT58 and L2-SL378-ST378-CT4349, and
159 L1-SL3-ST3-CT5507 and L1-SL288-ST330-CT5502).

160 There were 26 CTs (26/103, 25%) comprising more than one clinical case and 22 of which
161 spanning multiple years (Table 2), suggesting high prevalence of common CTs for several years,
162 which may imply unnoticed long-term outbreaks. There were five cgMLST types L1-SL87-
163 ST87-CT58 ($n=54$, 14%), L1-SL87-ST87-CT4373 ($n=53$, 10%), L1-SL5-ST5-CT4358 ($n=42$,
164 13%), L1-SL87-ST87-CT4374 ($n=28$, 7%) and L2-SL378-ST378-CT4349 ($n=57$, 13%)
165 accounting for 57% of human listeriosis, and spanned up to 6 years (Table 2 and Fig. 2B).
166 These CTs also predominated in CNS and maternal-neonatal infections in 2018-2019 for which
167 disease information was available (62.5%, Fig. S1D). While L1-SL87-ST87-CT58, L1-SL87-
168 ST87-CT4373 and L2-SL378-ST378-CT4349 were widely spread across regions and did not
169 exhibit a city-specific distribution, L1-SL5-ST5-CT4358 (odds ratio 8.274, $p < 0.0001$) and L1-
170 SL87-ST87-CT4374 (odds ratio 4.084, $p = 0.0024$) were highly associated to Taichung city (Fig.
171 S1E). In addition, 10 CTs (10/117, 8.5%) detected in this study have been previously reported
172 in the context of listeriosis surveillance in North America, Europe, Oceania and Asia (Table
173 S3).

174

175 **Virulence and stress resistance genetic traits**

176 Relevant traits of virulence and resistance were analyzed in all clinical and food isolates.
177 The presence of virulence traits was consistent with isolate phylogeny. All the isolates carry
178 intact *Listeria* Pathogenicity Island 1 (LIPI-1), which contains the virulence transcription
179 regulator *prfA*, the listeriolysin gene *hly* and multiple genes involved in intracellular vacuole
180 disruption (31) (Fig. 3). LIPI-3, which includes the listeriolysin S gene targeting intestinal
181 bacteria (32, 33), was found in 11% (55/493) of the isolates in SL1, SL3, SL219, SL224, SL288
182 and SL455 of lineage I (Fig. 3, Table S2). LIPI-4, which contributes to *L. monocytogenes*
183 placental and blood-CNS barrier crossing in CC4 (15), is present in 35% (175/493) of the
184 isolates in SL87 (CC87, serogroup IIb), SL219 (CC4, serogroup IVb) and SL455 (CC455,

185 serogroup IVb) (Fig. 3). Almost all the isolates (473/493, 96%) carry an *inlA* gene encoding
186 for full-length internalin, a *L. monocytogenes* invasion protein involved in the crossing of the
187 intestinal and placental barriers (14, 16, 17) (Fig. 3). In the remaining 4% isolates, mutations
188 in *inlA* leading to premature stop codons (PMSC) were detected, resulting in the expression of
189 a truncated InlA. InlA mutations identified concerned all SL9 isolates [(previously reported
190 PMSC_4 ($n=15$), PMSC_8 ($n=1$) and PMSC_11 ($n=2$)] (34, 35), and two SL3 isolates that
191 carry a new PMSC. The new *inlA* PMSC [termed PMSC_32, following the existing
192 nomenclature (8, 36)] consists of a C→A substitution at nucleotide position 1041, within the
193 leucine-rich repeat coding domain, leading to a shorter unanchored InlA protein of 346 amino
194 acids.

195 Acquired resistance genes towards antibiotics were found in 52 isolates, coding for resistance
196 towards tetracycline (*tetM*, $n=50$; 10.1%) and aminoglycosides (*aadA2*, $n=1$; *aacA4*, $n=1$) (Fig.
197 3). Genetic traits involved in resistance towards metals, antiseptics and/or biofilm formation
198 were also found. These included the arsenic resistance listeria genomic islands LGI-2 ($n=27$,
199 in SL1 and SL2) and LGI-3 ($n=21$, in SL101) involved in metal resistance, the stress survival
200 islet 1 (SSI-1, $n=157$, mainly in SL3, SL5, SL8, SL9, and SL155), and the resistance cassette
201 *bcrABC* conferring resistance to benzalkonium chloride ($n=4$, in SL9 and SL155) (37-40) (Fig.
202 3). Disrupted *comK*, proposed to be associated with biofilm formation on food-conditioned
203 surfaces (41), was found in 29% (143/493) of the isolates (Fig. 3), and was significantly more
204 prevalent in food isolates as compared to clinical isolates ($p<0.05$, Chi square test). All these
205 traits have been shown to confer tolerance to environmental stress in *L. monocytogenes*,
206 facilitating its long-term persistence in production environments (38, 39, 42, 43).

207

208 **Discussion**

209 Here we described the findings of a genomic nationwide study on clinical and food *L.*

210 *monocytogenes* isolates over a 6-year period, prior to and after mandatory notification of
211 human listeriosis cases. The incidence of listeriosis has sharply increased from 0.83 to 7 cases
212 per million population after the implementation of mandatory notification of listeriosis in 2018,
213 illustrating the effectiveness of mandatory notification. As a result, since July 2021, the Taiwan
214 Food and Drug Administration has extended food monitoring of *Listeria* from infant and dairy
215 products to refrigerated and ready-to-eat food products (<https://consumer.fda.gov.tw>). As
216 compared to the regulation before this date where food products were randomly sampled once,
217 an extended sampling plan (5 to 10 sampling times per location according to the guidelines on
218 sampling CAC GL 50-2004) is now in place for food microorganisms. This may help to identify
219 more contamination sources of *L. monocytogenes* in different food products.

220 As compared to Europe and North America where pregnancy-associated listeriosis accounts
221 for 9-14% of human listeriosis, only 2.8% of listeriosis cases in Taiwan were pregnancy-
222 associated (2, 44, 45). This might be due to the difference in placental barrier crossing ability
223 of circulating clones, and/or to the low total fertility rate in Taiwan, which is ranked the last in
224 the world (<https://www.cia.gov/the-world-factbook/field/total-fertility-rate/>). Median patient
225 age (67 years) in non-maternal-neonatal listeriosis was slightly lower than previously reported
226 in Europe and North America (67-77 years) (2, 44, 45). Here, all-cause mortality (11.9%, within
227 7 days) in non-maternal-neonatal listeriosis was similar to that in Germany (13%, within 7 days)
228 but lower than previously reported in France (30-46%, 90 days) and the United States (21%)
229 (2, 44, 45), although these differences might be due to varied follow-up periods. As observed
230 in our previous study of listeriosis cases in a medical center in Taiwan, the all-cause mortality
231 was 14.8% within 7 days and 25.2% within 30 days (20, 45). Thus, the mortality of listeriosis
232 is likely underestimated when considering a short follow-up period, as listeriosis can have an
233 impact on mortality at least up to three months (2).

234 While in Western countries serogroup IVb is the main cause of human listeriosis (9, 15, 18), in

235 Taiwan serogroups IIb (SL87/CC87, SL5/CC5, SL378/CC19) and IIa (SL155/CC155) are
236 responsible for the majority of human listeriosis cases (74.5%, 306 of 411). Previously,
237 SL87/CC87 (serogroup IIb) has been also reported to be responsible for 28-34% of clinical
238 infections in China and highly prevalent in raw meat products (46-48). These geographical
239 differences may result from different food habits in Asia, since serotype IVb (particularly
240 SL1/CC1) appears to be strongly associated to dairy products (15, 18) and dairy consumption
241 in Taiwan is relatively low across all age groups (49).

242 Whole genome sequencing of isolates collected prior to and after mandatory notification
243 allowed the identification of multiple local and nationwide clusters of cases spanning up to six
244 years. For instance, L1-SL87-ST87-CT58 (CC87, serogroup IIb) and L2-SL378-ST378-
245 CT4349 (CC19, serogroup IIa) comprised both clinical and food (chicken and pork) isolates
246 distributed across multiple geographic regions from 2014 to 2019. The presence of identical
247 cgMLST types across locations and along the years suggests long-term contamination of either
248 food-processing plants, meat refrigeration and/or large-scale distribution facilities, although
249 trace-back investigations are needed to confirm the source of the outbreaks. In contrast, the
250 region-associated distribution of L1-SL5-ST5-CT4358 to Taichung city suggests a local source
251 of contamination. Food sampling extended to abattoir environments and meat distribution
252 chains will help to identify the sources of contamination, if any (50). While up to 82% (96/117)
253 of cgMLST types in this study were exclusively seen in clinical isolates, a more extensive food
254 sampling will also benefit to identification of the sources of contamination, which remain
255 unknown for most clusters observed (82%, 96/117). The introduction of new regulation on food
256 microorganisms sampling in 2021 will help to identify the environmental niches of *Listeria* in
257 Taiwan.

258 The majority (91%) of cgMLST types found were unique to this study, reinforcing the
259 importance of comprehensive national surveillance programs to control *Listeria* transmission

260 (51). Nevertheless, 10 CTs have been previously reported within the scope of listeriosis
261 surveillance in other countries (Table S3), suggestive of punctual cross-country contaminations
262 or, in the case L1-SL87-ST87-CT58, of contamination sources with large scale distributions.
263 This study constitutes the first nationwide genome-based characterization of *L. monocytogenes*
264 genotypes circulating in Taiwan, and highlights the importance of mandatory declaration in
265 understanding listeriosis incidence. Continuous real-time genome-based surveillance coupled
266 with epidemiological investigations and regular surveillance of food products for *L.*
267 *monocytogenes* will allow to reducing contaminating sources and lower the disease burden in
268 Taiwan.

269

270 **Acknowledgments**

271 This work was supported by Ministry of Health and Welfare, Taiwan [grant numbers DOH102-
272 DC-2102 (2013), MOHW108-CDC-C-315-112520 (2019), MOHW109-CDC-C-315-122415
273 (2020)], Ministry of Science and Technology, Taiwan (grant number MOST 109-2320-B-010-
274 039), National Health Research Institutes, Taiwan (grant number NHRI-EX111-11120SC),
275 Institut Pasteur and Inserm.

276

277 **Potential conflicts of interest**

278 The authors have no conflicts to disclose.

279 **References**

- 280 1. Lecuit M. 2007. Human listeriosis and animal models. *Microbes Infect* 9:1216-25.
- 281 2. Charlier C, Perrodeau E, Leclercq A, Cazenave B, Pilmis B, Henry B, Lopes A,
282 Maury MM, Moura A, Goffinet F, Dieye HB, Thouvenot P, Ungeheuer MN,
283 Tourdjman M, Goulet V, de Valk H, Lortholary O, Ravaud P, Lecuit M, group Ms.
284 2017. Clinical features and prognostic factors of listeriosis: the MONALISA national
285 prospective cohort study. *Lancet Infect Dis* 17:510-519.
- 286 3. European Centre for Disease Prevention and Control. 2020. Listeriosis. In: ECDC.
287 Annual epidemiological report for 2017 doi:
288 [https://www.ecdc.europa.eu/en/publications-data/listeriosis-annual-epidemiological-](https://www.ecdc.europa.eu/en/publications-data/listeriosis-annual-epidemiological-report-2017)
289 [report-2017](https://www.ecdc.europa.eu/en/publications-data/listeriosis-annual-epidemiological-report-2017). ECDC, Stockholm.
- 290 4. Tack DM, Ray L, Griffin PM, Cieslak PR, Dunn J, Rissman T, Jervis R, Lathrop S,
291 Muse A, Duwell M, Smith K, Tobin-D'Angelo M, Vugia DJ, Zablotsky Kufel J,
292 Wolpert BJ, Tauxe R, Payne DC. 2020. Preliminary Incidence and Trends of
293 Infections with Pathogens Transmitted Commonly Through Food - Foodborne
294 Diseases Active Surveillance Network, 10 U.S. Sites, 2016-2019. *MMWR Morb*
295 *Mortal Wkly Rep* 69:509-514.
- 296 5. Ray LC, Collins JP, Griffin PM, Shah HJ, Boyle MM, Cieslak PR, Dunn J, Lathrop S,
297 McGuire S, Rissman T, Scallan Walter EJ, Smith K, Tobin-D'Angelo M, Wymore K,
298 Kufel JZ, Wolpert BJ, Tauxe R, Payne DC. 2021. Decreased Incidence of Infections
299 Caused by Pathogens Transmitted Commonly Through Food During the COVID-19
300 Pandemic - Foodborne Diseases Active Surveillance Network, 10 U.S. Sites, 2017-
301 2020. *MMWR Morb Mortal Wkly Rep* 70:1332-1336.
- 302 6. Doumith M, Buchrieser C, Glaser P, Jacquet C, Martin P. 2004. Differentiation of the
303 major *Listeria monocytogenes* serovars by multiplex PCR. *J Clin Microbiol* 42:3819-

- 304 22.
- 305 7. Yin Y, Yao H, Doijad S, Kong S, Shen Y, Cai X, Tan W, Wang Y, Feng Y, Ling Z,
306 Wang G, Hu Y, Lian K, Sun X, Liu Y, Wang C, Jiao K, Liu G, Song R, Chen X, Pan
307 Z, Loessner MJ, Chakraborty T, Jiao X. 2019. A hybrid sub-lineage of *Listeria*
308 *monocytogenes* comprising hypervirulent isolates. *Nat Commun* 10:4283.
- 309 8. Moura A, Criscuolo A, Pouseele H, Maury MM, Leclercq A, Tarr C, Bjorkman JT,
310 Dallman T, Reimer A, Enouf V, Larssonneur E, Carleton H, Bracq-Dieye H, Katz LS,
311 Jones L, Touchon M, Tourdjman M, Walker M, Stroika S, Cantinelli T, Chenal-
312 Francisque V, Kucerova Z, Rocha EP, Nadon C, Grant K, Nielsen EM, Pot B, Gerner-
313 Smidt P, Lecuit M, Brisse S. 2016. Whole genome-based population biology and
314 epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol* 2:16185.
- 315 9. Cossart P. 2011. Illuminating the landscape of host-pathogen interactions with the
316 bacterium *Listeria monocytogenes*. *Proc Natl Acad Sci U S A* 108:19484-91.
- 317 10. Halbedel S, Wilking H, Holzer A, Kleta S, Fischer MA, Luth S, Pietzka A, Huhulescu
318 S, Lachmann R, Krings A, Ruppitsch W, Leclercq A, Kamphausen R, Meincke M,
319 Wagner-Wiening C, Contzen M, Kraemer IB, Al Dahouk S, Allerberger F, Stark K,
320 Flieger A. 2020. Large Nationwide Outbreak of Invasive Listeriosis Associated with
321 Blood Sausage, Germany, 2018-2019. *Emerg Infect Dis* 26:1456-1464.
- 322 11. Luth S, Halbedel S, Rosner B, Wilking H, Holzer A, Roedel A, Dieckmann R, Vincze
323 S, Prager R, Flieger A, Al Dahouk S, Kleta S. 2020. Backtracking and forward
324 checking of human listeriosis clusters identified a multiclonal outbreak linked to
325 *Listeria monocytogenes* in meat products of a single producer. *Emerg Microbes Infect*
326 9:1600-1608.
- 327 12. Self JL, Conrad A, Stroika S, Jackson A, Whitlock L, Jackson KA, Beal J, Wellman
328 A, Fatica MK, Bidol S, Huth PP, Hamel M, Franklin K, Tschetter L, Kopko C, Kirsch

- 329 P, Wise ME, Basler C. 2019. Multistate Outbreak of Listeriosis Associated with
330 Packaged Leafy Green Salads, United States and Canada, 2015-2016. *Emerg Infect*
331 *Dis* 25:1461-1468.
- 332 13. Thomas J, Govender N, McCarthy KM, Erasmus LK, Doyle TJ, Allam M, Ismail A,
333 Ramalwa N, Sekwadi P, Ntshoe G, Shonhiwa A, Essel V, Tau N, Smouse S, Ngomane
334 HM, Disenyeng B, Page NA, Govender NP, Duse AG, Stewart R, Thomas T,
335 Mahoney D, Tourdjman M, Disson O, Thouvenot P, Maury MM, Leclercq A, Lecuit
336 M, Smith AM, Blumberg LH. 2020. Outbreak of Listeriosis in South Africa
337 Associated with Processed Meat. *N Engl J Med* 382:632-643.
- 338 14. Jacquet C, Doumith M, Gordon JI, Martin PM, Cossart P, Lecuit M. 2004. A
339 molecular marker for evaluating the pathogenic potential of foodborne *Listeria*
340 *monocytogenes*. *J Infect Dis* 189:2094-100.
- 341 15. Maury MM, Tsai YH, Charlier C, Touchon M, Chenal-Francisque V, Leclercq A,
342 Criscuolo A, Gaultier C, Roussel S, Brisabois A, Disson O, Rocha EPC, Brisse S,
343 Lecuit M. 2016. Uncovering *Listeria monocytogenes* hypervirulence by harnessing its
344 biodiversity. *Nat Genet* 48:308-313.
- 345 16. Lecuit M, Vandormael-Pournin S, Lefort J, Huerre M, Gounon P, Dupuy C, Babinet
346 C, Cossart P. 2001. A transgenic model for listeriosis: role of internalin in crossing the
347 intestinal barrier. *Science* 292:1722-5.
- 348 17. Disson O, Grayo S, Huillet E, Nikitas G, Langa-Vives F, Dussurget O, Ragon M, Le
349 Monnier A, Babinet C, Cossart P, Lecuit M. 2008. Conjugated action of two species-
350 specific invasion proteins for fetoplacental listeriosis. *Nature* 455:1114-8.
- 351 18. Maury MM, Bracq-Dieye H, Huang L, Vales G, Lavina M, Thouvenot P, Disson O,
352 Leclercq A, Brisse S, Lecuit M. 2019. Hypervirulent *Listeria monocytogenes* clones'
353 adaption to mammalian gut accounts for their association with dairy products. *Nat*

- 354 Commun 10:2488.
- 355 19. Fan Z, Xie J, Li Y, Wang H. 2019. Listeriosis in mainland China: A systematic review.
356 Int J Infect Dis 81:17-24.
- 357 20. Huang YT, Kuo YW, Lee MR, Tsai YH, Teng LJ, Tsai MS, Liao CH, Hsueh PR. 2021.
358 Clinical and molecular epidemiology of human listeriosis in Taiwan. Int J Infect Dis
359 doi:10.1016/j.ijid.2021.01.056.
- 360 21. Huang YT, Liao CH, Yang CJ, Teng LJ, Wang JT, Hsueh PR. 2011. Listeriosis,
361 Taiwan, 1996-2008. Emerg Infect Dis 17:1731-3.
- 362 22. Miya S, Takahashi H, Nakagawa M, Kuda T, Igimi S, Kimura B. 2015. Genetic
363 characteristics of Japanese clinical *Listeria monocytogenes* isolates. PLoS One
364 10:e0122902.
- 365 23. Okutani A, Okada Y, Yamamoto S, Igimi S. 2004. Nationwide survey of human
366 *Listeria monocytogenes* infection in Japan. Epidemiol Infect 132:769-72.
- 367 24. Huang YT, Ko WC, Chan YJ, Lu JJ, Tsai HY, Liao CH, Sheng WH, Teng LJ, Hsueh
368 PR. 2015. Disease Burden of Invasive Listeriosis and Molecular Characterization of
369 Clinical Isolates in Taiwan, 2000-2013. PLoS One 10:e0141241.
- 370 25. Thouvenot P, Vales G, Bracq-Dieye H, Tessaud-Rita N, Maury MM, Moura A, Lecuit
371 M, Leclercq A. 2018. MALDI-TOF mass spectrometry-based identification of
372 *Listeria* species in surveillance: A prospective study. J Microbiol Methods 144:29-32.
- 373 26. Criscuolo A, Brisse S. 2013. AlienTrimmer: a tool to quickly and accurately trim off
374 multiple short contaminant sequences from high-throughput sequencing reads.
375 Genomics 102:500-6.
- 376 27. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
377 Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,
378 Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and

- 379 its applications to single-cell sequencing. *J Comput Biol* 19:455-77.
- 380 28. Jolley KA, Maiden MC. 2010. BIGSdb: Scalable analysis of bacterial genome
381 variation at the population level. *BMC Bioinformatics* 11:595.
- 382 29. Robertson J, Nash JHE. 2018. MOB-suite: software tools for clustering,
383 reconstruction and typing of plasmids from draft assemblies. *Microb Genom* 4.
- 384 30. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER:
385 a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16-
386 21.
- 387 31. Vazquez-Boland JA, Dominguez-Bernal G, Gonzalez-Zorn B, Kreft J, Goebel W.
388 2001. Pathogenicity islands and virulence evolution in *Listeria*. *Microbes Infect*
389 3:571-84.
- 390 32. Cotter PD, Draper LA, Lawton EM, Daly KM, Groeger DS, Casey PG, Ross RP, Hill
391 C. 2008. Listeriolysin S, a novel peptide haemolysin associated with a subset of
392 lineage I *Listeria monocytogenes*. *PLoS Pathog* 4:e1000144.
- 393 33. Quereda JJ, Nahori MA, Meza-Torres J, Sachse M, Titos-Jimenez P, Gomez-Laguna
394 J, Dussurget O, Cossart P, Pizarro-Cerda J. 2017. Listeriolysin S Is a Streptolysin S-
395 Like Virulence Factor That Targets Exclusively Prokaryotic Cells In Vivo. *mBio* 8.
- 396 34. Rousseaux S, Olier M, Lemaitre JP, Piveteau P, Guzzo J. 2004. Use of PCR-restriction
397 fragment length polymorphism of *inlA* for rapid screening of *Listeria monocytogenes*
398 strains deficient in the ability to invade Caco-2 cells. *Appl Environ Microbiol*
399 70:2180-5.
- 400 35. Felicio MT, Hogg T, Gibbs P, Teixeira P, Wiedmann M. 2007. Recurrent and sporadic
401 *Listeria monocytogenes* contamination in alheiras represents considerable diversity,
402 including virulence-attenuated isolates. *Appl Environ Microbiol* 73:3887-95.
- 403 36. Gelbicova T, Kolackova I, Pantucek R, Karpiskova R. 2015. A novel mutation leading

- 404 to a premature stop codon in inlA of *Listeria monocytogenes* isolated from neonatal
405 listeriosis. *New Microbiol* 38:293-6.
- 406 37. Ortiz S, Lopez-Alonso V, Rodriguez P, Martinez-Suarez JV. 2016. The Connection
407 between Persistent, Disinfectant-Resistant *Listeria monocytogenes* Strains from Two
408 Geographically Separate Iberian Pork Processing Plants: Evidence from Comparative
409 Genome Analysis. *Appl Environ Microbiol* 82:308-17.
- 410 38. Ryan S, Begley M, Hill C, Gahan CG. 2010. A five-gene stress survival islet (SSI-1)
411 that contributes to the growth of *Listeria monocytogenes* in suboptimal conditions. *J*
412 *Appl Microbiol* 109:984-95.
- 413 39. Lee S, Ward TJ, Jima DD, Parsons C, Kathariou S. 2017. The Arsenic Resistance-
414 Associated *Listeria* Genomic Island LGI2 Exhibits Sequence and Integration Site
415 Diversity and a Propensity for Three *Listeria monocytogenes* Clones with Enhanced
416 Virulence. *Appl Environ Microbiol* 83.
- 417 40. Palma F, Brauge T, Radomski N, Mallet L, Felten A, Mistou MY, Brisabois A,
418 Guillier L, Midelet-Bourdin G. 2020. Dynamics of mobile genetic elements of
419 *Listeria monocytogenes* persisting in ready-to-eat seafood processing plants in France.
420 *BMC Genomics* 21:130.
- 421 41. Verghese B, Lok M, Wen J, Alessandria V, Chen Y, Kathariou S, Knabel S. 2011.
422 comK prophage junction fragments as markers for *Listeria monocytogenes* genotypes
423 unique to individual meat and poultry processing plants and a model for rapid niche-
424 specific adaptation, biofilm formation, and persistence. *Appl Environ Microbiol*
425 77:3279-92.
- 426 42. Hein I, Klinger S, Dooms M, Flekna G, Stessl B, Leclercq A, Hill C, Allerberger F,
427 Wagner M. 2011. Stress survival islet 1 (SSI-1) survey in *Listeria monocytogenes*
428 reveals an insert common to *Listeria innocua* in sequence type 121 *L. monocytogenes*

- 429 strains. Appl Environ Microbiol 77:2169-73.
- 430 43. Dutta V, Elhanafi D, Kathariou S. 2013. Conservation and distribution of the
431 benzalkonium chloride resistance cassette *bcrABC* in *Listeria monocytogenes*. Appl
432 Environ Microbiol 79:6067-74.
- 433 44. Centers for Disease C, Prevention. 2013. Vital signs: *Listeria* illnesses, deaths, and
434 outbreaks--United States, 2009-2011. MMWR Morb Mortal Wkly Rep 62:448-52.
- 435 45. Wilking H, Lachmann R, Holzer A, Halbedel S, Flieger A, Stark K. 2021. Ongoing
436 high incidence and case-fatality rates for invasive listeriosis, Germany, 2010-2019.
437 Emerg Infect Dis 27:2485-2488.
- 438 46. Lu B, Yang J, Gao C, Li D, Cui Y, Huang L, Chen X, Wang D, Wang A, Liu Y, Li Y,
439 Zhang Z, Jiao M, Xu H, Song Y, Fu B, Xu L, Yang Q, Ning Y, Wang L, Bao C, Luo G,
440 Wu H, Yang T, Li C, Tang M, Wang J, Guo W, Zeng J, Zhong W. 2021. Listeriosis
441 cases and genetic diversity of their *L. monocytogenes* isolates in China, 2008-2019.
442 Front Cell Infect Microbiol 11:608352.
- 443 47. Yin Y, Doijad S, Wang W, Lian K, Pan X, Korycinski I, Hu Y, Tan W, Ye S, Wang Z,
444 Pan Z, Chakraborty T, Jiao X. 2020. Genetic Diversity of *Listeria monocytogenes*
445 Isolates from Invasive Listeriosis in China. Foodborne Pathog Dis 17:215-227.
- 446 48. Zhang J, Cao G, Xu X, Allard M, Li P, Brown E, Yang X, Pan H, Meng J. 2016.
447 Evolution and Diversity of *Listeria monocytogenes* from Clinical and Food Samples
448 in Shanghai, China. Front Microbiol 7:1138.
- 449 49. Lee MS, Huang LY, Chen MC, Wahlqvist ML. 2009. The demography of food in
450 health security: current experience with dairy consumption in Taiwan. Asia Pac J Clin
451 Nutr 18:585-9.
- 452 50. Lin CH, Adams PJ, Huang JF, Sun YF, Lin JH, Robertson ID. 2021. Contamination of
453 chicken carcasses and the abattoir environment with *Listeria monocytogenes* in

454 Taiwan. Br Poult Sci 62:701-709.

455 51. Moura A, Lefrancq N, Wirth T, Leclercq A, Borges V, Gilpin B, Dallman TJ, Frey J,
456 Franz E, Nielsen EM, Thomas J, Pightling A, Howden BP, Tarr CL, Gerner-Smidt P,
457 Cauchemez S, Salje H, Brisse S, Lecuit M, Listeria CCSG. 2021. Emergence and
458 global spread of *Listeria monocytogenes* main clinical clonal complex. Sci Adv
459 7:eabj9805.

460

461 Table 1. Demographic data and *L. monocytogenes* clone distribution of the 327 human
 462 listeriosis cases between Jan. 1, 2018 and Dec. 31, 2019.

Disease	Bacteremia	CNS infection	Peritonitis & cholecystitis	Maternal-neonatal infection	All diseases
Total	278	32	7	9	327
Clinical information					
Female	145 (52%)	17 (53%)	3 (43%)	7 (77.8%)	173 (52.9%)
*Age (year)	68.5 (0, 96)	65.5 (20, 93)	60 (41, 87)	30 (26, 38) [†]	67 (0, 96) [†]
Diarrhea	18 (6.47%)	0 (0%)	0 (0%)	0 (0%)	19 (5.71%)
7-day mortality (%)	34 (12.2%)	4 (12.5%)	1 (14.3%)	0 [‡]	39 (11.9%) [‡]
Lineages					
L1	178 (64%)	27 (84.4%)	4 (57.1%)	6 (66.7%)	215 (66%)
L2	100 (36%)	5 (35.6%)	3 (42.9%)	3 (33.3%)	112 (34%)
Serogroups					
IVb	22 (7.9%)	3 (9.4%)	1 (14.2%)	0 (0%)	26 (7.95%)
IIb	156 (56.1%)	24 (75%)	3 (42.9%)	6 (66.7%)	189 (57.80%)
IIa	97 (34.9%)	5 (15.6)	3 (42.9%)	3 (33.3%)	109 (33.33%)
IIc	3 (1.1%)	0 (0%)	0 (0%)	0 (0%)	3 (0.92%)
Sublineages (Clonal Complexes)					
SL87 (CC87)	104 (37.4%)	12 (37.5%)	2 (28.6%)	4 (44.5%)	122 (37.31%)
SL5 (CC5)	37 (13.3%)	9 (28.13%)	1 (14.3%)	2 (22.2%)	49 (14.98%)
SL378 (CC19)	41 (14.7%)	2 (6.25%)	1 (14.3%)	1 (11.1%)	45 (13.76%)
SL155 (CC155)	22 (7.9%)	2 (6.25%)	0 (0%)	2 (22.2%)	27 (8.26%)
SL1 (CC1)	13 (4.7%)	1 (3.12%)	0 (0%)	0 (0%)	14 (4.28%)
Other SLs/CCs	61 (22.0%)	6 (18.75%)	3 (42.9%)	0 (0%)	70 (21.41%)

463 Data are represented as number (percentage) unless described.

464 *Age is presented as median (min, max).

465 [†] The age of neonatal infection was excluded.

466 [‡] Perinatal infections were excluded.

467 Table 2. cgMLST types detected in this study comprising more than 2 isolates ($n=33$ out of
 468 117). cgMLST types with 10 or more clinical isolates are highlighted in bold.

cgMLST type (clonal complex, serogroup)	No. isolates (%, $N=493$)	No. Clinical isolates (%, $n=411$)	No. Food isolates (food type*)	Isolation Years	No. Regions
<i>Lineage I</i>					
L1-SL87-ST87-CT58 (CC87, IIb)	77 (15.6%)	54 (13%)	23 (13 C, 10 P)	2014-19	17
L1-SL87-ST87-CT4373 (CC87, IIb)	53 (10.8%)	53 (13%)		2014-19	16
L1-SL5-ST5-CT4358 (CC5, IIb)	42 (8.5%)	42 (10%)		2014-19	9
L1-SL87-ST87-CT4374 (CC87, IIb)	28 (5.7%)	28 (7%)		2016-19	8
L1-SL1-ST1,ST1533-CT2384 (CC1, IVb)	11 (2.2%)	11 (3%)		2015, 2017-19	6
L1-SL5-ST5-CT4360 (CC5, IIb)	5 (1.0%)	5 (1%)		2016, 2018-19	3
L1-SL3-ST3-CT4369 (CC3, IIb)	4 (0.8%)	4 (1%)		2016,2018-19	3
L1-SL288-ST330-CT5502 (CC288, IIb)	3 (0.6%)	1 (0.2%)	2 (P)	2019	2
L1-SL87-ST87-CT9095 (CC87, IIb)	3 (0.6%)	3 (1%)		2018	3
L1-SL1-ST1-CT4353 (CC1, IVb)	2 (0.4%)	2 (0.5%)		2014	1
L1-SL288-ST330-CT5035 (CC288, IIb)	2 (0.4%)	2 (0.5%)		2018-19	1
L1-SL5-ST5-CT4359 (CC5, IIb)	2 (0.4%)	2 (0.5%)		2015, 2019	1
L1-SL5-ST5-CT5033 (CC5, IIb)	2 (0.4%)	2 (0.5%)		2018	2
L1-SL5-ST5-CT5501 (CC5, IIb)	2 (0.4%)	2 (0.5%)		2019	1
L1-SL87-ST87-CT4372 (CC87, IIb)	2 (0.4%)		2 (C)	2016	1
L1-SL87-ST87-CT4376 (CC87, IIb)	2 (0.4%)	2 (0.5%)		2015-19	2
<i>Lineage II</i>					
L2-SL378-ST378-CT4349 (CC19, IIa)	77 (15.6%)	57 (14%)	20 (7 C, 13 P)	2015-19	12
L2-SL101-ST101-CT4370 (CC101, IIa)	18 (3.7%)	18 (4%)		2016-19	7
L2-SL155-ST155-CT4362 (CC155, IIa)	10 (2.0%)	10 (2%)		2016-19	7
L2-SL9-ST9-CT4379 (CC9, IIc)	11 (2.2%)	1 (0.2%)	10 (1 C, 9 P)	2016, 2019	4
L2-SL177-ST177-CT5041 (CC177, IIa)	8 (1.6%)	8 (2%)		2018-19	6
L2-SL155-ST155-CT4363 (CC155, IIa)	7 (1.4%)	7 (2%)		2016-19	4
L2-SL155-ST155-CT4364 (CC155, IIa)	6 (1.2%)	2 (0.5%)	4 (P)	2014-15, 2019	4
L2-SL1081-ST1081-CT4378 (ST1081, IIa)	5 (1.0%)	5 (1%)		2016, 2018-19	3
L2-SL155-ST155-CT5049 (CC155, IIa)	5 (1.0%)	5 (1%)		2018-19	4
L2-SL8-ST551-CT4347 (CC8, IIa)	5 (1.0%)	1 (0.2%)	4 (C)	2016, 2019	2
L2-SL155-ST155-CT5050 (CC155, IIa)	4 (0.8%)	3 (1%)	1 (P)	2018-19	4
L2-SL155-ST1524-CT9039 (CC155, IIa)	3 (0.6%)	3 (1%)		2018-19	3
L2-SL155-ST155-CT4365 (CC155, IIa)	2 (0.4%)	2 (0.5%)		2015-16	2
L2-SL155-ST155-CT5496 (CC155, IIa)	2 (0.4%)		2	2019	2
L2-SL155-ST155-CT5539 (CC155, IIa)	2 (0.4%)		2	2019	1
L2-SL26-ST26-CT5754 (CC26, IIa)	2 (0.4%)	2 (0.5%)		2019	2
L2-SL9-ST9-CT5533 (CC9, IIc)	2 (0.4%)		2	2019	1

469

470 *Food type: C, chicken; P, pork.

Figure legends


Figure. 1. Incidence of listeriosis in Taiwan. A total of non-redundant 411 clinical isolates between January 2014 and December 2019 were collected. (A) Total no. of listeriosis cases and incidence per million population before (2014-2017) and after (2018-2019) mandatory notification in Taiwan. (B) Total no. of listeriosis cases in the different administrative regions of Taiwan before (2014-2017) and after (2018-2019) mandatory. The regions in white color indicates no reported case in the period.

Figure. 2. cgMLST analyses of the 493 *L. monocytogenes* isolates collected between January 2014 and December 2019. (A) Minimum spanning tree of isolates' core genome MLST allelic profiles (based on 1748 loci). cgMLST profiles are represented by circles proportional to the number of isolates and the number of allelic distances between profiles are indicated in the connecting lines. The dashed bar delimits *L. monocytogenes* phylogenetic lineages, dashed circles delimit sublineages (SL; i.e. isolates sharing 150 or fewer allelic differences) and grey shadows denote clusters of isolates within the same cgMLST type (CT; i.e. isolates sharing 7 or fewer allelic differences). CT names are shown for clusters with 2 or more isolates. CTs comprising 1 or more human isolates are shown in colors. Non-clinical isolates are labeled as E (environmental), P (pork) and C (chicken). (B) Prevalence of cgMLST types with 2 or more isolates according to phylogenetic lineage and isolation source.

Figure. 3. Genetic diversity of the 493 *L. monocytogenes* isolates. A) Single linkage clustering based on the cgMLST profiles. Branches are colored by phylogenetic lineage (L1, red; L2, blue). Sublineages (SL) with more than 5 isolates are labelled. Information on the serogroup, year of isolation and source are provided, using the color code indicated in the legend. The presence of virulence, acquired resistance traits, prophages and plasmids are shown as blue color-filled boxes.

FIGURE 1

A



B




FIGURE 2





FIGURE 3



SUPPLEMENTAL MATERIAL

Genomic surveillance of *Listeria monocytogenes* in Taiwan, 2014-2019

Yu-Huan Tsai^{1,*}, Alexandra Moura^{2,3,*}, Zi-Qi Gu¹, Jui-Hsien Chang⁴, Ying-Shu Liao⁴, Ru-Hsiou Teng⁴, Kuo-Yao Tseng¹, Dai-Ling Chang¹, Wei-Ren Liu¹, Yu-Tsung Huang^{5,6}, Alexandre Leclercq^{2,3}, Hsiu-Jung Lo^{7,8,9}, Marc Lecuit^{2,3,10,#}, Chien-Shun Chiou^{4,#}

¹ Laboratory of Host-Microbe Interactions and Cell Dynamics, Institute of Microbiology and Immunology, National Yang Ming Chiao Tung University, Taipei, Taiwan

² Institut Pasteur, Université Paris Cité, Inserm U1117, Biology of Infection Unit, Paris, France

³ Institut Pasteur, National Reference Centre and WHO Collaborating Centre *Listeria*, Paris, France

⁴ Center for Diagnostics and Vaccine Development, Centers for Disease Control, Taichung, Taiwan

⁵ Departments of Laboratory Medicine and Internal Medicine, National Taiwan University Hospital, National Taiwan University College of Medicine, Taipei, Taiwan

⁶ Graduate Institute of Clinical Laboratory Sciences and Medical Biotechnology, National Taiwan University, Taipei, Taiwan

⁷ National Institute of Infectious Disease and Vaccinology, National Health Research Institutes, Miaoli, Taiwan

⁸ School of Dentistry, China Medical University, Taichung, Taiwan

⁹ Department of Biological Science and Technology, National Yang Ming Chiao Tung University, Hsinchu, Taiwan

¹⁰ Necker-Enfants Malades University Hospital, Division of Infectious Diseases and Tropical Medicine, Institut Imagine, APHP, Paris, France

Table S1. Number of *Listeria monocytogenes*-positive food samples collected per region and food type in this study.

Product	Taipei	New Tapei	Taichung	Hsinchu	Total
Raw ground pork	18/27 (67%)	19/28 (68%)	32/62 (52%)	-	69/117 (59%)
Raw chicken	5/23 (22%)	8/27 (30%)	12/63 (19%)	-	25/113 (22%)
Mixed salads	-	-	-	0/50 (0%)	0/50 (0%)
Total	23/50 (46%)	27/55 (49%)	44/125 (35%)	0/50 (0%)	94/290 (32%)

Table S2. Characteristics of *Listeria monocytogenes* isolates retrieved in this study (N=493).

[extended excel file]

Table S3. Previously reported cgMLST types detected in this study ($n=10$).

cgMLST type (clonal complex, serogroup)	source(s) (this study)	source(s) (other studies)	countries (other studies)	source lab	Min. cgMLST similarity (%)
<i>Lineage 1</i>					
L1-SL1-ST1-CT2384 (CC1, IVb)	H	H,F,PE	US	FDA,CDC	99.6%
L1-SL224-ST845-CT21 (CC224, IIb)	H		MX,US	FDA,SENEICA	99.9%
L1-SL87-ST87-CT58 (CC87, IIb)	H,F	H,F,PE	AU,CA,CN,CO,DE,FR,SK,US	CDC,CFIA,DI,FDA,IP,RKI	99.6%
<i>Lineage 2</i>					
L2-SL101-ST101-CT4370 (CC101, IIa)	H	n.a.	n.a.	n.a.	99.7%
L2-SL155-ST155-CT4364 (CC155, IIa)	H,F	n.a.	n.a.	n.a.	99.6%
L2-SL16-ST16-CT5516 (CC8, IIa)	F	F	CA	CFIA	99.7%
L2-SL177-ST177-CT5041 (CC177, IIa)	H	H	KH	CDC-TW	99.8%
L2-SL378-ST378-CT4349 (CC19, IIa)	H,F	H	KH	CDC-TW	99.6%
L2-SL8-ST8-CT5167 (CC8, IIa)	H	n.a.	n.a.	n.a.	99.9%
L2-SL9-ST9-CT5537 (CC9,IIa, IIc)	F	H,F	FR	IP	99.8%

Source abbreviations: H, human; F, food; PE, food production environment.

Country abbreviations: AU, Australia, CA, Canada; CN, China; CO, Colombia; FR, France; KH, Cambodia; MX, Mexico; SK, South Korea; US, United States of America; n.a., not available.

Source lab abbreviations: CDC, Centers for Disease Control, United States; CDC-TW, Centers for Disease Control Taiwan; CFIA, Canadian Food Inspection Agency, Canada; DI, Doherty Institute, Australia; FDA, Food and Drug Administration, United States; IP, Institut Pasteur, France; RKI, Robert Koch Institut, Germany; SENESICA, Servicio Nacional de Sanidad Inocuidad y Calidad Agroalimentaria, Mexico; n.a., not available.




Figure S1. Distribution of *L. monocytogenes* major cgMLST sublineages (SLs) and types (CTs) in clinical and food sources in Taiwan. (A) No. isolates per SL and clinical form (2018-2019). (B) Percentage of larger SLs in clinical and food sources (2014-2019). (C) Percentage of larger SLs per food type (2014-2019). (D) Prevalence of cgMLST types in maternal-neonatal and CNS infections, for which the clinical information was available ($n=327$ isolates). (E) Distribution of major CTs of clinical isolates in different administrative regions of Taiwan. Abbreviations: CNS, central nervous system infection; MN, maternal-neonatal infections.