



**HAL**  
open science

## Alternative Transcription Start Site Usage and Functional Implications in Pathogenic Fungi

Thi Tuong Vi Dang, Jessie Colin, Guilhem Janbon

► **To cite this version:**

Thi Tuong Vi Dang, Jessie Colin, Guilhem Janbon. Alternative Transcription Start Site Usage and Functional Implications in Pathogenic Fungi. *Journal of Fungi*, 2022, 8 (10), pp.1044. 10.3390/jof8101044 . pasteur-03940436

**HAL Id: pasteur-03940436**

**<https://pasteur.hal.science/pasteur-03940436>**

Submitted on 16 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Alternative transcription start site usage and functional implications in pathogenic fungi

Thi Tuong Vi Dang <sup>1</sup>, Jessie Colin <sup>1,2</sup> and Guilhem Janbon <sup>1\*</sup>

<sup>1</sup> Université de Paris Cité, Institut Pasteur, Unité Biologie des ARN des Pathogènes Fongiques, Département de Mycologie, F-75015, Paris, France

<sup>2</sup> Ecole Pratique des Hautes Etudes, PSL Research university, F-75014, Paris, France

\* Correspondence: [guilhem.janbon@pasteur.fr](mailto:guilhem.janbon@pasteur.fr)

**Abstract:** Pathogenic fungi require delicate gene regulation mechanisms to adapt to diverse living environments and escape host immune systems. Recent advances in sequencing technology have exposed the complexity of the fungal genome, thus allowing the gradual disentanglement of multiple layers of gene expression control. Alternative transcription start site (aTSS) usage, previously reported to be prominent in mammals and to play important roles in physiopathology, is also present in fungi to fine-tune gene expression. Depending on the alteration in their sequences, RNA isoforms arising from aTSSs acquire different characteristics that significantly alter their stability and translational capacity as well as the properties and biologic functions of the resulting proteins. Disrupted control of aTSS usage has been reported to severely impair growth, virulence, and the infectious capacity of pathogenic fungi. Here, we discuss principle concepts, mechanisms, and the functional implication of aTSS usage in fungi.

**Keywords:** Alternative transcription start sites; fungi

## 1. Introduction

RNA molecules are synthesized in the cell by DNA- or RNA-dependent RNA polymerases and mature through various processes, including splicing, capping, polyadenylation, methylation, editing, and endonuclease and/or exonuclease digestion [1]. The nucleotide sequence of RNA molecules and the region surrounding the corresponding gene locus play critical roles in the control of these processes and significantly influence the transcriptome structure. Typically, eukaryotic RNA polymerase promoters consist of a core promoter and associated regulatory regions [2]. The core promoter can be defined as a genomic region harboring specific sequences that allow the recruitment and assembly of the pre-initiation complex (PIC) and that prime transcription at a basal level or upon stimulating signals [3,4]. Regulatory inputs from cis-acting elements are transduced at the core promoter to regulate RNA synthesis. After the formation of the PIC, the double-stranded DNA is melted to create the “transcription bubble” at the core promoter, allowing 5'-to-3' scanning of RNA polymerase II and transcription initiation at the transcription start site (TSS) [4]. One conserved characteristic is that transcription of a given RNA generally does not start at a unique nucleotide position. Rather, TSSs appear to form clusters in which TSS positions are located near one another. This represents the major TSS (mTSS) position, which is the most frequently used location for the initiation of RNA synthesis [4]. Each TSS cluster has a corresponding core promoter that spans the cluster and primes transcription initiation [4]. While core promoters are characterized by elements such as the “TATA box”, initiator element (Inr), and/or TFIIB recognition element (BRE), gene-specific regulatory elements are located upstream of the core promoter and control transcription [2–5].

The general transcription machinery is highly conserved across eukaryotes. Accordingly, fungi share similarities with mammals, including cofactors, cis-elements that con-

trol RNA pol II initiation, and other parts of the transcription process [2,3,6–8]. In *Saccharomyces cerevisiae*, the TATA box is typically located at positions from 40 to 120 bp upstream of the TSS [3,9]. This distance between the TATA box and the TSS seems to be organism-dependent. For example, this distance is shorter in *Schizosaccharomyces pombe* (40 to 70 bp) [10–13]. Core promoters of genes in many pathogenic fungi, such as *Ustilago maydis* *crg1* and *mig2-1* to *mig2-6*, *Nectria haematococca* *PDA1*, and *Coprinopsis cinerea* *clp1*, contain a TATA box located around 40 to 46 bp upstream of TSS [14–16]. Likewise, BRE is found in the *Magnaporthe grisea* *MPG1* gene, while analysis of 19 oomycete *Phytophthora infestans* genes revealed an Inr sequence that is highly similar to that of mammals [14,17,18]. In human fungal pathogens, a genome-wide analysis in *Candida glabrata* showed that the TATA box exists within the region from 200 to 1 bp upstream of the TSS [19]. In *Cryptococcus neoformans* and *Cryptococcus deneoformans*, the TATA box, when present, can be found 35–40 bp upstream of the mTSS (Dang and colleagues, manuscript in preparation).

Recently, the analysis of the distance between the TATA box and TSS in 12 yeast species allowed the definition of two classes of transcription initiation mechanisms [20]. The first mechanism, referred to as the “classic model” because it seems to be more widespread among eukaryotes, involves direct recognition of the TSS from factors recruited to the PIC. The other model, referred to as the scanning model, is based on the recruitment of RNAPII to the PIC, the opening of the transcription bubble, and scanning by RNAPII from the PIC downstream region towards the 3’ end to find an acceptable TSS. *Candida albicans* is more likely to use the scanning model, while *Cryptococcus* utilizes the classic model of transcription initiation.

The historical model of gene expression is that a gene is transcribed into a single mRNA that eventually serves as a template for the synthesis of a functional protein. The Beadle and Tatum “one gene, one enzyme” postulate imprinted the minds of scientists for decades before diverse transcriptional and post-transcriptional diversity was described. The textbook definition of a gene is often given as a transcription unit with one core promoter; thus, a gene would have only one TSS. However, recent advances in sequencing technology have allowed a closer look at the diversity of RNA molecules, revealing the spectacular plasticity of the eukaryotic transcriptome [21–24]. In mammals, usage of alternative splicing (aSpl), alternative TSSs (aTSSs), and alternative Poly(A) sites (aPAS) is prominent [25–29]. These processes are developmentally regulated [30], and defects in these processes are associated with a number of genetic diseases, such as thalassemia, retinitis pigmentosa, heart failure, and various types of cancer [31–38]. Interestingly, quantitative analysis of TSS usage suggested that aTSS and aPAS usage represents a major source of transcript isoform diversity in human tissues [39]. Thus, aTSS usage can have consequences on RNA stability, localization, and coding potential. aSpl, aTSS, and aPAS appear to be common in fungi [40]. However, these processes mostly result in unproductive transcripts, which have a limited impact on proteome diversity and thus the processes function more as a way to regulate gene expression [40–44]. Due to technical difficulties associated with their study, knowledge on aTSS was, up to recently, restricted to the model yeasts *S. cerevisiae* and *S. pombe*. Although less is known in pathogenic fungi, existing data suggest that these mechanisms might contribute into fine-tuning gene expression during infection [45–47]. Here, we review recent literature on aTSS usage and regulation and the consequences on fungal biology. We focus on fungal pathogens and discuss the potential and the mostly unexplored consequences of these regulations on virulence.

## 2. Identification of alternative transcription start sites

An aTSS is a TSS cluster that differs from the reference TSS most frequently used to initiate RNA synthesis under various environmental conditions. In the scanning model, the PIC may assemble at a single location but initiates transcription at different aTSS clusters [48,49]. For clarity, we will consider aTSS to have a similar meaning to alternative TSS

cluster, alternative promoter, or promoter switch, and the term is not related to the heterogeneity of TSSs within a TSS cluster as mentioned above. More than half of human genes have alternative promoters, and a human gene has an average of 4 TSSs [50]. This implies that a single human gene can harbor multiple core promoters that drive the transcription of RNA polymerase at distinct TSSs. Recently developed techniques, such as CAGE, TIF-seq, TL-seq, TSS-seq, and STRIPE-seq, provide insight into the TSS position at a single-nucleotide resolution and have confirmed the model of multiple TSS genes in eukaryotes [51–55]. Kimura and colleagues reported that at least 7674 genes, accounting for 52% of total human annotated genes, were regulated by putative alternative promoters [56]. In *Drosophila*, data from 5' cap read sequencing revealed 34,664 discrete TSS clusters associated with 8577 genes, suggesting comparable complexity of aTSS usage in the fruit fly compared to humans [57]. The situation appears to be similar in fungi. For instance, 1773 *S. pombe* genes have at least 2 core promoters [58]. In *S. cerevisiae*, 56% of the genes have at least 2 TSS clusters, and alternative core promoter usage by a gene is widespread in response to changing environments [59]. Nevertheless, information about aTSS usage in pathogenic fungi is limited due to both the scarcity of genome-wide studies and caveats in data integration. Results from 5'-end-cDNA sequencing of *C. glabrata* showed that out of 4316 coding genes, only 10% (n = 435) were transcribed from a single TSS, but it is unclear as to how many TSS clusters these TSSs belong [19]. A study in *Aspergillus nidulans* grouped 18,817,969 TSS positions into 17,992 putative TSS clusters, but further analysis of alternative usage of these clusters in various growth conditions has yet to be conducted [60]. Nonetheless, evidence of aTSS usage in pathogenic and non-pathogenic fungi has been published. In *Neurospora crassa*, the TSS positions of the circadian clock gene *frq* are distributed into two alternative major clusters depending on light exposure of the fungus [61]. In *C. neoformans*, aTSS usage results in the production of two alternative isoforms of *PUM1* regulated by sexual development growth phase [62]. Our recent TSS-seq data analysis in two *Cryptococcus* species revealed the existence of more TSS clusters than coding genes, suggesting widespread aTSS usage in these pathogenic yeasts [63]. While global studies will provide more understanding of the extent and dynamics of aTSS usage in various fungal species, the biological functions and mechanism of aTSS in pathogenic fungi is of particular interest.

### 3. Biological consequences of aTSS in fungi

During the 2000s, promoter switching events associated with mammalian development and disease etiology have been the subject of reviews that have cited results from hundreds of studies [64–67]. aTSS usage is expected to have a similar impact on the fungal transcriptome and proteome. Although aTSS likely affects lncRNAs in addition to mRNAs, we focus on mRNAs and discuss the consequences of aTSS usage on fungal mRNA level, mRNA diversity, and proteome diversity.

#### 3.1. Alternative transcript isoforms share the same ORF but differ in leader (5'UTR) sequence

aTSSs can be found upstream or downstream of the annotated TSS, which modulates the 5'UTR length. This modulation can result in the inclusion or exclusion of sequences containing important regulatory elements (Figure 1). For instance, the presence of one or more upstream open-reading frames (uORFs) within this region can dramatically alter mRNA stability and translation efficiency. In mammals, an uORF-linked regulatory mechanism has been illustrated in the mouse with the cyclin-dependent kinase inhibitor (p18INK4c) and peroxisome proliferator-activated receptor (PPAR) beta/delta [68,69]. In *S. cerevisiae*, 791 mRNAs contain uORFs in their transcript leader sequence [70], and 252 genes contain conserved uORFs [71]. Although the function of most uORFs has not been explored in detail, examples of uORF-dependent-regulation associated with alternative TSS usage exist. For instance, alternative promoter usage of the *NDC80* gene produces either a nonfunctional long transcript bearing several uORFs or a short translatable

mRNA, allowing versatile management of protein expression [72]. The *NDC80* gene encodes a subunit of the outer kinetochore, which plays a key role in meiosis. The controlled expression of two *NDC80* mRNA isoforms are responsible for the inactivation and reactivation of the kinetochore during cell division through Ndc80 protein level. The translatable short isoform is highly expressed in vegetative growth, but at the beginning of meiosis, 100% of cells express only the nonfunctional long isoform, which harbors 9 uORFs. Hence, no Ndc80 protein is produced at the meiotic prophase, thereby ensuring proper removal of the kinetochore at this stage. Mutating all 9 AUGs led to an increase in Ndc80 protein during prophase, confirming the inhibitory effect of uORFs on the translation of the main coding sequence. This abnormal expression of *NDC80* from the mutated long mRNA isoform led to abnormal chromosome segregation during cell division. In pathogenic fungi, uORFs in the 5'UTR contribute to the control of translation and RNA stability in response to environmental cues [73,74]. In *C. neoformans*, a robust, transient transcriptome modification is triggered upon exposure to reactive oxygen species [75], which is explained by translation inhibition through Gcn2-mediated eIF2 $\alpha$  phosphorylation [73]. This likely favors the efficient translation of the oxidative stress response gene *ERG110* and *GCN4*, which both contain uORFs in the transcript leader (TL) of their mRNA [73,76]. In addition, the production of an alternative transcript isoform containing an uORF has been reported for the *Metarhizium robertsii* Mr-OPY2 gene [43]. The plant- and fly-infecting fungus *M. robertsii* requires precise regulation of the Mr-OPY2 membrane anchor protein during saprophyte-to-insect pathogen transition. During saprophytic growth, Mr-OPY2 mRNA contains 2 uORFs that impede the translation of the main coding sequence. The elevated level of Mr-OPY2 protein during infection stages is achieved through the production of a short mRNA isoform that does not contain the 2 uORFs. Mutant strains that fail to express Mr-OPY2 proteins were impaired in appressorial formation and infection capacity. Artificial constitutive expression of the major open reading frame (ORF) during saprophytic growth results in the fluffy phenotype, aberrant conidiophores, and significantly reduced conidial production. These findings illustrate that aTSS- and uORF-associated regulatory strategies are exploited by both fungi and mammals as a flexible tool to regulate protein levels.

Genome-wide data analyses of TL structure performed in *Saccharomyces*, *Cryptococcus*, *Aspergillus*, *Candida*, and *Neurospora* species confirmed the repressive effect of uORFs on translation efficiency [53,60,63,77]. The translation efficiency of annotated ORFs is negatively correlated with the number of uORFs present in the 5'UTR. This is in agreement with the observation in mouse fibroblasts that uORF-containing longer mRNA isoforms are enriched in the monosome fraction, which is indicative of a lack of translation of the main ORF [78]. Analysis of Ribo-seq data revealed some examples of such regulation in *Cryptococcus*. At the CNAG\_06246 and CNAG\_03140 loci, only uORF translation was observed in the tested conditions, whereas the main ORF translation was completely abolished, suggesting tight regulation of gene expression [63].


Although the general view is that uORF translation would eventually trigger mRNA to degrade via nonsense-mediated mRNA decay (NMD), as suggested in *Cryptococcus* and *S. cerevisiae* [53,63], some uORF-based regulation appears to be more complex [79]. A classic example is the *S. cerevisiae* *GCN4* gene, which has 4 small uORFs in the 5' leader sequence of its mRNA [80]. In amino acid-rich media, ribosomes are dissociated after passing a GC-rich region at the end of the fourth uORF within *GCN4* mRNA. Therefore, ribosome re-initiation scanning is hindered after translation of the fourth uORF, and no protein is synthesized from the main ORF.

Given the small length of uORF-born peptides, one question is whether these short peptides possess any biological function. Examples of functional small peptides (<100 amino acids in length) are available in vertebrates and *Drosophila* [81]. In fungi, *N. crassa* arg-2 mRNA contains an evolutionarily conserved uORF coding a 24-amino acid peptide called arginine attenuator peptide (AAP), which participates in the control of ribosome movement [82]. Nevertheless, the features of repressive uORFs are evolutionarily maintained rather than their amino acid sequence [83]. This suggests that conserved uORFs are

mostly involved in regulation of translation of the downstream major ORF rather than encoding functional protein [84].

A longer 5' leader sequence of transcripts resulting from an upstream aTSS can potentially form a secondary structure that interferes with translation initiation [85]. Once assembled, the 43S complex scans the TL sequence to identify the correct translation start site [86]. Though the ribosome has the ability to process helicase activity, RNA secondary structures, such as the stem-loop hairpin structure, can block translation initiation via blockage of the scanning process [87,88]. For instance, under hypoxic conditions, *S. pombe* transcribes a translationally silent transcript from an upstream promoter at the *TCO1* locus [89]. The additional 751-nucleotide sequence of this alternative transcript is computationally predicted to form a stable stem-loop structure thought to block translation initiation by impairing 43S scanning.

Additional cis-regulatory sequences included within the 5' leader sequence upon aTSS usage can alter mRNA translation potential. An interesting example is the Pumilio protein family *PUM1* gene in *C. neoformans* [62]. Pum1 is an RNA-binding protein required for hyphal formation and is known to indirectly enhance the mRNA stability of the master filamentation regulator Znf2 [90]. In yeast cells, the Pum1 protein binds to its own transcript leader sequence, blocking ribosome scanning and shutting off mRNA translation [62]. In filaments, *C. neoformans* utilizes an alternative downstream TSS to produce a shorter isoform lacking the Pum1 binding site [62]. Thus, aTSS usage prevents auto-inhibition, thereby allowing full expression of *PUM1* and activation of *ZNF2*.




**Figure 1.** aTSS usage regulates gene expression without impacting protein diversity. aTSS usage includes or excludes regulatory elements within the transcript leader sequence, such as the uORF or secondary structure. This regulates the translation, stability, or subcellular targeting of the produced mRNA.

### 3.2. Alternative transcripts are translated into N-termini-truncated proteins

Although aTSS usage maintains an intact ORF in 60% to 80% of studied cases in mammals, it can also be a source of proteome diversity [65]. Thus, the usage of TSS clusters within coding sequences potentially results in novel protein isoforms. The translationally active downstream ATG (dATG) is typically found in-frame with the annotated ATG (aATG), so the alternative protein is a N-terminus-truncated version of the annotated one. This shorter polypeptide can maintain the same functional domains but lacks sequences


critical for protein localization (Figure 2). For instance, the plant *Arabidopsis thaliana* growing in shaded conditions uses aTSS to produce a cytosolic isoform of *GLYK*, a photorespiration enzyme believed to solely reside in the chloroplast [91]. Such aTSS regulation of protein localization by N-terminus truncation has been described for some *S. cerevisiae* tRNA synthetase genes. In these examples, instead of having a dedicated gene for cytosolic and mitochondrial tRNA-synthetase, a single gene generates both enzymes [92]. Both cytoplasmic and mitochondrial versions of histidinyI, valinyI, and cysteinyI tRNA synthetases are encoded by a single gene (*HST1*, *VAS1* and *CRS1*, respectively) [92–94]. At these loci, aTSSs generate either a long or short transcript isoform leading to alternative protein versions of Hts1, Vas1, and Crs1. The truncated versions lack the mitochondria targeting signal (MTS) at the N terminus and remain in the cytosol, while the full-length versions are translocated into the mitochondria. In the fungal pathogen *C. neoformans*, both tRNA synthetase activities are encoded by a single gene for most amino acids [63]. The expression of the 2 protein isoforms can be controlled by aTSS usage to produce a long and a short transcript, and the long isoform can produce both cytosolic and mitochondrial enzymes. Another layer of regulation is the selection of the translation initiation position, which is based on the consensus level of the associated Kozac context of the long transcript [63]. However, aTSS-dependent protein localization has been reported for other *C. neoformans* genes. For instance, *C. neoformans* synthesizes a long *UVE1* mRNA isoform which codes a protein, following UV exposure, specifically targeted to the mitochondria [95]. This long Uve1 protein is functionally active as a DNA damage repair endonuclease and protects the mitochondrial genome from potentially lethal UV-induced DNA damage. Similarly, *C. neoformans* employs aTSSs as a novel layer of regulation of superoxide dismutase activities. The genes *SOD1* and *SOD2* encode genes with cytoplasmic Cu-dependent and mitochondrial manganese-dependent superoxide dismutase activities, respectively [96]. Upon Cu shortage, an aTSS is used to regulate both gene products: a translationally repressed *SOD1* mRNA is produced while a shorter version of *SOD2* mRNA produces a functional cytosolic version of manganese-dependent superoxide dismutase Sod2. Thus, during infection, a condition in which copper is limited in host cells, the *Cryptococcus* manganese-dependent superoxide dismutase typically located in the mitochondria is translocated to the cytosol to maintain the cytosolic redox equilibrium against oxidative stress [96]. TSS switch-driven subcellular localization of proteins is not restricted to mitochondrial and cytosolic targeting. For instance, aTSS usage regulates the alternative production of the secreted, glycosylated version of the *S. cerevisiae* invertase *SUC2* in glucose-rich culture instead of its constitutive intracellular, non-glycosylated form [97].



**Figure 2.** Alternative transcripts leading to N-terminus-truncated proteins with different localization to that of the canonical protein. Condition 1: mRNA is transcribed from TSS1. This mRNA is translated into a protein with a localization target signal (mitochondria targeting signal [MTS] in this example) at its N terminus. The protein is targeted to the mitochondria. Condition 2: mRNA is transcribed from TSS2. The shorter transcript isoform is translated into an N-terminus-truncated protein lacking the MTS. The resulting alternative protein localizes to cytosol.

Alternative transcription initiation can be used to produce a shorter protein that loses important functional domains (apart from the localization signal) and in turn acts as an inhibitor of its full-length counterpart (Figure 3). An example in fungi is the *ZEB2* gene of the cereal-infecting fungus *Fusarium graminearum* in which a TSS not only impacts protein localization but also alters its function [98]. The full-length Zeb2 protein (Zeb2L) localizes exclusively to the nucleus. This long isoform has a basic leucine zipper (bZIP) DNA-binding domain and functions as a transcription factor that induces the production of the polyketide mycotoxin zearalenone (ZEA), a chemical compound leading to hyperestrogenic syndrome in infected cereals. ZEA accumulation triggers the synthesis of an N-terminally truncated protein (Zeb2S) that lacks the bZIP domain and exists in both the nucleus and cytoplasm. The short isoform forms heterodimers with Zeb2L and thus impedes the DNA-binding activity of Zeb2L in an autoregulatory process [98]. Similarly, in *S. pombe* wtf genes encode killer meiotic drivers which are selfish DNA sequence. Interestingly, the wtf drivers use alternative TSS to produce two protein isoforms [99]. Here, the long protein isoform Wtf4<sup>antidote</sup> neutralizes the short protein isoform Wtf4<sup>poison</sup> via heterodimerization. Moreover, the expression of both isoforms is tightly controlled in timing and localization to give the driver a transmission advantage into the next generation [99].






**Figure 3.** Alternative TSS usage leads to the production of an N-terminal-truncated protein with an alternative function. Condition 1: The mRNA is transcribed from the TSS1. The produced protein contains the functional domain (green). Condition 2: The mRNA is transcribed from the TSS2. The aTSS (TSS2) is located within the annotated coding sequence. The resulting protein does not contain the functional domain and might function as an inhibitor of the full-length protein, as in the case of the *Fusarium graminearum* protein Zeb2.

aTSS can result in the production of alternative protein isoforms encoded by alternative ORFs. To our knowledge, there are only 2 characterized examples in mammals: cyclin-dependent kinase inhibitor 2A gene (CDKN2A) and p21 in humans [100,101]. Each encodes 2 protein isoforms with different reading frames resulting from an alternative promoter coupled with different splicing patterns. Until recently, reports of similar aTSS-induced out-of-frame proteins have been unavailable in fungi. However, genome-wide proteomics approaches that sequence the N-terminal peptide have enabled studies on the impact on the proteome and the discovery of out-of-frame peptides [102–104]. These putative detected out-of-frame peptides are clearly not degradation products of annotated proteins that have a different reading frame. Rather, they likely result from bona fide proteins that are translated starting from an out-of-frame dATG or an in-frame dATG in the presence of alternative splicing. It is likely that such alternative proteins exist in pathogenic fungi, but their impact on the biology and virulence of these organisms remains to be studied.

### 3.3. Transcript isoforms are similar in coding sequence and translational efficiency

In some cases, aTSS usage does not impact protein output or translation efficiency (Figure 4). Possible explanations of multiple promoters for essential genes include to ensure expression level regardless of diverse initiation environments, such as the available transcription factor pool, or to diminish the fatal effects of mutations within the promoter [66]. For example, in the fungus *Aspergillus oryzae*, most glycolysis-reversible enzyme-coding genes have multiple TSSs even though they are constitutively expressed [105]. Carbon source-dependent aTSS usage is only observed in two genes: those encoding enolase (enoA) and fructose-bisphosphate aldolase (fbaA), where 2 promoters are alternatively used in response to nutritional signals. Thus, with the exceptions of enoA and fbaA, the existence of multiple TSSs in other glycolysis/gluconeogenesis genes probably functions

to maintain a constant level of protein in any environmental condition. On the other hand, *enoA* is an interesting case because the use of the two promoters depends on the carbon source. This gene is transcribed into two mRNAs that differ only in the 5'UTR; the short isoform is specific to glycolytic conditions, and the long transcript is highly expressed in gluconeogenic conditions. Replacing the 5' UTR of one *enoA* mRNA isoform with another 5' UTR of the reporter gene does not alter translational activity. However, mutations abolishing transcription from the upstream TSS prevent cell growth in acetate, while transcription from the downstream TSS is important for cell proliferation in glucose. Thus, *A. oryzae* requires transcription of *enoA* from distinct TSS/promoters for environmental adaptation, but no clear functional differences between the two mRNA isoforms can be observed. Here, *A. oryzae* might use aTSSs to adapt to the different availability of some transcription factors so as to satisfy the different demands of an important reversible enzyme in glycolysis and gluconeogenesis.



**Figure 4.** Alternative transcript isoforms code identical proteins and possess the same translation capacity. Condition 1: Canonical mRNA from TSS1 is produced. Condition 2: Alternative mRNA from TSS2 is transcribed. The alternative TSS results in transcript isoforms that differ in the 5' UTR but have the same coding sequence. This alternative isoform does not significantly impact stability or translational efficiency.

#### 4. Mechanism of alternative TSS usage control

As discussed in examples above, some transcripts produced from alternative TSSs are functional and critical to the cell. Fungi actively control their synthesis by employing both cis-elements and trans-factors to accurately produce the correct mRNA isoforms at the correct time.

##### 4.1. Transcription factors regulate alternative TSS usage

Transcription can be initiated from aTSS via binding of transcription factors (TFs) to its dedicated promoter, with evidence available both in mammals and yeasts. The promoter switch of the rodent gene *satb1* during T-cell development is controlled by the transcription factor TCF1 [85]. In *S. pombe*, transcription from the upstream aTSS of the *tco1+* gene in hypoxic conditions is triggered by the oxygen-sensitive TF Sre1 [89]. *S. cerevisiae* Gcn4 is a striking example of a TF that induces unconventional transcription: 546 Gcn4 ChIP-seq peaks were enriched during amino acid starvation vs control nutrient-rich me-

dia. Nearly 60% of Gcn4 genic binding sites are not located at the canonical promoter region, but rather are found inside coding sequences (CDSs) [106]. Many of these Gcn4 signals within CDSs are adjacent to induced TATA-binding protein peaks, suggesting the existence of cryptic internal promoters. Indeed, Gcn4 binds to its site within the ORF of *POS1*, *SNX41*, *SPO21*, and *COG1* and induces both the noncanonical antisense and sense transcriptions starting around 100 bp upstream or downstream. The H3 histone occupancy profile suggests that Gcn4 does not bind to the pre-existing nucleosome-depleted region. Rather, Gcn4 binding is more likely to stimulate gentle histone removal at the surrounding region, suggesting that Gcn4 actively provokes transcription initiation. This implies active regulatory activities of Gcn4 on aTSS usage.

TFs function not only as inducers but also as repressors of alternative promoters. Examples in mammals include the transcription factor Chx10, which negatively regulates the synthesis of the 2 transcript isoforms (H- and D-Mitf), but not 8 other isoforms of MIFT gene, in the retina during murine eye development [107]. Examples of TFs that negatively control the transcription of a particular alternative mRNA isoform have yet to be discovered in yeast. However, *S. cerevisiae* transcription factor Rap1 is reported to bind at the promoters of highly expressed genes and repress pervasive transcription from these promoters, possibly by contributing to nucleosome positioning [108,109].

TFs are reported to mediate feedback regulation on aTSS usage at their encoding gene. In humans, the pluripotent stem cell transcription factor NANOG binds to and auto-upregulates the usage of its own proximal promoter [110]. In *C. neoformans*, the RNA-binding protein Pum1 autoregulates the expression of its encoding gene by binding *ZNF2* mRNA, thus regulating the expression of this TF [62]. In turn, Znf2p binds to the *PUM1* proximal promoter, inducing the synthesis of a short isoform that lacks the sequence required for translational repression as in the long isoform [62].


Identification of the TFs regulating aTSS usage is suggested by the analysis of the sequence surrounding these TSSs and confirmed through genetic and/or ChIP-based analyses [85,111,112]. For instance, the binding site of 2 TFs in *A. oryzae* (AcuK and AcuM) are detected at the upstream promoter of the *enoA* gene, suggesting that these TFs could regulate the expression of the long transcript of the *enoA* gene in glucose starvation conditions. Accordingly, mutating the potential binding motif of AcuK and AcuM at the upstream promoter of the *enoA* gene reduces expression of the long transcript in acetate culture conditions but not in glucose-rich conditions [105].

Some TFs can affect aTSS usage of multiple genes, as observed in mammals and plants [91,113]. In *S. cerevisiae*, the TFs Ume6 and Ime1 form a heterodimer to activate transcription from alternative promoters upstream of meiotic genes specifically during meiotic prophase. This allows the downregulation of these meiotic specific genes through the combined act of translational and transcriptional repression. Thus, the production of 5' extended transcripts down-regulates the expression of the short isoforms through transcriptional interference of the downstream promoters. Whereas, these long mRNA isoforms are enriched in uORFs, leading to translational repression of the main ORF [114]. Accordingly, Ume6 binding sites are highly enriched at the promoter of these non-canonical transcripts. Similarly, Gcn4 likely functions as the master regulator controlling the use of a number of aTSSs at different loci in response to amino acid limitation [106]. In addition, Zap1 regulates aTSS usage at the *RTC4* and *RAD27* loci through zinc-responsive elements (ZREs) [115]. In *C. neoformans*, the TF Cuf1 controls aTSS usage at the *SOD1* and *SOD2* loci via Cu-responsive elements (CuREs) [96].

#### 4.2. *cis* regulatory transcription activity and chromatin conformation remodeling control alternative transcription

Either alternative promoters are independently activated by different pathways via distinct transcription factors, or their usages are intertwined with each other through transcriptional interference. In *S. cerevisiae*, transcription through a promoter can impede its activity [72,116] (Figure 5). For instance, transcription of the short functional *NDC80*

mRNA isoform is inhibited by the transcription event starting upstream that generates a 5' extended isoform [72]. This repression likely results from co-transcriptional chromatin remodeling [117,118]. During transcription from the upstream TSS of *NDC80*, the histone methyl transferases Set1 and Set2 regulate H3 lysine 4 dimethylation (H3K4me2) and H3 lysine 36 trimethylation (H3K36me3) levels, respectively [119]. Global mapping of epigenetic imprinting showed that H3K36me3 and, to a lesser extent, H3K4me2 are strong predictors of transcription repression mediated by upstream transcription [114]. The repressive histone marks H3K4me2 and H3K36me3 are read by histone deacetylases Set3 and Rpd3S, respectively, to promote histone deacetylation at these regions [119]. Histone deacetylation enhances the electrostatic interaction between histones and DNA, thus promoting nucleosome occupancy and inhibiting local transcription initiation [120]. Transcription activity from upstream TSSs is likely associated with low-level histone acetylation followed by a tighter wrapping of the histone core by the DNA string at the gene body. The converse can be true, where limited upstream transcription activity brings about permissive histone marks at the gene body, thereby allowing downstream transcription. For instance, low transcriptional activity of *S. cerevisiae* lncRNA *IRT2* induces the acetylation of histone H3 lysine 56 (H3K56ac). The associated relaxed chromatin conformation allows binding of the TF Rme1 at the promoter of lncRNA *IRT1*, which is located downstream and facilitates its transcription [121]. However, not every gene expressing 5' extended transcripts is associated with a reduction in the abundance of the short transcripts [114]. Further, the effect of expressing distal promoters on proximal ones can range from repression to activation [122]. Interestingly, this effect appears to be dependent on the distance between the distal and proximal TSS. Enhanced transcription from the upstream TSS can repress transcription from the downstream TSS if located more than 80 bp from one another, while a smaller distance is linked to positive correlation of their usage [122]. It is possible that a downstream TSS distantly separated from the upstream TSS is likely to have its own promoter, so its activity is regulated by transcriptional interference. Nevertheless, this cis regulation through transcriptional interference of a TSS usage remains to be described in pathogenic fungi.



**Figure 5.** Regulatory action in cis and chromatin conformation control alternative transcription. Condition 1: There is no transcription from TSS1. Acetylated histones (orange circle) at the gene body downstream TSS1 are associated with a relaxed chromatin state. The transcription factor Y, which activates transcription at the TSS2, can bind. Condition 2: Transcription from TSS1 leads to histone deacetylation. This reduction in histone acetylation leads to a more condensed chromatin conformation that occludes the binding of transcription factor Y, thereby repressing transcription from TSS2.

In many situations discussed above, it is puzzling that the cell uses so many resources on synthesizing a translationally deficient mRNA. A possible explanation for this “unnecessary” energy consumption resides on the regulatory cis action of alternative transcription as discussed above. In *S. cerevisiae*, aTSS usage driven by alternative promoters has cis regulatory activities, mostly via transcriptional activation. However, even if the RNA species produced from these aTSS are not coding/translated, we cannot exclude that at least some may have additional trans regulatory activities. In metazoans, several examples exist of trans-acting lncRNA acting on chromatin through binding chromatin modifiers or remodelers [123]. Thus, it is possible that this regulatory activity could occur in fungi as well.

Though not yet reported in yeast and fungi, several lines of evidence suggest the involvement of DNA methylation in the regulation of aTSS usage in mammals. Cap-targeted and bisulfite genomic sequencing data reveal variation in methylation patterns among alternative promoters of tissue-biased genes [124]. In human and viruses, the activity of alternative promoters is negatively correlated with hypermethylation of its CpG islands [124–128]. In conditions in which demethylation of aTSS is observed, the active promoter-associated histone modification H3K4me3 is detected spanning the promoter, implying genuine transcriptional activities [124,125]. How DNA methylation and TFs interfere with each other, and impact promoter expression is still controversial. In the human gene *garp*, an *in vitro* DNA-protein binding assay demonstrated an inhibitory effect of methylated CpG flanking the alternative promoter on the binding capacity of the 2 positive transcriptional regulators NFAT and Foxp3 [124]. Thus, steric hindrance of the methyl groups associated with DNA methylation at alternative promoters can repress TF binding to the corresponding cis-elements [129]. Conversely, TF binding can also affect local DNA methylation patterns and induce transcription through passive demethylation of the bound region [129,130]. Although cytosine methylation has been identified in at least 16 yeast species [131], the impact of DNA methylation on transcription requires clarification in model yeasts and pathogenic fungi.

## 5. Conclusions and remaining challenges

aTSS usage is a major mechanism regulating gene expression and proteome diversity in eukaryotes. Use of alternative promoters results in alterations in mRNA that can significantly modify transcript stability and translational efficiency, as well as protein sequence localization and function. Thus, aTSS has a potential critical impact on cell growth, differentiation, and adaptation. This type of regulation of both the transcriptome and the proteome structure is still poorly studied in pathogenic fungi. However, data produced in fungal model organisms, such as *N. crassa*, *S. cerevisiae*, and *S. pombe*, suggest that regulation of gene expression via aTSS may be common in pathogenic fungi as well. Accordingly, genome-wide analysis in 2 species of *Cryptococcus* and in *C. albicans* identified several aTSS clusters associated with coding genes, although the regulation of their expression and the associated molecular mechanisms remain to be explored [20,63]. Fungal pathogens need to adapt to diverse conditions and efficiently produce virulence factors to escape the host immune system and potentially acquire drug resistance. This suggests sophisticated and precise gene regulation mechanisms, such as aTSS usage, to regulate their biology and virulence.

The analysis of aTSS usage in model organisms revealed a significant number of multiple-promoter genes showing no promoter switch in different conditions. In this case, it is possible that transcript isoforms are regulated under unknown conditions. They can also be co-regulated. Cells uniformly express both alternative transcripts across conditions, but a population of cells could express a specific isoform while the alternative transcript would be expressed in the other part of the cell population. In the latter scenario, analysis of the whole population at once does not capture the true picture of aTSS usage [132]. Thus, single-cell TSS analysis would allow more accurate findings. However, this type of analysis is challenging in fungi. TSS likely plays major role in host-pathogen interactions given the heterogeneity of *in vivo* and *in vitro* fungal populations [133].

---

Overall, the handful of studies on aTSSs in pathogenic fungi suggest wide usage and should prompt exploration of the mechanisms regulating the expression of virulence factors, drug resistance, or in vivo fungal cell biology. Although a limited number of sequence datasets have been produced, no genome-wide analysis of the regulation of aTSS usage has been performed in any pathogenic fungi.

**Author Contributions:** TTV.D. contributed to the conception and literature survey of the study and to the first draft and revision of the manuscript. J.C. and G.J. contributed to the revision of the manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding:**

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** We thank Cecelia Shertz Wall for editing the manuscript. Thi Tuong Vi Dang was supported by the Pasteur - Paris University (PPU) International PhD Program

**Conflicts of Interest:** The authors declare no conflict of interest

## References

1. Brown, T.A. Synthesis and Processing of RNA. **2002**. 522  
523
2. Haberle, V.; Stark, A. Eukaryotic Core Promoters and the Functional Basis of Transcription Initiation. *Nature Reviews Molecular Cell Biology* **2018**, *19*, 621–637, doi:10.1038/s41580-018-0028-8. 524  
525
3. Hampsey, M. Molecular Genetics of the RNA Polymerase II General Transcriptional Machinery. *Microbiology and Molecular Biology Reviews* **1998**, *62*, 465, doi:10.1128/MMBR.62.2.465-503.1998. 526  
527
4. Sandelin, A.; Carninci, P.; Lenhard, B.; Ponjavic, J.; Hayashizaki, Y.; Hume, D.A. Mammalian RNA Polymerase II Core Promoters: Insights from Genome-Wide Studies. *Nature Reviews Genetics* **2007**, *8*, 424–436, doi:10.1038/nrg2026. 528  
529  
530
5. Lenhard, B.; Sandelin, A.; Carninci, P. Metazoan Promoters: Emerging Characteristics and Insights into Transcriptional Regulation. *Nature Reviews Genetics* **2012**, *13*, 233–245, doi:10.1038/nrg3163. 531  
532
6. Rhee, H.S.; Pugh, B.F. Genome-Wide Structure and Organization of Eukaryotic Pre-Initiation Complexes. *Nature* **2012**, *483*, 295–301, doi:10.1038/nature10799. 533  
534
7. Thomas, M.C.; Chiang, C.M. The General Transcription Machinery and General Cofactors. *Critical reviews in biochemistry and molecular biology* **2006**, *41*, 105–178, doi:10.1080/10409230600648736. 535  
536
8. Petrenko, N.; Struhl, K. Comparison of Transcriptional Initiation by RNA Polymerase II across Eukaryotic Species. *eLife* **10**, e67964, doi:10.7554/eLife.67964. 537  
538
9. Murakami, K.; Mattei, P.J.; Davis, R.E.; Jin, H.; Kaplan, C.D.; Kornberg, R.D. Uncoupling Promoter Opening from Start-Site Scanning. *Molecular Cell* **2015**, *59*, 133–138, doi:10.1016/J.MOLCEL.2015.05.021. 539  
540  
541
10. Huisinga, K.L.; Pugh, B.F. A Genome-Wide Housekeeping Role for TFIID and a Highly Regulated Stress-Related Role for SAGA in *Saccharomyces Cerevisiae*. *Molecular Cell* **2004**, *13*, 573–585, doi:10.1016/S1097-2765(04)00087-5. 542  
543  
544
11. Basehoar, A.D.; Zanton, S.J.; Pugh, B.F. Identification and Distinct Regulation of Yeast TATA Box-Containing Genes. *Cell* **2004**, *116*, 699–709, doi:10.1016/S0092-8674(04)00205-3. 545  
546
12. Li, Y.; Flanagan, P.M.; Tschochner, H.; Kornberg, R.D. RNA Polymerase II Initiation Factor Interactions and Transcription Start Site Selection. *Science* **1994**, *263*, 805–807, doi:10.1126/science.8303296. 547  
548  
549
13. Yang, C.; Ponticelli, A.S. Evidence That RNA Polymerase II and Not TFIIB Is Responsible for the Difference in Transcription Initiation Patterns between *Saccharomyces Cerevisiae* and *Schizosaccharomyces Pombe*. *Nucleic Acids Research* **2012**, *40*, 6495–6507, doi:10.1093/nar/gks323. 550  
551  
552
14. Basse, C.W.; Farfsing, J.W. Promoters and Their Regulation in *Ustilago Maydis* and Other Phytopathogenic Fungi. *FEMS Microbiology Letters* **2006**, *254*, 208–216, doi:10.1111/j.1574-6968.2005.00046.x. 553  
554  
555
15. Ruan, Y.; Straney, D.C.; Ruan, Y.; Straney, D.C. *Identification of Elements in the PDAI Promoter of Nectria Haematococca Necessary for a High Level of Transcription in Vitro*; 1996; Vol. 250, pp. 29–38;. 556  
557
16. Inada, K.; Morimoto, Y.; Arima, T.; Murata, Y.; Kamada, T. The Clp1 Gene of the Mushroom *Coprinus Cinereus* Is Essential for A-Regulated Sexual Development. *Genetics* **2001**, *157*, 133–140. 558  
559
17. Soanes, D.M.; Kershaw, M.J.; Cooley, R.N.; Talbot, N.J. *Regulation of the MPG1 Hydrophobin Gene in the Rice Blast Fungus Magnaporthe Grisea*; 2002; Vol. 15, pp. 1253–1267;. 560  
561
18. McLeod, A.; Smart, C.D.; Fry, W.E. Core Promoter Structure in the Oomycete *Phytophthora Infestans*. *Eukaryotic Cell* **2004**, *3*, 91–99, doi:10.1128/EC.3.1.91-99.2004. 562  
563

- 
19. Aoyama, T.; Nakayama, H.; Ueno, K.; Inukai, T.; Tanabe, K.; Nagi, M.; Bard, M.; Chibana, H. Genome-Wide Survey of Transcriptional Initiation in the Pathogenic Fungus, *Candida Glabrata*. *Genes to Cells* **2014**, *19*, 478–503, doi:10.1111/gtc.12147. 564–566
  20. Lu, Z.; Lin, Z. The Origin and Evolution of a Distinct Mechanism of Transcription Initiation in Yeasts. *Genome Research* **2020**, *31*, gr.264325.120, doi:10.1101/GR.264325.120. 567–568
  21. Kodzius, R.; Kojima, M.; Nishiyori, H.; Nakamura, M.; Fukuda, S.; Tagami, M.; Sasaki, D.; Imamura, K.; Kai, C.; Harbers, M.; et al. Cage: Cap Analysis of Gene Expression. *Nature Methods* **2006**, *3*, 211, doi:10.1038/nmeth0306-211. 569–571
  22. Pelechano, V.; Wei, W.; Steinmetz, L.M. Extensive Transcriptional Heterogeneity Revealed by Isoform Profiling. *Nature* **2013**, *497*, 127–131, doi:10.1038/nature12121. 572–573
  23. Tian, B.; Hu, J.; Zhang, H.; Lutz, C.S. A Large-Scale Analysis of MRNA Polyadenylation of Human and Mouse Genes. *Nucleic Acids Research* **2005**, *33*, 201–212, doi:10.1093/nar/gki158. 574–575
  24. Hoque, M.; Ji, Z.; Zheng, D.; Luo, W.; Li, W.; You, B.; Park, J.Y.; Yehia, G.; Tian, B. Analysis of Alternative Cleavage and Polyadenylation by 3' Region Extraction and Deep Sequencing. *Nature Methods* **2013**, *10*, 133–139, doi:10.1038/nmeth.2288. 576–578
  25. Wang, E.T.; Sandberg, R.; Luo, S.; Khrebtkova, I.; Zhang, L.; Mayr, C.; Kingsmore, S.F.; Schroth, G.P.; Burge, C.B. Alternative Isoform Regulation in Human Tissue Transcriptomes. *Nature* **2008**, *456*, 470–476, doi:10.1038/nature07509. 579–581
  26. Nilsen, T.W.; Graveley, B.R. Expansion of the Eukaryotic Proteome by Alternative Splicing. *Nature* **2010**, *463*, 457–463, doi:10.1038/nature08909. 582–583
  27. Carninci, P.; Sandelin, A.; Lenhard, B.; Katayama, S.; Shimokawa, K.; Ponjavic, J.; Semple, C.A.M.; Taylor, M.S.; Engström, P.G.; Frith, M.C.; et al. Genome-Wide Analysis of Mammalian Promoter Architecture and Evolution. *Nature Genetics* **2006**, *38*, 626–635, doi:10.1038/ng1789. 584–586
  28. Lianoglou, S.; Garg, V.; Yang, J.L.; Leslie, C.S.; Mayr, C. Ubiquitously Transcribed Genes Use Alternative Polyadenylation to Achieve Tissue-Specific Expression. *Genes and Development* **2013**, *27*, 2380–2396, doi:10.1101/gad.229328.113. 587–589
  29. Ji, Z.; Lee, J.Y.; Pan, Z.; Jiang, B.; Tian, B. Progressive Lengthening of 3' Untranslated Regions of MRNAs by Alternative Polyadenylation during Mouse Embryonic Development. *Proceedings of the National Academy of Sciences of the United States of America* **2009**, *106*, 7028–7033, doi:10.1073/pnas.0900028106. 590–593
  30. Pal, S.; Gupta, R.; Kim, H.; Wickramasinghe, P.; Baubet, V.; Showe, L.C.; Dahmane, N.; Davuluri, R.V. Alternative Transcription Exceeds Alternative Splicing in Generating the Transcriptome Diversity of Cerebellar Development. *Genome Research* **2011**, *21*, 1260–1272, doi:10.1101/gr.120535.111. 594–596
  31. Ley, T.J.; Anagnou, N.P.; Pepe, G.; Nienhuis, A.W. RNA Processing Errors in Patients with Beta-Thalassemia. *Proceedings of the National Academy of Sciences of the United States of America* **1982**, *79*, 4775–4779, doi:10.1073/pnas.79.15.4775. 597–599
  32. Buskin, A.; Zhu, L.; Chichagova, V.; Basu, B.; Mozaffari-Jovin, S.; Dolan, D.; Droop, A.; Collin, J.; Bronstein, R.; Mehrotra, S.; et al. Disrupted Alternative Splicing for Genes Implicated in Splicing and Ciliogenesis Causes PRPF31 Retinitis Pigmentosa. *Nature Communications* **2018**, *9*, 1–19, doi:10.1038/s41467-018-06448-y. 600–603



33. Li, C.-Q.; Pang, B.; Kiziltepe, T.; Trudel, L.J.; Engelward, B.P.; Dedon, P.C.; Wogan, G.N. Threshold Effects of Nitric Oxide-Induced Toxicity and Cellular Responses in Wild-Type and P53-Null Human Lymphoblastoid Cells. *Chemical Research in Toxicology* **2006**, *19*, 399–406, doi:10.1021/tx050283e.
34. Agarwal, V.R.; Bulun, S.E.; Leitch, M.; Rohrich, R.; Simpson, E.R. Use of Alternative Promoters to Express the Aromatase Cytochrome P450 (CYP19) Gene in Breast Adipose Tissues of Cancer-Free and Breast Cancer Patients. *The Journal of Clinical Endocrinology & Metabolism* **1996**, *81*, 3843–3849, doi:10.1210/jcem.81.11.8923826.
35. Mayr, C.; Bartel, D.P. Widespread Shortening of 3'UTRs by Alternative Cleavage and Polyadenylation Activates Oncogenes in Cancer Cells. *Cell* **2009**, *138*, 673–684, doi:10.1016/j.cell.2009.06.016.
36. Tan, S.; Li, H.; Zhang, W.; Shao, Y.; Liu, Y.; Guan, H.; Wu, J.; Kang, Y.; Zhao, J.; Yu, Q.; et al. NUDT21 Negatively Regulates PSMB2 and CXXC5 by Alternative Polyadenylation and Contributes to Hepatocellular Carcinoma Suppression. *Oncogene* **2018**, *37*, 4887–4900, doi:10.1038/s41388-018-0280-6.
37. Masamha, C.P.; Xia, Z.; Yang, J.; Albrecht, T.R.; Li, M.; Shyu, A.B.; Li, W.; Wagner, E.J. CFIm25 Links Alternative Polyadenylation to Glioblastoma Tumour Suppression. *Nature* **2014**, *510*, 412–416, doi:10.1038/nature13261.
38. Gacita, A.M.; Dellefave-Castillo, L.; Page, P.G.T.; Barefield, D.Y.; Wasserstrom, J.A.; Puckelwartz, M.J.; Nobrega, M.A.; McNally, E.M. Altered Enhancer and Promoter Usage Leads to Differential Gene Expression in the Normal and Failed Human Heart. *Circulation: Heart Failure* **2020**, doi:10.1161/CIRCHEARTFAILURE.120.006926.
39. Reyes, A.; Huber, W. Alternative Start and Termination Sites of Transcription Drive Most Transcript Isoform Differences across Human Tissues. *Nucleic Acids Research* **2018**, *46*, 582–592, doi:10.1093/nar/gkx1165.
40. Grutzmann, K.; Szafranski, K.; Pohl, M.; Voigt, K.; Petzold, A.; Schuster, S. Fungal Alternative Splicing Is Associated with Multicellular Complexity and Virulence: A Genome-Wide Multi-Species Study. *DNA Research* **2014**, *21*, 27–39, doi:10.1093/dnares/dst038.
41. Hoppins, S.C.; Go, N.E.; Klein, A.; Schmitt, S.; Neupert, W.; Rapaport, D.; Nargang, F.E. Alternative Splicing Gives Rise to Different Isoforms of the *Neurospora Crassa* Tob55 Protein That Vary in Their Ability to Insert  $\beta$ -Barrel Proteins Into the Outer Mitochondrial Membrane. *Genetics* **2007**, *177*, 137–149, doi:10.1534/genetics.107.075051.
42. Muzafar, S.; Sharma, R.D.; Shah, A.H.; Gaur, N.A.; Dasgupta, U.; Chauhan, N.; Prasad, R. Identification of Genomewide Alternative Splicing Events in Sequential, Isogenic Clinical Isolates of *Candida albicans* Reveals a Novel Mechanism of Drug Resistance and Tolerance to Cellular Stresses. *mSphere* **2020**, *5*, doi:10.1128/msphere.00608-20.
43. Guo, N.; Qian, Y.; Zhang, Q.; Chen, X.; Zeng, G.; Zhang, X.; Mi, W.; Xu, C.; St Leger, R.J.; Fang, W. Alternative Transcription Start Site Selection in Mr-OPY2 Controls Lifestyle Transitions in the Fungus *Metarhizium robertsii*. *Nature Communications* **2017**, *8*, 1–13, doi:10.1038/s41467-017-01756-1.
44. Franceschetti, M.; Bueno, E.; Wilson, R.A.; Tucker, S.L.; Gómez-Mena, C.; Calder, G.; Sesma, A. Fungal Virulence and Development Is Regulated by Alternative Pre-mRNA 3'End Processing in *Magnaporthe oryzae*. *PLoS Pathogens* **2011**, *7*, e1002441, doi:10.1371/journal.ppat.1002441.

- 
45. Jin, L.; Li, G.; Yu, D.; Huang, W.; Cheng, C.; Liao, S.; Wu, Q.; Zhang, Y. Transcriptome Analysis Reveals the Complexity of Alternative Splicing Regulation in the Fungus *Verticillium dahliae*. *BMC Genomics* **2017**, *18*, 130, doi:10.1186/s12864-017-3507-y. 644-646
46. Sieber, P.; Voigt, K.; Kämmer, P.; Brunke, S.; Schuster, S.; Linde, J. Comparative Study on Alternative Splicing in Human Fungal Pathogens Suggests Its Involvement during Host Invasion. *Frontiers in Microbiology* **2018**, *9*, doi:10.3389/fmicb.2018.02313. 647-649
47. Ibrahim, H.M.M.; Kusch, S.; Didelon, M.; Raffaele, S. Genome-wide Alternative Splicing Profiling in the Fungal Plant Pathogen *Sclerotinia Sclerotiorum* during the Colonization of Diverse Host Families. *Molecular Plant Pathology* **2021**, *22*, 31–47, doi:10.1111/mpp.13006. 650-652
48. Thiebaut, M.; Colin, J.; Neil, H.; Jacquier, A.; Séraphin, B.; Lacroute, F.; Libri, D. Futile Cycle of Transcription Initiation and Termination Modulates the Response to Nucleotide Shortage in *S. Cerevisiae*. *Mol Cell* **2008**, *31*, 671–682, doi:10.1016/j.molcel.2008.08.010. 653-655
49. Kuehner, J.N.; Brow, D.A. Regulation of a Eukaryotic Gene by GTP-Dependent Start Site Selection and Transcription Attenuation. *Mol Cell* **2008**, *31*, 201–211, doi:10.1016/j.molcel.2008.05.018. 656-657
50. Xu, C.; Park, J.K.; Zhang, J. Evidence That Alternative Transcriptional Initiation Is Largely Nonadaptive. *PLoS Biology* **2019**, *17*, doi:10.1371/JOURNAL.PBIO.3000197. 658-659
51. Takahashi, H.; Lassmann, T.; Murata, M.; Carninci, P. 5' End-Centered Expression Profiling Using Cap-Analysis Gene Expression and next-Generation Sequencing. *Nature Protocols* **2012**, *7*, 542–561, doi:10.1038/nprot.2012.005. 660-662
52. Pelechano, V.; Wei, W.; Jakob, P.; Steinmetz, L.M. Genome-Wide Identification of Transcript Start and End Sites by Transcript Isoform Sequencing. *Nature Protocols* **2014**, *9*, 1740–1759, doi:10.1038/nprot.2014.121. 663-665
53. Arribere, J.A.; Gilbert, W.V. Roles for Transcript Leaders in Translation and mRNA Decay Revealed by Transcript Leader Sequencing. *Genome Res.* **2013**, *23*, 977–987, doi:10.1101/gr.150342.112. 666-667
54. Malabat, C.; Feuerbach, F.; Ma, L.; Saveanu, C.; Jacquier, A. Quality Control of Transcription Start Site Selection by Nonsense-Mediated-mRNA Decay. *eLife* **2015**, *2015*, 1–59, doi:10.7554/eLife.06722. 668-669
55. Policastro, R.A.; Raborn, R.T.; Brendel, V.P.; Zentner, G.E. Simple and Efficient Profiling of Transcription Initiation and Transcript Levels with STRIPE-Seq. *Genome Research.* **2020**, doi:10.1101/gr.261545.120. 670-672
56. Kimura, K.; Wakamatsu, A.; Suzuki, Y.; Ota, T.; Nishikawa, T.; Yamashita, R.; Yamamoto, J.I.; Sekine, M.; Tsuritani, K.; Wakaguri, H.; et al. Diversification of Transcriptional Modulation: Large-scale identification and characterization of putative alternative promoters of human genes. *Genome Research* **2006**, *16*, 55–65, doi:10.1101/gr.4039406. 673-676
57. Ni, T.; Corcoran, D.L.; Rach, E.A.; Song, S.; Spana, E.P.; Gao, Y.; Ohler, U.; Zhu, J. A Paired-end sequencing strategy to map the complex landscape of transcription initiation. *Nature Methods* **2010**, *7*, 521–527, doi:10.1038/nmeth.1464. 677-679
58. Li, H.; Hou, J.; Bai, L.; Hu, C.; Tong, P.; Kang, Y.; Zhao, X.; Shao, Z. Genome-wide analysis of core promoter structures in *Schizosaccharomyces pombe* with DeepCAGE. *RNA Biology* **2015**, *12*, 525–537, doi:10.1080/15476286.2015.1022704. 680-682
59. Lu, Z.; Lin, Z. Pervasive and dynamic transcription initiation in *Saccharomyces cerevisiae*. *Genome Research* **2019**, *29*, 1198–1210, doi:10.1101/gr.245456.118. 683-684

- 
60. Galagan, J.E.; Calvo, S.E.; Cuomo, C.; Ma, L.J.; Wortman, J.R.; Batzoglou, S.; Lee, S.I.; Baştürkmen, M.; Spevak, C.C.; Clutterbuck, J.; et al. Sequencing of *Aspergillus nidulans* and Comparative Analysis with *A. Fumigatus* and *A. Oryzae*. *Nature* **2005**, *438*, 1105–1115, doi:10.1038/nature04341. 685–687
61. Colot, H.V.; Loros, J.J.; Dunlap, J.C. Temperature-Modulated Alternative Splicing and Promoter Use in the Circadian Clock Gene Frequency. *Molecular Biology of the Cell* **2005**, *16*, 5563–5571, doi:10.1091/mbc.E05-08-0756. 688–690
62. Kaur, J.N.; Panepinto, J.C. Morphotype-Specific Effector Functions of *Cryptococcus neoformans* PUM1. *Scientific Reports* **2016**, *6*, doi:10.1038/srep23638. 691–692
63. Wallace, E.W.J.; Maufrais, C.; Sales-Lee, J.; Tuck, L.R.; De Oliveira, L.; Feuerbach, F.; Moyrand, F.; Natarajan, P.; Madhani, H.D.; Janbon, G. Quantitative Global Studies Reveal Differential Translational Control by Start Codon Context across the Fungal Kingdom. *Nucleic Acids Research* **2020**, *48*, 2312–2331, doi:10.1093/nar/gkaa060. 693–696
64. Davuluri, R.V.; Suzuki, Y.; Sugano, S.; Plass, C.; Huang, T.H.M. The Functional Consequences of Alternative Promoter Use in Mammalian Genomes. *Trends in Genetics* **2008**, *24*, 167–177, doi:10.1016/j.tig.2008.01.008. 697–699
65. Landry, J.R.; Mager, D.L.; Wilhelm, B.T. Complex Controls: The Role of Alternative Promoters in Mammalian Genomes. *Trends in Genetics* **2003**, *19*, 640–648, doi:10.1016/j.tig.2003.09.014. 700–701
66. Ayoubi, T. Alternative Promoter Usage. In *eLS*; John Wiley & Sons, Ltd, 2005. 702
67. Pankratova, E.V. Alternative Promoters in Expression of Genetic Information. *Molecular Biology* **2008**, *42*, 371–380, doi:10.1134/S0026893308030047. 703–704
68. Phelps, D.E.; Hsiao, K.-M.; Li, Y.; Hu, N.; Franklin, D.S.; Westphal, E.; Lee, E.Y.-H.P.; Xiong, Y. Coupled Transcriptional and Translational Control of Cyclin-Dependent Kinase Inhibitor P18INK4c Expression during Myogenesis. *Molecular and Cellular Biology* **1998**, *18*, 2334–2343, doi:10.1128/mcb.18.4.2334. 705–708
69. Larsen, L.K.; Amri, E.Z.; Mandrup, S.; Pacot, C.; Kristiansen, K. Genomic Organization of the Mouse Peroxisome Proliferator-Activated Receptor  $\beta/\delta$  Gene: Alternative Promoter Usage and Splicing Yield Transcripts Exhibiting Differential Translational Efficiency. *Biochemical Journal* **2002**, *366*, 767–775, doi:10.1042/BJ20011821. 709–712
70. Spealman, P.; Naik, A.W.; May, G.E.; Kuersten, S.; Freeberg, L.; Murphy, R.F.; McManus, J. Conserved Non-AUG UORFs Revealed by a Novel Regression Analysis of Ribosome Profiling Data. *Genome Research* **2018**, *28*, 214–222, doi:10.1101/GR.221507.117/-/DC1. 713–715
71. Cvijović, M.; Dalevi, D.; Bilsland, E.; Kemp, G.J.L.; Sunnerhagen, P. Identification of Putative Regulatory Upstream ORFs in the Yeast Genome Using Heuristics and Evolutionary Conservation. *BMC Bioinformatics* **2007**, *8*, 295, doi:10.1186/1471-2105-8-295. 716–718
72. Chen, J.; Tresenrider, A.; Chia, M.; McSwiggen, D.T.; Spedale, G.; Jorgensen, V.; Liao, H.; Van Werven, F.J.; Ünal, E. Kinetochore Inactivation by Expression of a Repressive mRNA. *eLife* **2017**, *6*, doi:10.7554/eLife.27417. 719–721
73. Leipheimer, J.; Bloom, A.L.M.; Campomizzi, C.S.; Salei, Y.; Panepinto, J.C. Translational Regulation Promotes Oxidative Stress Resistance in the Human Fungal Pathogen *Cryptococcus neoformans*. *mBio* **2019**, *10*, doi:10.1128/MBIO.02143-19. 722–724

- 
74. Sundaram, A.; Grant, C.M. A Single Inhibitory Upstream Open Reading Frame (UORF) Is Sufficient to Regulate *Candida Albicans* GCN4 Translation in Response to Amino Acid Starvation Conditions. *RNA* **2014**, *20*, 559–567, doi:10.1261/rna.042267.113.
75. Upadhyaya, R.; Campbell, L.T.; Donlin, M.J.; Aurora, R.; Lodge, J.K. Global Transcriptome Profile of *Cryptococcus Neoformans* during Exposure to Hydrogen Peroxide Induced Oxidative Stress. *PLoS One* **2013**, *8*, e55110, doi:10.1371/journal.pone.0055110.
76. Stovall, A.K.; Knowles, C.M.; Kalem, M.C.; Panepinto, J.C. A Conserved Gcn2-Gcn4 Axis Links Methionine Utilization and the Oxidative Stress Response in *Cryptococcus Neoformans*. *Frontiers in Fungal Biology* **2021**, *2*.
77. Zhang, H.; Wang, Y.; Wu, X.; Tang, X.; Wu, C.; Lu, J. Determinants of Genome-Wide Distribution and Evolution of UORFs in Eukaryotes. *Nature Communications* **2021**, *12*, 1–17, doi:10.1038/s41467-021-21394-y.
78. Wang, X.; Hou, J.; Quedenau, C.; Chen, W. Pervasive Isoform-specific Translational Regulation via Alternative Transcription Start Sites in Mammals. *Molecular Systems Biology* **2016**, *12*, 875, doi:10.15252/msb.20166941.
79. Vilela, C.; McCarthy, J.E.G. Regulation of Fungal Gene Expression via Short Open Reading Frames in the MRNA 5'untranslated Region. *Molecular Microbiology* **2003**, *49*, 859–867, doi:10.1046/j.1365-2958.2003.03622.x.
80. Abastado, J.P.; Miller, P.F.; Jackson, B.M.; Hinnebusch, A.G. Suppression of Ribosomal Reinitiation at Upstream Open Reading Frames in Amino Acid-Starved Cells Forms the Basis for GCN4 Translational Control. *Mol Cell Biol* **1991**, *11*, 486–496, doi:10.1128/mcb.11.1.486-496.1991.
81. Andjus, S.; Morillon, A.; Wery, M. From Yeast to Mammals, the Nonsense-Mediated MRNA Decay as a Master Regulator of Long Non-Coding RNAs Functional Trajectory. *ncRNA* **2021**, *7*, 44, doi:10.3390/ncrna7030044.
82. Wang, Z.; Fang, P.; Sachs, M.S. The Evolutionarily Conserved Eukaryotic Arginine Attenuator Peptide Regulates the Movement of Ribosomes That Have Translated It. *Molecular and Cellular Biology* **1998**, *18*, 7528, doi:10.1128/MCB.18.12.7528.
83. Johnstone, T.G.; Bazzini, A.A.; Giraldez, A.J. Upstream ORFs Are Prevalent Translational Repressors in Vertebrates. *EMBO J* **2016**, *35*, 706–723, doi:10.15252/embj.201592759.
84. Orr, M.W.; Mao, Y.; Storz, G.; Qian, S.-B. Alternative ORFs and Small ORFs: Shedding Light on the Dark Proteome. *Nucleic Acids Res* **2020**, *48*, 1029–1042, doi:10.1093/nar/gkz734.
85. Patta, I.; Madhok, A.; Khare, S.; Gottimukkala, K.P.; Verma, A.; Giri, S.; Dandewad, V.; Seshadri, V.; Lal, G.; Misra-Sen, J.; et al. Dynamic Regulation of Chromatin Organizer SATB1 via TCR-Induced Alternative Promoter Switch during T-Cell Development. *Nucleic acids research* **2020**, *48*, 5873–5890, doi:10.1093/nar/gkaa321.
86. Hinnebusch, A.G. Molecular Mechanism of Scanning and Start Codon Selection in Eukaryotes. *Microbiol Mol Biol Rev* **2011**, *75*, 434–467, doi:10.1128/MMBR.00008-11.
87. Takyar, S.; Hickerson, R.P.; Noller, H.F. MRNA Helicase Activity of the Ribosome. *Cell* **2005**, *120*, 49–58, doi:10.1016/j.cell.2004.11.042.
88. Leppék, K.; Das, R.; Barna, M. Functional 5' UTR MRNA Structures in Eukaryotic Translation Regulation and How to Find Them. *Nature Reviews Molecular Cell Biology* **2018**, *19*, 158–174, doi:10.1038/nrm.2017.103.

- 
89. Sehgal, A.; Hughes, B.T.; Espenshade, P.J. Oxygen-Dependent, Alternative Promoter Controls Translation of Tco1+ in Fission Yeast. *Nucleic Acids Research* **2008**, *36*, 2024–2031, doi:10.1093/nar/gkn027. 767  
768  
769
90. Wang, L.; Tian, X.; Gyawali, R.; Upadhyay, S.; Foyle, D.; Wang, G.; Cai, J.J.; Lin, X. Morphotype Transition and Sexual Reproduction Are Genetically Associated in a Ubiquitous Environmental Pathogen. *PLoS Pathog* **2014**, *10*, e1004185, doi:10.1371/journal.ppat.1004185. 770  
771  
772
91. Ushijima, T.; Hanada, K.; Gotoh, E.; Yamori, W.; Kodama, Y.; Tanaka, H.; Kusano, M.; Fukushima, A.; Tokizawa, M.; Yamamoto, Y.Y.; et al. Light Controls Protein Localization through Phytochrome-Mediated Alternative Promoter Selection. *Cell* **2017**, *171*, 1316–1325.e12, doi:10.1016/j.cell.2017.10.018. 773  
774  
775
92. Chatton, B.; Walter, P.; Ebel, J.P.; Lacroute, F.; Fasiolo, F. The Yeast VAS1 Gene Encodes Both Mitochondrial and Cytoplasmic Valyl-TRNA Synthetases. *Journal of Biological Chemistry* **1988**, *263*, 52–57, doi:10.1016/S0021-9258(19)57354-9. 776  
777  
778
93. Natsoulis, G.; Hilger, F.; Fink, G.R. The HTS1 Gene Encodes Both the Cytoplasmic and Mitochondrial Histidine TRNA Synthetases of *S. Cerevisiae*. *Cell* **1986**, *46*, 235–243, doi:10.1016/0092-8674(86)90740-3. 779  
780  
781
94. Nishimura, A.; Nasuno, R.; Yoshikawa, Y.; Jung, M.; Ida, T.; Matsunaga, T.; Morita, M.; Takagi, H.; Motohashi, H.; Akaike, T. Mitochondrial Cysteinyl-TRNA Synthetase Is Expressed via Alternative Transcriptional Initiation Regulated by Energy Metabolism in Yeast Cells. *Journal of Biological Chemistry* **2019**, *294*, 13781–13788, doi:10.1074/jbc.RA119.009203. 782  
783  
784  
785
95. Verma, S.; Idnurm, A. The Uve1 Endonuclease Is Regulated by the White Collar Complex to Protect *Cryptococcus Neoformans* from UV Damage. *PLoS Genetics* **2013**, *9*, e1003769, doi:10.1371/journal.pgen.1003769. 786  
787  
788
96. Smith, A.D.; Garcia-Santamarina, S.; Ralle, M.; Loiselle, D.R.; Haystead, T.A.; Thiele, D.J. Transcription Factor-Driven Alternative Localization of *Cryptococcus neoformans* Superoxide Dismutase. *Journal of Biological Chemistry* **2021**, *296*, doi:10.1016/j.jbc.2021.100391. 789  
790  
791
97. Carlson, M.; Botstein, D. Two Differentially Regulated MRNAs with Different 5' Ends Encode Secreted and Intracellular Forms of Yeast Invertase. *Cell* **1982**, *28*, 145–154, doi:10.1016/0092-8674(82)90384-1. 792  
793
98. Park, A.R.; Son, H.; Min, K.; Park, J.; Goo, J.H.; Rhee, S.; Chae, S.K.; Lee, Y.W. Autoregulation of ZEB2 Expression for Zearalenone Production in *Fusarium Graminearum*. *Molecular Microbiology* **2015**, *97*, 942–956, doi:10.1111/mmi.13078. 794  
795  
796
99. Nuckolls, N.L.; Srinivasa, A.N.; Mok, A.C.; Núñez, M.A.B.; Lange, J.J.; Gallagher, T.J.; Seidel, C.W.; Zanders, S.E. *S. Pombe* Wtf Genes Use Dual Transcriptional Regulation and Selective Protein Exclusion from Spores to Cause Meiotic Drive 2021, 2021.09.30.462505. 797  
798  
799
100. Ouelle, D.E.; Zindy, F.; Ashmun, R.A.; Sherr, C.J. Alternative Reading Frames of the INK4a Tumor Suppressor Gene Encode Two Unrelated Proteins Capable of Inducing Cell Cycle Arrest. *Cell* **1995**, *83*, 993–1000, doi:10.1016/0092-8674(95)90214-7. 800  
801  
802
101. Nozell, S.; Chen, X. P21B, a Variant of P21waf1/Cip1, Is Induced by the P53 Family. *Oncogene* **2002**, *21*, 1285–1294, doi:10.1038/sj.onc.1205191. 803  
804
102. Kleifeld, O.; Doucet, A.; Prudova, A.; Auf Dem Keller, U.; Gioia, M.; Kizhakkedathu, J.N.; Overall, C.M. Identifying and Quantifying Proteolytic Events and the Natural N Terminome by Terminal Amine Isotopic Labeling of Substrates. *Nature Protocols* **2011**, *6*, 1578–1611, doi:10.1038/nprot.2011.382. 805  
806  
807

- 
103. Fournier, C.T.; Cherny, J.J.; Truncali, K.; Robbins-Pianka, A.; Lin, M.S.; Krizanc, D.; Weir, M.P. Amino Termini of Many Yeast Proteins Map to Downstream Start Codons. *Journal of Proteome Research* **2012**, *11*, 5712–5719, doi:10.1021/pr300538f. 808  
809  
810
104. Wei, W.; Hennig, B.P.; Wang, J.; Zhang, Y.; Piazza, I.; Sanchez, Y.P.; Chabbert, C.D.; Adjalley, S.H.; Steinmetz, L.M.; Pelechano, V. Chromatin-Sensitive Cryptic Promoters Putatively Drive Expression of Alternative Protein Isoforms in Yeast. *Genome Research* **2019**, *29*, 1974–1984, doi:10.1101/gr.243378.118. 811  
812  
813  
814
105. Inoue, T.; Toji, H.; Tanaka, M.; Takama, M.; Hasegawa-Shiro, S.; Yamaki, Y.; Shintani, T.; Gomi, K. Alternative Transcription Start Sites of the Enolase-Encoding Gene EnoA Are Stringently Used in Glycolytic/Gluconeogenic Conditions in *Aspergillus Oryzae*. *Current Genetics* **2020**, *66*, 729–747, doi:10.1007/s00294-020-01053-3. 815  
816  
817  
818
106. Rawal, Y.; Chereji, R.V.; Valabhoju, V.; Qiu, H.; Ocampo, J.; Clark, D.J.; Hinnebusch, A.G. Gcn4 Binding in Coding Regions Can Activate Internal and Canonical 5' Promoters in Yeast. *Molecular Cell* **2018**, *70*, 297–311.e4, doi:10.1016/j.molcel.2018.03.007. 819  
820  
821
107. Bharti, K.; Liu, W.; Csermely, T.; Bertuzzi, S.; Arnheiter, H. Alternative Promoter Use in Eye Development: The Complex Role and Regulation of the Transcription Factor MITF. *Development* **2008**, *135*, 1169–1178, doi:10.1242/dev.014142. 822  
823  
824
108. Wu, A.C.K.; Patel, H.; Chia, M.; Moretto, F.; Frith, D.; Snijders, A.P.; van Werven, F.J. Repression of Divergent Noncoding Transcription by a Sequence-Specific Transcription Factor. *Molecular Cell* **2018**, *72*, 942–954.e7, doi:10.1016/j.molcel.2018.10.018. 825  
826  
827
109. Challal, D.; Barucco, M.; Kubik, S.; Feuerbach, F.; Candelli, T.; Geoffroy, H.; Benaksas, C.; Shore, D.; Libri, D. General Regulatory Factors Control the Fidelity of Transcription by Restricting Non-Coding and Ectopic Initiation. *Molecular Cell* **2018**, *72*, 955–969.e7, doi:10.1016/j.molcel.2018.11.037. 828  
829  
830
110. Das, S. Transcriptional Regulation of Human NANOG by Alternate Promoters in Embryonic Stem Cells. *Journal of Stem Cell Research & Therapy* **2012**, *01*, doi:10.4172/2157-7633.s10-009. 831  
832
111. Guske, K.; Schmitz, B.; Schelleckes, M.; Duning, K.; Kremerskothen, J.; Pavenstädt, H.J.; Brand, S.M.; Brand, E. Tissue-Specific Differences in the Regulation of KIBRA Gene Expression Involve Transcription Factor TCF7L2 and a Complex Alternative Promoter System. *Journal of Molecular Medicine* **2014**, *92*, 185–196, doi:10.1007/s00109-013-1089-y. 833  
834  
835  
836
112. Lutze, P.; Wanka, H.; Bäumgen, I.; Staar, D.; Grunow, B.; Peters, J. An Alternative Promoter in Intron1 of the Renin Gene Is Regulated by Glucose Starvation via Serum Response Factor. *Cellular Physiology and Biochemistry* **2017**, *42*, 1447–1457, doi:10.1159/000479209. 837  
838  
839
113. Feng, G.; Tong, M.; Xia, B.; Luo, G.; Wang, M.; Xie, D.; Wan, H.; Zhang, Y.; Zhou, Q.; Wang, X. Ubiquitously Expressed Genes Participate in Cell-specific Functions via Alternative Promoter Usage. *EMBO reports* **2016**, *17*, 1304–1313, doi:10.15252/embr.201541476. 840  
841  
842
114. Tresenrider, A.; Morse, K.; Jorgensen, V.; Chia, M.; Liao, H.; Jacobus van Werven, F.; Ünal, E. Integrated Genomic Analysis Reveals Key Features of Long Undecoded Transcript Isoform-Based Gene Repression. *Molecular Cell* **2021**, doi:10.1016/j.molcel.2021.03.013. 843  
844  
845
115. Taggart, J.; MacDiarmid, C.W.; Haws, S.; Eide, D.J. Zap1-Dependent Transcription from an Alternative Upstream Promoter Controls Translation of RTC4 mRNA in Zinc-Deficient *Saccharomyces Cerevisiae*. *Molecular Microbiology* **2017**, *106*, 678–689, doi:10.1111/mmi.13851. 846  
847  
848

- 
116. Moretto, F.; Wood, N.E.; Kelly, G.; Doncic, A.; Van Werven, F.J. A Regulatory Circuit of Two LncRNAs and a Master Regulator Directs Cell Fate in Yeast. *Nature Communications* **2018**, *9*, doi:10.1038/s41467-018-03213-z. 849–851
117. Hampsey, M.; Reinberg, D. Tails of Intrigue: Phosphorylation of RNA Polymerase II Mediates Histone Methylation. *Cell* **2003**, *113*, 429–432, doi:10.1016/S0092-8674(03)00360-X. 852–853
118. Ard, R.; Allshire, R.C. Transcription-Coupled Changes to Chromatin Underpin Gene Silencing by Transcriptional Interference. *Nucleic Acids Research* **2016**, *44*, 10619–10630, doi:10.1093/nar/gkw801. 854–855
119. Chia, M.; Tresenrider, A.; Chen, J.; Spedale, G.; Jorgensen, V.; Ünal, E.; van Werven, F.J. Transcription of a 5' Extended mRNA Isoform Directs Dynamic Chromatin Changes and Interference of a Downstream Promoter. *eLife* **2017**, *6*, doi:10.7554/eLife.27420. 856–858
120. De Ruijter, A.J.M.; Van Gennip, A.H.; Caron, H.N.; Kemp, S.; Van Kuilenburg, A.B.P. Histone Deacetylases (HDACs): Characterization of the Classical HDAC Family. *Biochemical Journal* **2003**, *370*, 737–749, doi:10.1042/BJ20021321. 859–861
121. Moretto, F.; Wood, N.E.; Chia, M.; Li, C.; Luscombe, N.M.; van Werven, F.J. Transcription Levels of a Noncoding RNA Orchestrate Opposing Regulatory and Cell Fate Outcomes in Yeast. *Cell Reports* **2021**, *34*, 108643, doi:10.1016/j.celrep.2020.108643. 862–864
122. Chia, M.; Li, C.; Marques, S.; Pelechano, V.; Luscombe, N.M.; van Werven, F.J. High-Resolution Analysis of Cell-State Transitions in Yeast Suggests Widespread Transcriptional Tuning by Alternative Starts. *Genome Biology* **2021**, *22*, 34, doi:10.1186/s13059-020-02245-3. 865–867
123. Statello, L.; Guo, C.-J.; Chen, L.-L.; Huarte, M. Gene Regulation by Long Non-Coding RNAs and Its Biological Functions. *Nat Rev Mol Cell Biol* **2021**, *22*, 96–118, doi:10.1038/s41580-020-00315-9. 868–869
124. Haupt, S.; Söntgerath, V.S.A.; Leipe, J.; Schulze-Koops, H.; Skapenko, A. Methylation of an Intragenic Alternative Promoter Regulates Transcription of GARP. *Biochimica et Biophysica Acta - Gene Regulatory Mechanisms* **2016**, *1859*, 223–234, doi:10.1016/j.bbagr.2015.11.003. 870–872
125. Wiesner, T.; Lee, W.; Obenauf, A.C.; Ran, L.; Murali, R.; Zhang, Q.F.; Wong, E.W.P.; Hu, W.; Scott, S.N.; Shah, R.H.; et al. Alternative Transcription Initiation Leads to Expression of a Novel ALK Isoform in Cancer. *Nature* **2015**, *526*, 453–457, doi:10.1038/nature15258. 873–875
126. Bujko, M.; Kober, P.; Rusetska, N.; Wakula, M.; Goryca, K.; Grecka, E.; Matyja, E.; Neska, J.; Mandat, T.; Bonicki, W.; et al. Aberrant DNA Methylation of Alternative Promoter of DLC1 Isoform 1 in Meningiomas. *Journal of Neuro-Oncology* **2016**, *130*, 473–484, doi:10.1007/s11060-016-2261-3. 876–878
127. Arenas, F.; Hervías, I.; Sáez, E.; Melero, S.; Prieto, J.; Parés, A.; Medina, J.F. Promoter Hypermethylation of the AE2/SLC4A2 Gene in PBC. *JHEP Reports* **2019**, *1*, 145–153, doi:10.1016/j.jhepr.2019.05.006. 879–881
128. Strassheim, S.; Gennart, I.; Muylkens, B.; André, M.; Rasschaert, D.; Laurent, S. Oncogenic Marek's Disease Herpesvirus Encodes an Isoform of the Conserved Regulatory Immediate Early Protein ICP27 Generated by Alternative Promoter Usage. *Journal of General Virology* **2016**, *97*, 2399–2410, doi:10.1099/jgv.0.000547. 882–885
129. Li, E.; Zhang, Y. DNA Methylation in Mammals. *Cold Spring Harbor Perspectives in Biology* **2014**, *6*, 19133–19134, doi:10.1101/cshperspect.a019133. 886–887
130. Hsieh, C.L. Dynamics of DNA Methylation Pattern. *Current Opinion in Genetics and Development* **2000**, *10*, 224–228, doi:10.1016/S0959-437X(00)00064-2. 888–889

- 
131. Tang, Y.; Gao, X.-D.; Wang, Y.; Yuan, B.-F.; Feng, Y.-Q. Widespread Existence of Cytosine Methylation in Yeast DNA Measured by Gas Chromatography/Mass Spectrometry. *Anal Chem* **2012**, *84*, 7249–7255, doi:10.1021/ac301727c. 890  
891  
892
132. Karlsson, K.; Lönnerberg, P.; Linnarsson, S. Alternative TSSs Are Co-regulated in Single Cells in the Mouse Brain. *Molecular Systems Biology* **2017**, *13*, 930, doi:10.15252/msb.20167374. 893  
894
133. Hogan, D.A.; Gladfelter, A.S. Editorial Overview: Heterogeneity in Fungal Cells, Populations, and Communities. *Current opinion in microbiology* **2015**, *26*, vii, doi:10.1016/J.MIB.2015.07.003. 895  
896  
897  
898  
899