



HAL
open science

New geographical records for tick-borne pathogens in ticks collected from cattle in Benin and Togo

Roland Eric Yessinou, Cristina Daniela Cazan, Luciana Cătălina Panait, Eyabana Mollong, Abel S Biguezoton, Sarah Irène Bonnet, Souaïbou Farougou, Martin H Groschup, Andrei Daniel Mihalca

► To cite this version:

Roland Eric Yessinou, Cristina Daniela Cazan, Luciana Cătălina Panait, Eyabana Mollong, Abel S Biguezoton, et al.. New geographical records for tick-borne pathogens in ticks collected from cattle in Benin and Togo. *Veterinary Medicine and Science*, 2023, 9 (1), pp.345 - 352. 10.1002/vms3.1022 . pasteur-04008651

HAL Id: pasteur-04008651

<https://pasteur.hal.science/pasteur-04008651v1>

Submitted on 28 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

New geographical records for tick-borne pathogens in ticks collected from cattle in Benin and Togo

Roland Eric Yessinou¹  | Cristina Daniela Cazan^{2,3} | Luciana Cătălina Panait² | Eyabana Mollong⁴ | Abel S. Biguezoton⁵ | Sarah Irène Bonnet^{6,7} | Souaïbou Farougou¹ | Martin H. Groschup⁸ | Andrei Daniel Mihalca²

¹Communicable Disease Research Unit (URMaT), University of Abomey-Calavi, Cotonou, Benin

²Department of Parasitology and Parasitic Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Cluj-Napoca, Romania

³CDS-9, Molecular Biology and Veterinary Parasitology Unit, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Cluj-Napoca, Romania

⁴Laboratory of Applied Entomology, Section: Agro-Resources Protection, Faculty of Sciences, University of Lome, Lome, Togo

⁵Unité Maladies à Vecteurs et Biodiversité (UMaVeB), Centre International de Recherche-Développement sur l'Élevage en zone Subhumide (CIRDES), Bobo-Dioulasso, Burkina Faso

⁶Functional Genetics of Infectious Diseases Unit, Institut Pasteur, CNRS UMR 2000, Université de Paris, Paris 7, France

⁷Animal Health Department, Nouzilly, France

⁸Friedrich-Loeffler-Institut (FLI), Federal Research Institute for Animal Health, Institute of Novel and Emerging Infectious Diseases, Greifswald-Insel Riems, Germany

Correspondence

Roland Eric Yessinou, Communicable Disease Research Unit (URMaT), University of Abomey-Calavi, 01 P.O. Box: 2009, Cotonou, Benin.
Email: eric.yessinou@gmail.com

Funding information

Francophony University Agency (AUF) for Postdoctoral Fellowship Eugen Ionescu

Abstract

Background: Ticks are obligate hematophagous arthropods capable of transmitting a great variety of endemic and emerging pathogens causing diseases in animals and humans.

Objectives: The aim of this study was to investigate the presence of *Bartonella* spp., *Rickettsia* spp., *Borrelia burgdorferi* sensu lato (s.l.) and *Anaplasma phagocytophilum* in ticks collected from cattle in Benin and Togo.

Methods: Overall, 396 (148 males, 205 females and 43 nymphs) ticks were collected from cattle in 17 districts (Benin and Togo) between 2019 and 2020. Ticks were pooled into groups of 2–6 ticks per pool according to individual host, location, species and developmental stage. The DNA of each pool was extracted for molecular screening.

Results: PCR results revealed that 20 tick pools were positive for *Bartonella* spp. (Benin and Togo) and 23 tick pools positive for *Rickettsia* spp. (Benin), while all pools were negative for *A. phagocytophilum* and *B. burgdorferi* s.l. Sequence analysis of positive *Rickettsia* samples revealed the presence of *Rickettsia aeschlimannii*.

Conclusions: The present study highlights the presence of zoonotic agents in ticks collected from cattle in Benin and Togo. This information will raise awareness of tick-borne diseases among physicians and veterinarians, stimulate further studies to

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Veterinary Medicine and Science* published by John Wiley & Sons Ltd.

monitor these pathogens, and advise on necessary measures to control the spread of these zoonoses.

KEYWORDS

bacteria, public health, *Rickettsia aeschlimannii*, tick, West Africa

1 | INTRODUCTION

Ticks are known to be important disease vectors, capable of transmitting different pathogens such as protozoa, bacteria, and viruses to a wide range of hosts, including humans (Dantas-Torres et al., 2012). The distribution of tick-borne pathogens is influenced by a series of climatic and environmental factors, as well as density and diversity of the host species (Ogden, 2017; Wikel, 2018). Among the tick-borne pathogens, *Rickettsia*, *Anaplasma* or *Bartonella* species are of high importance in cattle (Kasaija et al., 2021). *Rickettsia* spp. are obligate intracellular alpha-proteobacteria, with a worldwide distribution and transmitted to humans and animals by various arthropod vectors, including ticks (Parola et al., 2013). They have a high host adaptability, and the infections are important causes of morbidity and mortality worldwide (Diop et al., 2019; Saravanan et al., 2020). *Rickettsia* spp. are considered emerging zoonotic pathogens that are responsible for conditions such as the spotted fevers and typhus (Bhengri et al., 2016). The continuous identification of new species or genotypes of *Rickettsia* (Parola et al., 2013; Portillo et al., 2015) in ticks brings up new questions on their true diversity, ecology and biology and the role of climate change in their distribution (Dantas-Torres et al., 2012). This study provides information on these bacteria in West Africa, including Benin and Togo.

Bartonella spp. (Rhizobiales) are intracellular Gram-negative bacteria. About 30 *Bartonella* species were described in a great variety of domestic and wild mammals worldwide (Kosoy et al., 2012). *Bartonella* spp. cause endocarditis in humans and domestic animals, including cattle (Erol et al., 2013). Detection of *Bartonella* spp. in bats was reported from Ghana and Nigeria (Billeter et al., 2012; Kamani et al., 2014). It has also been detected in rodents and their ectoparasites in Nigeria (Kamani et al., 2013). Although several *Bartonella* spp. were identified in ticks, their vectorial competence is still under discussion (Telford & Wormser, 2010). Despite the potential public and veterinary health importance, no studies were conducted on these bacteria in Benin and Togo.

Borrelia burgdorferi sensu lato (s.l.) complex comprises more than 20 genospecies and was described worldwide in a great variety of animals and vectors including ticks (Margos et al., 2019). *Borrelia burgdorferi* s.l. is one of the most important tick-borne zoonotic agents in humans (Lefeuvre et al., 2020). They are maintained in nature by the interaction that exists with its vector ticks vertebrate reservoir hosts. *Borrelia burgdorferi* s.l. is transmitted by hard ticks and has been identified in Europe (Marchant et al., 2017), America (Scott et al., 2018) and Asia (Pukhovskaya et al., 2019), but few studies reported this zoonotic agent in Africa (Chitanga et al., 2014). While there are some reports of the presence of *B. burgdorferi* s.l. in Northern Africa (Elhelw et al., 2021), in Western Africa, mainly *B. crocidurae* and other species responsible

for relapsing fever than Lyme borreliosis have been detected (Margos et al., 2019).

Anaplasma phagocytophilum (Anaplasmataceae) is an obligate intracellular bacterium known as a zoonotic tick-borne pathogen and poses significant public health importance (Dumler & Walker, 2001). *Anaplasma phagocytophilum* was reported in several species of wild and domestic mammals worldwide, causing the human granulocytic anaplasmosis (HGA). In animals, it is responsible for reduction of milk production, abortion and also death (Stuen, 2007). *Anaplasma phagocytophilum* has been reported in various countries from Africa including Morocco, Zimbabwe, Tunisia, Algeria and South Africa (El Hamiani Khatat et al., 2021; Kelly et al., 2014; Nakayima et al., 2014). Currently, very few confirmed cases of *A. phagocytophilum* infection were reported in West Africa (Djiba et al., 2013).

In Africa, human tick-borne diseases are underestimated and tick bites in humans go unreported due to the lack of awareness, knowledge of the risk of TBDs and the failure of the epidemiological surveillance system (N'koué Sambiéni et al., 2015; Rapp, 2014; Tutin, 2000). However, accurate identification of pathogens circulating between wild and domestic animals, ticks, and humans in a region is essential to facilitate diagnosis and treatment regimens, which depend on the pathogen involved. The lack of epidemiological knowledge on tick-borne diseases induces confusion, with wrong diagnoses, and therefore, wrong treatment. The purpose of this study was to identify *Bartonella* spp., *Rickettsia* spp., *B. burgdorferi* s.l. and *A. phagocytophilum* in ticks collected from cattle in Benin and Togo (West Africa).

2 | MATERIALS AND METHODS

2.1 | Study area and tick collection

A total of 396 ticks were collected from cattle in 2019–2020 in Benin and Togo (Figure 1). Ticks were preserved in 70% ethanol and transported to the Department of Parasitology and Parasitic Diseases of the University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca. Identification of tick species was done under a stereomicroscope by using morphological keys (Estrada-Peña et al., 2004; J. J. Walker et al., 2000; A. R. Walker et al., 2003).

2.2 | Molecular assessment

Ticks were pooled into groups of 2–6 ticks per pool according to individual host, location, species and developmental stage. The DNA of each pool was extracted using ISOLATE II Genomic DNA Kit




FIGURE 1 Geographical distribution of tick collection areas in Benin and Togo

(Bioline Meridian Bioscience, Luckenwalde, Germany), according to the manufacturer's instructions and stored at -20°C .

Genomic DNA samples were screened for the presence of *Bartonella* spp., *Rickettsia* spp., *B. burgdorferi* s.l. and *A. phagocytophilum* by conventional and nested PCRs (Table 1), targeting the citrate synthase (*gltA*) gene of *Rickettsia* spp. and *Bartonella* spp., the flagellin B (*flaB*) gene of *B. burgdorferi* s.l. and the 16S rRNA gene of *A. phagocytophilum*.

The PCR amplifications were performed in $25\ \mu\text{l}$ reaction volume, containing $12.5\ \mu\text{l}$ Green PCR Mastermix (Roalab GmbH, Teltow, Germany), $6.5\ \mu\text{l}$ of ultra-pure water, $1\ \mu\text{l}$ ($10\ \text{pmol}/\mu\text{l}$) of each primer (Table 1), and $4\ \mu\text{l}$ of isolated DNA aliquot. One negative control (ultra-pure water) as well as a positive control were included. For the nested PCR targeting the *flaB* gene of *B. burgdorferi* s.l. and 16S gene of *A. phagocytophilum*, a second reaction was performed in $25\ \mu\text{l}$ reaction volume, containing $12.5\ \mu\text{l}$ Green PCR Mastermix (Roalab GmbH), $9.5\ \mu\text{l}$ of ultra-pure water, $1\ \mu\text{l}$ ($10\ \text{pmol}/\mu\text{l}$) of each of the two primers (Table 1), and $1\ \mu\text{l}$ aliquot of the first PCR reaction product.

The PCR was performed using the T1000™ Thermal Cycler (Bio-Rad, London, UK). Amplification products were visualized by electrophoresis on 1.5% agarose gel stained with ECO Safe 20,000 × Nucleic Acid Staining Solution (Pacific Image Electronics, New Taipei, Taiwan) and their molecular weight was assessed by comparison to a molecular marker (O'GeneRuler™ 100 bp DNA Ladder, Thermo Fisher Scientific Inc., Waltham, MA, USA). PCR products were purified using

the ISOLATE II PCR and Gel Kit (Bioline Meridian Bioscience) and sent for sequencing (Macrogen Europe, Amsterdam, the Netherlands).

2.3 | Sequencing

All sequences were analyzed and edited using Geneious® 4.85 software. Basic Local Alignments Tool (BLAST) analyses (<https://blast.ncbi.nlm.nih.gov>) were conducted to compare all the obtained sequences with the ones deposited in the GenBank™ database.

3 | RESULTS

3.1 | Molecular detection

A total of 396 (148 males, 205 females and 43 nymphs) ticks were collected from cattle in 17 districts (Benin, Togo) (Figure 1) and divided into 96 pools. Overall, PCR results showed that 20 tick pools were positive for *Bartonella* spp. and 23 pools for *Rickettsia* spp.

In Benin, pools were positive for *Bartonella* spp. (25.39%) and *Rickettsia* spp. (30.15%). *Bartonella* spp. and *Rickettsia* spp. have been detected in *Amblyomma variegatum*, *Hyalomma rufipes* and *Rhipicephalus microplus* (Table 3). The PCR targeting the *gltA* gene allowed identifying

TABLE 1 Primers and polymerase chain reaction conditions used for the detection of pathogens in ticks

Target species	Gene (~amplicon length)	Forward and reverse primers (5'-3') (Reference)	Temperature and duration of:					Number of cycles
			Initial denaturation	Denaturation	Annealing	Extension	Final extension	
<i>Bartonella</i> spp.	<i>gltA</i> (380–400 bp)	bart781: GGG GAC CAG CTC ATG GTG G bart1137: AAT GCA AAA AGA ACA GTA AAC A (Norman et al., 1995)	95°C, 5 m	95°C, 30 s	52°C, 30 s	72°C, 30 s	72°C, 5 m	35
<i>Rickettsia</i> spp.	<i>gltA</i> (381 bp)	Rsf877: GGG GGC CTG CTC ACG GCG G Rsf1258: ATT GCA AAA AGT ACA GTG AAC A (Regnery et al., 1991)	95°C, 2 m	95°C, 30 s	58°C, 30 s	72°C, 30 s	72°C, 2 m	35
<i>B. burgdorferi</i> s.l. (nested PCR)	<i>flaB</i> (650 bp)	FlaLL: ACA TAT TCA GAT GCA GAC AGA GGT FlaRL: TGT TAG ACG TTA CCG ATA CTA ACG (Barbour et al., 1996)	95°C, 5 m	95°C, 30 s	55°C, 30 s	72°C, 45 s	72°C, 5 m	35
	<i>flaB</i> (350–400 bp)	FlaLS: AAC AGC TGA AGA GCT TGG AAT G FlaRS: CGA TAA TCT TAC TAT TCA CTA GTT TC (Clark et al., 2013)	95°C, 5 m	95°C, 30 s	59°C, 30 s	72°C, 30 s	72°C, 5 m	35
<i>A. phagocytophilum</i> (nested PCR)	16S rRNA (945 bp)	Ge3a: CAC ATG CAA GTC GAA CGG ATT ATT C Ge10r: TTC CGT TAA GAA GGA TCT AAT CTC C (Massung et al., 1998)	95°C, 2 m	94°C, 30 s	60°C, 30 s	72°C, 1 m	72°C, 5 m	40
	16S rRNA (570 bp)	Ge9f: AAC GGA TTA TTC TTT ATA GCT TGC T Ge2: GGC AGT ATT AAA AGC AGC TCC AGG (Massung et al., 1998)	95°C, 2 m	94°C, 30 s	55°C, 30 s	72°C, 1 m	72°C, 5 m	30

Abbreviations: m, minute; s, second.

Bartonella spp. and *Rickettsia* spp. in several districts in Benin, such as Abomey-Calavi, Bassila Dassa-Zoumè, Grand-Popo, Kétou, Natitingou, Pehunco, Tchaourou and Zangnanando (Table 2).

In Togo, *gltA* rickettsial DNA was detected in four out of 33 tested tick pools, including *A. variegatum*, *R. microplus*, *Rhipicephalus muhsamae*, and *Rhipicephalus sulcatus* pools. The DNA of *Rickettsia* spp. was identified in *A. variegatum*, *Hyalomma truncatum*, *Rhipicephalus annulatus* and *R. microplus* (Table 3). *Bartonella* spp. were identified in Kamina and Tabligbo, while *Rickettsia* spp. were found in Kamina, Kékékopé and Tabligbo. All samples tested for *B. burgdorferi* s.l. and *A. phagocytophilum* in this study were negative.

3.2 | Sequencing

The *gltA* gene sequences carried out from *Bartonella* spp. obtained from two pools of *R. microplus* in Benin were closely related to a *Bartonella* strain from China (98.23% and 90.76% identity) (accession No. KX354203). Analysis of the *gltA* gene from *Rickettsia* spp. revealed of sequences which had 97.29%–100% identity to various *Rickettsia*

aeschlimannii sequences: accession No. MH267736, MH932014, MH932013; MH675642–MH675648, MK608659–MK608560; KJ663742; KY233219; KU961540 obtained from China, France, Italy, Lebanon and Crimean Peninsula. *Rickettsia aeschlimannii* was detected in seven tick pools including *A. variegatum*, *R. microplus* and *H. rufipes* collected in Benin.

4 | DISCUSSION

Data on the distribution of tick-borne pathogens are necessary for designing appropriate control measures and future approaches to more comprehensive surveillance (Gondard et al., 2017). Very few data exist on this topic in West Africa where a high diversity of tick species was recorded in Benin and Togo (Yessinou et al., 2022). Their presence and distribution can be explained by ecological and climatic conditions that are favourable or unfavourable for the development of ticks in certain countries. All these ticks identified in the study area are capable of playing a role in the transmission and/or maintenance of *Bartonella* spp. and *Rickettsia* spp. (Dantas-Torres et al., 2012). In this study, five

TABLE 2 *Bartonella* spp. and *Rickettsia* spp. detected in different study sites

Country	District	Pools (n)	<i>Bartonella</i> spp.			<i>Rickettsia</i> spp.		
			M	F	N	M	F	N
Benin	Abomey-Calavi	6 (26)	-	1/6	-	1/6	1/6	-
	Adjara	4 (19)	-	-	-	-	-	-
	Bassila	3 (8)	-	-	-	-	1/3	-
	Dassa-Zoumè	7 (36)	1/7	2/7	-	1/7	1/7	-
	Djidja	4 (21)	-	-	-	-	-	-
	Grand-Popo	6 (20)	1/6	1/6	-	-	1/6	-
	Kandi	4 (19)	-	-	-	-	-	-
	Kétou	3 (18)	-	-	-	1/3	1/3	-
	Natitingou	4 (23)	1/4	2/4	-	1/4	2/4	-
	Pehunco	4 (19)	1/4	1/3	-	1/4	3/4	-
	Tanguiéta	5 (20)	-	-	-	-	-	-
	Tchaourou	6 (32)	-	1/6	-	1/6	-	-
	Zangnanando	7 (30)	1/7	1/7	-	2/7	1/7	-
	Subtotal	63 (291)	5/63	9/63	-	8/63	11/63	-
Togo	Kamina	10 (27)	-	1/10	-	-	1/1-	-
	Kékékopé	6 (23)	-	-	-	-	1/6	-
	Adétikopé	9 (25)	-	-	-	-	-	-
	Tabligbo	8 (30)	3/8	-	-	-	2/8	-
	Subtotal	33 (105)	3/33	1/33	-	-	4/33	-
Total	96/396	8/96	10/96	-	8/96	15/96	-	

tick species collected from cattle were positive for the presence of *Bartonella* DNA, namely *A. variegatum*, *R. microplus*, *H. rufipes*, *R. muhsamae*, and *R. sulcatu*s. Several species of *Bartonella* have been reported in ticks, mammals and humans so far (Tsai et al., 2010; Vayssier-Taussat et al., 2015). Our findings represent the first detection of *Bartonella* spp. in hard ticks in Benin and Togo. *Bartonella* spp. were previously also detected in rodents and ticks from Nigeria (Kamani et al., 2013).

Sequencing results also confirmed that *A. variegatum*, *R. microplus* and *H. rufipes* were infected with *R. aeschlimannii*, a tick-borne pathogen, reported in 1997 in *Hyalomma marginatum* ticks collected in Morocco (Beati et al., 1997). For the first time, the presence of *R. aeschlimannii* in ticks collected from cattle in Benin was shown in this study. In Senegal, Mediannikov et al. (2010) reported *R. aeschlimannii* in *H. (m) rufipes*, *H. truncatum* and *Rhipicephalus evertsi evertsi* ticks collected from cows, donkeys, sheep, goats and horses. Others studies in Niger, Mali, Nigeria Ivory-Coast and Burkina-Faso identified *R. aeschlimannii* in *Hyalomma* spp. (Ehounoud et al., 2016; Kamani et al., 2015; Parola et al., 2001; Tomassone et al., 2016), but also in Europe (Fernández Soto et al., 2003; Punda-Polic et al., 2003). *Hyalomma marginatum* appears to serve as a vector but also as a reservoir of *R. aeschlimannii* (Parola et al., 2005). *Rickettsia aeschlimannii* was reported in other tick genera including *Amblyomma*, *Dermacentor*, *Ixodes* and *Rhipicephalus* (Orkun et al., 2014a; Rumer et al., 2011; Toma et al., 2014). This pathogen is implicated in several cases of fever in humans but also reported in several places in animals (Orkun et al., 2014b; Tosoni

et al., 2016). The symptoms following an infection are similar to those of Mediterranean spotted fever caused by *Rickettsia conorii* (Rovero & Raoult, 2008). Raoult et al. (2002) reported cases of human infection caused by *R. aeschlimannii* in France.

New tick-borne pathogens infecting humans have increased in recent years, thus playing an important role in public health. In sub-Saharan Africa, *R. aeschlimannii* is known as a potentially important pathogen and was identified in humans, animals and ticks (Parola et al., 2013). In these regions, febrile illnesses in breeders, farmers and animal health professionals may be caused by rickettsiosis (Moumouni et al., 2016). Mediannikov et al. (2010) showed that tick-borne rickettsioses are among the causes of acute nonmalaria febrile diseases. Traditionally, in West Africa, almost all febrile conditions are generally considered to be linked to malaria and are treated as such without further investigations, but other tick-borne pathogens might be involved. Thus, clinical manifestations of Rickettsioses should be considered in the differential diagnosis of the malaria in patients mainly in rural areas or locations of ruminants rearing.

It should also be noted that the identification of pathogen DNA in a tick does not imply that it is necessarily a biological vector. Studies must be conducted to prove the vector competence of the tick species. The growth of the world population, the transformation of natural habitats, the global changes and the practices of use of the fauna are some of the factors that modify and facilitate the interactions between wild and anthropic environments (Cable et al., 2017). These growing contacts

TABLE 3 *Bartonella* spp. and *Rickettsia* spp. detected in tick species

Country	Tick species	Number of tested ticks			Pools (n)	PCR positive pools	
		Male	Female	Nymph		<i>Bartonella</i> spp.	<i>Rickettsia</i> spp.
Benin	<i>Amblyomma variegatum</i>	73	62	12	29 (147)	24.13 7/29	48.27 14/29
	<i>Hyalomma rufipes</i>	13	8	-	9 (21)	33.33 3/9	33.33 3/9
	<i>Rhipicephalus microplus</i>	15	77	31	25 (123)	24 6/25	8 2/25
	Subtotal	101	147	43	63 (291)	25.39 16/63	30.15 19/63
Togo	<i>Amblyomma variegatum</i>	24	16	-	8 (40)	12.5 1/8	12.5 1/8
	<i>Hyalomma truncatum</i>	6	5	-	7 (11)	-	14.28 1/7
	<i>Rhipicephalus annulatus</i>	2	6	-	4 (5)	-	25 1/4
	<i>Rhipicephalus decoloratus</i>	-	3	-	1 (3)	-	-
	<i>Rhipicephalus lunulatus</i>	-	3	-	2 (3)	-	-
	<i>Rhipicephalus microplus</i>	4	21	-	5 (26)	20 1/5	20 1/5
	<i>Rhipicephalus muhsamae</i>	-	2	-	1 (2)	100 1/1	-
	<i>Rhipicephalus sanguineus</i>	8	2	-	4 (10)	-	-
	<i>Rhipicephalus sulcatus</i>	3	-	-	1 (3)	100 1/1	-
Subtotal	47	58	0	33 (105)	12.12 4/33	12.12 4/33	
Total					20.83 20/96	23.95 23/96	

between wildlife and domestic animals, humans and ticks gradually promote the exchange of pathogens that can have harmful health consequences on the animals and humans.

5 | CONCLUSION

This study reports the identification and distribution of pathogens in hard ticks collected from cattle in West Africa. These results suggest the need to include bartonellosis and rickettsiosis among the causes of febrile illnesses among breeders, para-veterinarians, veterinarians and travellers in West Africa. Doctors should also consider these tick-borne illnesses as a differential diagnostic with malaria. On the other hand, the role of domestic and wild animals in the epidemiology of diseases transmitted by ticks requires further investigation.

AUTHOR CONTRIBUTIONS

Conceptualization, data curation, formal analysis, investigation, methodology, and writing—original draft: Roland Eric Yessinou. *Conceptualization, formal analysis, and methodology:* Cristina Daniela Cazan. *Conceptualization, formal analysis, and methodology:* Luciana Cătălina Panait. *Investigation:* Eyabana Mollong. *Conceptualization and methodology:* Abel S. Biguezoton. *Conceptualization, methodology, and writing—original draft:* Sarah Irène Bonnet. *Conceptualization:* Souaïbou Farougou.

ACKNOWLEDGEMENT
We are grateful to the Francophony University Agency (AUF) for Post-doctoral Fellowship EUGEN IONESCU and Department of Parasitology and Parasitic Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Romania for supporting this research.

ACKNOWLEDGEMENT

CONFLICT OF INTEREST
The authors declare no conflict of interest.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ETHICS STATEMENT

No ethical approval was required, as this study does not involve clinical trials or experimental procedures. The cattle's are still alive and used

for milk and meat production. This study did not involve endangered or protected species.

ORCID

Roland Eric Yessinou  <https://orcid.org/0000-0003-1743-318X>

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1002/vms3.1022>.

REFERENCES

- Barbour, A. G., Maupin, G. O., Teltow, G. J., Carter, C. J., & Piesman, J. (1996). Identification of an uncultivable *Borrelia* species in the hard tick *Amblyomma americanum*: Possible agent of a Lyme disease-like illness. *Journal of Infectious Diseases*, 173, 403–409.
- Beati, L., Meskini, M., Thiers, B., & Raoult, D. (1997). *Rickettsia aeschlimannii* sp. nov., a new spotted fever group rickettsia associated with *Hyalomma marginatum* ticks. *International Journal of Systematic and Evolutionary Microbiology*, 47, 548–554.
- Bhengsi, S., Baggett, H. C., Edouard, S., Dowell, S. F., Dasch, G. A., Fisk, T. L., Raoult, D., & Parola, P. (2016). *Sennetsu neorickettsiosis*, spotted fever group, and typhus group rickettsioses in three provinces in Thailand. *American Journal of Tropical Medicine and Hygiene*, 95, 43–49.
- Billeter, S. A., Hayman, D. T. S., Peel, A. J., Baker, K., Wood, J. L. N., Cunningham, A., Suu-Ire, R., Dittmar, K., & Kosoy, M. Y. (2012). Bartonellaspaces in bat flies (Diptera: Nycteribiidae) from western Africa. *Parasitology*, 139(3), 324–329. <https://doi.org/10.1017/S0031182011002113>
- Cable, J., Barber, I., Boag, B., Ellison, A. R., Morgan, E. R., Murray, K., Pascoe, E. L., Sait, S. M., Wilson, A. J., & Booth, M. (2017). Global change, parasite transmission and disease control: Lessons from ecology. *Philosophical Transactions of The Royal Society B Biological Sciences*, 372, 20160088.
- Chitanga, S., Gaff, H., & Mukaratirwa, S. (2014). Tick-borne pathogens of potential zoonotic importance in the southern African Region. *Journal of the South African Veterinary Association*, 85, 01–03.
- Clark, K. L., Leydet, B., & Hartman, S. (2013). Lyme borreliosis in human patients in Florida and Georgia, USA. *International Journal of Medical Sciences*, 10, 915.
- Dantas-Torres, F., Chomel, B. B., & Otranto, D. (2012). Ticks and tick-borne diseases: A One Health perspective. *Trends in Parasitology*, 28, 437–446.
- Diop, A., Raoult, D., & Fournier, P.-E. (2019). Paradoxical evolution of rickettsial genomes. *Ticks and Tick-borne Diseases*, 10, 462–469.
- Djiba, M. L., Mediannikov, O., Mbengue, M., Thiongane, Y., Molez, J.-F., Seck, M. T., Fenollar, F., Raoult, D., & Ndiaye, M. (2013). Survey of Anaplasmataceae bacteria in sheep from Senegal. *Tropical Animal Health and Production*, 45, 1557–1561.
- Dumler, J. S., & Walker, D. H. (2001). Tick-borne ehrlichioses. *The Lancet Infectious Diseases*, 1, 21–28.
- Ehounoud, C. B., Yao, K. P., Dahmani, M., Achi, Y. L., Amanzougaghene, N., Kacou N'Douba, A., N'Guessan, J. D., Raoult, D., Fenollar, F., & Mediannikov, O. (2016). Multiple pathogens including potential new species in tick vectors in Côte d'Ivoire. *PLoS Neglected Tropical Diseases*, 10, e0004367.
- El Hamiani Khatat, S., Daminet, S., Duchateau, L., Elhachimi, L., Kachani, M., & Sahibi, H. (2021). Epidemiological and clinicopathological features of *Anaplasma phagocytophilum* infection in dogs: A systematic review. *Frontiers in Veterinary Science*, 8, 686644. <https://doi.org/10.3389/fvets.2021.686644>
- Elhelw, R., Elhariri, M., Hamza, D., Abuowarda, M., Ismael, E., & Farag, H. (2021). Evidence of the presence of *Borrelia burgdorferi* in dogs and associated ticks in Egypt. *BMC Veterinary Research*, 17, 1–9.
- Erol, E., Jackson, C., Bai, Y., Sells, S., Locke, S., & Kosoy, M. (2013). *Bartonella bovis* isolated from a cow with endocarditis. *Journal of Veterinary Diagnostic Investigation*, 25, 288–290.
- Estrada-Peña, A., Bouattour, A., Camicas, J. L., & Walker, A. R. (2004). *Ticks of domestic animals in the Mediterranean region: A guide to identification of species*. University of Zaragoza.
- Fernández Soto, P., Grandes, A. E., & Pérez Sánchez, R. (2003). *Rickettsia aeschlimannii* in Spain: Molecular evidence in *Hyalomma marginatum* and five other tick species that feed on humans. *Emerging Infectious Diseases*, 9(7), 889–890.
- Gondard, M., Cabezas-Cruz, A., Charles, R. A., Vayssier-Taussat, M., Albina, E., & Moutailler, S. (2017). Ticks and tick-borne pathogens of the Caribbean: Current understanding and future directions for more comprehensive surveillance. *Frontiers in Cellular and Infection Microbiology*, 7, 490.
- Kamani, J., Baneth, G., Apanaskevich, D. A., Mumcuoglu, K. Y., & Harrus, S. (2015). Molecular detection of *Rickettsia aeschlimannii* in *Hyalomma* spp. ticks from camels (*Camelus dromedarius*) in Nigeria, West Africa. *Medical and Veterinary Entomology*, 29, 205–209. <https://doi.org/10.1111/mve.12094>
- Kamani, J., Morick, D., Mumcuoglu, K. Y., & Harrus, S. (2013). Prevalence and diversity of *Bartonella* species in commensal rodents and ectoparasites from Nigeria, West Africa. *PLoS Neglected Tropical Diseases*, 7, e2246.
- Kasajja, P. D., Estrada-Peña, A., Contreras, M., Kirunda, H., & de la Fuente, J. (2021). Cattle ticks and tick-borne diseases: A review of Uganda's situation. *Ticks and Tick-borne Diseases*, 12(5), 101756.
- Kelly, P., Marabini, L., Dutlow, K., Zhang, J., Loftis, A., & Wang, C. (2014). Molecular detection of tick-borne pathogens in captive wild felids, Zimbabwe. *Parasites & Vectors*, 7, 1–6.
- Kosoy, M., Hayman, D. T., & Chan, K. - S. (2012). *Bartonella* bacteria in nature: Where does population variability end and a species start? *Infection, Genetics and Evolution*, 12, 894–904.
- Lefevre, B., Cantero, P., Ehret-Sabatier, L., Lenormand, C., Barthel, C., Po, C., Parveen, N., Grillon, A., Jaulhac, B., & Boulanger, N. (2020). Effects of topical corticosteroids and lidocaine on *Borrelia burgdorferi* sensu lato in mouse skin: Potential impact to human clinical trials. *Scientific Reports*, 10, 1–13.
- Marchant, A., Coupance, A. L., Joly, C., Perthame, E., Sertour, N., Garnier, M., Godard, V., Ferquel, E., & Choumet, V. (2017). Infection of *Ixodes ricinus* by *Borrelia burgdorferi* sensu lato in peri-urban forests of France. *PLoS One*, 12, e0183543.
- Margos, G., Fingerle, V., & Reynolds, S. (2019). *Borrelia bavariensis*: Vector switch, niche invasion, and geographical spread of a tick-borne bacterial parasite. *Frontiers in Ecology and Evolution*, 7(401), 1–20.
- Massung, R. F., Slater, K., Owens, J. H., Nicholson, W. L., Mather, T. N., Solberg, V. B., & Olson, J. G. (1998). Nested PCR assay for detection of granulocytic ehrlichiae. *Journal of Clinical Microbiology*, 36, 1090–1095.
- Mediannikov, O., Diatta, G., Fenollar, F., Sokhna, C., Trape, J.-F., & Raoult, D. (2010). Tick-borne rickettsioses, neglected emerging diseases in rural Senegal. *PLoS Neglected Tropical Diseases*, 4, e821.
- Moumouni, P. F. A., Terkawi, M. A., Jirapattharasate, C., Cao, S., Liu, M., Nakao, R., Umemiya-Shirafuji, R., Yokoyama, N., Sugimoto, C., & Fujisaki, K. (2016). Molecular detection of spotted fever group rickettsiae in *Amblyomma variegatum* ticks from Benin. *Ticks and Tick-borne Diseases*, 7, 828–833.
- Nakayima, J., Hayashida, K., Nakao, R., Ishii, A., Ogawa, H., Nakamura, I., Moonga, L., Hang'ombe, B. M., Mweene, A. S., & Thomas, Y. (2014). Detection and characterization of zoonotic pathogens of free-ranging non-human primates from Zambia. *Parasites & Vectors*, 7, 1–7.
- N'koué Sambiéni, E., Danko, N., & Ridde, V. (2015). La Fièvre Hémorragique à Virus Lassa au Bénin en 2014 en contexte d'Ebola : une épidémie révélatrice de la faiblesse du système sanitaire. *Anthropologie et Santé*, 11, 16. <https://doi.org/10.4000/anthropologiesante.1772>

- Norman, A. F., Regnery, R., Jameson, P., Greene, C., & Krause, D. C. (1995). Differentiation of *Bartonella*-like isolates at the species level by PCR-restriction fragment length polymorphism in the citrate synthase gene. *Journal of Clinical Microbiology*, 33, 1797–1803.
- Ogden, N. H. (2017). Climate change and vector-borne diseases of public health significance. *FEMS Microbiology Letters*, 364, fnx186.
- Orkun, Ö., Karaer, Z., Çakmak, A., & Nalbantoğlu, S. (2014a). Spotted fever group rickettsiae in ticks in Turkey. *Ticks and Tick-borne Diseases*, 5, 213–218. <https://doi.org/10.1016/j.ttbdis.2012.11.018>
- Orkun, Ö., Karaer, Z., Çakmak, A., & Nalbantoğlu, S. (2014b). Identification of tick-borne pathogens in ticks feeding on humans in Turkey. *PLoS Neglected Tropical Diseases*, 8, e3067. <https://doi.org/10.1371/journal.pntd.0003067>
- Parola, P., Inokuma, H., Camicas, J.-L., Brouqui, P., & Raoult, D. (2001). Detection and identification of spotted fever group Rickettsiae and Ehrlichiae in African ticks. *Emerging Infectious Diseases*, 7, 1014–1017.
- Parola, P., Paddock, C. D., & Raoult, D. (2005). Tick-borne rickettsioses around the world: Emerging diseases challenging old concepts. *Clinical Microbiology Reviews*, 18, 719–756.
- Parola, P., Paddock, C. D., Socolovschi, C., Labruna, M. B., Mediannikov, O., Kernif, T., Abdad, M. Y., Stenos, J., Bitam, I., & Fournier, P.-E. (2013). Update on tick-borne rickettsioses around the world: A geographic approach. *Clinical Microbiology Reviews*, 26, 657–702.
- Portillo, A., Santibáñez, S., García-Álvarez, L., Palomar, A. M., & Oteo, J. A. (2015). Rickettsioses in Europe. *Microbes and Infection*, 17, 834–838.
- Pukhovskaya, N. M., Morozova, O. V., Vysochina, N. P., Belozerova, N. B., & Ivanov, L. I. (2019). Prevalence of *Borrelia burgdorferi* sensu lato and *Borrelia miyamotoi* in ixodid ticks in the Far East of Russia. *International Journal for Parasitology: Parasites and Wildlife*, 8, 192–202.
- Punda-Polic, V., Petrovec, M., Trilar, T., Duh, D., Bradaric, N., Klismanic, Z., & Avsic-Zupanc, T. (2003). Detection and identification of spotted fever group rickettsiae in ticks collected in southern Croatia. *Ticks and tick-borne pathogens* (pp.169–176). Springer.
- Raoult, D., Fournier, P.-E., Abboud, P., & Caron, F. (2002). First documented human *Rickettsia aeschlimannii* infection. *Emerging Infectious Diseases*, 8, 748. <https://doi.org/10.3201/eid0807.010480>
- Rapp, C. (2014). Epidémie à virus Ebola en Afrique de l'Ouest: réalités et perspectives. *Médecine Santé Trop*, 24, 229–231.
- Regnery, R. L., Spruill, C. L., & Plikaytis, B. D. (1991). Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *Journal of Bacteriology*, 173, 1576–1589.
- Rovero, C., & Raoult, D. (2008). Mediterranean spotted fever. *Infectious Disease Clinics of North America*, 22, 515–530. <https://doi.org/10.1016/j.idc.2008.03.003>
- Rumer, L., Graser, E., Hillebrand, T., Talaska, T., Dautel, H., Mediannikov, O., Roy-Chowdhury, P., Sheshukova, O., Mantke, O. D., & Niedrig, M. (2011). *Rickettsia aeschlimannii* in *Hyalomma marginatum* ticks, Germany. *Emerging Infectious Diseases*, 17, 325.
- Saravanan, P., Nagaraj, M. V., & Somashekhar, C. (2020). Rickettsial fever in tertiary care hospital in rural Bengaluru: Clinical profile and complications. *Indian Journal of Maternal & Child Health*, 7, 385–387.
- Scott, J. D., Clark, K. L., Foley, J. E., Bierman, B. C., & Durden, L. A. (2018). *Far-reaching dispersal of Borrelia burgdorferi sensu lato-infected blacklegged ticks by migratory songbirds in Canada*. Healthcare. Multidisciplinary Digital Publishing Institute.
- Stuen, S. (2007). *Anaplasma phagocytophilum*—The most widespread tick-borne infection in animals in Europe. *Veterinary Research Communications*, 31, 79–84.
- Telford III, S. R., & Wormser, G. P. (2010). *Bartonella* spp. transmission by ticks not established. *Emerging Infectious Diseases*, 16, 379.
- Toma, L., Mancini, F., Di Luca, M., Cecere, J. G., Bianchi, R., Khoury, C., Quarchioni, E., Manzia, F., Rezza, G., & Ciervo, A. (2014). Detection of microbial agents in ticks collected from migratory birds in central Italy. *Vector-Borne and Zoonotic Diseases*, 14, 199–205.
- Tomassone, L., De Meneghi, D., Adakal, H., Rodighiero, P., Pressi, G., & Grego, E. (2016). Detection of *Rickettsia aeschlimannii* and *Rickettsia africae* in ixodid ticks from Burkina Faso and Somali Region of Ethiopia by new real-time PCR assays. *Ticks and Tick-borne Diseases*, 7, 1082–1088.
- Tosoni, A., Mirijello, A., Ciervo, A., Mancini, F., Rezza, G., Damiano, F., & Addolorato, G. (2016). Human *Rickettsia aeschlimannii* infection: First case with acute hepatitis and review. *European Review for Medical and Pharmacological Sciences*, 20, 2630–2633.
- Tsai, Y.-L., Chuang, S.-T., Chang, C.-C., Kass, P. H., & Chomel, B. B. (2010). *Bartonella* species in small mammals and their ectoparasites in Taiwan. *American Journal of Tropical Medicine and Hygiene*, 83, 917–923.
- Tutin, C. E. G. (2000). Ecologie et organisation sociale des primates de la forêt tropicale africaine: aide à la compréhension de la transmission des retrovirus. *Bulletin de la Société de Pathologie Exotique*, 93, 3–157.
- Vayssier-Taussat, M., Kazimirova, M., Hubalek, Z., Hornok, S., Farkas, R., Cosson, J.-F., Bonnet, S., Vourch, G., Gasqui, P., Mihalca, A. D., Plantard, O., Silaghi, C., Cutler, S., & Rizzoli, A. (2015). Emerging horizons for tick-borne pathogens: From the 'one pathogen–one disease' vision to the pathobiome paradigm. *Future Microbiology*, 10, 2033–2043. <https://doi.org/10.2217/fmb.15.114>
- Walker, A. R., Bouattour, A. A., Camicas, J. L., & Estrada-Pena, I. G. (2003). *Ticks of domestic animals in Africa: A guide to identification of species*. Bioscience Reports Edinburgh.
- Walker, J. B., Keirans, J. E., & Horak, I. G. (2000). The genus *Rhipicephalus* (Acari, Ixodidae): A guide to the brown ticks of the world. *Tropical Animal Health and Production*, 32, 417.
- Wikel, S. K. (2018). Ticks and tick-borne infections: Complex ecology, agents, and host interactions. *Journal of Veterinary Science*, 5, 60.
- Yessinou, R. E., Cazan, C. D., Bonnet, S. I., Farougou, S., & Mihalca, A. D. (2022). Geographical distribution of hard ticks (Acari:Ixodidae) and tick-host associations in Benin, Burkina-Faso, Ivory-Coast and Togo. *Acta Tropica*, 232, 106510.

How to cite this article: Yessinou, R. E., Cazan, C. D., Panait, L. C., Mollong, E., Biguezoton, A. S., Bonnet, S. I., Farougou, S., Groschup, M. H., & Mihalca, A. D. (2022). New geographical records for tick-borne pathogens in ticks collected from cattle in Benin and Togo. *Veterinary Medicine and Science*, 1–8. <https://doi.org/10.1002/vms3.1022>