



HAL
open science

Don't forget to be picky – selective autophagy of protein aggregates in neurodegenerative diseases

Anne Simonsen, Thomas Wollert

► To cite this version:

Anne Simonsen, Thomas Wollert. Don't forget to be picky – selective autophagy of protein aggregates in neurodegenerative diseases. *Current Opinion in Cell Biology*, 2022, 75, pp.102064. 10.1016/j.cecb.2022.01.009 . pasteur-04047116

HAL Id: pasteur-04047116

<https://pasteur.hal.science/pasteur-04047116>

Submitted on 27 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Don't forget to be picky – selective autophagy of protein aggregates in neurodegenerative diseases

Anne Simonsen^{a,b} and Thomas Wollert^c

Abstract

The homeostasis of cells depends on the selective degradation of damaged or superfluous cellular components. Autophagy is the major pathway that recognizes such components, sequesters them in *de novo* formed autophagosomes and delivers them to lysosomes for degradation. The recognition of specific cargo and the biogenesis of autophagosomes involve a dedicated machinery of autophagy related (ATG) proteins. Intense research over the past decades has revealed insights into the function of autophagy proteins and mechanisms that govern cargo recognition. Other aspects including the molecular mechanisms involved in the onset of human diseases are less well understood. However, autophagic dysfunctions, caused by age related decline in autophagy or mutations in ATG proteins, are directly related to a large number of human pathologies including neurodegenerative disorders. Here, we review most recent discoveries and breakthroughs in selective autophagy and its relationship to neurodegeneration.

Addresses

^a Department of Molecular Medicine, Institute of Basic Medical Sciences and Centre for Cancer Cell Reprogramming, Institute of Clinical Medicine, University of Oslo, 0372 Oslo, Norway

^b Department of Molecular Cell Biology, The Norwegian Radium Hospital Montebello, 0379, Oslo, Norway

^c Membrane Biochemistry and Transport, Institut Pasteur, UMR3691 CNRS, 75015, Paris, France

Corresponding authors: Simonsen, Anne (anne.simonsen@medisin.uio.no); Wollert, Thomas (thomas.wollert@pasteur.fr)

Current Opinion in Cell Biology 2022, 75:102064

This review comes from a themed issue on **Membrane Trafficking 2022**

Edited by **Chiara Zurzolo** and **Benjamin Glick**

For complete overview of the section, please refer the article collection - [Membrane Trafficking 2022](#)

Available online 28 February 2022

<https://doi.org/10.1016/j.ceb.2022.01.009>

0955-0674/© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Autophagy was discovered more than 60 years ago as a mechanism by which the cell captures cytoplasm in double membrane vesicles, termed autophagosomes, upon starvation. It took much longer to reveal that

autophagy also degrades damaged or superfluous cytoplasmic components, including protein aggregates and mitochondria, in a highly selective manner [1]. An important milestone was the discovery of a direct relationship of selective autophagy and human diseases, including cancer and neurodegeneration [2]. Notably, proteins that are prone to aggregate into amyloid-like filaments, including Huntingtin, α -Synuclein, Tau and the Amyloid-Precursor-Protein fragment A β , are causative agents for neurodegenerative diseases and also substrates of selective autophagy [3]. Mechanistic studies of the formation of such aggregates and their degradation by selective autophagy (aggrephagy) have revealed the importance of membrane-less organelles (MLO) and liquid–liquid phase separation (LLPS) [4]. Moreover, autophagy receptors have been found to regulate the formation of LLPs and initiate selective autophagy by recruiting core autophagy components to cargo. The goal of this review is to summarize most recent advances in our understanding of how selective autophagy counteracts amyloid-like aggregation and neurodegeneration.

Recognition of autophagic substrates by the autophagy machinery

Selective autophagy is orchestrated by a dedicated machinery (Box 1) that assembles in a spatiotemporally well-defined order to enable sequestration of cytoplasmic cargo by a phagophore membrane. The process starts with the recruitment of autophagy receptors to cargo, which can occur in an ubiquitin (Ub) dependent or independent manner [1]. Examples of Ub-dependent autophagy receptors include the Sequestosome-like receptors SQSTM1/p62, NBR1, OPTN, NDP52 and well as human CUET family proteins [5]. These receptors possess canonical Ub-binding motifs such as UBA, UBZ and UBA domains and their recruitment depends on cargo ubiquitination by upstream acting E3-ligases [6]. Ubiquitin-independent autophagy receptors are often integral membrane proteins being activated in response to organellar stress or dysfunction. Examples include mitochondrial receptors BNIP3 and BNIP3L or ER receptors FAM134B and RTNL3 [1]. Common to all autophagy receptors is the presence of an LC3 interacting region (LIR) that mediates their interaction with ATG8 proteins to promote tethering of cargo to the phagophore membrane by multivalent interactions of


receptors and membrane-bound ATG8 proteins [7]. The autophagy-related kinases ULK1 and TBK1 can phosphorylate residues within and surrounding the core LIR motif to regulate the affinity and specificity of cargo receptors for ATG8 proteins [8,9]. In addition to their interaction with cargo and ATG8 proteins, several autophagy receptors have recently been found to facilitate recruitment of the early autophagy machinery (Box 1), leading to *de novo* phagophore formation and LC3 lipidation in a cargo-dependent manner [10]. However, the specificity and redundancy of cargo receptors in different types of selective autophagy are just starting to be unraveled [11].

Role of cargo receptors in liquid–liquid phase separation and aggrephagy

Misfolded ubiquitinated proteins are substrates for proteasomal degradation, but can also form phase-separated condensates that become substrates for selective autophagy in a process involving cargo receptors

(Figure 1). p62 is key to this process, as it contains a PB1 domain that promotes its self-polymerization to form filamentous polymers [12–14] that again can assemble into gel-like liquid droplets [15], referred to as p62 bodies or p62 gels. Elegant *in vitro* studies showed that p62 is sufficient to phase separate ubiquitinated proteins into condensates [16,17]. p62 recruits NBR1 via PB1-PB1 interactions, which further promotes condensate formation by high affinity binding to ubiquitin and recruitment of TAX1BP1 [18]. These three cargo receptors are all able to interact with the ULK1 complex subunit FIP200, leading to recruitment of the downstream autophagy machinery and *de novo* phagophore formation around the p62 bodies. This may happen independently of ATG8 lipidation [19], although an LIR-dependent interaction of p62 with ATG8 proteins was found to facilitate engulfment of whole p62-gels by autophagosomes [20]. TAX1BP1 further recruits the kinase TBK1 to p62 bodies, leading to phosphorylation of p62 to enhance its binding to ubiquitin [21].

Figure 1



Aggrephagy facilitates clearance of misfolded ubiquitinated proteins. Environmental and cellular stress can cause protein misfolding and their ubiquitination. The autophagy receptor protein p62 contains a self-polymerizing PB1 domain and an UBA domain that together with the autophagy receptors NBR1 and TAX1BP1 and the kinase TBK1 mediate liquid–liquid phase separation of ubiquitinated proteins to form p62 bodies. These autophagy receptors further promote phagophore formation by recruitment of the ULK complex subunit FIP200 and ATG8 proteins, leading to autophagic clearance of parts of or whole p62 bodies.

It is not clear what triggers the oligomerization of p62, but it has been found that oxidative stress leads to oxidation of p62 at cysteine residues (105 and 1113), enhancing its oligomerization and condensation [22]. The ability of p62 to oligomerize and phase separate can also be regulated by chaperones and post-translational modifications of its UBA and PB1 domain. Expression of the prefoldin-like chaperone UXT promoted p62 clustering and autophagic clearance of p62 bodies and delayed SOD1(A4V)-induced degeneration of motor neurons in a *Xenopus* model system [23]. It has also been suggested that the type of ubiquitin chain linked to cargo regulates p62 oligomerization and aggrephagy. Importantly, p62 oligomerization is physiologically important. Upon oxidative stress, the p62-interacting protein Keap1 is recruited to p62 gels, leading to activation of the transcription factor Nrf2 important for the anti-oxidative response [20]. This is further stimulated by NBR1, which is induced in response to oxidative stress and mediates p62 body formation [24]. Impaired turnover of p62-gels leads to *in vivo* Nrf2 hyperactivation and liver injury [20]. Moreover, several mutations of TBK1 or p62 linked to amyotrophic lateral sclerosis (ALS) and fronto-temporal dementia (FTD) disrupt p62 oligomerization and its binding to ubiquitinated cargo, as well as the recruitment of KEAP1 to p62 bodies under oxidative stress, leading to formation of TDP-43 associated stress granules implicated in ALS-FTD neurotoxicity [22]. Moreover, premature lipofuscin accumulation and ubiquitin positive aggregates were detected in the brain of young TAX1BP1 knockout mice, showing *in vivo* relevance of aggrephagy [25].

A recent physical model shows that the surface tension of p62 bodies supports the formation of autophagosomal membrane sheets and that the bending of such sheets can lead to sequestration of a piece of a droplet or an entire droplet [26]. In line with this, aggrephagy cargo segregated by liquid–liquid phase separation in p62 bodies seems to be highly mobile and undergo dynamic exchange with the surrounding cytosol. The exact mechanisms involved in *de novo* phagophore biogenesis during aggrephagy still remains largely unknown, but knowledge from other types of selective autophagy indicates that autophagy receptors share some general functions, including receptor-mediated recruitment of FIP200 to the cargo. Similar to p62, NBR1 and TAX1BP, the cargo receptor NDP52 recruits FIP200 to ubiquitinated cargo [27]. Elegant structural characterization shows that NDP52 promotes membrane-binding of the ULK1 complex by allosteric regulation of the membrane-binding portion of the FIP200 coiled coil domain [28] and that its LIR domain directly interacts with the claw domain of FIP200 [29], indicating that FIP200 compete with ATG8 proteins for binding to autophagy receptors [30]. ATG8 proteins can also be recruited to cargo in a LIR-independent manner to further mediate ubiquitin-independent recruitment of cargo receptors via the

LIR motif, thereby amplifying selective autophagy [31]. Another mitophagy receptor, OPTN, forms a complex with vesicles containing ATG9A [32], a lipid scramblase required for phagophore biogenesis [33]. The function of OPTN in Parkin-dependent mitophagy is regulated by its phosphorylation by both TBK1 and ULK1 complex activity [34]. The high structural similarity between the N-terminal domain of FIP200 and the dimerization domain of TBK1 suggest they may function in a similar manner [35]. In addition to ubiquitin, cargo-specific “eat me” signals such as mitochondrial NIPSNAP proteins [36] can contribute to recruitment of cargo specific receptors. Thus, complex multivalent interactions of autophagy receptors with several core autophagy proteins seem to drive *de novo* synthesis of autophagosomal membranes during selective autophagy [31].

Interestingly, phase separated condensates are not only formed by p62, but also by many amyloid-like proteins. Revealing how these condensates influence each other and whether co-condensation of p62 and cargo occurs is of particular interest, since phase condensates act as nucleation centers for the formation of amyloid-like fibrils that are the hallmark of neurodegeneration and proteinopathies [37].

Membrane-less organelles in neurodegenerative diseases

Liquid–liquid phase separation (LLPS) is a fundamental physical property of macromolecules that is based on dehydration, that is unmixing of an aqueous solution by replacing macromolecule–water interactions by inter-macromolecular interactions [38]. Recent advances revealed that many cellular structures that have initially been described as puncta are in fact biological condensates or membrane-less organelles (MLO). Thus, LLPS emerges as a more general phenomenon that helps structuring the cellular content to facilitate mRNA processing, local translation, and chromatin compaction [39]. MLOs can also serve very distinct and specialized functions. Examples include neurons, which rely heavily on MLOs to structure pre- and postsynaptic densities and to regulate neural plasticity [40].

Most proteins that undergo spontaneous LLPS contain intrinsically disordered regions (IDR) of low complexity. The major driving force for the formation of MLOs are multivalent IDR–IDR, IDR-domain or IDR-nucleic acid interactions based on short linear binding motifs [41]. Posttranslational modifications or mutations within these binding regions can influence their physical properties, leading to a changed propensity to undergo LLPS with sometime detrimental consequences for cells. Tau represents a prominent and well characterized example [42]. Phosphorylation of Tau or the mutation of proline 301 to leucine are associated with the onset of Alzheimer’s disease (AD), Parkinson’s disease (PD) and

frontotemporal dementia with parkinsonism-17 (FTDP-17) in humans [43]. Both modifications trigger LLPS of Tau, which promotes the nucleation of Tau fibrils [44]. LLPS plays also an important role in the pathogenesis of Huntington's disease (HD), a neurodegenerative disease caused by poly-Q expansion in the exon 1 of Huntingtin. The poly-Q tract induces phase separation of Huntingtin and the resulting liquid droplets serve as nucleation centers for fibrillar Huntingtin [45].

Autophagic clearance of disease related liquid phases and aggregates

Many neurodegenerative diseases are caused or accompanied by the formation of protein aggregates. The process usually starts in a restricted area of the brain from which aggregates spread in a prion-like manner to healthy neighboring cells [46]. The high connectivity of neural tissues facilitates spreading of toxic protein fibrils, which correlates with the progression of pathology [47]. Autophagy degrades toxic protein aggregates, counteracts fibrillization and limits the spread of toxic

fibrils in neural tissue, indicating that autophagy plays an essential role in protecting neurons from accumulating toxic aggregates [48]. Accordingly, several mutations in autophagy factors are related to an increased risk of developing neurodegenerative disorders (Table 1).

Although the molecular mechanisms remain to be established, autophagic clearance appears to be most efficient early in neurodegeneration since elongated fibrils can escape autophagy by interfering with the autophagy-lysosomal pathway [49]. How autophagy counteracts fibrillization of proteins is currently unknown, but recent experimental evidence suggests that LLPS is an important factor in this process [50].

Tau, the causative agent of tauopathies, is extensively regulated by posttranslational modifications that impact on its proteasomal and autophagic clearance and its tendency to form insoluble aggregates [51,52]. For example, phosphorylation of Tau at Ser214, Ser262 and Ser305 prevents its aggregation [53], while

Table 1

Disease causing mutations in autophagy and cargo proteins. The table summarizes mutations that are directly related to the onset of neurodegenerative diseases. Data have been compiled using the Human Gene Mutation Database (Cooper et al., Institute of Medical Genetics, Cardiff University, Cardiff, UK) and The Human Variants Database Ganesan K, Kulandaisamy A, Binny Priya S, Gromiha MM (2019) HuVarBase: A human variant database with comprehensive information at gene and protein levels. PLoS ONE 14(1): e0210475.

Protein	Disease	Mutation
Tau	Frontotemporal dementia Pick's disease	R5H, E659V, L583V, G589V, L596K, P618S, S622N, K634M, V654M K574T, K686I, S637F
	Parkinson's Disease	N613del
	Alzheimer's Disease	A90V, A152T
α -Synuclein	Parkinson's Disease	A30P, H50Q, G51T, A53T, N203S, S222L, K230E, M711V, A714T, R806H, K904E, S912R
	Lewy body dementia	E46K
APP	Alzheimer's Disease	D678N, E682K, K687Q, F690_V695del, A692G, E693del, E693G, T714A, T715A, I716F, V717I, T719N, M722K, L723P
Parkin	Parkinson's Disease	V15M, R33Q, R42H, A46P, K161N, M192L, C212Y, A230V, I236V, V148E, T240R, C253Y, N254S, R256C, D280N, G284R, C289G, Q311R, G328E, T351P, R366W, A405T, T415N, G430D, C431F, C418R, C441R, R455H
Pink1	Parkinson's Disease	G30R, Q115L, A168P, A217D, A232V, H271Q, R279H, L308Q, G309D, T313M, A340T, L347P, Y431H, N521T, K526N
p62/SQSTM1	Frontotemporal dementia/ALS	A16V, D80E, V90M, R107W, V153I, R212W, G219V, S226P, P228L, P232T, K238del, D258N, S318P, D329G, S370P, A381V, T430P
	Amyotrophic Lateral Sclerosis	M1K, R12del, E20del
	Alzheimer's Disease	P29S, D260A, L268V, R393W
WIP14/WDR45	Bata-propeller protein-associated neurodegeneration	M1I, V34M, L37S, N61K, N84G, L98P, F100S, L160R, deletions at various positions
OPTN	Amyotrophic Lateral Sclerosis	H3Y, P16A, S73L, R83C, A93P, A136V, V161M, K258N, T282P, V295P, Q314L, I451T, N478G, L494W, L500P, L557T, D564H, T567A, L568S, H571N
TBK1	Amyotrophic Lateral Sclerosis	T4A, V23I, G26D, T31A, R47H, L94S, D118N, G121D, R143C, I207 T/V, G217R, G244V, I246T, R308N, T320I, H322R, R358 C/H/W, K401D, Y424E, R444N, M559R, M598V, M662T, R724C
	Frontotemporal dementia Alzheimer's Disease (late)	T4A, G26D, R143C, G244V, T320I, I418V, M662T D534H

phosphorylation at positions Ser202, Thr205 and Ser208 inhibits the Ub/proteasome system and promotes aggregation of Tau [54]. Moreover, hyperphosphorylated Tau has an increased tendency to phase separate and is hyperubiquitinated by the E3-ligase CHIP. Both mechanisms promote the formation of Tau fibrils, the building blocks for neurofibrillary tangles [55]. However, over-expressing CHIP in a Tau-BiFC mouse model, in which the progression of Tau oligomers can be spatiotemporally visualized based on the fluorescent protein [56], leads to a reduction of oligomeric Tau and phosphorylated Tau species in an autophagy dependent manner [57]. Since polyubiquitinated cargo stimulates LLPS of p62 [12] one could speculate that hyperphosphorylated and hyperubiquitinated Tau aggregates are included in p62 bodies to be degraded by selective autophagy. To what extent the formation of co-condensates influence the fate of Tau, notably whether Tau is degraded by autophagy or forms larger Tau-tangles, remains to be investigated.

Box 1. The core autophagy machinery in selective autophagy.

Local activation of the ULK kinase complex (consisting of ULK1/2, FIP200, ATG13 and ATG1010) is the earliest event upon initiation of autophagy by various specific and pathway dependent stimuli. ULK1 kinase activity is suppressed through its phosphorylation at Ser757 by mTOR1. Thus, inactivation of mTOR1, for example upon nitrogen starvation, leads to activation of ULK1, while glucose starvation activates AMPK, which activates ULK1 by phosphorylating Ser317 and Ser777 [58]. During selective autophagy, ULK1 needs to be activated although its activity is constantly suppressed by mTOR1. This is facilitated by cargo-bound autophagy receptors, which recruit the ULK1 complex through a direct interaction with its subunit FIP200 [27,30,59]. The ULK1 complex further promotes the recruitment of ATG9 vesicles and the PtdIns-3 kinase complex to cargo, leading to the nucleation of the phagophore membrane. ATG9 functions as a lipid scramblase to promote the translocation of phospholipids from the cytoplasmic to the luminal leaflet of the expanding phagophore membrane [33] and together with the lipid transfer protein ATG2 [60,61], being recruited to ER-associated membranes, it provides lipids and proteins needed for *de novo* autophagosome biogenesis [62]. Following its recruitment by the ULK1 kinase complex subunits ATG13 and ATG101, the PtdIns-3 kinase complex enriches the phagophore with PtdIns-3-phosphate which is recognized by the WD-domain containing protein WIPI2 [63]. The latter initiates the lipidation of ATG8 proteins to the phagophore by recruiting ATG16L1 [64]. The conjugation of ATG8 proteins to lipids is catalyzed by an autophagy specific Ub-like conjugation system, comprising the E1-like enzyme ATG7, E2-like ATG3 and the E3-like ligase ATG12–ATG5–ATG16L1 [65]. Based on their sequence, ATG8 proteins can be grouped into the GABARAP (Gamma-aminobutyric acid receptor-associated protein) and L3C (Microtubule-associated proteins 1A/1B light chain 3) subfamilies. ATG8 proteins fulfil various essential functions in autophagy. They tether cargo to phagophores by binding cargo receptors, promote the maturation of phagophores and coordinate fusion of autophagosomes with lysosomes [5,7,11], but are dispensable for some types of selective autophagy, including the autophagic degradation of mitochondria by mitophagy [19,66]. The final closure of the phagophore membrane is facilitated by the endosomal sorting complexes required for transport (ESCRT) machinery [67].

Conclusion

The application of physical methods and concepts in cell biology has increased our understanding of fundamental biological processes tremendously. This is particularly true for the field of membrane trafficking and autophagy, where many phenomena and observations could only be explained by combining traditional cell biological and physical techniques. Liquid phase separation is a physical process that all of us know well from daily experience. However, its contribution to structure the complex cellular environment just begin to unfold. In contrast to membrane bound organelles, phase separated structures exchange components with the cytoplasm in a highly dynamic and much less constrained manner, providing another level of subcellular organization and regulation. Several processes in autophagy, including cargo sequestration as well as initiation and expansion of phagophores, involve LLPS. Understanding the consequences of perturbed phase separation or changes in the composition of phase separated structures in autophagy will be key to reveal insights into the origin of many human pathologies including neurodegeneration. Future studies of dynamic and structural aspects of phase separation in autophagy bear the potential to develop new therapeutical concepts to treat neurodegeneration and other human diseases.

Conflict of interest statement

Nothing declared.

Acknowledgement

We would like to thank Laura Rodriguez de la Ballina for help with the figure. This work was supported by the Research Council of Norway through its Centres of Excellence funding scheme (Project: 262652) and FRIPRO grant (Project: 314684) to A.S. and funding of the Institut Pasteur to T.W.

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
 - ** of outstanding interest
1. Lamark T, Johansen T: **Mechanisms of selective autophagy.** *Annu Rev Cell Dev Biol* 2021, **37**:1–27.
 2. Mizushima N, White E, Rubinsztein DC: **Breakthroughs and bottlenecks in autophagy research.** *Trends Mol Med* 2021, <https://doi.org/10.1016/j.molmed.2021.06.012>.
 3. Conway O, Akpinar HA, Rogov VV, Kirkin V: **Selective autophagy receptors in neuronal health and disease.** *J Mol Biol* 2020, **432**:2483–2509.
 4. Noda NN, Wang Z, Zhang H: **Liquid–liquid phase separation in autophagy.** *J Cell Biol* 2020, **219**.
 5. Kirkin V, Rogov VV: **A diversity of selective autophagy receptors determines the specificity of the autophagy pathway.** *Mol Cell* 2019, **76**:268–285.
 6. Stolz A, Ernst A, Dikic I: **Cargo recognition and trafficking in selective autophagy.** *Nat Cell Biol* 2014, **16**:495–501.

7. Wirth M, Zhang W, Razi M, Nyoni L, Joshi D, O'Reilly N, Johansen T, Tooze SA, Mouilleron S: **Molecular determinants regulating selective binding of autophagy adapters and receptors to ATG8 proteins.** *Nat Commun* 2019, **10**:2055.
- This study reveals the structural bases for the selective interaction of LIRs with GABARAPs by exploring the contribution and specificity of residues in the LIR binding region of GABARAPs and in LIRs of interaction partners in a systematic fashion.
8. Wirth M, Mouilleron S, Zhang W, Sjøttem E, Abudu YP, Jain A, Olsvik HL, Bruun J-A, Razi M, Jefferies HBJ, et al.: **Phosphorylation of the LIR domain of SCOC modulates ATG8 binding affinity and specificity.** *J Mol Biol* 2021, **433**:166987.
9. Poole LP, Bock-Hughes A, Berardi DE, Macleod KF: **ULK1 promotes mitophagy via phosphorylation and stabilization of BNIP3.** *Sci Rep* 2021, **11**:20526.
10. Chang C, Shi X, Jensen LE, Yokom AL, Fracchiolla D, Martens S, Hurley JH: **Reconstitution of cargo-induced LC3 lipidation in mammalian selective autophagy.** *Sci Adv* 2021, **7**, eabg4922.
11. Zellner S, Schifferer M, Behrends C: **Systematically defining selective autophagy receptor-specific cargo using autophagosome content profiling.** *Mol Cell* 2021, **81**:1337–1354.e8.
- This study combined protease protection and APEX2 proximity proteomics to reveal the selectivity of autophagy receptors for certain cargos under basal and aggrephagy inducing conditions.
12. Jakobi AJ, Huber ST, Mortensen SA, Schultz SW, Palara A, Kuhm T, Shrestha BK, Lamark T, Hagen WJH, Wilmanns M, et al.: **Structural basis of p62/SQSTM1 helical filaments and their role in cellular cargo uptake.** *Nat Commun* 2020, **11**:440.
13. Lamark T, Perander M, Outzen H, Kristiansen K, Øvervatn A, Michaelsen E, Bjørkøy G, Johansen T: **Interaction codes within the family of mammalian Phox and Bem1p domain-containing proteins*.** *J Biol Chem* 2003, **278**:34568–34581.
14. Ciuffa R, Lamark T, Tarafder AK, Guesdon A, Rybina S, Hagen WJH, Johansen T, Sachse C: **The selective autophagy receptor p62 forms a flexible filamentous helical scaffold.** *Cell Rep* 2015, **11**:748–758.
15. Berkamp S, Mostafavi S, Sachse C: **Structure and function of p62/SQSTM1 in the emerging framework of phase separation.** *FEBS J* 2021, <https://doi.org/10.1111/febs.15672>.
16. Sun D, Wu R, Zheng J, Li P, Yu L: **Polyubiquitin chain-induced p62 phase separation drives autophagic cargo segregation.** *Cell Res* 2018, **28**:405–415.
17. Zaffagnini G, Savova A, Danieli A, Romanov J, Tremel S, Ebner M, Peterbauer T, Sztacho M, Trapannone R, Tarafder AK, et al.: **p62 filaments capture and present ubiquitinated cargos for autophagy.** *EMBO J* 2018, **37**, e98308.
18. Turco E, Savova A, Gere F, Ferrari L, Romanov J, Schuschnig M, Martens S: **Reconstitution defines the roles of p62, NBR1 and TAX1BP1 in ubiquitin condensate formation and autophagy initiation.** *Nat Commun* 2021, **12**:5212.
- This study uses an in vitro reconstitution system in combination with cell biology to dissect the roles of p62, NBR1 and TAX1BP1 in the formation and clearance of ubiquitin condensates.
19. Ohnstad AE, Delgado JM, North BJ, Nasa I, Kettenbach AN, Schultz SW, Shoemaker CJ: **Receptor-mediated clustering of FIP200 bypasses the role of LC3 lipidation in autophagy.** *EMBO J* 2020, <https://doi.org/10.15252/embj.2020104948>.
20. Kageyama S, Gudmundsson SR, Sou Y-S, Ichimura Y, Tamura N, Kazuno S, Ueno T, Miura Y, Noshiro D, Abe M, et al.: **p62/SQSTM1-droplet serves as a platform for autophagosome formation and anti-oxidative stress response.** *Nat Commun* 2021, **12**:16.
- This study shows that p62-bodies serves as platforms to initiate the anti-oxidative stress responses. The p62- and Nrf2-interacting protein Keap1 translocates to p62-bodies upon oxidative stress, leading to activation of Nrf2 and transcription of genes important the anti-oxidative response. Impaired clearance of p62-bodies leads to Nrf2 hyper-activation *in vivo* and liver injury.
21. Schlütermann D, Berleth N, Deitersen J, Wallot-Hieke N, Friesen O, Wu W, Stuhldreier F, Sun Y, Berning L, Friedrich A, et al.: **FIP200 controls the TBK1 activation threshold at SQSTM1/p62-positive condensates.** *Sci Rep* 2021, **11**:13863.
22. Carroll B, Otten EG, Manni D, Stefanatos R, Menzies FM, Smith GR, Jurk D, Kenneth N, Wilkinson S, Passos JF, et al.: **Oxidation of SQSTM1/p62 mediates the link between redox state and protein homeostasis.** *Nat Commun* 2018, **9**:256.
23. Yoon MJ, Choi B, Kim EJ, Ohk J, Yang C, Choi Y-G, Lee J, Kang C, Song HK, Kim YK, et al.: **UXT chaperone prevents proteotoxicity by acting as an autophagy adaptor for p62-dependent aggrephagy.** *Nat Commun* 2021, **12**:1955.
- This study identify the prefoldin-like chaperone UXT as an interacting protein of p62 and shows that UXT increases p62 clustering and p62 body formation and clearance. Importantly, UXT expression delays motor neuron degeneration in a SOD1(A4V)-induced model, demonstrating a relationship between molecular chaperones and the aggrephagy machinery.
24. Sánchez-Martín P, Sou Y, Kageyama S, Koike M, Waguri S, Komatsu M: **NBR1-mediated p62-liquid droplets enhance the Keap1-Nrf2 system.** *EMBO Rep* 2020, **21**, e48902.
25. Sarraf SA, Shah HV, Kanfer G, Pickrell AM, Holtzclaw LA, Ward ME, Youle RJ: **Loss of TAX1BP1-directed autophagy results in protein aggregate accumulation in the brain.** *Mol Cell* 2020, **80**:779–795.e10.
- This study provides a physical model demonstrating that autophagosome-like membrane sheets form at the surface of p62-containing droplets in vitro through partial wetting. The model predicts that bending of such sheets can divide droplets for piecemeal sequestration or sequester entire droplets.
26. Agudo-Canalejo J, Schultz SW, Chino H, Migliano SM, Saito C, Koyama-Honda I, Stenmark H, Brech A, May AI, Mizushima N, et al.: **Wetting regulates autophagy of phase-separated compartments and the cytosol.** *Nature* 2021, **591**:142–146.
27. Vargas JNS, Wang C, Bunker E, Hao L, Maric D, Schiavo G, Randow F, Youle RJ: **Spatiotemporal control of ULK1 activation by NDP52 and TBK1 during selective autophagy.** *Mol Cell* 2019, **74**:347–362.e6.
28. Shi X, Chang C, Yokom AL, Jensen LE, Hurley JH: **The autophagy adaptor NDP52 and the FIP200 coiled-coil allosterically activate ULK1 complex membrane recruitment.** *Elife* 2020, **9**, e59099.
- This study uses negative stain electron microscopy (NSEM) to characterize the structure of full-length FIP200 and hydrogen–deuterium exchange mass spectrometry analysis (HDX-MS) to elucidate how the ULK1 complex interacts with FIP200 in the absence or presence of the cargo receptor NDP52. Using GUVs they further investigate how NDP52 affects membrane recruitment of the ULK1 complex.
29. Fu T, Zhang M, Zhou Z, Wu P, Peng C, Wang Y, Gong X, Li Y, Wang Y, Xu X, et al.: **Structural and biochemical advances on the recruitment of the autophagy-initiating ULK and TBK1 complexes by autophagy receptor NDP52.** *Sci Adv* 2021, **7**, eabi6582.
30. Turco E, Witt M, Abert C, Bock-Bierbaum T, Su M-Y, Trapannone R, Sztacho M, Danieli A, Shi X, Zaffagnini G, et al.: **FIP200 claw domain binding to p62 promotes autophagosome formation at ubiquitin condensates.** *Mol Cell* 2019, **74**.
31. Padman BS, Nguyen TN, Uoselis L, Skulsupaisarn M, Nguyen LK, Lazarou M: **LC3/GABARAPs drive ubiquitin-independent recruitment of Optineurin and NDP52 to amplify mitophagy.** *Nat Commun* 2019, **10**:408.
32. Yamano K, Kikuchi R, Kojima W, Hayashida R, Koyano F, Kawawaki J, Shoda T, Demizu Y, Naito M, Tanaka K, et al.: **Critical role of mitochondrial ubiquitination and the OPTN–ATG9A axis in mitophagy.** *J Cell Biol* 2020, **219**, e201912144.
33. Matoba K, Kotani T, Tsutsumi A, Tsuji T, Mori T, Noshiro D, Sugita Y, Nomura N, Iwata S, Ohsumi Y, et al.: **Atg9 is a lipid scramblase that mediates autophagosomal membrane expansion.** *Nat Struct Mol Biol* 2020, **27**:1185–1193.
34. Harding O, Evans CS, Ye J, Cheung J, Maniatis T, Holzbaur ELF: **ALS- and FTD-associated missense mutations in TBK1 differentially disrupt mitophagy.** *Proc Natl Acad Sci Unit States Am* 2021, **118**, e2025053118.
35. Shi X, Yokom AL, Wang C, Young LN, Youle RJ, Hurley JH: **ULK complex organization in autophagy by a C-shaped FIP200 N-terminal domain dimer.** *J Cell Biol* 2020, **219**, e201911047.

36. Abudu YP, Pankiv S, Mathai BJ, Lystad AH, Bindsbøll C, Brenne HB, Ng MYW, Thiede B, Yamamoto A, Nthiga TM, *et al.*: **NIPSNAP1 and NIPSNAP2 act as “eat me” signals for mitophagy**. *Dev Cell* 2019, **49**:509–525.e12.
37. Ray S, Singh N, Kumar R, Patel K, Pandey S, Datta D, Mahato J, Panigrahi R, Navalkar A, Mehra S, *et al.*: **α -Synuclein aggregation nucleates through liquid–liquid phase separation**. *Nat Chem* 2020, **12**:705–716.
- This study demonstrate that phase separated condensates of α -Synuclein are nucleation centers for the *de novo* generation of disease-related α -Synuclein fibrils.
38. Fare CM, Villani A, Drake LE, Shorter J: **Higher-order organization of biomolecular condensates**. *Open Biol* 2021, **11**:210137.
39. Alberti S, Hyman AA: **Biomolecular condensates at the nexus of cellular stress, protein aggregation disease and ageing**. *Nat Rev Mol Cell Biol* 2021, **22**:196–213.
40. Ryan VH, Fawzi NL: **Physiological, pathological, and targetable membraneless organelles in neurons**. *Trends Neurosci* 2019, **42**:693–708.
41. Riback JA, Zhu L, Ferrolino MC, Tolbert M, Mitrea DM, Sanders DW, Wei M-T, Kriwacki RW, Brangwynne CP: **Composition-dependent thermodynamics of intracellular phase separation**. *Nature* 2020, **581**:209–214.
- This study revealed the thermodynamic basis of multi-component LLPS in human cells and established the physical basis for assembly and vectorial transport of ribonucleoprotein complexes.
42. Ambadipudi S, Biernat J, Riedel D, Mandelkow E, Zweckstetter M: **Liquid-liquid phase separation of the microtubule-binding repeats of the Alzheimer-related protein Tau**. *Nat Commun* 2017, **8**:275.
43. Lewis J, McGowan E, Rockwood J, Melrose H, Nacharaju P, Sleightenhorst MV, Gwinn-Hardy K, Murphy MP, Baker M, Yu X, *et al.*: **Neurofibrillary tangles, amyotrophy and progressive motor disturbance in mice expressing mutant (P301L) Tau protein**. *Nat Genet* 2000, **25**:402–405.
44. Savastano A, Flores D, Kadavath H, Biernat J, Mandelkow E, Zweckstetter M: **Disease-associated tau phosphorylation hinders tubulin assembly within tau condensates**. *Angew Chem Int Ed* 2021, **60**:726–730.
45. Peskett TR, Rau F, O’Driscoll J, Patani R, Lowe AR, Saibil HR: **A liquid to solid phase transition underlying pathological Huntingtin Exon1 aggregation**. *Mol Cell* 2018, **70**:588–601.e6.
46. Chastagner P, Loria F, Vargas JY, Tois J, Diamond MI, Okafo G, Brou C, Zurzolo C: **Fate and propagation of endogenously formed Tau aggregates in neuronal cells**. *EMBO Mol Med* 2020, **12**, e12025.
47. Cornblath EJ, Li HL, Changolkar L, Zhang B, Brown HJ, Gathagan RJ, Olufemi MF, Trojanowski JQ, Bassett DS, Lee VMY, *et al.*: **Computational modeling of Tau pathology spread reveals patterns of regional vulnerability and the impact of a genetic risk factor**. *Sci Adv* 2021, **7**, eabg6677.
48. Gao J, Perera G, Bhadbhade M, Halliday GM, Dzamko N: **Autophagy activation promotes clearance of α -synuclein inclusions in fibril-seeded human neural cells**. *J Biol Chem* 2019, **294**:14241–14256.
49. Riera-Tur I, Schäfer T, Hornburg D, Mishra A, Padilha M da S, Fernández-Mosquera L, Feigenbutz D, Auer P, Mann M, Baumeister W, *et al.*: **Amyloid-like aggregating proteins cause lysosomal defects in neurons via gain-of-function toxicity**. *Life Sci Alliance* 2022, **5**, e202101185.
50. Silva JM, Rodrigues S, Sampaio-Marques B, Gomes P, Neves-Carvalho A, Dioli C, Soares-Cunha C, Mazuik BF, Takashima A, Ludovico P, *et al.*: **Dysregulation of autophagy and stress granule-related proteins in stress-driven Tau pathology**. *Cell Death Differ* 2019, **26**:1411–1427.
51. Arakhamia T, Lee CE, Carlomagno Y, Duong DM, Kundinger SR, Wang K, Williams D, DeTure M, Dickson DW, Cook CN, *et al.*: **Posttranslational modifications mediate the structural diversity of tauopathy strains**. *Cell* 2020, **180**:633–644.e12.
- This study established structural differences in Tau fibrils form brains of patients with Alzheimer’s disease and corticobasal degeneration depending on posttranslational modifications using a combination of proteomics and cryo-electron microscopy.
52. Wesseling H, Mair W, Kumar M, Schlaffner CN, Tang S, Beerepoot P, Fatou B, Guise AJ, Cheng L, Takeda S, *et al.*: **Tau PTM profiles identify patient heterogeneity and stages of Alzheimer’s disease**. *Cell* 2020, **183**:1699–1713.e13.
53. Strang KH, Sorrentino ZA, Riffe CJ, Gorion K-MM, Vijayaraghavan N, Golde TE, Giasson BI: **Phosphorylation of serine 305 in Tau inhibits aggregation**. *Neurosci Lett* 2019, **692**:187–192.
54. Xia Y, Prokop S, Gorion K-MM, Kim JD, Sorrentino ZA, Bell BM, Manaolis AN, Chakrabarty P, Davies P, Giasson BI: **Tau Ser208 phosphorylation promotes aggregation and reveals neuro-pathology diversity in Alzheimer’s disease and other tauopathies**. *Acta Neuropathologica Commun* 2020, **8**:88.
55. Kim JH, Lee J, Choi WH, Park S, Park SH, Lee JH, Lim SM, Mun JY, Cho H-S, Han D, *et al.*: **CHIP-mediated hyper-ubiquitylation of Tau promotes its self-assembly into the insoluble Tau filaments**. *Chem Sci* 2021, **12**:5599–5610.
56. Shin S, Kim D, Song JY, Jeong H, Hyeon SJ, Kowall NW, Ryu H, Pae AN, Lim S, Kim YK: **Visualization of soluble Tau oligomers in TauP301L-BiFC transgenic mice demonstrates the progression of tauopathy**. *Prog Neurobiol* 2020, **187**:101782.
57. Subramanian M, Hyeon SJ, Das T, Suh YS, Kim YK, Lee J-S, Song EJ, Ryu H, Yu K: **UBE4B, a microRNA-9 target gene, promotes autophagy-mediated Tau degradation**. *Nat Commun* 2021, **12**:3291.
58. Kim J, Kundu M, Viollet B, Guan K-L: **AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1**. *Nat Cell Biol* 2011, **13**, 1–22.
59. Ravenhill BJ, Boyle KB, Muhlinen N von, Ellison CJ, Masson GR, Otten EG, Foeglein A, Williams R, Rando F: **The cargo receptor NDP52 initiates selective autophagy by recruiting the ULK complex to cytosol-invading bacteria**. *Mol Cell* 2019, **74**:320–329.e6.
60. Maeda S, Otomo C, Otomo T: **The autophagic membrane tether ATG2A transfers lipids between membranes**. *Elife* 2019, **8**, e45777.
61. Osawa T, Kotani T, Kawaoka T, Hirata E, Suzuki K, Nakatogawa H, Ohsumi Y, Noda NN: **Atg2 mediates direct lipid transfer between membranes for autophagosome formation**. *Nat Struct Mol Biol* 2019, **26**:281–288.
62. Judith D, Jefferies HBJ, Boeing S, Frith D, Sniijders AP, Tooze SA: **ATG9A shapes the forming autophagosome through Arfaptin 2 and phosphatidylinositol 4-kinase III β** . *J Cell Biol* 2019, **114**, jcb.201901115.
63. Nishimura T, Tooze SA: **Emerging roles of ATG proteins and membrane lipids in autophagosome formation**. *Cell Discov* 2020, **6**:32.
64. Dooley HC, Razi M, Polson HEJ, Girardin SE, Wilson MI, Tooze SA: **WIPI2 links LC3 conjugation with PI3P, autophagosome formation, and pathogen clearance by recruiting Atg12–5–16L1**. *Mol Cell* 2014, **55**:238–252.
65. Mizushima N: **The ATG conjugation systems in autophagy**. *Curr Opin Cell Biol* 2020, **63**:1–10.
66. Nguyen TN, Padman BS, Zellner S, Khuu G, Uoselis L, Lam WK, Skulsuppaisarn M, Lindblom RJS, Watts EM, Behrends C, *et al.*: **ATG4 family proteins drive phagophore growth independently of the LC3/GABARAP lipidation system**. *Mol Cell* 2021, **81**:2013–2030.e9.
67. Zhou F, Wu Z, Zhao M, Murtazina R, Cai J, Zhang A, Li R, Sun D, Li W, Zhao L, *et al.*: **Rab5-dependent autophagosome closure by ESCRT**. *J Cell Biol* 2019, **111**, jcb.201811173.
- This study revealed that ESCRT-III proteins catalyze the closure of phagophores to generate fully matured and sealed autophagosomes that are competent to fuse with lysosomal compartments.