



HAL
open science

Challenges in the Biotechnological Implementation of Oral RNA Interference as an Antiviral Strategy in *Aedes aegypti*

Ottavia Romoli, Annabelle Henrion-Lacritick, Hervé Blanc, Lionel Frangeul, Maria-Carla Saleh

► **To cite this version:**

Ottavia Romoli, Annabelle Henrion-Lacritick, Hervé Blanc, Lionel Frangeul, Maria-Carla Saleh. Challenges in the Biotechnological Implementation of Oral RNA Interference as an Antiviral Strategy in *Aedes aegypti*. 2024. pasteur-04376142

HAL Id: pasteur-04376142

<https://pasteur.hal.science/pasteur-04376142>

Preprint submitted on 6 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 Challenges in the Biotechnological Implementation of Oral RNA Interference as an 2 Antiviral Strategy in *Aedes aegypti*

3 Authors:

4 Ottavia Romoli^{1,*}, Annabelle Henrion-Lacritick¹, Hervé Blanc¹, Lionel Frangeul¹ and Maria-Carla
5 Saleh^{1,*}

6 Affiliation:

7 ¹ Institut Pasteur, Université Paris Cité, CNRS UMR3569, Viruses and RNAi Unit, F-75015 Paris, France.

8 * correspondence to: Ottavia Romoli (ottavia.romoli@pasteur.fr) and Maria-Carla Saleh
9 (carla.saleh@pasteur.fr)

10 Abstract

11 Mosquitoes, particularly *Aedes aegypti*, are critical vectors for globally significant pathogenic
12 viruses. This study examines the limitations of oral RNA interference (RNAi) as a strategy to disrupt
13 viral transmission by *Ae. aegypti*. We hypothesized that double-stranded RNA (dsRNA) targeting the
14 Zika virus (ZIKV) or chikungunya virus (CHIKV) genomes produced by engineered bacterial symbionts
15 could trigger an antiviral response. Mosquitoes mono-colonized with *Escherichia coli* producing
16 dsZIK or dsCHIK did not display reduced viral titers following exposure to virus-contaminated
17 bloodmeals and failed to generate dsZIK- or dsCHIK-derived small interfering RNAs. To address
18 potential limitations of bacterial dsRNA release, we explored dsRNA inoculation via feeding and
19 injection. While viral replication was impeded in mosquitoes injected with dsZIK or dsCHIK, no
20 antiviral effect was observed in dsRNA-fed mosquitoes. These findings highlight complexities of
21 implementing oral RNAi as an antiviral strategy in *Ae. aegypti* and warrant further exploration of
22 local and systemic RNAi mechanisms.

23 Keywords

24 *Aedes aegypti*, arbovirus, oral RNA interference, small RNA, paratransgenesis

25

26 Introduction

27 Mosquitoes are vectors for several arthropod-borne viruses (or arboviruses), which are responsible
28 for multiple outbreaks worldwide. Over the past decade, the emergence or re-emergence of viruses
29 such as dengue (DENV), Zika (ZIKV), or chikungunya (CHIKV) viruses, together with the current lack of
30 vaccines for most of these viruses and the emergence of insecticide resistant mosquito populations
31 has necessitated the development of new strategies to control the spread of these pathogens and
32 their vectors.

33 Mosquito vector competence (*i.e.*, the ability of a mosquito to acquire and transmit a virus) is
34 dependent on efficient viral replication and dissemination from the gut to the salivary glands, where
35 the virus must enter the saliva to be transmitted to the next vertebrate host during subsequent blood-
36 feeding. One of the mechanisms that mosquitoes use to control viral replication is RNA interference
37 (RNAi), which is triggered by viral double-stranded RNA (dsRNA) intermediates produced during RNA
38 virus replication. The RNase Dicer-2 binds dsRNAs and processes them into 21 nucleotide (nt) small-
39 interfering RNAs (siRNAs)¹. These are used by Argonaute-2 as sequence-specific guides to target
40 complementary viral RNAs, thereby hindering viral replication²⁻⁵. Key studies in *Aedes aegypti*
41 mosquitoes showed that a protective RNAi-mediated antiviral response could be triggered by
42 exposure to dsRNA molecules corresponding to portions of the DENV or ZIKV genomes either by
43 transgenic production of dsDENV or by injection of dsZIKV^{6,7}, validating the potential use of exogenous
44 dsRNA molecules to limit arbovirus infection.

45 During viral infection through a bloodmeal, the mosquito midgut represents the first mechanical
46 and immunological barrier encountered by the virus and crossing this barrier is critical to establish
47 an infection in the mosquito. Thus, we reasoned that blocking viral replication in the midgut could
48 be an effective way to limit viral transmission and we hypothesized that bacterial symbionts in the
49 mosquito gut could be used as a platform to produce and deliver dsRNA to mosquitoes and
50 stimulate a prophylactic RNAi-based antiviral response. Indeed, a recent study in honeybees showed
51 the possibility of reducing viral replication through colonization with an antiviral dsRNA-producing
52 symbiont⁸. When it comes to mosquitoes, numerous studies have demonstrated the potential for
53 gene silencing by introducing dsRNA through feeding or using microorganisms that produce dsRNA⁹⁻
54 ¹⁴. While a handful of studies have pointed out challenges like low efficiency and variability in oral
55 RNA interference in mosquitoes^{15,16}, it's noteworthy that none of these investigations have explored
56 the concept of microbiota-mediated oral delivery of antiviral dsRNA.

57 Paratransgenesis is the genetic modification of a symbiotic microorganism that is associated with a
58 disease vector with the aim of disrupting the transmission of pathogens carried by the vector. This
59 approach offers the advantage of potential self-sustainability, as the targeted microorganism is
60 typically selected to undergo effective horizontal (between individuals) and vertical (from parents
61 to progeny) transmission¹⁷. Various paratransgenic methods have been proposed to reduce
62 *Plasmodium* transmission by *Anopheles* mosquitoes, including the use symbiotic bacteria
63 engineered to produce anti-*Plasmodium* molecules or mosquito immune factors¹⁸⁻²⁰. Additionally,
64 other paratransgenic strategies have focused on interfering with mosquito development or

65 fertility^{10,12,21}. However, there are currently no developed strategies to impede virus transmission
66 by *Aedes* mosquitoes through manipulation of their microbiota.

67 Here, we investigated the potential targeted induction of antiviral RNAi against ZIKV and CHIKV in
68 *Ae. aegypti* mosquitoes through colonization with bacteria capable of producing antiviral dsRNA.
69 We used *Escherichia coli* as a proof-of-concept and found that mosquitoes colonized by dsZIK- and
70 dsCHIKV-expressing bacteria did not exhibit reduced viral titers when infected with the
71 corresponding virus through a bloodmeal. Additionally, there was no discernible production of
72 siRNAs induced by the ingestion of bacteria-produced dsRNA. When we fed mosquitoes with naked
73 dsRNA, we did not observe enhanced antiviral immunity, likely due to the absence of an effective
74 antiviral RNAi-based response in the midgut and insufficient activation of systemic RNAi.

75 **Results**

76 *Colonization of mosquitoes with dsRNA-producing Escherichia coli*

77 To generate *E. coli* strains expressing antiviral dsRNA, we took advantage of the HT115 strain, which
78 has historically been used in *Caenorhabditis elegans* to induce RNAi. This strain is suitable for dsRNA
79 production as it possesses an isopropylthio- β -galactoside (IPTG)-inducible *T7 polymerase* gene and
80 has a mutation in the *rnc* allele encoding RNase III, resulting in reduced dsRNA degradation²². This
81 allows the synthesis of the target sequence by transcription from a DNA template containing the T7
82 promoter. To induce the production of dsRNA for ZIKV or CHIKV, we cloned the target viral
83 sequences between two directionally opposed T7 promoters and terminators in the T444T plasmid
84 to produce dsRNA by convergent transcription²³ (**Figure 1A, Supplementary Figure 1A**). To test
85 whether a unidirectionally-transcribed RNA with a hairpin secondary structure would result in a
86 more stable and effective dsRNA production, an inverted repeat of the CHIKV target sequence was
87 cloned in a pUC18 plasmid downstream from a T7 promoter. The inverted repeat consisted of the
88 CHIKV target sequence in both polarities separated by a non-viral spacer sequence (**Supplementary**
89 **Figure 1B**). The target viral sequences were selected for dsRNA production based on the following
90 criteria: (i) we excluded sequences that could produce 21mers with high similarity to any sequence
91 in the *Ae. aegypti* genome in order to avoid potential interference of endogenous gene expression;
92 (ii) we targeted regions corresponding to only partial coding regions of viral structural genes to
93 prevent translation of the target sequences into functional viral proteins; (iii) we avoided sequences
94 corresponding to noncoding subgenomic flaviviral RNA or viral proteins shown to repress mosquito
95 immunity^{24–26} (**Figure 1B, Supplementary Figure 1C**). A T444T plasmid containing a portion of the
96 GFP coding sequence instead of a target viral sequence was produced as a control.

97 To confirm the production of dsRNA by bacteria, RNA was extracted from overnight cultures of *E.*
98 *coli* HT115-T444T-ZIK1 induced or not with IPTG for 24 h. Northern blot with ZIK1 sense and
99 antisense probes was performed on bacterial total RNA and on dsZIK1 synthesized *in vitro* as a
100 positive control. The agarose gel showed a major 480 nt band for *in vitro* synthesized dsZIK1,
101 although a higher ~900 nt band was visible (**Supplementary Figure 2A**). In the corresponding
102 northern blot, ZIK1 sense and antisense probes bound to both 480 and ~900 nt bands
103 (**Supplementary Figure 2B**), suggesting that the larger band is composed by dsRNA multimers or

104 higher order assemblies. These molecules are commonly found in both *in vitro* and *in vivo*
105 synthesized dsRNA with unknown effect on RNAi activity^{27,28}. A signal at ~350 nt was detected by
106 sense and antisense probes in uninduced bacteria (**Supplementary Figure 2B**). This might represent
107 a truncated RNA transcribed by the T7 RNA Polymerase, whose transcription is directed by *lac*
108 operon control elements. This system is commonly known to be leaky and to show basal
109 transcription in absence of IPTG induction, especially in stationary phase cultures^{29,30}. Bacteria
110 induced with IPTG showed a major ~650 nt band in the agarose gel (**Supplementary Figure 2A**),
111 which was recognized by both ZIK1 probes together with other transcription products spanning
112 from ~250 to ~900 nt (**Supplementary Figure 2B**). The major band at ~650 nt represents the main
113 ZIK1 transcript from the T444T-ZIK1 plasmid, which is 186 nt longer than the *in vitro* transcript due
114 to additional plasmid sequences flanking the *NotI* and *AgeI* cloning sites. The ZIK1 sense probe
115 showed high specificity for the target ZIKV sequence, while no significant signal was visible in a
116 northern blot of total RNA extracted from overnight cultures of *E. coli* HT115-T444T-CHIK1
117 (**Supplementary Figure 2C-D**). Altogether the northern blot analysis revealed abundant production
118 of dsRNA by the modified bacteria.

119 To test the mosquito colonization properties of *E. coli* HT115, the transstadial transmissibility of *E.*
120 *coli* HT115 (from larvae to adults), and the stability of the T444T plasmid *in vivo* in the absence of
121 antibiotic selection, we produced *Ae. aegypti* larvae mono-colonized by *E. coli* HT115-T444T-ZIK1
122 (**Figure 1C**). Mono-colonization at the larval stage was used to maximize the exposure of mosquitoes
123 to dsRNA. *E. coli* HT115 was added to axenic larvae obtained from surface sterilized eggs³¹. The
124 HT115 strain was able to support standard larval development, as previously described for other *E.*
125 *coli* strains^{31,32}. To assess the transstadial transmission of *E. coli* HT115, we measured bacterial loads
126 in adult mosquitoes originating from mono-colonized larvae kept in non-sterile conditions after
127 adult emergence. When mosquitoes were provided only with sterile sucrose, *E. coli* HT115 carrying
128 the T444T-ZIK1 plasmid could not be isolated from whole mosquitoes collected one or four days
129 post-emergence, showing the low transstadial transmissibility of this strain (**Figure 1D**). When
130 mosquitoes were provided with a sucrose solution contaminated with *E. coli* HT115 carrying the
131 T444T-ZIK1 plasmid, bacteria could be isolated from whole mosquitoes at both tested time points,
132 with 100 % of mosquitoes colonized four days post-emergence. Bacterial loads were comparable
133 when measured on antibiotic-free plates or plates containing ampicillin to select for bacteria
134 carrying the T444T-ZIK1 plasmid (Wilcoxon test: sterile sugar day 1, $p = \text{NA}$; sterile sugar day 4, $p =$
135 0.38; bacteria supplemented day 1, $p = 1$; bacteria supplemented day 4, $p = 0.71$), indicating that *E.*
136 *coli* HT115 could stably colonize adult mosquitoes, it represented most of the bacteria present at
137 the adult stage, and it could maintain the plasmid *in vivo* for at least four days.

138 *Mosquitoes colonized with dsRNA-producing bacteria do not show reduced viral titers or production*
139 *of dsRNA-specific siRNAs*

140 To assess the effect of the bacterial-produced dsRNA on viral infections, we exposed mosquitoes to
141 a ZIKV- or CHIKV-containing infectious bloodmeal. We reasoned that mosquitoes mono-colonized
142 at the larval stage with the dsRNA-producing bacteria would be the best model to work with, as
143 they were exposed to the dsRNA during their entire development. We produced larvae mono-

144 colonized with *E. coli* HT115 carrying the T444T-GFP, T444T-ZIK1, T444T-CHIK1, or pUC18-CHIK1
145 plasmids and we provided the same bacteria to the adult mosquitoes that developed from these
146 larvae (**Figure 2A, Supplementary Figure 1D**). To activate transcription of the dsRNA from the T7
147 promoter(s), IPTG was added to the bacterial culture medium, the larval water, and the sucrose
148 solution provided to adults. Mosquitoes were offered bacteria-contaminated sucrose from their
149 emergence onward. Four- to five-day-old mosquitoes were blood-fed with a ZIKV- or CHIKV-
150 containing infectious bloodmeal and provided with bacteria-contaminated sucrose after the
151 bloodmeal. For ZIKV, viral titers were measured in the midguts, carcasses, and heads of mosquitoes
152 at six and ten days post-infection to assess infection and dissemination rates. We could not detect
153 a significant difference in ZIKV titers for mosquitoes colonized by dsZIK1-producing *E. coli* compared
154 to mosquitoes colonized by dsGFP-producing *E. coli* in any tissue at any time-point (**Figure 2B, C**,
155 Wilcoxon test on log transformed viral titers: day 6 midguts $p = 0.54$, carcasses $p = 0.72$, heads $p =$
156 0.87 ; day 10 midguts $p = 0.09$, carcasses $p = 0.29$, heads $p = 0.16$). The prevalence of infection was
157 also similar between the two conditions (Wilcoxon test: day 6 midguts $p = \text{NA}$, carcasses $p = 0.71$,
158 heads $p = 0.63$; day 10 midguts $p = 0.55$, carcasses $p = 0.35$, heads $p = 0.29$). These data indicate that
159 bacteria-produced dsRNA did not induce a protective RNAi-based antiviral response.

160 For CHIKV, viral titers were measured in the bodies and heads of mosquitoes at two and five days
161 post-infection. The results indicated that CHIKV titers were not specifically reduced by the exposure
162 of mosquitoes to dsCHIKV. Contrary to our hypothesis that dsRNA-producing bacteria could induce
163 a sequence-specific prophylactic antiviral response, CHIKV titers were higher in the bodies and
164 heads of mosquitoes colonized with bacteria carrying the T444T-CHIK1 or pUC18-CHIK1 plasmids
165 than those of mosquitoes colonized with bacteria carrying the T444T-GFP plasmid at day two post-
166 infection, while viral loads were similar in mosquitoes colonized with the three types of bacteria at
167 day five post-infection (**Supplementary Figure 1E-F**, Kruskal-Wallis test on log transformed viral
168 titers: day 2 body $p < 0.001$; day 2 head $p = 0.004$; day 5 body $p = 0.3$, ns; day 5 head $p = 0.8$, ns;
169 Kruskal-Wallis test on prevalence day 2 body $p = \text{NA}$; day 2 head $p = 0.003$; day 5 body $p = \text{NA}$; day
170 5 head $p = 0.5$, ns; see Supplementary Table 2 for individual comparisons). Taken together, these
171 results suggest that colonization of mosquitoes with dsRNA-producing bacteria did not trigger an
172 efficient antiviral response against flaviviruses or alphaviruses.

173 To verify whether the mosquito RNAi machinery could produce siRNAs from the dsRNA produced
174 by bacteria, we sequenced small RNAs from adult mosquitoes mono-colonized for two days with *E.*
175 *coli* HT115 carrying the T444T-ZIK1 plasmid (**Figure 3A**). Reads mapping to the ZIKV target sequence
176 on both positive and negative strands were obtained, with an enrichment of positive strand-reads
177 (**Figure 3B**). This further confirmed that dsRNA is effectively transcribed *in vivo* by *E. coli* HT115.
178 Most of the reads mapping to the target viral sequence fell within the expected range of 20 to 28
179 nt, but no enrichment at 21 nt was observed, suggesting that siRNAs were not produced from the
180 bacterial-produced dsRNA (**Figure 3C**). As our mosquito colony is persistently infected with the
181 insect specific virus Phasi-Charoen Like Virus (PCLV), we used this natural infection as a quality
182 control for our small RNA sequencing. We observed that 21 nt reads were significantly enriched
183 among all small RNA reads from 18 to 33 nt, suggesting that PCLV-specific siRNAs were produced in

184 these mosquitoes and were detected by our small RNA sequencing pipeline (**Supplementary Figure**
185 **3A-F**).

186 To assess whether different cloning strategies would result in an improved dsRNA-induced
187 response, we colonized mosquitoes with *E. coli* HT115 bacteria carrying the pUC18 hairpin-
188 producing plasmid containing either the 481 bp ZIK1 sequence (**Figure 3D**) or a shorter (175 bp) ZIK1
189 sequence (**Figure 3G**). We did not find evidence for siRNA production from dsRNA derived from
190 either the long (**Figure 3E-F**) or short construct (**Figure 3H-I**).

191 To assess whether the observed lack of an antiviral RNAi-based response was due to poor release
192 of dsRNA from live bacteria, we sequenced small RNAs from mosquitoes supplemented with the
193 same *E. coli* strains heat-killed before supplementation, as heat-killed bacteria were previously used
194 to induce oral RNAi in mosquitoes²¹. We did not find evidence for target-specific siRNA production
195 in any of the samples produced from mosquitoes supplemented with heat-killed bacteria
196 (**Supplementary Figure 4A-I**). We conclude that dsRNA produced by bacteria failed to interact with
197 the mosquito RNAi machinery and to induce the production of target-specific siRNAs.

198 *Oral supplementation of in vitro synthesized dsRNA does not induce an antiviral response in adult*
199 *mosquitoes*

200 The lack of an interaction between the bacterial-produced dsRNA and the mosquito RNAi machinery
201 could be due to several factors, such as lack of internalization of dsRNA by mosquito gut cells or low
202 stability of dsRNA in the mosquito gut. To investigate whether dsRNA could be internalized by the
203 mosquito gut epithelium and to overcome the potential limiting factor of insufficient release of
204 dsRNA from bacteria, we produced Cy3-labeled dsZIK1 (dsZIK1-Cy3) *in vitro*. Due to the low quantity
205 of dsZIK1-Cy3 that we obtained, we tested whether mosquitoes could feed on a 200 μ L sucrose
206 solution provided on a microtube cap (to prevent dilution of dsZIK1-Cy3 in a standard feeding
207 volume). To determine the rate of feeding with this setup, we provided mosquitoes with a sucrose
208 solution containing a blue dye. Mosquitoes that fed on the solution showed a readily visible blue
209 abdomen (**Figure 4A**). We dissected these mosquitoes after one or two days of feeding and verified
210 the presence of blue dye in the crop (where the sugar is stored before digestion) and midgut. While
211 after one day of feeding 67 % (10/15) of mosquitoes showed a blue crop and 80 % (12/15) a blue
212 midgut, after two days all mosquitoes (15/15) were found to have fed on the sucrose solution and
213 showed both a blue crop and midgut. dsZIK1-Cy3 was thus provided to mosquitoes for 48 h in a
214 sucrose solution with the setup described above, but without the blue dye that could potentially
215 interfere with fluorescence detection. Adult mosquitoes fed with dsZIK1-Cy3 were dissected and
216 guts (including crop and hindgut) were observed under a confocal microscope. Mosquitoes fed on
217 sucrose containing dsZIK1-Cy3 showed bright red fluorescence in all gut compartments (**Figure 4B-**
218 **C**). Interestingly, cells displaying red fluorescence in both the cytoplasm and nucleus were observed
219 in the midgut, which is the region in which blood is stored during digestion and that is primarily
220 infected by viruses ingested with blood (**Figure 4D**). These data indicate that at least some mosquito
221 gut epithelial cells can internalize dsRNA.

222 To explore if *in vitro*-synthesized dsRNA could induce an antiviral RNAi-based response when orally
223 administered to mosquitoes, we fed mosquitoes with non-labeled dsZIK1 and, as a control, with
224 dsGFP (**Figure 4E**). To confirm that the same dsRNA could induce an antiviral effect when introduced
225 systemically into the mosquito body, we injected mosquitoes with dsZIK1 or dsGFP. Four days after
226 the oral supplementation or the injection, we provided mosquitoes with a ZIKV-containing
227 infectious bloodmeal and measured viral titers in whole mosquitoes by plaque assays at seven and
228 14-days post-infection. We found that oral supplementation of dsZIK1 did not decrease viral titers
229 in mosquitoes collected at either time point, while injection of dsZIK1 completely blocked viral
230 replication at both timepoints (**Figure 4F-G**, Kruskal-Wallis test on log transformed viral titers: day
231 7 $p < 0.001$; day 14 $p < 0.001$; Kruskal-Wallis test on prevalence: day 7 $p < 0.001$; day 14 $p < 0.001$;
232 see Supplementary Table 2 for individual comparisons). In line with these results, mosquitoes fed
233 with dsCHIK1 did not show any significant reduction in viral titers after infection with CHIKV, while
234 mosquitoes injected with dsCHIK1 showed reduced viral titers in whole bodies two days post-
235 infection and marginally significant reduced titers five days post-infection (**Supplementary Figure**
236 **5A-C**, Kruskal-Wallis test on log transformed viral titers: day 2 $p < 0.001$; day 5 $p = 0.04$; Kruskal-
237 Wallis test on prevalence: day 2 $p = 0.18$; day 5 $p = 0.07$; see Supplementary Table 2 for individual
238 comparisons).

239 To verify that siRNAs were produced from dsRNA introduced into mosquitoes by injection or
240 feeding, we performed small RNA sequencing of RNA from whole mosquito samples. We found that
241 21 nt siRNAs were enriched in RNA from whole bodies of mosquitoes fed or injected with dsRNA
242 four days before sampling (**Figure 4H-K**).

243 To investigate the localization and integrity of dsRNA in fed and injected mosquitoes, RNA was
244 extracted from midguts, carcasses, and heads of mosquitoes collected two and four days after
245 feeding/injection (**Supplementary Figure 6A**). Northern blot analysis of samples collected at day
246 two indicated that RNA corresponding to the dsZIK1 sequence was detectable in the midgut,
247 carcass, and head of mosquitoes fed with dsRNA, while only in the carcass and head of mosquitoes
248 injected with dsRNA (**Supplementary Figure 6C**). At day four, a signal from dsZIK1 was detected in
249 the carcass of both fed and injected mosquitoes, and in the midgut of fed mosquitoes
250 (**Supplementary Figure 6C**). It is important to mention that, while the crop was not included in any
251 sample because of the presumable high concentration of dsRNA in this organ, carcass samples
252 contained Malpighian tubules and hindguts, and head samples included the anterior part of the
253 digestive system. Therefore, we cannot determine whether the dsZIK1 signal visible in RNA samples
254 extracted from fed mosquitoes derives from dsRNA that is circulating in the haemocoel and entering
255 other organs or if it represents what is ingested by mosquitoes that fed on dsRNA. However, these
256 results confirmed that dsRNA was not degraded by midgut RNases in fed mosquitoes.

257 Discussion

258 In mosquitoes, targeted RNAi can be activated by introduction of dsRNA corresponding to target
259 sequences. This stimulates the production of target-specific siRNAs from the introduced dsRNA,
260 which are used to guide degradation of complementary RNA. When the introduced dsRNA is
261 designed to target viral RNA, introduction of dsRNA prior to infection can limit subsequent viral

262 replication in a sequence-specific manner. In this paper we tested whether bacteria colonizing the
263 mosquito midgut could be used as a platform to produce virus-specific dsRNA and stimulate an
264 enhanced antiviral response against arboviruses. We showed that the *E. coli* HT115 strain, originally
265 used in *C. elegans* to induce RNAi, is poorly transmitted from mono-colonized larvae to adult
266 mosquitoes but can stably colonize the mosquito gut when provided to adult mosquitoes through
267 sucrose supplementation (**Figure 1D**). However, we did not detect any significant reduction on ZIKV
268 or CHIKV titers in mosquitoes colonized by dsZIK- or dsCHIK-producing bacteria (**Figure 2B-C** and
269 **Supplementary Figure 1E-F**). Moreover, our small RNA sequencing results did not show evidence
270 for the production of 21 nt siRNAs specific for the viral target sequence in mosquitoes provided with
271 live or heat-killed dsRNA-producing bacteria (**Figure 3, Supplementary Figure 4**). We identified at
272 least three potential issues that may have limited the interaction of the bacterial-produced dsRNA
273 with the mosquito RNAi machinery: (i) insufficient production or release of dsRNA by bacteria in the
274 mosquito gut, (ii) low stability of dsRNA in the mosquito gut, and/or (iii) limited uptake of dsRNA
275 from mosquito gut cells.

276 Although we confirmed dsRNA production *in vivo* by bacteria (**Supplementary Figure 2**), we cannot
277 confirm whether dsRNA was released by bacteria in the mosquito gut. In previous experiments in *C.*
278 *elegans*, *E. coli* HT115 was used as a source of food for the nematode and undergoes cell lysis during
279 ingestion and release dsRNA^{22,33}. It is unknown to what extent bacterial cells are lysed in the
280 mosquito gut. Live bacteria are needed at the larval stage to support mosquito development and
281 are found in the larval gut by two hours post-hatching, but dead bacteria can be found in the larval
282 gastric caeca and midgut by 16 hours post-hatching³⁴. This suggests that, at least at the larval stage,
283 some bacteria undergo cell lysis in the mosquito gut and potentially release dsRNA. Live, culturable
284 bacteria have been isolated from the adult gut from both lab-reared and field-collected
285 mosquitoes³⁵⁻³⁷ and an expansion of some members of the microbiota is observed 24 h after a
286 bloodmeal^{38,39}. However, the relative proportions of live and dead bacteria in adult mosquitoes are
287 unknown.

288 To test whether production or release of dsRNA from bacteria was a limiting factor in our system,
289 we decided to orally administer naked dsRNA to mosquitoes. Northern blot analysis of different
290 mosquito tissues confirmed that dsRNA molecules were, at least in part, intact after being provided
291 to mosquitoes via oral supplementation for two days, indicating that orally acquired dsRNA is not
292 totally degraded by gut RNases (**Supplementary Figure 6C**). Although oral acquisition of dsZIK
293 induced production of dsZIK-derived siRNAs in mosquitoes (**Figure 4H-I**), it failed to trigger an
294 antiviral RNAi response and limit ZIKV replication (**Figure 4F-G**). In contrast, the same dsZIK provided
295 systemically through injection completely blocked viral replication (**Figure 4F-G**). This suggests that
296 dsRNA provided via oral supplementation is not able to induce an RNAi response capable of
297 inhibiting viral replication. Downstream from siRNA production, RNAi has been shown to have a
298 limited role in restricting DENV or ZIKV infection in the *Ae. aegypti* midgut, predominantly due to
299 the lack of expression of the dsRNA-binding protein, Loqs2, in this tissue⁴⁰. This may explain why
300 siRNA produced after oral administration of dsRNA failed to produce an effective RNAi-based
301 response. To overcome this limitation, further applications involving oral RNAi should explore the
302 co-expression of Loqs2 in the mosquito midgut.

303 Cy3-labeled dsRNA provided orally to mosquitoes could be visualized in the crop, diverticula, and
304 some midgut cells, where fluorescence could be detected in both the cytoplasm and nucleus (**Figure**
305 **4B-D**). However, we did not investigate whether orally supplemented dsRNA could be found in other
306 tissues distal from the digestive system. A recent study described a similar strong fluorescence signal
307 in the alimentary canal of *Ae. aegypti* larvae and adults fed with fluorescently-labeled dsRNA⁴¹.
308 Interestingly, this study showed that adult mosquitoes fed with fluorescently-labeled dsRNA did not
309 display fluorescence in other tissues, while mosquitoes intrathoracically injected with dsRNA
310 showed strong fluorescence in hemocytes, pericardial cells, and ovaries. Uptake of dsRNA and
311 amplification of the RNAi signal by hemocytes is essential for effective antiviral RNAi in *Drosophila*
312 *melanogaster*⁴². We can therefore speculate that the failure of orally acquired dsRNA to inhibit viral
313 replication in *Ae. aegypti* may be due to incorrect processing of this dsRNA, which may be
314 internalized by midgut cells, but not hemocytes or cells in other tissues relevant for an antiviral
315 response.

316 The only previous report of successful microbiota-mediated antiviral RNAi in insects was in
317 honeybees infected with deformed wing virus, where bees colonized by viral dsRNA-producing
318 microbiota were found to exhibit longer lifespans and reduced viral replication compared to bees
319 carrying dsGFP-producing bacteria⁸. However, mosquitoes and bees might have physiological
320 differences that could limit this application against arboviruses. For instance, the honeybee
321 microbiota almost exclusively colonizes the hindgut⁴³, while mosquitoes have high bacterial loads
322 mainly in the crop and midgut⁴⁴. Differences in viral infection dynamics and/or dsRNA uptake by
323 different tissues might explain the different results obtained with *Apis mellifera* and *Ae. aegypti*.

324 Over the last ten years, oral RNAi targeting essential mosquito genes has been proposed as a vector
325 control strategy to inhibit or block mosquito development and/or reproduction. Several methods
326 have been developed to deliver dsRNA to mosquito larvae or adults, including feeding of naked
327 dsRNA, colonization with microorganisms producing dsRNA, or feeding with nanoparticle-dsRNA
328 complexes (reviewed in ²¹). However, some studies have reported high variability or low gene
329 silencing efficiency when dsRNA was provided via oral feeding^{15,16} and it is likely that other studies
330 with contradictory or negative results were left unpublished. As emphasized in a recent discussion
331 by Devang Mehta, the importance of showcasing negative results for the advancement of science
332 cannot be overstated⁴⁵. It is key to acknowledge that when negative results go unpublished, other
333 scientists can't learn from them and end up repeating failed experiments, leading to a waste of
334 public funds and a delay in genuine progress. Our results point to clear limitations of oral RNAi as a
335 strategy to disrupt viral transmission by mosquitoes and should be considered when developing
336 new strategies for vector control.

337 **Materials and Methods**

338 *Bacteria, viruses, cells and mosquitoes*

339 *E. coli* HT115(DE3, Genotype: F-, mcrA, mcrB, IN(rrnD-rrnE)1, rnc14::Tn10) was obtained from the
340 Caenorhabditis Genetics Center (CGC, University of Minnesota). For all experiments, bacteria were
341 inoculated from fresh colonies in Luria-Bertani (LB) broth supplemented with 20 µg/mL tetracycline.

342 The growth medium was supplemented with 100 µg/mL ampicillin for bacteria carrying the T444T
343 or pUC18 plasmids.

344 The ZIKV strain used in this study is the prototype African MR766 strain, derived from a previously
345 described infectious clone⁴⁶. The CHIKV strain used is the Caribbean strain, Asian genotype originally
346 described in⁴⁷. Viral stocks were produced in Vero E6 cells.

347 Vero E6 cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Life Technologies)
348 supplemented with 10 % fetal bovine serum (Invitrogen) and 1 % Penicillin/Streptomycin (P/S, Life
349 Technologies) at 37 °C with 5 % CO₂.

350 *Ae. aegypti* mosquitoes belonged to the 28th–31st generation of a colony created in 2013 from
351 mosquitoes collected in the Kamphaeng Phet Province, Thailand⁴⁸. They were maintained under
352 standard insectary conditions (27 °C, 70 % relative humidity, 12h:12h light/dark cycle). Mosquitoes
353 are routinely blood-fed with rabbit blood with an Hemotek system (Hemotek Ltd). Conventionally
354 reared larvae were reared in 30x40 cm plastic trays with 1.5 L of dechlorinated tap water and ground
355 TetraMin fish food. Pupae were transferred in plastic cages (BugDorm) for emergence and adults
356 were provided with autoclaved 10 % sucrose solution.

357 *Selection of viral target sequences for dsRNA production*

358 Sequences of the ZIKV (MR766 strain) or CHIKV (Caribbean strain) genome spanning regions of the
359 polyprotein coding sequence corresponding to different mature proteins were selected and used as
360 a blastn⁴⁹ query against the *Ae. aegypti* genome in Vectorbase and all *Aedes* sequences in Genbank
361 to verify that they did not target any *Aedes* sequence. To exclude any interference by mosquito
362 small RNAs on the target viral sequence, small RNA library data previously produced in the lab from
363 *Ae. aegypti* uninfected mosquitoes were mapped on the ZIKV and CHIKV sequences. Any mosquito
364 small RNA was found to map on the selected viral sequences.

365 *Cloning*

366 The T444T plasmid was a gift from Tibor Vellai (Addgene plasmid #113081; RRID:
367 Addgene_113081²³). GFP and target viral sequences were cloned into T444T using *NotI* and *AgeI*
368 restriction sites and standard cloning techniques. The NEBuilder® HiFi DNA Assembly Kit (New
369 English Biolabs) was used to insert the following elements into the pUC18 plasmid: the T7 promoter,
370 *NotI* and *AgeI* restriction sites, a spacer sequence corresponding to a 300 bp portion of an intron of
371 the *Drosophila melanogaster actin* gene (GenBank: M18829.1), *NheI* and *KpnI* restriction sites and
372 the T7 terminator sequence. The final pUC18-spacer plasmid sequence and map are available upon
373 request. The same primers used for the cloning in T444T were used to clone GFP and target viral
374 sequences in pUC18-spacer using *NotI* and *AgeI* restriction sites. To clone the inverted repeat, *NheI*
375 and *KpnI* restriction sites were used. A shorter ZIK1 sequence (175 bp) was also cloned in pUC18-
376 spacer using the same cloning strategy. All primers used are listed in Supplementary Table 1. After
377 cloning, plasmid sequences were confirmed via Sanger sequencing from both sides of the insert.

378 *Mono-colonization of Ae. aegypti larvae with E. coli HT115*

379 Egg sterilization and larval mono-colonization were performed in sterile conditions working inside
380 a microbiological safety cabinet. *Ae. aegypti* eggs were surface sterilized with subsequent 5-minute
381 washes in 70 % ethanol, 1 % bleach, and 70 % ethanol. After three rinses in sterile Milli-Q water,
382 eggs were transferred to a 25 mL cell culture flask with vented cap overnight for hatching. On the
383 same day, *E. coli* HT115 carrying the selected plasmid was used to inoculate LB supplemented with
384 40 μ M IPTG, 20 μ g/mL tetracycline and 100 μ g/mL ampicillin, and grown for \sim 16 h at 37 $^{\circ}$ C, 200 rpm
385 shaking. The following day, the bacterial culture was centrifuged and resuspended in sterile Milli-Q
386 water at a final concentration of 10^8 CFUs/mL. First instar larvae were placed in plastic boxes cleaned
387 with ethanol (200 larvae per box), together with 500 mL of bacterial suspension, 40 μ M IPTG and 5
388 mL of sterile fish food (TetraMin baby, 50 mg/mL). Boxes were closed with filtered covers and kept
389 at 27 $^{\circ}$ C, 12h:12h light/dark cycle. Pupae were transferred into sterile boxes for emergence as in³².
390 Larval water was plated on LB agar and LB agar supplemented with 100 μ g/mL ampicillin to confirm
391 the mono-colonization of mosquitoes. The same number of colonies and the same colony
392 morphology were observed in both media.

393 To monitor the transstadial transmission of *E. coli* HT115-T444T-ZIK1 from larvae to adults, adult
394 mosquitoes originating from mono-colonized larvae were either provided with 10 % sterile sucrose
395 or with bacteria resuspended in 10 % sterile sucrose at a final concentration of 10^9 CFUs/mL. At days
396 one and four post-emergence, mosquitoes were collected, surface sterilized for 3 minutes in 70 %
397 ethanol, rinsed in sterile PBS, and individually homogenized in 200 μ L LB. Mosquito homogenates
398 were serially diluted in LB and plated on LB agar and LB agar supplemented with 100 μ g/mL
399 ampicillin. Plates were placed overnight at 37 $^{\circ}$ C and the number of colonies was counted to
400 estimate whole mosquito bacterial loads.

401 *Infection of mono-colonized mosquitoes with ZIKV or CHIKV*

402 Larvae mono-colonized by the selected strain of *E. coli* HT115 were produced as described above.
403 Larval water was supplemented with 40 μ M IPTG to induce the transcription of the dsRNA by
404 bacteria. Adult mosquitoes were provided with a 10 % sterile sucrose solution supplemented with
405 10^9 CFUs/mL bacteria and 40 μ M IPTG for four days. The day before the infectious bloodmeal,
406 mosquitoes were deprived of the sucrose/bacteria/IPTG solution for starvation. Mosquitoes were
407 transferred to a biosafety level 3 insectary and offered an infectious bloodmeal on a Hemotek
408 system (Hemotek Ltd) consisting of washed rabbit blood supplemented with 10^6 PFUs/mL of ZIKV
409 or CHIKV and 10 mM ATP. Fully engorged mosquitoes were kept at 27 $^{\circ}$ C, 70 % relative humidity,
410 12h:12h light/dark cycle until sampling and provided with fresh bacteria and sucrose every three
411 days. At the selected time points, mosquitoes were dissected and tissues were placed in 200 μ L PBS.
412 Samples were kept at -80 $^{\circ}$ C until processing.

413 *Plaque assay*

414 The day before the assay, Vero-E6 cells were seeded in 24-well plates at a density of 1.75×10^5
415 cells/well in DMEM with 10 % FBS and 1 % P/S. The day of the assay, mosquito samples were
416 homogenized in a bead beater for 30 sec at 5000 rpm. Ten-fold dilutions of mosquito homogenates
417 or of an aliquot of the bloodmeal were prepared in DMEM. Two-hundred μ L of each dilution were

418 added to the cell monolayer and incubated for 1 h at 37 °C with 5 % CO₂. Cells were covered with
419 DMEM with 2 % FBS, 1 % P/S and 0.8 % agarose. After seven (ZIKV) or three (CHIKV) days, cells were
420 fixed with 4 % paraformaldehyde. Plaques were counted after 0.2 % crystal violet staining.

421 *Small RNA sequencing*

422 RNA was extracted from pools of five mosquitoes using TRIzol. Small RNAs of 18-33 nt in length were
423 purified from a 15% acrylamide/bisacrylamide (37.5:1), 7 M urea gel as described previously⁵⁰.
424 Libraries were prepared using the NEBNext® Small RNA Library Prep Set for Illumina® (New England
425 BioLabs), with the Universal miRNA Cloning Linker (New England BioLabs) as the 3' adaptor and in-
426 house designed indexed primers. Libraries were diluted to 4 nM and sequenced using an Illumina
427 NextSeq 500 High Output kit v2 (75 cycles) on an Illumina NextSeq 500 platform.

428 *Bioinformatics analysis of small RNA libraries*

429 FastQC⁵¹ was used to assess the quality of fastq file. Low quality portions and adaptors were
430 trimmed from each read using cutadapt⁵². Only reads with a phred score ≥20 were kept. FastQC was
431 used on the fastq files created by cutadapt to evaluate the overall quality. Mapping was performed
432 using Bowtie⁵³ with the -v 1 option (one mismatch between the read and its target) using as
433 reference sequences the ZIKV, CHIKV, or GFP target sequences, or the PCLV genome. Bowtie
434 generates results in sam format. All sam files were analyzed by different tools of the package
435 samtools⁵⁴ to produce bam indexed files. To analyze these bam files, different kind of graphs were
436 generated using home-made R scripts with several Bioconductor libraries such as Rsamtools and
437 Shortreads⁵⁵.

438 *Synthesis and labelling of dsRNA*

439 RNA corresponding to the same ZIK1, CHIK1 and GFP sequences cloned in the T444T plasmid was
440 transcribed *in vitro* using the T7 MEGAscript Kit (Invitrogen) following the manufacturer's
441 instructions. A PCR amplicon generated with both forward and reverse primers containing the T7
442 promoter sequence (Supplementary Table 1) was used as a template for the *in vitro* transcription.
443 A DNase treatment was performed after transcription to eliminate the DNA template. The annealing
444 of the two strands of dsRNA was allowed by heating the RNA at 98 °C for 5 min and leaving it to cool
445 down slowly to room temperature. After quantification with Nanodrop, RNA was stored at -80 °C
446 until use. The Silencer™ siRNA Labeling Kit with Cy™3 dye (Invitrogen) was used to label dsRNA
447 following the manufacturer's instructions. An aliquot of the labeled dsRNA was run on a 1 % agarose
448 gel to visualize a shift in the migration and thus confirm the labeling.

449 *Oral feeding of mosquitoes with labeled dsRNA, sample collection and visualization*

450 One-to-three day old conventionally reared female mosquitoes were starved for one day and then
451 provided with a 10 % sucrose solution containing 200 ng/μL of the labeled dsRNA. After two days of
452 feeding, midguts were dissected and fixed overnight in 4 % methanol-free paraformaldehyde in PBS.
453 After three washes with PBS, samples were permeabilized in 1 % BSA, 0.1 % Triton X-100 in PBS for
454 1 h at room temperature. After three washes with PBS, midguts were incubated for 30 minutes with

455 DAPI and Oregon Green 488 Phalloidin (Invitrogen) dyes to stain DNA and actin, respectively. After
456 three additional washes in PBS, midguts were mounted on slides with Vectashield H-1000 (Vector
457 Laboratories) and visualized on a Zeiss LSM 780 confocal microscope at 100X magnification. To
458 reconstruct mosquito full digestive tract images, multiple flanking images were acquired using the
459 “tile” function and subsequently assembled using the ZEN Microscopy Software.

460 *Infection of mosquitoes fed or injected with dsRNA*

461 One-to-three day old conventionally reared female mosquitoes were starved for one day and then
462 provided with a 10 % sucrose solution containing 200 ng/ μ L of dsRNA and 1 mg/mL of the blue dye,
463 Erioglaucine disodium salt (Sigma-Aldrich). The control consisted of mosquitoes fed on blue-dyed
464 sucrose without dsRNA. All mosquitoes were allowed to feed on the sucrose solution for three days
465 and then only mosquitoes showing a blue abdomen were selected for infection. In parallel, one-to-
466 three day old conventionally reared female mosquitoes were intrathoracically injected with 250 ng
467 of dsRNA using a Nanoject III system (Drummond). They were left to recover for three days. The day
468 before the infectious bloodmeal, mosquitoes were deprived of sucrose for starvation. Mosquitoes
469 were infected with 10^6 PFUs/mL of ZIKV MR766 or CHIKV as described above. At the selected time
470 points, whole mosquitoes were collected in 2 mL screwcap tubes containing 200 μ L of sterile PBS
471 and glass beads. Samples were kept at -80 °C until processing. Viral titers in whole mosquitoes were
472 measured by plaque assay as described above.

473 *Northern blot*

474 ZIK1 sense and antisense RNA probes labeled with digoxigenin were synthesized using the DIG
475 Northern Starter Kit (Roche) following the manufacturer’s instructions. The template consisted of a
476 PCR product obtained with primers containing the T7 (sense) or T3 (antisense) promoter sequence
477 (Supplementary Table 1). Serial dilutions of the two probes were visualized on a dot blot for probe
478 quantification.

479 *E. coli* HT115-T444T-ZIK1 or -CHIK1 (probe specificity control) was inoculated in 5 mL LB
480 supplemented with 20 μ g/mL tetracycline and 100 μ g/mL ampicillin and grown overnight at 37 °C
481 shaking at 200 rpm. The following day, 100 μ L of the overnight culture were used to inoculate 5 mL
482 of LB with 20 μ g/mL tetracycline and 100 μ g/mL ampicillin with or without 40 μ M IPTG. Bacteria
483 were grown overnight at 37 °C shaking at 200 rpm. After 24 h, 1 mL of bacterial culture was
484 centrifuged and the bacterial pellet was resuspended in 100 μ L of 0.1 % SDS. Bacterial extracts were
485 incubated for 2 min at 95 °C for cell lysis and RNA was extracted using TRIzol.

486 Mosquitoes fed or injected with dsRNA were dissected after two or four days of feeding with 200
487 ng/ μ L dsZIK or after injection with 250 ng/mosquito of dsZIK1. For each time-point/condition, 15 to
488 28 mosquitoes were dissected and their midguts, carcasses (excluding crop, including hindgut and
489 Malpighian tubules), and heads were pooled and stored at -80 °C until processing. Total RNA was
490 extracted from each sample pool using TRIzol.

491 For northern blot analysis, RNA samples were mixed with a glyoxal-containing loading dye,
492 incubated at 50 °C for 30 min for RNA denaturation, and loaded on a 1.2 % agarose gel. Five μ g of

493 total RNA were loaded on the gel for all samples (including *in vitro* synthesized dsRNA) except for
494 midguts at day 4 (injected) and heads (fed/injected) at day 4, for which 1.5 µg of RNA were loaded
495 due to low RNA yields. After gel electrophoresis, RNA was transferred to a Nytran Super Charge
496 membrane (Whatman) using the NorthernMax™-Gly kit (Invitrogen) following the manufacturer's
497 instructions. After transfer and crosslinking with UV light, the membrane was incubated overnight
498 with 0.2 ng/µL of the selected probe. Membranes were incubated with an Alkaline Phosphatase-
499 conjugated Anti-Digoxigenin antibody and the chemiluminescent signal was revealed using a
500 ChemiDoc MP imaging system (Biorad).

501 *Statistics*

502 All graphs were created using GraphPad Prism (version 9.5.1) or R (version 4.2.1). Statistical analyses
503 were performed in R (version 4.2.1). The normality of data was tested with a Shapiro test. A
504 Wilcoxon Signed Rank test was used to compare mosquito bacterial loads, prevalence and log₁₀
505 transformed viral titers of mosquitoes colonized by dsZIK-producing bacteria. A Kruskal-Wallis one-
506 way analysis of variance followed by a Dunn test with Bonferroni correction was used to compare
507 log₁₀ transformed viral titers in mosquitoes colonized by dsCHIK producing bacteria, or in
508 mosquitoes fed or injected with dsZIK/dsCHIK. Detailed results of statistical analyses are shown in
509 Supplementary Table 2. R codes are available upon request.

510

511 **Data availability**

512 The small RNA sequencing data were deposited to the Sequence Read Archive under accession
513 number PRJNA1034089.

514 **Acknowledgments**

515 We are grateful to Jared Nigg and Louis Lambrechts for critically reading of the manuscript. We
516 thank Catherine Lallemand for technical assistance in mosquito rearing, and Madeleine Lausted for
517 assistance in dsRNA feeding.

518 This work was supported by funding from the French Government's Investissement d'Avenir
519 program, Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases (grant ANR-
520 10-LABX-62-IBEID), Agence Nationale de la Recherche (grant ANR-22-CE35-0001, MAMMAMIA),
521 and Fondation iXcore - iXlife - iXblue Pour La Recherche to M.C.S., the Pasteur-Roux-Cantarini
522 fellowship of Institut Pasteur to O.R. We acknowledge the kind financial support to the Photonic
523 BioImaging (UTechS PBI) platform by the Institut Pasteur (Paris), the France-BioImaging
524 infrastructure network supported by the Agence Nationale de la Recherche (ANR-10-INBS-04,
525 Investments for the future), and the Région Ile-de-France (program DIM-Malinf).

526 **Authors contributions**

527 O.R. and M.-C.S. designed the experiments; O.R., A.H.-L., and H.B. performed the experiments; O.R.
528 and L.F. analyzed the data; O.R. wrote the paper with input from co-authors. All authors reviewed
529 and approved the final version of the manuscript.

530

531 **Declaration of interests**

532 The authors declare no competing interests.

533

534 **Figures legends**

535 **Figure 1. Colonization of mosquitoes with dsRNA producing *E. coli*.** (A) Representation of the T444T
536 plasmid. The target viral sequence was cloned between two directionally opposed T7 promoters to
537 allow convergent transcription of a sense and antisense RNAs that form dsRNA upon annealing. (B)
538 Position of the target viral sequence on the ZIKV genome. The selected sequence is 481 nt and spans
539 the coding sequences for the Capsid (Cap) and the Pre-membrane (Pr) protein coding genes. The
540 other components of the ZIKV genome are: M (Membrane), E (Envelope) NS1-5 (Non-Structural
541 proteins). (C) Experimental set up for mosquito mono-colonization with *E. coli* HT115. Sterile eggs
542 were obtained by subsequent washes in 70 % ethanol and 1 % bleach. An overnight culture of *E. coli*
543 HT115 carrying the T444T-ZIK1 plasmid was added to axenic larvae hatched from sterile eggs to
544 obtain mono-colonized larvae. Adults emerging from mono-colonized larvae were provided with
545 either sterile sucrose or sucrose supplemented with *E. coli* HT115. Colony forming units (CFUs) were
546 determined in whole adults to evaluate the efficiency of bacterial colonization. (D) Bacterial loads

547 (mean) and plasmid stability *in vivo*. CFUs were determine one and four days-post-emergence in
548 adult mosquitoes mono-colonized with *E. coli* HT115-T444T-ZIK1 at the larval stage, fed with either
549 sterile (black) or bacterial supplemented (yellow) sucrose. To determine plasmid stability in bacteria
550 during mosquito colonization, CFUs were determined on LB plates supplemented (full dots) or not
551 (empty dots) with ampicillin. Bacterial loads were compared using a Wilcoxon test.

552 **Figure 2. Mosquitoes colonized with dsZIK-producing bacteria do not show reduced viral titers.**
553 (A) Experimental protocol for ZIKV infection of mosquitoes colonized by dsRNA-producing *E. coli*.
554 Mosquitoes originating from mono-colonized larvae were kept on bacteria-supplemented sucrose
555 for four days. After one day of starvation, they were offered a ZIKV infectious bloodmeal and
556 subsequently kept with bacteria-supplemented sucrose until sampling at six and ten days post-
557 infection. At each time point midguts, carcasses, and heads of individual mosquitoes were collected.
558 (B-C) Viral titers (mean \pm SEM) in midguts, carcasses, and heads of individual mosquitoes colonized
559 by *E. coli* HT115 expressing dsGFP (grey) or dsZIK1 (yellow) and collected six (B) or ten (C) days post-
560 infection. Each plot shows the log of plaque forming units (PFUs) measured in individual mosquitoes
561 in each tissue. Pie charts show the proportion of infected (grey/yellow) and uninfected (white)
562 mosquitoes. Numbers below pie charts represent the number of infected over uninfected
563 mosquitoes per tissue/time-point. Viral loads were compared using a Wilcoxon test.

564 **Figure 3. Mosquitoes colonized with dsZIK-producing bacteria do not show production of dsZIK-**
565 **specific siRNAs.** Small RNAs were sequenced from pools of whole mosquitoes colonized for two
566 days with the *E. coli* HT115 strain carrying either the T444T-ZIK1 plasmid (A), or the pUC18-ZIK1
567 plasmid with the 481 nt (D) or 175 nt (G) ZIKV sequence. For each condition, the number of reads
568 mapping to each position of the positive strand (grey) or negative strand (black) of the target
569 sequence is displayed (B, E, H). The size distribution of the reads shown in (B, E, H) is shown in (C, F,
570 I), respectively.

571 **Figure 4. A protective antiviral response is induced by systemic injection of dsRNA in adult**
572 **mosquitoes, but not by feeding.** (A) A blue sucrose solution containing dsRNA was provided to
573 mosquitoes for two days. Blue dye in the crop and midgut is visible through the cuticle and allows
574 selection of fed mosquitoes. (B-D) Confocal images of the digestive tract of a mosquito fed for 48 h
575 with dsZIK1-Cy3. In (B) different dyes indicate DNA (blue, DAPI), actin (green, Phalloidin), and
576 dsZIK1-Cy3 (red, Cy3), while in (C) only the signal of dsZIK1-Cy3 is shown. In (C) the different
577 components of the mosquito digestive tract are indicated. (D) represents a higher magnification
578 image of a region of the same midgut shown in (B-C). Scale bars correspond to 500 μ m (B-C) or 50
579 μ m (D). (E) Experimental protocol for ZIKV infection of mosquitoes fed or injected with dsZIK1 or
580 dsGFP. At day 0, mosquitoes were fed with a 200 ng/ μ L solution of dsZIK1 or dsGFP, or injected with
581 250 ng dsZIK1 or dsGFP/mosquito. After four days, they were offered a ZIKV infectious bloodmeal.
582 At seven and 14 days after infection, whole individual mosquitoes were collected. (F-G) Viral titers
583 (mean \pm SEM) in individual mosquitoes (whole bodies) fed on sucrose (black) or sucrose containing
584 dsGFP (full grey) or dsZIK1 (full yellow), or injected with dsGFP (empty grey) or dsZIK1 (empty
585 yellow). Mosquitoes were collected seven (F) or 14 (G) days post-infection. Each plot shows the log
586 of plaque forming units (PFUs) measured in individual mosquitoes. Pie charts show the proportion

587 of infected (black/grey/yellow) and uninfected (white) mosquitoes. Numbers below pie charts
588 represent the number of infected over uninfected mosquitoes per tissue/time-point. Viral loads
589 were compared using a Kruskal-Wallis one-way ANOVA followed by a Dunn test with Bonferroni
590 correction. The global ANOVA p value is displayed in the top-right corner of each plot and
591 significance groups ($p < 0.05$) are denoted by letters. (H-K) Small RNAs were sequenced from pools
592 of five whole mosquitoes fed with dsZIK1 for two days (H, I) or injected with dsZIK1 two days before
593 sampling (J, K). (H, J) For each condition, the number of reads mapping to each position of the
594 positive strand (grey) or negative strand (black) of the target sequence is displayed. The size
595 distribution of the reads shown in (H, J) is shown in (I, K), respectively.

596 **Supplementary figures and tables legends**

597 **Supplementary Figure 1. Mosquitoes colonized with dsCHIK-producing bacteria do not show**
598 **reduced viral titers.** (A) Representation of the T444T plasmid. The target viral sequence was cloned
599 between two directionally opposed T7 promoters to allow convergent transcription of a sense and
600 antisense RNAs that form dsRNA upon annealing. (B) Representation of the pUC18 plasmid. An
601 inverted repeat of the target viral sequence was cloned downstream from a T7 promoter, allowing
602 the formation of an RNA hairpin after transcription. (C) Position of the target viral sequence on the
603 CHIKV genome. The selected sequence is 557 nt long and spans the coding sequences for the Capsid
604 (Cap) and the Envelope (E3) peptide coding genes. The other components of the CHIKV genome are
605 NSPs (Non-Structural Proteins). (D) Experimental protocol for CHIKV infection of mosquitoes
606 colonized by dsRNA-producing *E. coli*. Mosquitoes originating from mono-colonized larvae were
607 kept on bacteria-supplemented sucrose for four days. After one day of starvation, they were offered
608 a CHIKV-containing infectious bloodmeal and subsequently kept with bacteria-supplemented
609 sucrose until sampling at two and five days post-infection. At each time point, bodies and heads of
610 individual mosquitoes were collected. (E-F) Viral titers (mean \pm SEM) in bodies and heads of
611 individual mosquitoes colonized by *E. coli* HT115 carrying the T444T-GFP (grey), T444T-CHIK1 (dark
612 blue), or pUC18-CHIK1 (light blue) plasmid and collected two (E) or five (F) days post-infection. Each
613 plot shows the log of plaque forming units (PFUs) measured in individual mosquitoes in each tissue.
614 Pie charts show the proportion of infected (grey/blue) and uninfected (white) mosquitoes. Numbers
615 below pie charts represent the number of infected over uninfected mosquitoes per tissue/
616 time-point. Viral loads were compared using a Kruskal-Wallis one-way ANOVA followed by a Dunn test
617 with Bonferroni correction. The global ANOVA p value is displayed in the top-right corner of each
618 plot and significance groups ($p < 0.05$) are denoted by letters.

619 **Supplementary Figure 2. Northern blot analysis of *E. coli* HT115-T444T-ZIK1 total RNA.** (A, B)
620 Ethidium bromide-stained agarose gel (A) and northern blot (B) of dsZIK1 synthesized *in vitro* and
621 total RNA extracted from a 24 h bacterial culture without and with 40 μ M IPTG induction. For (B),
622 the same samples were loaded in duplicate on the same agarose gel and transferred on the same
623 membrane. The membrane was cut and incubated with either a sense (left) or antisense (right) 345
624 nt probe annealing to the ZIK1 sequence. (C, D) Specificity of the ZIK1 sense probe. Ethidium
625 bromide-stained agarose gel (C) and northern blot (D) of the same samples showed in (A, B) loaded
626 together with dsCHIK1 synthesized *in vitro* and total RNA extracted from a 24 h culture of *E. coli*

627 HT115-T444T-CHIK1 without and with 40 μ M IPTG induction. Northern blot membrane was
628 incubated with the sense ZIK1 probe to show binding specificity to dsZIK1 RNA.

629 **Supplementary Figure 3. Sequencing quality control of small RNAs mapping on the Phasi-Charoen**
630 **Like Virus (PCLV) genome.** Small RNAs were sequenced from the pools of whole mosquitoes shown
631 in Figure 3 and mapped to the PCLV genomic segments S (A-B), M (C-D), and L (E-F). The number of
632 reads mapping to each position of the positive strand (grey) or negative strand (black) of the PCLV
633 genome is shown (A, C, E). The size distribution of the reads shown in (A, C, E) is shown in (B, D, F),
634 respectively.

635 **Supplementary Figure 4. Mosquitoes colonized with heat-killed dsZIK-producing bacteria do not**
636 **show production of dsZIK-specific siRNAs.** Small RNAs were sequenced from pools of whole
637 mosquitoes supplemented for two days with sucrose containing heat-killed *E. coli* HT115 (heated
638 for 5 min at 98 °C) carrying either the T444T-ZIK1 plasmid (A), or the pUC18-ZIK1 plasmid with the
639 481 nt (D) or 175 nt (G) ZIKV sequence. For each condition, the number of reads mapping to each
640 position of the positive strand (grey) or negative strand (black) of the target sequence is displayed
641 (B, E, H). The size distribution of the reads shown in (B, E, H) is shown in (C, F, I), respectively.

642 **Supplementary Figure 5. Effect of oral supplementation or injection of *in vitro* synthesized dsRNA**
643 **on CHIKV infection.** (A) Experimental protocol for CHIKV infection of mosquitoes fed or injected
644 with dsCHIK1 or dsGFP. At day 0, mosquitoes were fed with a 200 ng/ μ L solution of dsCHIK1 or
645 dsGFP, or injected with 250 ng dsCHIK1 or dsGFP/mosquito. After four days, they were offered a
646 CHIKV infectious bloodmeal. At two and five days after infection, whole individual mosquitoes were
647 collected. (B-C) Viral titers (mean \pm SEM) in individual mosquitoes (whole bodies) fed on sucrose
648 (black) or sucrose containing dsGFP (full grey) or dsCHIK1 (full blue), or injected with dsGFP (empty
649 grey) or dsCHIK1 (empty blue). Mosquitoes were collected two (B) or five (C) days post-infection.
650 Each plot shows the log of plaque forming units (PFUs) measured in individual mosquitoes. Pie
651 charts show the proportion of infected (black/grey/blue) and uninfected (white) mosquitoes.
652 Numbers below pie charts represent the number of infected over uninfected mosquitoes per
653 tissue/time-point. Viral loads were compared using a Kruskal-Wallis one-way ANOVA followed by a
654 Dunn test with Bonferroni correction. The global ANOVA *p* value is displayed in the top-right corner
655 of each plot and significance groups (*p* < 0.05) are denoted by letters.

656 **Supplementary Figure 6. Northern blot analysis of total RNA extracted from mosquitoes fed or**
657 **injected with dsZIK1.** (A) Experimental protocol for sample collection of mosquitoes fed or injected
658 with dsZIK1. At day 0, mosquitoes were fed with a 200 ng/ μ L solution of dsZIK1 or injected with 250
659 ng dsZIK1/mosquito. After two and four days, midguts, carcasses, and heads of individual
660 mosquitoes were collected and pooled. RNA was extracted from pools of 15 to 28 mosquitoes. (B-
661 C) Ethidium bromide-stained agarose gel (B) and northern blot (C) of RNA samples extracted from
662 pools of midguts (M), carcasses (C), or heads (H) of mosquitoes fed or injected with dsZIK1 collected
663 two or four days after feeding/injection. The membrane was incubated with a sense 345 nt probe
664 annealing to the ZIK1 sequence. dsZIK1 synthesized *in vitro* was loaded on the same gel as a control.

665 **Supplementary Table 1. List of primers used in this study.** For each primer, the following
666 information is indicated: name, sequence (5'-3'), GenBank reference sequence, position on the
667 reference sequence, amplicon length and application.

668 **Supplementary Table 2. Results of statistical analyses.**

669

670 **References**

- 671 1. Galiana-Arnoux, D., Dostert, C., Schneemann, A., Hoffmann, J. A. & Imler, J. L. Essential function in
672 vivo for Dicer-2 in host defense against RNA viruses in drosophila. *Nature Immunology* 2006 7:6 **7**,
673 590–597 (2006).
- 674 2. Rand, T. A., Ginalski, K., Grishin, N. V. & Wang, X. Biochemical identification of Argonaute 2 as the
675 protein required for RNA-induced silencing complex activity. *Proc Natl Acad Sci U S A* **101**, 14385–
676 14389 (2004).
- 677 3. Van Rij, R. P. *et al.* The RNA silencing endonuclease Argonaute 2 mediates specific antiviral immunity
678 in *Drosophila melanogaster*. *Genes Dev* **20**, 2985–2995 (2006).
- 679 4. Sánchez-Vargas, I. *et al.* Dengue Virus Type 2 Infections of *Aedes aegypti* Are Modulated by the
680 Mosquito’s RNA Interference Pathway. *PLoS Pathog* **5**, e1000299 (2009).
- 681 5. Blair, C. D. A Brief History of the Discovery of RNA-Mediated Antiviral Immune Defenses in Vector
682 Mosquitos. *Microbiology and Molecular Biology Reviews* (2022) doi:10.1128/MMBR.00191-21.
- 683 6. Olson, K. E. *et al.* Genetically Engineered Resistance to Dengue-2 Virus Transmission in Mosquitoes.
684 *Science (1979)* **272**, 884–886 (1996).
- 685 7. Magalhaes, T. *et al.* Induction of RNA interference to block Zika virus replication and transmission in
686 the mosquito *Aedes aegypti*. *Insect Biochem Mol Biol* **111**, 103169 (2019).
- 687 8. Leonard, S. P. *et al.* Engineered symbionts activate honey bee immunity and limit pathogens. *Science*
688 (1979) **367**, 573–576 (2020).
- 689 9. Van Ekert, E., Powell, C. A., Shatters, R. G. & Borovsky, D. Control of larval and egg development in
690 *Aedes aegypti* with RNA interference against juvenile hormone acid methyl transferase. *J Insect*
691 *Physiol* **70**, 143–150 (2014).
- 692 10. Whyard, S. *et al.* Silencing the buzz: A new approach to population suppression of mosquitoes by
693 feeding larvae double-stranded RNAs. *Parasit Vectors* **8**, 1–11 (2015).
- 694 11. Mysore, K. *et al.* Yeast interfering RNA larvicides targeting neural genes induce high rates of
695 *Anopheles* larval mortality. *Malar J* **16**, 1–12 (2017).
- 696 12. Lopez, S. B. G. *et al.* RNAi-based bioinsecticide for *Aedes* mosquito control. *Scientific Reports* 2019
697 9:1 **9**, 1–13 (2019).
- 698 13. Taracena, M. L., Hunt, C. M., Benedict, M. Q., Pennington, P. M. & Dotson, E. M. Downregulation of
699 female doublesex expression by oral-mediated RNA interference reduces number and fitness of
700 *Anopheles gambiae* adult females. *Parasit Vectors* **12**, 1–11 (2019).
- 701 14. Ding, J. *et al.* Engineered Gut Symbiotic Bacterium-Mediated RNAi for Effective Control of *Anopheles*
702 Mosquito Larvae. *Microbiol Spectr* **11**, (2023).
- 703 15. Dhandapani, R. K., Gurusamy, D., Howell, J. L. & Palli, S. R. Development of CS-TPP-dsRNA
704 nanoparticles to enhance RNAi efficiency in the yellow fever mosquito, *Aedes aegypti*. *Sci Rep* **9**,
705 (2019).


- 706 16. Giesbrecht, D. *et al.* RNA Interference Is Enhanced by Knockdown of Double-Stranded RNases in the
707 Yellow Fever Mosquito *Aedes aegypti*. *Insects* 2020, Vol. 11, Page 327 **11**, 327 (2020).
- 708 17. Mancini, M. V. *et al.* Paratransgenesis to control malaria vectors: a semi-field pilot study. *Parasit*
709 *Vectors* **9**, (2016).
- 710 18. Riehle, M. A., Moreira, C. K., Lampe, D., Lauzon, C. & Jacobs-Lorena, M. Using bacteria to express
711 and display anti-Plasmodium molecules in the mosquito midgut. *Int J Parasitol* **37**, 595–603 (2007).
- 712 19. Wang, S. *et al.* Fighting malaria with engineered symbiotic bacteria from vector mosquitoes. *Proc*
713 *Natl Acad Sci U S A* **109**, 12734–12739 (2012).
- 714 20. Wang, S. & Jacobs-Lorena, M. Paratransgenesis Applications: Fighting Malaria With Engineered
715 Mosquito Symbiotic Bacteria. *Arthropod Vector: Controller of Disease Transmission* **1**, 219–234
716 (2017).
- 717 21. Wiltshire, R. M. & Duman-Scheel, M. Advances in oral RNAi for disease vector mosquito research
718 and control. *Curr Opin Insect Sci* **40**, 18–23 (2020).
- 719 22. Timmons, L., Court, D. L. & Fire, A. Ingestion of bacterially expressed dsRNAs can produce specific
720 and potent genetic interference in *Caenorhabditis elegans*. *Gene* **263**, 103–112 (2001).
- 721 23. Sturm, Á., Saskoï, É., Tibor, K., Weinhardt, N. & Vellai, T. Highly efficient RNAi and Cas9-based auto-
722 cloning systems for *C. elegans* research. *Nucleic Acids Res* **46**, (2018).
- 723 24. Pompon, J. *et al.* Dengue subgenomic flaviviral RNA disrupts immunity in mosquito salivary glands to
724 increase virus transmission. *PLoS Pathog* **13**, e1006535 (2017).
- 725 25. Qiu, Y. *et al.* Flavivirus induces and antagonizes antiviral RNA interference in both mammals and
726 mosquitoes. *Sci Adv* **6**, (2020).
- 727 26. Slonchak, A. *et al.* Zika virus noncoding RNA suppresses apoptosis and is required for virus
728 transmission by mosquitoes. *Nature Communications* 2020 11:1 **11**, 1–14 (2020).
- 729 27. Gholamalipour, Y., Karunanayake Mudiyansele, A. & Martin, C. T. 3' end additions by T7 RNA
730 polymerase are RNA self-templated, distributive and diverse in character—RNA-Seq analyses.
731 *Nucleic Acids Res* **46**, 9253 (2018).
- 732 28. Nwokeoji, A. O. *et al.* Analysis of long dsRNA produced in vitro and in vivo using atomic force
733 microscopy in conjunction with ion-pair reverse-phase HPLC †. *Cite this: Analyst* **144**, 4985 (2019).
- 734 29. Grossman, T. H., Kawasaki, E. S., Punreddy, S. R. & Osburne, M. S. Spontaneous cAMP-dependent
735 derepression of gene expression in stationary phase plays a role in recombinant expression
736 instability. *Gene* **209**, 95–103 (1998).
- 737 30. Nielsen, B. L., Willis, V. C. & Lin, C. Y. Western blot analysis to illustrate relative control levels of the
738 lac and ara promoters in *Escherichia coli*. *Biochemistry and Molecular Biology Education* **35**, 133–
739 137 (2007).
- 740 31. Coon, K. L., Vogel, K. J., Brown, M. R. & Strand, M. R. Mosquitoes rely on their gut microbiota for
741 development. *Mol Ecol* **23**, 2727 (2014).

- 742 32. Romoli, O., Schönbeck, J. C., Hapfelmeier, S. & Gendrin, M. Production of germ-free mosquitoes via
743 transient colonisation allows stage-specific investigation of host–microbiota interactions. *Nat*
744 *Commun* **12**, (2021).
- 745 33. Xiao, R., Chun, L., Liu, J., Shawn, X. Z. & Correspondence, X. RNAi Interrogation of Dietary
746 Modulation of Development, Metabolism, Behavior, and Aging in *C. elegans*. *CellReports* **11**,
747 1123–1133 (2015).
- 748 34. Coon, K. L. *et al.* Bacteria-mediated hypoxia functions as a signal for mosquito development. *Proc*
749 *Natl Acad Sci U S A* **114**, E5362–E5369 (2017).
- 750 35. Tuanudom, R., Yurayart, N., Rodkhum, C. & Tiawsirisup, S. Diversity of midgut microbiota in
751 laboratory-colonized and field-collected *Aedes albopictus* (Diptera: Culicidae): A preliminary study.
752 *Heliyon* **7**, e08259 (2021).
- 753 36. Berhanu, A. *et al.* Isolation and identification of microflora from the midgut and salivary glands of
754 *Anopheles* species in malaria endemic areas of Ethiopia. *BMC Microbiol* **19**, 1–8 (2019).
- 755 37. Ranasinghe, K., Gunathilaka, N., Amarasinghe, D., Rodrigo, W. & Udayanga, L. Diversity of midgut
756 bacteria in larvae and females of *Aedes aegypti* and *Aedes albopictus* from Gampaha District, Sri
757 Lanka. *Parasit Vectors* **14**, (2021).
- 758 38. Oliveira, J. H. M. *et al.* Blood Meal-Derived Heme Decreases ROS Levels in the Midgut of *Aedes*
759 *aegypti* and Allows Proliferation of Intestinal Microbiota. *PLoS Pathog* **7**, 1001320 (2011).
- 760 39. Gendrin, M. *et al.* Antibiotics in ingested human blood affect the mosquito microbiota and capacity
761 to transmit malaria. *Nature Communications* **6**:1 **6**, 1–7 (2015).
- 762 40. Olmo, R. P. *et al.* Control of dengue virus in the midgut of *Aedes aegypti* by ectopic expression of the
763 dsRNA-binding protein Loqs2. *Nature Microbiology* **2018 3:12** **3**, 1385–1393 (2018).
- 764 41. Airs, P. M., Kudrna, K. E., Lubinski, B., Phanse, Y. & Bartholomay, L. C. A Comparative Analysis of
765 RNAi Trigger Uptake and Distribution in Mosquito Vectors of Disease. (2023) doi:10.3390/insects.
- 766 42. Tassetto, M., Kunitomi, M. & Andino, R. Circulating Immune Cells Mediate a Systemic RNAi-Based
767 Adaptive Antiviral Response in *Drosophila*. *Cell* **169**, 314–325 (2017).
- 768 43. Martinson, V. G., Moy, J. & Moran, N. A. Establishment of characteristic gut bacteria during
769 development of the honeybee worker. *Appl Environ Microbiol* **78**, 2830–2840 (2012).
- 770 44. Strand, M. R. Composition and functional roles of the gut microbiota in mosquitoes. *Curr Opin Insect*
771 *Sci* **28**, 59 (2018).
- 772 45. Mehta, D. Highlight negative results to improve science. *Nature* (2019) doi:10.1038/D41586-019-
773 02960-3.
- 774 46. Schwarz, M. C. *et al.* Rescue of the 1947 Zika Virus Prototype Strain with a Cytomegalovirus
775 Promoter-Driven cDNA Clone. (2016) doi:10.1128/mSphere.00246-16.
- 776 47. Stapleford, K. A. *et al.* Whole-Genome Sequencing Analysis from the Chikungunya Virus Caribbean
777 Outbreak Reveals Novel Evolutionary Genomic Elements. (2016) doi:10.1371/journal.pntd.0004402.

- 778 48. Lequime, S., Fontaine, A., Ar Gouilh, M., Moltini-Conclois, I. & Lambrechts, L. Genetic Drift, Purifying
779 Selection and Vector Genotype Shape Dengue Virus Intra-host Genetic Diversity in Mosquitoes. *PLoS*
780 *Genet* **12**, e1006111 (2016).
- 781 49. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J*
782 *Mol Biol* **215**, 403–410 (1990).
- 783 50. Suzuki, Y. *et al.* Non-retroviral Endogenous Viral Element Limits Cognate Virus Replication in *Aedes*
784 *aegypti* Ovaries. *Current Biology* **30**, 3495-3506.e6 (2020).
- 785 51. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data.
786 <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- 787 52. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J*
788 **17**, 10–12 (2011).
- 789 53. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of
790 short DNA sequences to the human genome. *Genome Biol* **10**, 1–10 (2009).
- 791 54. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
792 (2009).
- 793 55. Bioconductor - Home. <https://www.bioconductor.org/>.
- 794


A

Cloning strategy




B

Target viral sequence







C




D

Bacterial colonization







A

T444T-ZIK1 (live bacteria)


B

Coverage of sRNAs


C

Length distribution of sRNAs


D

pUC18-ZIK1 (live bacteria)


E

Coverage of sRNAs


F

Length distribution of sRNAs


G



pUC18-ZIK1-short (live bacteria)

H


Coverage of sRNAs

I


Length distribution of sRNAs




F Viral titers - Day 7 post-infection




G Viral titers - Day 14 post-infection




H Coverage of sRNAs



I Length distribution of sRNAs



J Coverage of sRNAs



K Length distribution of sRNAs

