



**HAL**  
open science

## Clinical and genomic features of a *Listeria monocytogenes* fatal case of meningitis in Madagascar

Saïda Rasoanandrasana, Mamitina Alain Noah Rabenandrasana, Lucia Mélanie Ravaoharisoa, Narindra Randrianaivo, Vonintsoa Lalaina Rahajamanana, Zafindrasoa Domoina Rakotovao-Ravahatra, Alexandra Moura, Marc Lecuit, Andriamiadana Luc Rakotovao

### ► To cite this version:

Saïda Rasoanandrasana, Mamitina Alain Noah Rabenandrasana, Lucia Mélanie Ravaoharisoa, Narindra Randrianaivo, Vonintsoa Lalaina Rahajamanana, et al.. Clinical and genomic features of a *Listeria monocytogenes* fatal case of meningitis in Madagascar. *Access Microbiology*, 2024, 6 (6), pp.000764.v3. 10.1099/acmi.0.000764.v2 . pasteur-04632587

**HAL Id: pasteur-04632587**

**<https://pasteur.hal.science/pasteur-04632587>**

Submitted on 2 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Clinical and genomic features of a *Listeria monocytogenes* fatal case of meningitis in Madagascar

Saïda Rasoanandrasana<sup>1,†</sup>, Mamitina Alain Noah Rabenandrasana<sup>2,\*†</sup>, Lucia Mélanie Ravaoharisoa<sup>1</sup>, Narindra Randrianaivo<sup>3</sup>, Vonintsoa Lalaina Rahajamanana<sup>4</sup>, Zafindraso Domoïna Rakotovoao-Ravahatra<sup>1</sup>, Alexandra Moura<sup>5,6</sup>, Marc Lecuit<sup>5,6,7</sup> and Andriamiadana Luc Rakotovoao<sup>1</sup>

## Abstract

Listeriosis constitutes a significant public health threat due to its high mortality rate. This study investigates the microbiological and genomic characteristics of *Listeria monocytogenes* isolates in Madagascar, where listeriosis is a notifiable disease. The analysis focuses on a fatal case of meningeal listeriosis in a 12-year-old child. Genomic analysis revealed a novel cgMLST type (L2-SL8-ST8-CT11697; CC8, serogroup lia) with typical virulence and antibiotic resistance profiles. These isolates, unique to Madagascar, formed an independent clade in the phylogenetic tree. This study presents the first genomic characterization of *Listeria* isolates in Madagascar, highlighting the necessity of ongoing genomic surveillance to strengthen listeriosis prevention and control strategies in the region.

## Impact Statement

This study's significant findings on *Listeria monocytogenes* infections in Madagascar highlight the elevated risk among specific populations, particularly pregnant women from sub-Saharan Africa and the Maghreb region. The research reveals previously undocumented manifestations of the disease, expanding our understanding of its clinical presentation. Crucially, genomic analysis has identified a novel strain unique to Madagascar, providing valuable insights for tracing the pathogen's origins and transmission. The presence of specific virulence genes and antibiotic resistance profiles underscores the need for tailored prevention strategies. This pioneering genomic characterization lays the groundwork for targeted interventions and enhances public health initiatives, ultimately contributing to the prevention of listeriosis in the region.

*Access Microbiology* is an open research platform. Pre-prints, peer review reports, and editorial decisions can be found with the online version of this article. Received 05 January 2024; Accepted 04 June 2024; Published 27 June 2024

**Author affiliations:** <sup>1</sup>Bacteriology Laboratory, CHU Joseph Raseta Befelatanana, Antananarivo, Madagascar; <sup>2</sup>Experimental Bacteriology Unit, Institut Pasteur of Madagascar, Antananarivo, Madagascar; <sup>3</sup>Pediatric Service, CHU Joseph Raseta Befelatanana, Antananarivo, Madagascar; <sup>4</sup>Bacteriology Laboratory, Centre Hospitalier Universitaire Mère Enfant Tsaralàlana, Antananarivo, Madagascar; <sup>5</sup>Institut Pasteur, Biology of Infection Unit, Université Paris Cité, Inserm U1117, Paris, 75015, France; <sup>6</sup>Institut Pasteur, National Reference Centre and WHO Collaborating Centre *Listeria*, Paris, France; <sup>7</sup>Division of Infectious Diseases and Tropical Medicine, Institut Imagine, APHP, Necker-Enfants Malades University Hospital, Paris, France.

\*Correspondence: Mamitina Alain Noah Rabenandrasana, rabalainnoah@gmail.com

**Keywords:** fatal case; genomic characterization; listeriosis; Madagascar; meningitis.

**Abbreviations:** BDAL, Bruker Daltonics Library; BIGSdb, bacterial isolate genome sequence database; cgMLST, core genome multi-locus sequence typing; CSF, cerebrospinal fluid; IVD, in vitro diagnostic; MALDI-TOF MS, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry; MLST, multi-locus sequence typing; MSP, main spectral profiles; PCR, polymerase chain reaction. The assemblies and SRA data are deposited on NCBI under the bioproject PRJNA1032442.

†These authors contributed equally to this work

A supplementary table is available with the online version of this article.

000764.v3 © 2024 The Authors



This is an open-access article distributed under the terms of the Creative Commons Attribution License. The Microbiology Society waived the open access fees for this article.

## DATA SUMMARY

The assemblies and SRA data are deposited on NCBI under the bioproject PRJNA1032442 (all accession numbers are listed in the supplementary table, available with the online version of this article).

## INTRODUCTION


Listeriosis is one of the most serious food-borne illnesses, caused by the Gram-positive bacteria *Listeria monocytogenes*. It is a relatively rare disease, with 0.1 to 10 cases per million inhabitants per year, depending on the country and region. Although the number of listeriosis cases is low, the high mortality rate associated with this infection makes it a worrying public health issue [1]. In pregnant women, listeriosis is higher in populations from sub-Saharan Africa and the Maghreb than in those from Latin America and the Caribbean. Among the 107 cases of maternal-neonatal listeriosis, 35 (33%) originated from the Maghreb or sub-Saharan Africa, a rate three times higher than that observed in the general pregnant population in France in 2010, as indicated by national records (88 000 [11%] out of 832 000 women,  $P < 0.0001$ ) [2].

Listeriosis is a notifiable disease in Madagascar. A recent study [3] documented the case of a 75-year-old man with an aortic bioprosthesis, who was admitted with polyarthritis in an afebrile state. Blood cultures confirmed the presence of *L. monocytogenes*, leading to the diagnosis of *Listeria* endocarditis and spondylodiscitis, both uncommon manifestations of listeriosis. The patient showed improvement following antibiotic therapy [3]. Here we describe a case of invasive listeriosis at the CHU Befelatanana in Madagascar.

## CASE PRESENTATION

A 12-year-old patient was admitted for meningeal syndrome, with a history of hyperthermic convulsive seizure at age 3. The patient is the first child in a sibling group of three, with an up-to-date immunization schedule. There was no similar case in the family. We received a sample of the cerebrospinal fluid and blood cultures. Direct examination of the CSF revealed hyperleukocytosis with a predominance of polynuclear neutrophils, hyperproteinorrachia, and hypoglycorrachia. Solid and liquid cultures were positive for a Gram-positive bacillus, which was identified as *L. monocytogenes* using the BioFire FilmArray Meningitis/Encephalitis (ME) Panel (Biomérieux, Marcy-l'Étoile, France). Concurrently, the blood culture also grew a Gram-positive bacillus with a 48 h growth period. Unfortunately, despite appropriate treatment according to the Common Infections Guidelines in Madagascar, 2019 edition [4] with ceftriaxone, gentamicin, and vancomycin, the patient did not survive. The antimicrobial susceptibility testing (AST) is not performed at the time of patient admission due to cost constraints for patients, especially considering the urgency of promptly treating bacterial meningitis, particularly in children. We used methods such as Gram-staining and Matrix-assisted laser desorption/ionization – time of flight mass spectrometry (MALDI-TOF MS) using the MALDI-TOF Biotyper System with the BDAL spectra database containing 8 468 MSPs, and the IVD containing 8 326 MSPs, for bacterial identification and the extraction method with 70% formic acid and acetonitrile. The results indicated the presence of Gram-positive bacteria (Fig. 1), with the identification as *L. monocytogenes* with a score of  $> 2.0$  (Table 1). We notified the appropriate authorities; however, as of our current understanding, the origin remains undetermined.


Isolate characterization was performed by whole genome sequencing with Illumina technology and genomic analysis with TORMES pipeline [5] and VirulenceFinder [6, 7]. PCR-serogrouping, multilocus sequencing typing (MLST), core genome MLST (cgMLST) and virulence and resistance profiles were performed using BIGSdb-*Listeria* [6, 8–14]. We conducted a phylogenetic analysis using core genome alignments that were not purged for genetic recombination, obtained with Parsnp



**Fig. 1.** Gram-staining of *Listeria monocytogenes* strains (a) isolated from cerebrospinal fluid (CSF) and blood cultures (b).

**Table 1.** Identification results on the MALDI-TOF MS for the two strains of *Listeria monocytogenes*. If the score is between: 2.00–3.00, the identification is performed with a high confidence level; 1.70–1.99, the identification is performed with a low confidence level; 0.00–1.69, no identification is attributed to the organism

Strains origins	Corresponding reference mass spectra	Score	NCBI identifier
Cerebrospinal fluid	<i>Listeria monocytogenes</i> CCUG 32843 CCUG	2.14	1639
	<i>Listeria monocytogenes</i> CCUG 59664 CCUG	2.14	1639
	<i>Listeria monocytogenes</i> CCUG 31527 CCUG	2.09	1639
	<i>Listeria monocytogenes</i> CCUG 35751 CCUG	2.08	1639
	<i>Listeria monocytogenes</i> DSM 20600T DSM_1	2.07	1639
	<i>Listeria monocytogenes</i> CCUG 61052 CCUG	2.07	1639
	<i>Listeria monocytogenes</i> Mb19348_1 CHB_1	2.06	1639
	<i>Listeria monocytogenes</i> CCUG 49753 CCUG	2.03	1639
	<i>Listeria monocytogenes</i> ser4B ATCC 19115 THL_1	2.02	1639
	<i>Listeria monocytogenes</i> CCUG 32964A CCUG	2.02	1639
Blood culture	<i>Listeria monocytogenes</i> CCUG 59664 CCUG	2.21	1639
	<i>Listeria monocytogenes</i> CCUG 32843 CCUG	2.12	1639
	<i>Listeria monocytogenes</i> CCUG 49753 CCUG	2.12	1639
	<i>Listeria monocytogenes</i> CCUG 35751 CCUG	2.11	1639
	<i>Listeria monocytogenes</i> CCUG 31527 CCUG	2.1	1639
	<i>Listeria monocytogenes</i> CCUG 61052 CCUG	2.1	1639
	<i>Listeria monocytogenes</i> CCUG 33548 CCUG	2.09	1639
	<i>Listeria monocytogenes</i> CCUG 32964A CCUG	2.09	1639
	<i>Listeria monocytogenes</i> Mb19348_1 CHB_1	2.08	1639
	<i>Listeria monocytogenes</i> CCUG 61052 CCUG	2.03	1639



**Fig. 2.** Phylogenetic tree based on the core genome alignment of Malagasy isolates and 100 publicly available CC8 sublineage SL8 genomes from BIGSdb-Listeria (as of 9 September 2023). From the innermost to the outermost circle, the first circle represents the distribution by ST, the second represents the distribution by continent, and the last one represents the isolation sources (b) with a zoom on the clades containing strains from Madagascar (a).

[15], incorporating publicly available strains from the BIGSdb-*Listeria* database belonging to clonal complex CC8 (as of 9 September 2023).

The two isolates of *L. monocytogenes* isolates belonged to the same cgMLST type (L2-SL8-ST8-CT11697; CC8, serogroup Iia), which is new in BIGSdb-*Listeria* database. The typical *L. monocytogenes* virulence genes were present and included the pathogenic island LIPI-1 [16], as well the *inlAB* operon necessary for gut and brain invasion [16]. Intrinsic antibiotic resistance (*fosX*, *lin*, *norB*, *sul*, *pbp*) genes were present, and no acquired resistance was detected [16]. Both isolates carried the stress island SSI-1 [16], which favours adaptation to low pH and high salt concentration [16] and LGI-3 [16].

This new type differs by 12 allelic differences from L2-SL8-ST8-CT750, which has been found in Europe (Denmark, Poland, Estonia) and Latin America (Chile). Our phylogenetic analysis (Fig. 2) confirms that the Malagasy isolates exhibit a high degree of genetic similarity and form an independent clade, distinct from those originating in other countries.

## DISCUSSION

To the best of our knowledge, this work represents the first genomic characterization of *Listeria* isolates from Madagascar. Further sequencing efforts will help to identify sources of contamination and improve listeriosis surveillance and prevention efforts in our region. The discovery of a novel strain specific to Madagascar highlights the uniqueness of *Listeria* circulating in the local population. This groundbreaking research sets the stage for further sequencing efforts, aiming to identify contamination sources and enhance listeriosis surveillance and prevention strategies within Madagascar. The findings underscore the importance of integrating genomic data into public health initiatives, enabling targeted interventions, swift outbreak responses, and tailored patient care. Ultimately, this study serves as a crucial foundation for comprehensive research and collaborative efforts to mitigate the impact of listeriosis in Madagascar.

### Funding information

This study was supported by internal funding. The authors have received no specific grant from any funding agency.

### Acknowledgements

We thank the National Reference Centre and WHO Collaborating Centre *Listeria* at Institut Pasteur in Paris (France) and the Experimental Bacteriology Unit at the Institut Pasteur of Madagascar (Madagascar) for their collaboration and support. We also acknowledge the Mutualised Microbiology Platform (P2M) at Institut Pasteur in Paris (France) for conducting the complete genome sequencing of the strains.

### Author contributions

S.R.: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Writing - original draft, Writing - review & editing. M.A.N.R.: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Writing - original draft, Writing - review & editing. L.M.R.: Investigation, Methodology, Writing - original draft. N.R.: Investigation, Methodology, Writing - original draft. V.L.R.: Investigation, Methodology, Writing - original draft. Z.D.R.R.: Investigation, Methodology, Writing - original draft. A.M.: Data curation, Formal analysis, Methodology, Validation, Writing - original draft, Writing - review & editing. M.L.: Funding acquisition, Supervision, Validation, Writing - original draft, Writing - review & editing. A.L.R.: Conceptualization, Funding acquisition, Supervision, Validation, Writing - original draft, Writing - review & editing. The corresponding author has provided authorship verification on behalf of co-author, N. R.

### Conflicts of interest

The authors declare that there are no conflicts of interest.

### Consent to publish

Consent for publishing this case report has been obtained from the parent, given the patient's age being under 18 years.

### References

- World Health Organization African Region. Listeriosis; 2023. <https://www.afro.who.int/fr/health-topics/listeriose>
- Charlier C, Perrodeau É, Leclercq A, Cazenave B, et al, Group M. Clinical features and prognostic factors of listeriosis: the MONALISA national prospective cohort study. *Lancet Infect Dis* 2017;17:510–519.
- Randrianarisoa RMF, Ramanandafy H, Mania A, Benelhadj AD, Clément M, et al. *Listeria* endocarditis and spondylodiscitis: A case report and review of the literature. *Clin Case Rep* 2022;10:e05899.
- F.M. Société de Pathologie Infectieuse de Madagascar (SPIM), Société Malgache de Biologie (SOMABIO), Société Malgache de Pédiatrie (SOMAPED) and F. de Médecine. In: *Guide Des Infections Courantes à Madagascar*, 2019th edn. Antananarivo, Madagascar, 2019.
- Quijada NM, Rodríguez-Lázaro D, Eiros JM, Hernández M. TORMES: an automated pipeline for whole bacterial genome analysis. *Bioinformatics* 2019;35:4207–4212.
- Moura A, Tourdjman M, Leclercq A, Hamelin E, Laurent E, et al. Real-time whole-genome sequencing for surveillance of *Listeria monocytogenes*, France. *Emerg Infect Dis* 2017;23:1462–1470.
- Malberg Tetzschner AM, Johnson JR, Johnston BD, Lund O, Scheutzd F. In silico genotyping of *Escherichia coli* isolates for sequencing data. *J Clin Microbiol* 2020;58:e01269–20.
- Moura A, Criscuolo A, Pouseele H, Maury MM, Leclercq A, et al. Whole genome-based population biology and epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol* 2016;2:16185.
- Maury MM, Tsai Y, Charlier C, Touchon M, Chenal-Francisque V, et al. Uncovering *Listeria monocytogenes* hypervirulence by harnessing its biodiversity. *Nat Genet* 2016;48:308–313.
- Chenal-Francisque V, Lopez J, Cantinelli T, Caro V, Tran C, et al. Worldwide distribution of major clones of *Listeria monocytogenes*. *Emerg Infect Dis* 2011;17:1110–1112.
- Ragon M, Wirth T, Hollandt F, Lavenir R, Lecuit M, et al. A new perspective on *Listeria monocytogenes* evolution. *PLoS Pathog* 2008;4:e1000146.

12. Doumith M, Buchrieser C, Glaser P, Jacquet C, Martin P. Differentiation of the major *Listeria monocytogenes* serovars by multiplex PCR. *J Clin Microbiol* 2004;42:3819–3822.
13. Jolley KA, Maiden MCJ. BIGSdb: scalable analysis of bacterial genome variation at the population level. *BMC Bioinformatics* 2010;11:595.
14. Maiden MCJ, Bygraves JA, Feil E, Morelli G, Russell JE, *et al.* Multi-locus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc Natl Acad Sci U S A* 1998;95:3140–3145.
15. Treangen TJ, Ondov BD, Koren S, Phillippy AM. The harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol* 2014;15:1–15.
16. Moura M, Leclercq A, Valès G, Tessaud-Rita N, Bracq-Dieye H, *et al.* Phenotypic and genotypic antimicrobial resistance of *Listeria monocytogenes* in France. 2023. DOI: 10.2139/ssrn.4401708.

**The Microbiology Society is a membership charity and not-for-profit publisher.**

**Your submissions to our titles support the community – ensuring that we continue to provide events, grants and professional development for microbiologists at all career stages.**

**Find out more and submit your article at [microbiologyresearch.org](https://microbiologyresearch.org)**