



HAL
open science

Complete genome sequence of a vampire bat-related rabies virus obtained by metagenomics from a patient with encephalitis of unknown etiology, French Guiana

Véronique Hourdel, Charlotte Balière, Jessica Vanhomwegen, Angela Brisebarre, Quentin Grassin, Jean-Claude Manuguerra, Hatem Kallel, Magalie Demar, Laurent Dacheux, Valérie Caro

► To cite this version:

Véronique Hourdel, Charlotte Balière, Jessica Vanhomwegen, Angela Brisebarre, Quentin Grassin, et al.. Complete genome sequence of a vampire bat-related rabies virus obtained by metagenomics from a patient with encephalitis of unknown etiology, French Guiana. *Microbiology Resource Announcements*, 2024, 13 (11), pp.e00514-24. 10.1128/mra.00514-24. pasteur-04724112

HAL Id: pasteur-04724112

<https://pasteur.hal.science/pasteur-04724112v1>

Submitted on 7 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Complete genome sequence of a vampire bat-related rabies virus obtained by metagenomics from a patient with encephalitis of unknown etiology, French Guiana

Véronique Hourdel,¹ Charlotte Balière,¹ Jessica Vanhomwegen,¹ Angela Brisebarre,¹ Quentin Grassin,¹ Jean-Claude Manuguerra,¹ Hatem Kallel,² Magalie Demar,³ Laurent Dacheux,¹ Valérie Caro¹

AUTHOR AFFILIATIONS See affiliation list on p. 3.

ABSTRACT We report the complete genome sequence of a rabies virus obtained by direct metagenomics from the cerebellum of a gold panner who died of unknown encephalitis in French Guiana. Phylogenetic analysis exhibited a close genetic relationship with vampire bat-related isolates, confirming the second case of human rabies identified in this territory.

KEYWORDS rabies, French Guiana, complete genome, metagenomics, encephalitis, diagnosis, Illumina, next generation sequencing, bat, human

Rabies remains a major but still neglected tropical disease, with an estimated number of 59,000 fatal human cases worldwide each year (1). Rabies virus (RABV) is the principal etiological agent, belonging to the species *Rabies lyssavirus* within the genus *Lyssavirus*, family *Rhabdoviridae* (order *Mononegavirales*) (2). Apart from domestic dogs, other mammals such as bats can act as natural hosts and vectors. In French Guiana, an overseas territory localized in the northern part of South America, rabies transmitted by hematophagous bats (mainly the *Desmodus rotundus* species) remains endemic. Regular cases are detected in cattle, domestic carnivores, or even bats (3, 4). In 2008, the first confirmed human case of vampire rabies was detected (5). We describe here the RABV genome sequence of the second human Guyanese case, whose diagnosis was based on direct metagenomics.

A post-mortem cerebellum biopsy (lab ref 2024-6-10) was collected from a deceased patient presenting with encephalitis of unknown etiology in late February 2024 in Cayenne Hospital, French Guiana. This patient first developed symptoms while working on an illegal gold mining site in the Amazon rainforest. Nucleic acids were extracted with a DNA/RNA Microprep plus kit (Zymo Research). Total RNA was used for reverse transcription using random hexamer primers and the SuperScript IV First-Strand cDNA Synthesis kit (ThermoFisher Scientific) and amplified with the Quantitect Whole Transcriptome kit (Qiagen). All steps were performed according to the manufacturer's instructions. A sequencing-ready library was prepared using the TruSeq DNA PCR-Free prep kit (Illumina) and sequenced on a NextSeq2000 system (Illumina) generating nearly 14.4 million of 300-bp single-end length reads (88.2% of unique reads and 12.8% of duplicate reads).

Nearly 77 million contigs were generated and identified against the RVDB database (v25.0) using Microseek, retaining only contigs with a length of 100 nucleotides (6). Among them, a contig (9,010 nt) related to *Lyssavirus rabies* species was identified. A second dedicated workflow built on Galaxy@Pasteur (a Galaxy platform hosted by Institut Pasteur) was subsequently performed, combining cleaning, *de novo* assembly, and mapping (CLC Assembly Cell, Qiagen, v5.1.10) (7–10). A final consensus was

Editor Jelle Matthijnsens, Katholieke Universiteit, Leuven, Belgium

Address correspondence to Laurent Dacheux, laurent.dacheux@pasteur.fr, or Valérie Caro, valerie.caro@pasteur.fr.

Véronique Hourdel and Charlotte Balière contributed equally to this article. Author order was determined in order of decreasing seniority.

The authors declare no conflict of interest.

Received 23 May 2024

Accepted 17 September 2024

Published 4 October 2024

Copyright © 2024 Hourdel et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).




FIG 1 Phylogenetic analysis of the RABV strain (2024-6-10) obtained from the cerebellum of a deceased patient presenting with encephalitis of unknown etiology in French Guiana and different representative American strains. The tree was based on the multiple sequence alignment corresponding to the concatenation of the five ORFs (10,815–10,818 nt), performed with ClustalW (v2.1) (14) implemented in Galaxy@Pasteur (9). The tree was constructed using the ML approach based on the generalized time-reversible model proportion of invariable sites plus gamma-distributed rate heterogeneity (GTR+I+ Γ 4) utilizing Subtree Prune-and-Regraft (SPR) branch-swapping, as estimated in PhyML 3.0 (12) with Smart Model Selection, also implemented in Galaxy@Pasteur. The robustness of individual nodes was estimated using 100 bootstrap replicates. All tools were run with default parameters unless otherwise specified. The ML phylogenetic tree was visualized with FigTree (<http://tree.bio.ed.ac.uk/>). Only bootstrap values ≥ 80 are indicated for the major branches. The scale bar indicates nucleotide substitutions per site. Each sequence denomination contains the GenBank accession number, the nature of the infected species, and the country and year of isolation, where applicable. Rabies virus isolates associated with vampire bats are indicated by an open circle. The isolate (2024_6_10_RABV_GUY) described in this study is indicated in bold, with a black circle. RAC-SK, raccoon–skunk (rabies virus associated with the raccoon and skunk host animals).

generated with the nearly 8 million reads retained and manually edited using SnapGene (v7.2) before visualization using Tablet (11).

This complete RABV genome was 11,922 nucleotides in length, with 294.80 \times coverage on average and 18,078 (0.22%) mapped reads. The five canonical genes were present, encoding the nucleoprotein (N, 1,353 nt, 450 aa), phosphoprotein (P, 894 nt, 297 aa), matrix protein (M, 609 nt, 202 aa), glycoprotein (G, 1,575 nt, 524 aa), and RNA polymerase (L, 6,387 nt, 2,128 aa). The sequences of leader and trailer extremities were complete, with 58 and 69 nucleotides in length, respectively (checking done after alignment with genetically close and available complete genomes). The transcription initiation signal AACA and the transcription termination polyadenylation (TTP) sequence

TGA7 were observed for all the genes, except for the G gene, which presented the AGA7 motif for TTP.

Maximum likelihood (ML) phylogenetic analysis was performed on the five concatenated open reading frames (ORFs) with different representative American strains using PhyML (12), and genetic analysis confirmed that the virus clustered together with vampire bat-related isolates within the Bat Clade (Fig. 1) (13).

ACKNOWLEDGMENTS

We thank the other members of the Laboratory for Urgent Response to Biological Threats (CIBU) for their technical help, especially Séverine Matheus for her network. We are also grateful to the Biomics Core Facility of Institut Pasteur for access to their facilities.

This work was funded through the convention with the Direction Générale de la Santé (for CIBU), with the support of the DURABLE EU4Health program under grant agreement 101102733 (for ERI unit).

AUTHOR AFFILIATIONS

¹Institut Pasteur, Université Paris Cité, Laboratory for Urgent Response to Biological Threats (CIBU), Environment and Infectious Risks (ERI) Unit, Paris, France

²Service de Réanimation Polyvalente, Centre Hospitalier de Cayenne, Cayenne, French Guiana

³Laboratoire de Biologie Médicale, Centre Hospitalier de Cayenne, Cayenne, French Guiana

AUTHOR ORCID*s*

Laurent Dacheux  <http://orcid.org/0000-0002-4401-3706>

Valérie Caro  <http://orcid.org/0000-0001-8477-5708>

AUTHOR CONTRIBUTIONS

Véronique Hourdel, Data curation, Formal analysis, Writing – original draft, Writing – review and editing | Charlotte Balière, Data curation, Formal analysis, Methodology, Writing – original draft, Writing – review and editing | Jessica Vanhomwegen, Methodology, Supervision, Validation, Writing – review and editing | Angela Brisebarre, Formal analysis, Writing – review and editing | Quentin Grassin, Formal analysis, Writing – review and editing | Jean-Claude Manuguerra, Funding acquisition, Writing – review and editing | Hatem Kallel, Investigation, Resources, Writing – review and editing | Magalie Demar, Investigation, Resources, Writing – original draft, Writing – review and editing | Laurent Dacheux, Conceptualization, Data curation, Formal analysis, Funding acquisition, Methodology, Project administration, Supervision, Validation, Writing – original draft, Writing – review and editing | Valérie Caro, Conceptualization, Methodology, Supervision, Validation, Writing – original draft, Writing – review and editing

DATA AVAILABILITY

The complete genome sequence of the vampire bat-related rabies virus from French Guiana was deposited in GenBank under accession number [PP764242](#) and BioProject accession number [PRJNA1108161](#).

ETHICS APPROVAL

The lab investigations described in this article were carried out as part of the diagnosis activities of the Laboratory for Urgent Response to Biological Threats (CIBU). After the patient's death, the relatives were informed about the use of the post-mortem cerebellum biopsy for genetic characterization of the rabies strain, after the diagnosis had been established, and consent was obtained.

REFERENCES

1. Hampson K, Coudeville L, Lembo T, Sambo M, Kieffer A, Attlan M, Barrat J, Blanton JD, Briggs DJ, Cleaveland S, et al. 2015. Estimating the global burden of endemic canine rabies. *PLoS Negl Trop Dis* 9:4. <https://doi.org/10.1371/journal.pntd.0003709>
2. Walker PJ, Freitas-Astúa J, Bejerman N, Blasdell KR, Breyta R, Dietzgen RG, Fooks AR, Kondo H, Kurath G, Shi PL, Stone M, Tesh DM, Tordo RB, Vasilakis N, Whitfield N. 2022. ICTV virus taxonomy profile: *Rhabdoviridae* 2022. *J Gen Virol* 103. <https://doi.org/10.1099/jgv.0.001689>
3. Lavergne A, Darcissac E, Bourhy H, Tirera S, de Thoisy B, Lacoste V. 2016. Complete genome sequence of a vampire bat rabies virus from French Guiana. *Genome Announc* 4:e00188-16. <https://doi.org/10.1128/genomeA.00188-16>
4. de Thoisy B, Bourhy H, Delaval M, Pontier D, Dacheux L, Darcissac E, Donato D, Guidez A, Larrous F, Lavenir R, Salmier A, Lacoste V, Lavergne A. 2016. Bioecological drivers of rabies virus circulation in a neotropical bat community. *PLoS Negl Trop Dis* 10:e0004378. <https://doi.org/10.1371/journal.pntd.0004378>
5. Meynard J-B, Flamand C, Dupuy C, Mahamat A, Eltges F, Queuche F, Renner J, Fontanella J-M, Hommel D, Dussart P, Grangier C, Djossou F, Dacheux L, Goudal M, Berger F, Ardillon V, Krieger N, Bourhy H, Spiegel A. 2012. First human rabies case in French Guiana, 2008: epidemiological investigation and control. *PLoS Negl Trop Dis* 6:e1537. <https://doi.org/10.1371/journal.pntd.0001537>
6. Pérot P, Bigot T, Temmam S, Regnault B, Eloit M. 2022. Microseek: a protein-based metagenomic pipeline for virus diagnostic and discovery. *Viruses* 14:1990. <https://doi.org/10.3390/v14091990>
7. Luo D-S, Li B, Shen X-R, Jiang R-D, Zhu Y, Wu J, Fan Y, Bourhy H, Hu B, Ge X-Y, Shi Z-L, Dacheux L. 2021. Characterization of novel rhabdoviruses in Chinese bats. *Viruses* 13:64. <https://doi.org/10.3390/v13010064>
8. Pallandre L, Luo D, Feuvrier C, Liefbrig F, Pozet F, Dacheux L, Bigarré L. 2020. Revisiting the classification of percid perhabdoviruses using new full-length genomes. *Viruses* 12:649. <https://doi.org/10.3390/v12060649>
9. Mareuil F, D-A O, Ménager H. 2017. A public galaxy platform at pasteur used as an execution engine for web services. *F1000Res* 1030. <https://doi.org/10.7490/f1000research.1114334.1>
10. Dacheux L, Dommergues L, Chouanibou Y, Doméon L, Schuler C, Bonas S, Luo D, Maufrais C, Cetre-Sossah C, Cardinale E, Bourhy H, Métras R. 2019. Co-circulation and characterization of novel African arboviruses (genus *Ephemerovirus*) in cattle, Mayotte island, Indian Ocean, 2017. *Transbound Emerg Dis* 66:2601–2604. <https://doi.org/10.1111/tbed.13323>
11. Milne I, Stephen G, Bayer M, Cock PJA, Pritchard L, Cardle L, Shaw PD, Marshall D. 2013. Using Tablet for visual exploration of second-generation sequencing data. *Brief Bioinform* 14:193–202. <https://doi.org/10.1093/bib/bbs012>
12. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59:307–321. <https://doi.org/10.1093/sysbio/syq010>
13. Troupin C, Dacheux L, Tanguy M, Sabeta C, Blanc H, Bouchier C, Vignuzzi M, Duchene S, Holmes EC, Bourhy H. 2016. Large-scale phylogenomic analysis reveals the complex evolutionary history of rabies virus in multiple carnivore hosts. *PLOS Pathog* 12:e1006041. <https://doi.org/10.1371/journal.ppat.1006041>
14. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007. Clustal W and clustal X version 2.0. *Bioinformatics* 23:2947–2948. <https://doi.org/10.1093/bioinformatics/btm404>