

### Horseshoe bats from Southeast Asia host a high diversity of Sarbecoviruses, including close ancestors of SARS-CoV-2

Sarah Temmam, Khamsing Vongphayloth, Tran Cong Tu, Béatrice Regnault, Massimiliano Bonomi, Eduard Baquero, Cécile Hérate, Sandie Munier, Delphine Chrétien, Flora Donati, et al.

#### ▶ To cite this version:

Sarah Temmam, Khamsing Vongphayloth, Tran Cong Tu, Béatrice Regnault, Massimiliano Bonomi, et al.. Horseshoe bats from Southeast Asia host a high diversity of Sarbecoviruses, including close ancestors of SARS-CoV-2. 8th World One Health Congress, Sep 2024, Cape Town / Le Cap, South Africa. pasteur-04727508

#### HAL Id: pasteur-04727508 https://pasteur.hal.science/pasteur-04727508v1

Submitted on 9 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Horseshoe bats from Southeast Asia host a high diversity of Sarbecoviruses, including close ancestors of SARS-CoV-2.

**Sarah Temmam**<sup>1</sup>, Khamsing Vongphayloth<sup>2</sup>, Tran Cong Tu<sup>3</sup>, Béatrice Regnault<sup>1</sup>, Massimiliano Bonomi<sup>4</sup>, Eduard Baquero<sup>5</sup>, Cécile Hérate<sup>6</sup>, Sandie Munier<sup>7</sup>, Flora Donati<sup>8</sup>, Delphine Chrétien<sup>1</sup>, Thomas Bigot<sup>9</sup>, Léa Vendramini<sup>1</sup>, Philippe Pérot<sup>1</sup>, Yasaman Karami<sup>4</sup>, Océane Dehan<sup>8</sup>, Faustine Amara<sup>7</sup>, Bounsavane Douangboubpha<sup>10</sup>, Daosavanh Sanamxay<sup>10</sup>, Vilakhan Xayaphet<sup>10</sup>, Phetphoumin Paphaphanh<sup>10</sup>, Somphavanh Somlor<sup>2</sup>, Khaithong Lakeomany<sup>2</sup>, Nothasin Phommavanh<sup>2</sup>, Mikael Attia<sup>7</sup>, Laurine Conquet<sup>11</sup>, Grégory Jouvion<sup>12</sup>, Juliana Pipoli Da Fonseca<sup>13</sup>, Thomas Cokelaer<sup>13</sup>, Tran Nhu Duong<sup>3</sup>, Tran Vu Phong<sup>3</sup>, Nguyen Thi Yen<sup>3</sup>, Hoang Ngoc Anh<sup>3</sup>, Tran Hai Son<sup>3</sup>, Pham Tuan Anh<sup>3</sup>, Vu Dinh Thong<sup>14</sup>, Francis Relouzat<sup>6</sup>, Thibaut Naninck<sup>6</sup>, Julien Lemaitre<sup>6</sup>, Nathalie Derreudre-Bosquet<sup>6</sup>, Quentin Pascal<sup>6</sup>, Michael Nilges<sup>4</sup>, Vincent Lacoste<sup>2</sup>, Xavier Montagutelli<sup>11</sup>, Vu Sinh Nam<sup>3</sup>, Roger Le Grand<sup>6</sup>, Félix A. Rey<sup>5</sup>, Sylvie van der Werf<sup>7</sup>, Paul T. Brey<sup>2</sup>, Marc Eloit<sup>1</sup>.

## contact: sarah.temmam@pasteur.fr / marc.eloit@pasteur.fr

1. Institut Pasteur, Université de Paris, Pathogen Discovery Laboratory, Paris, France. 8. Institut Pasteur, Université Paris Cité, National Reference Center for Respiratory Viruses, Paris, France. 2. Institut Pasteur du Laos, Vientiane, Lao People's Democratic Republic. 9. Bioinformatic and Biostatistic Hub - Computational Biology Department, Institut Pasteur, Université Paris Cité, Paris, France. 3. National Institute of Hygiene and Epidemiology, Hanoi 100000, Vietnam. 10. Faculty of Environmental Sciences, National University of Laos, Vientiane, Lao People's Democratic Republic. 11. Institut Pasteur, Université Paris Cité, Mouse Genetics Laboratory, Paris, France. 4. Institut Pasteur, Université de Paris, CNRS UMR 3528, Structural Bioinformatics Unit, Paris, France. 5. Institut Pasteur, Université de Paris, CNRS UMR 3569, Structural Virology Unit, Paris, France. 12. Ecole Nationale Vétérinaire d'Alfort, Unité d'Histologie et d'Anatomie Pathologique, Maisons-Alfort, France. 6. Center for Immunology of Viral, Auto-immune, Hematological and Bacterial Diseases (IMVA-HB/IDMIT), Université Paris-Saclay, Inserm, CEA, Fontenay-aux-Roses, France. 13. Biomics Platform, C2RT, Institut Pasteur, Université Paris Cité, Paris, France. 7. Institut Pasteur, Université de Paris, CNRS UMR 3569, Molecular Genetics of RNA Viruses Unit, Paris, France. 14. Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology (VAST), Hanoi 70072, Vietnam.

## Background

> Bats are **reservoirs** of many alpha- & beta-CoVs

## **Objectives**

- Understanding of the origins, natural history & dissemination of *Sarbecovirus*
- Identify **natural reservoirs** of *Sarbecovirus*
- > 4000 CoVs identified in 14 bat species Ruiz-Aravena et al., 2022 Nat. Rev. Microbiol. DOI: 10.1038/s41579-021-00652-2
- > SE Asia: huge richness of horseshoe bat species
- > Many outbreaks due to CoVs, Sarbecovirus at the Keusch et al., 2022 PNAS. DOI: 10.1073/pnas.2202871119 Zhou et al., 2021 Cell. DOI: 10.1016/j.cell.2021.06.008 origin of the last human pandemic Sanchez et al., 2022 Nat. Comm. DOI: 10.1038/s41467-022-31860-w
- Decipher **interspecies transmission** (bat-to-bat, bat-to-mammals, bat-to-humans)
- Study the **pathogenicity** of bat *Sarbecovirus* in human-relevant models
- Detect past CoV infections in humans 3.
  - Identify early circulation of *Sarbecovirus* before global spread



# Results

- Identification of **19 novel** *Sarbecovirus* in 6 *Rhinolophus* sp. bat species: (1.)
  - *R. siamensis* & *R. thomasi* in Vietnam
  - *R. marshalli, R. malayanus, R. pusillus* & R. *coelophyllus* in Laos
    - Rhinolophus sp. are the natural reservoirs of Sarbecovirus
    - > No Sarbecovirus identified in other bat species
    - **No furin site** in Spike protein of bat *Sarbecovirus*

#### **Recombination** and **cross-species transmission** are major drivers of virus evolution: (2.)

Concomitant circulation of  $\neq$  lineages in same cave  $\rightarrow$  risk of recombination





RtVN21-193 RtVN21-194 (#) RtVN21-197 **RtVN21-200** 

**RtVN21-192** 





- Same *Sarbecovirus* strain identified in  $\neq$  bat species  $\rightarrow$  cross-species transmission between bats that live in **sympatry**
- **Host range** of bat *Sarbecovirus* partly driven by ability to interact with ACE2 receptor: (3.
  - Genomic signature of Receptor-Binding Domain in Spike protein
    - Deletion in RBD = inability to interact with ACE2 (RsVN21-195, RtVN21-29 to -201 + BANAL-20-116/247)
    - **Conservation** of contact residues = **ability** to interact with ACE2 (**BANAL-20-52/103/236, -21-606**)
  - Interaction with hACE2 receptor
    - > No interaction & no entry in human cells (RsVN21-195, RtVN21-29 to -201 + BANAL-20-116/247)  $\rightarrow$  involvement of another receptor?
    - High affinity & ACE2-dependent entry in human cells (BANAL-20-52/103/236, -21-606) → risk of spillover?

**4.** Pathogenicity of bat *Sarbecovirus* in human-relevant models (e.g. BANAL-20-236):

- <u>Tg mice</u>: **no significant body weight loss** & virus load in lungs << SARS-CoV-2 (Wuhan strain) **cross-protection** against SARS-CoV-2 <sup>D</sup>614<sup>G</sup> challenge
- <u>NHP: no clinical signs & transient virus load in the enteric route compared to SARS-CoV-2</u>



## <u>Conclusion & perspectives</u>

- **Ecological factors** contributing to a possible emergence of *Sarbecovirus*:
  - 1. Bat species richness  $\rightarrow$  horseshoe bats are natural reservoirs
  - 2. Bat lifestyle  $\rightarrow$  colonies with mixed species more prone to exchange viruses through

- (5.) **Past** *Sarbecovirus* **exposure** of humans highly exposed to bat feces
  - Lack of serological detection in bat guano collectors (Laos)
  - Only cross-reactive Ab detected for COVID-19 patients



**Enterotropism** of bat Sarbecovirus > No/few circulation of bat Sarbecovirus in humans

→ Acquisition of a furin site & transmission by aerosol route in intermediate host?

- feco-oral route
- 3. Co-circulation of different virus strains in same ecosystem  $\rightarrow$  risk of recombination
- 4. Human exposure  $\rightarrow 7$  risk of contact with bat feces for guano collectors, religious communities, tourists visiting caves
- **Emergence of SARS-CoV-2**: questions to be solved
  - 1. Origin of furin cleavage site  $\rightarrow$  likely acquired through recombination
    - Origin of donor sequence (virus genome, host transcript, etc.)?
    - Host for recombination (bat, intermediate host, etc.)?
  - 2. Route of transmission to human populations
    - Furin acquisition = unique event to explain shift from enteric to respiratory tropism?
- Proof-of-concept to study the emergence of CoVs in Indochinese peninsula

**Perspectives**: New collections from Vietnam, at the border with China

# References

Temmam S. et al., Bat coronaviruses related to SARS-CoV-2 and infectious for human cells. Nature. 2022 Apr;604(7905):330-336.

doi: 10.1038/s41586-022-04532-4

Temmam S. et al.. SARS-CoV-2-related bat virus behavior in human-relevant models sheds light on the origin of COVID-19. EMBO Rep. 2023 Apr 5;24(4):e56055.

Temmam S. et al., Genotype and Phenotype haracterization of *Rhinolophus* sp. Sarbecoviruses from Vietnam: Implications for Coronavirus Emergence. Viruses. 2023 Sep 8;15(9):1897.

doi: 10.3390/v15091897



doi: 10.15252/embr.202256055