



HAL
open science

Escape of SARS-CoV-2 Variants KP.1.1, LB.1, and KP.3.3 From Approved Monoclonal Antibodies

Delphine Planas, Isabelle Staropoli, Cyril Planchais, Emilie Yab, Banujaa Jeyarajah, Yannis Rahou, Matthieu Prot, Florence Guivel-Benhassine, Frederic Lemoine, Vincent Enouf, et al.

► **To cite this version:**

Delphine Planas, Isabelle Staropoli, Cyril Planchais, Emilie Yab, Banujaa Jeyarajah, et al.. Escape of SARS-CoV-2 Variants KP.1.1, LB.1, and KP.3.3 From Approved Monoclonal Antibodies. *Pathogens and Immunity*, 2024, 10 (1), pp.1-11. 10.20411/pai.v10i1.752 . pasteur-04748459

HAL Id: pasteur-04748459

<https://pasteur.hal.science/pasteur-04748459v1>

Submitted on 22 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Published September 30, 2024

ESCAPE OF SARS-CoV-2 VARIANTS KP.1.1, LB.1, AND KP.3.3 FROM APPROVED MONOCLONAL ANTIBODIES

AUTHORS

Delphine Planas^{1,2}, Isabelle Staropoli¹, Cyril Planchais³, Emilie Yab⁴, Banujaa Jeyarajah⁴, Yannis Rahou⁴, Matthieu Prot⁵, Florence Guivel-Benhassine¹, Frederic Lemoine⁶, Vincent Enouf⁴, Etienne Simon-Loriere^{5*}, Hugo Mouquet^{3*}, Marie-Anne Rameix-Welti^{4,7*}, Olivier Schwartz^{1,2*}

AFFILIATED INSTITUTIONS

¹ Virus and Immunity Unit, Institut Pasteur, Université Paris Cité, CNRS UMR3569, Paris, France

² Vaccine Research Institute, Créteil, France

³ Humoral Immunology Unit, Institut Pasteur, Université Paris Cité, Paris, France

⁴ National Reference Center for Respiratory Viruses, Molecular Mechanisms of Multiplication of Pneumoviruses Unit, Institut Pasteur, Université Paris Cité, Paris, France

⁵ G5 Evolutionary Genomics of RNA Viruses, Institut Pasteur, Université Paris Cité, Paris, France

⁶ Bioinformatics and Biostatistics Hub, Paris, France

⁷ Molecular Mechanisms of Multiplication of Pneumoviruses, Université Paris-Saclay, Université de Versailles St. Quentin, UMR 1173 (2I), INSERM; Assistance Publique des Hôpitaux de Paris, Paris, France

*co-last authors

CORRESPONDING AUTHOR

Olivier Schwartz

email: olivier.schwartz@pasteur.fr

DOI

10.20411/pai.v10i1.752

SUGGESTED CITATION

Planas D, Staropoli I, Planchais C, Yab E, Jeyarajah B, Rahou Y, Prot⁵, Guivel-Benhassine F, Lemoine F, Enouf V, Simon-Loriere E, Mouquet H, Rameix-Welti MA, Schwartz O. Escape of SARS-CoV-2 Variants KP.1.1, LB.1, and KP.3.3 From Approved Monoclonal Antibodies. *Pathogens and Immunity*. 2024;10(1):1–11. doi: 10.20411/pai.v10i1.752

ABSTRACT

Background: First-generation anti-SARS-CoV-2 monoclonal antibodies (mAbs) used for prophylaxis or therapeutic purposes in immunocompromised patients have been withdrawn because of the emergence of resistant Omicron variants. In 2024, 2 novel mAbs, VYD222/Pemivibart and AZD3152/Sipavibart, were approved by health authorities, but their activity against contemporary JN.1 sublineages is poorly characterized.

Methods: We isolated authentic JN.1.1, KP.1.1, LB.1, and KP.3.3 viruses and evaluated their sensitivity to neutralization by these mAbs in 2 target cell lines.

Results: Compared to ancestral strains, VYD222/Pemivibart remained moderately active against JN.1 subvariants, with a strong increase of 50% Inhibitory Concentration (IC₅₀), reaching up to 3 to 15 µg/mL for KP.3.3. AZD3152/Sipavibart neutralized JN.1.1 but lost antiviral efficacy against KP.1.1, LB.1, and KP.3.3.


Conclusions: Our results highlight the need for a close clinical monitoring of VYD222/Pemivibart and raise concerns about the clinical efficacy of AZD3152/Sipavibart.

KEYWORDS

SARS-CoV-2 variants; monoclonal antibodies; Sipavibart; Pemivibart; neutralization

INTRODUCTION

The JN.1 lineage arose in late 2023 and rapidly outcompeted previous SARS-CoV-2 variants [1]. Since then, JN.1 continued its evolution, with the appearance of sublineages carrying convergent mutations in the Spike (S) protein, notably F456L or R346T, and more recently S31del [2, 3] (Figure 1A,B). Sublineage specific mutations also appeared, such as Q493E. As of August 2024, the KP.1, KP.2, LB.1, and KP.3 variants, that carry various combinations of these substitutions, represented about 80% of sequenced circulating strains (Figure 1B,C). These mutations are collectively responsible for increased immune escape from previously infected and vaccinated populations [2, 3].







Figure 1. Sequence and proportion of SARS-CoV-2 variants circulating in mid-2024. (A) Spike mutations of SARS-CoV-2 variants relative to the spike domains of the BA.2.86.1 parental strain. JN.1, KP.1.1, LB.1, and KP.3.3 were further studied. The mutations in KP.2, JN.1.18, and JN.1.16 are also indicated. (B) Evolution of the prevalence of the main SARS-CoV-2 lineages from January to August 4, 2024, showing the expansion of JN.1-derived lineages. The star indicates the presence of different sublineages. The variants highlighted with a frame are analyzed in this study. Data from the GISAID EpiCoV database. (C) Comparison of spike mutations of selected SARS-CoV-2 variants using Wuhan_Hu-1 (NC_045512) as reference. The color scale reflects the frequency of the mutations within lineages based on the data available on the GISAID EpiCoV database.

The sensitivity of KP.1, LB.1, and KP.3 variants to monoclonal antibodies (mAbs) developed for clinical use is poorly characterized. First-generation anti-S mAbs, previously approved by the Food and Drug Administration (FDA), the European Medicines Agency (EMA), or other agencies, had their authorization withdrawn after the emergence of Omicron variants, because of escape mutations in the receptor binding domain (RBD) of Spike protein. In 2024, novel mAbs, efficient against Omicron variants, have been tested in clinical trials or are available for clinical use in some countries [4]. These include AZD3152/Sipavibart, VYD222/Pemivibart, and SA55 that belong to different anti-RBD antibody classes and target distinct epitopes.

AZD3152 (or SipavibartTM) [5] was authorized in July 2024 by the EMA, for pre-exposure prophylaxis in patients with immunocompromising conditions and at high risk of developing severe COVID-19 [6]. However, AZD3152/Sipavibart-resistant viruses, carrying escape mutations at RBD positions 415, 456, and 458, have been described in pre-JN.1 lineages [5]. VYD222 (also termed Pemivibart or PempardaTM) obtained an emergency use authorization by the FDA in March 2024, for pre-exposure prophylaxis in patients who are moderately to severely immunocompromised [7]. VYD222/Pemivibart preserved *in vitro* efficacy against pre-JN.1 strains carrying the F456L mutation [7]. SA55 was isolated from a SARS-CoV-1 infected, SARS-CoV-2 vaccinated individual and displays a broad sarbecovirus neutralization profile, including JN.1, when used alone or in combination with another mAb (SA58) [8, 9]. SA55 has been tested in a clinical trial initiated in 2023 in China, in patients with hematological disorders who are persistently positive for SARS-CoV-2 [10].

Here, we isolated the main SARS-CoV-2 variants circulating in mid-2024 and tested their sensitivity to neutralization by a panel of mAbs.

METHODS

Virus Isolation

Viral strains were amplified through 1 or 2 passages on Vero E6 TMPRSS2 cells and 1 passage on IGROV-1 cells. Cells were plated in T75 flasks and cultivated in culture media (Dulbecco's Modified Eagle Medium (DMEM), 10% fetal calf serum, and 1% Penicillin/Streptomycin) at 37°C, 5% CO₂. Supernatants were harvested 2 or 3 days after viral exposure. Viral supernatants were sequenced directly from nasopharyngeal swabs and after isolation and amplification on IGROV-1 cells to confirm identity, the presence of specific mutations in the spike protein, and the absence of cell culture-derived mutations. The titration of viral stocks was performed on S-Fuse cells [1, 11, 12].

The D614G and JN.1 strains have been described [1, 13]. The KP.1.1 (hCoV-19/France/IDF-RELAB-IPP05044/2024), LB.1 (hCoV-19/France/GES-RELAB-IPP04736/2024), and KP.3.3 strains (hCoV-19/France/BFC-IPP06087/2024) were isolated and amplified by the National Reference Center for Respiratory Viruses hosted by Institut Pasteur.

Monoclonal Antibodies

S309 (or Sotrovimab) was previously described [14]. Codon-optimized synthetic DNA fragments coding for the immunoglobulin variable domains of SA55 (BD55-5514) [8], AZ3152/Sipavibart [5], and VYD222/Pemivibart were synthesized (GeneArt, Thermo Fisher Scientific) and cloned into human IgG1 expression vectors as previously described [15]. Recombinant IgG1 antibodies were

produced by transient co-transfection of Freestyle™ 293-F suspension cells (Thermo Fisher Scientific) using the PEI-precipitation method and purified from culture supernatants by affinity chromatography using Protein G Sepharose® 4 Fast Flow (GE Healthcare) as previously described [15].

Cell Lines

IGROV-1 and S-Fuse (U20S) cells were previously described [1, 11]. Cells were regularly tested to confirm they were negative for mycoplasma.

Virus Titration for Neutralizing Assay

Titration of viral stocks was performed on S-Fuse and IGROV-1 cells. Neutralization assays were conducted using a multiplicity of infection sufficient to produce about 200 syncytia/well with S-Fuse cells and achieve 40% of infected IGROV-1 cells.

S-Fuse Neutralization Assay

U2OS-ACE2 GFP1-10 and GFP11 cells, also termed S-Fuse cells, become GFP+ when they are productively infected by SARS-CoV-2 [11, 13]. Cells were mixed (ratio 1:1) and plated overnight at 12×10^3 per well in a μ Clear 96-well plate (Greiner Bio-One). The indicated SARS-CoV-2 strains were incubated with serially diluted monoclonal antibodies (mAbs) for 15 minutes at room temperature and added to S-Fuse cells. After 18 hours, cells were fixed with 2% PFA (Electron Microscopy Sciences, cat# 15714-S), washed, and stained with Hoechst (dilution of 1:1,000, Invitrogen, cat# H3570). Images were acquired using an Opera Phenix high-content confocal microscope (PerkinElmer). The number of GFP syncytia and the number of nuclei were quantified using Harmony software (PerkinElmer). The percentage of neutralization was calculated using the number of syncytia with the following formula: $100 \times (1 - [\text{value with mAb} - \text{value in "non-infected"}] / [\text{value in "no mAb"} - \text{value in "non-infected"}])$.

For each mAb, the half-maximal inhibitory concentration (IC₅₀) in ng/mL was calculated with a reconstructed curve using the percentage of neutralization at each concentration.

IGROV-1 Neutralization Assay

Sixteen hours before infection, 30×10^3 cells per well were seeded in a μ Clear black 96-well plate (Greiner Bio-One). The indicated SARS-CoV-2 strains were incubated with serially diluted mAbs for 15 minutes at room temperature and added to IGROV-1 cells. After 24 hours, cells were fixed with 2% PFA (Electron Microscopy Sciences, cat# 15714-S). The cells were then intracellularly stained with anti-SARS-CoV-2 nucleoprotein (N) antibody NCP-1 (0.1 μ g/mL) as described [1]. The staining was carried out in PBS with 0.05% saponin 1% BSA, and 0.05% sodium azide for 1 hour. Cells were then washed twice with PBS and stained with anti-IgG Alexa Fluor 488 (dilution 1:500, Invitrogen; cat# A11029) for 30 minutes before being washed twice with PBS. Hoechst 33342 (Invitrogen, cat# H3570) was added during the final PBS wash. Images were captured using an Opera Phenix high-content confocal microscope (PerkinElmer). The N-positive area and the number of nuclei were quantified using Harmony Software v4.9 (PerkinElmer). The percentage of neutralization was calculated using the N-positive area with the following formula: $100 \times (1 - [\text{value with mAb} - \text{value in "non-infected"}] / [\text{value in "no mAb"} - \text{value in "non-infected"}])$. For each mAb, the half-maximal inhibitory concentration (IC₅₀) in ng/mL was calculated with a reconstructed curve using the percentage of neutralization at each concentration.

Statistical Analysis

Figures were generated using Prism 9 (GraphPad Software). Statistical analysis was conducted using GraphPad Prism 9. Data are mean \pm SD of 3 independent experiments.

Lineage Monitoring

To visualize the evolution of the frequency of SARS-CoV-2 lineages, we analyzed the viral genomic surveillance data deposited in the GISAID database (<https://www.gisaid.org>; metadata downloaded on July 7, 2024) [16, 17]. The hierarchical relationships between lineages were retrieved from the pangolin GitHub repository (<https://github.com/cov-lineages/pango-designation>). We analyzed SARS-CoV-2 data collected from January 1, 2024, to August 4, 2024, using R 4.3 and ggplot 3.4.3. Mutations that are common and specific to lineages of interest were computed using the outbreak.info R package (<https://outbreak-info.github.io/R-outbreak-info>) [18].

Data Availability

All data supporting the findings of this study are available within the article or from the corresponding author upon reasonable request without any restrictions. The sequencing data generated in this study have been deposited in the GISAID EpiCoV database.

RESULTS

We examined the sensitivity of SARS-CoV-2 variants JN.1.1, KP.1.1, LB.1, and KP.3.3 to VYD222/Pemivibart, AZD3152/Sipavibart, and SA55. We included the ancestral D614G strain as control. We isolated KP.1.1, LB.1, and KP.3.3 (which carries the same S protein as KP.3) variants from nasal swabs of individuals with sequence-diagnosed infections. Sequences of outgrown viruses confirmed the identity of the variants (Figure 1A). The mAbs were not commercially available for research purposes. Therefore, we retrieved their sequences from public databases and produced biosimilar molecules. As an additional control, we used SotrovimabTM that neutralizes several Omicron strains but not JN.1, and is no longer approved [1].


We measured the sensitivity of the viral isolates to mAbs first using S-Fuse cells as targets [12] (Figure 2A). These cells were engineered to express ACE2 and are thus sensitive to SARS-CoV-2 [1, 12]. The four mAbs efficiently neutralized D614G, with EC50s of 18-39 ng/mL, corresponding to those described in the literature. As expected, S309/Sotrovimab lost any activity against the 4 JN.1-derived strains. AZD3152/Sipavibart inhibited JN.1.1, with an EC50 of 198 ng/mL, but no longer neutralized KP.1.1, LB.1, and KP.3.3 (Figure 1B). The F456L substitution present in the 3 variants likely mediates this resistance. VYD222/Pemivibart was poorly active against JN.1.1 and displayed a decreased antiviral activity against KP.1.1, LB.1, and KP.3.3 (Fig. 2A). The EC50s reached up to 16,000 ng/mL, corresponding to up to 888-fold reduction of potency against the 4 variants compared to D614G. The antiviral activity of SA55 was preserved against the variants, with EC50s that remained remarkably low (7 to 23 ng/mL) (Figure 1B).

We did not isolate a KP.2 variant, but its profile of resistance is likely similar to KP.1, since their S only differ at position 1086, outside the RBD (Figure 1A).


We then sought to confirm these results using another cell line. We selected IGROV-1 cells, because they naturally express ACE2 and are highly sensitive to SARS-CoV-2, including Omicron

and JN.1 variants [1]. The profile of neutralization of the 5 SARS-CoV-2 strains was similar in S-Fuse (Figure 2A) and IGROV-1 cells (Figure 2B). The IC50 were also in the same range in the 2 cell types (Figure 2C).

A



B



C

	S309/Sotrovimab		VYD222/Pemivibart		AZD3152/Sipavibart		SA55	
	S-Fuse	IGROV-1	S-Fuse	IGROV-1	S-Fuse	IGROV-1	S-Fuse	IGROV-1
D614G	39	26	18	13	27	38	23	29
JN.1.1	-	-	3114	1136	198	118	16	6
KP.1.1	-	-	7265	2950	-	-	7	7
LB.1	-	-	16387	9297	-	-	23	11
KP.3.3	-	-	14963	3668	-	-	11	3

IC50 (ng/mL)

Figure 2. Neutralization of SARS-CoV-2 variants circulating in mid-2024 by mAbs. A. Neutralization curves of mAbs in S-Fuse cells. Dose-response analysis of neutralization of the indicated variants by S309/Sotrovimab, VYD222/Pemivibart, AZD3152/Sipavibart, and SA55. Data are presented as mean ± standard deviation of 2-3 independent experiments. B. Neutralization curves of mAbs in IGROV-1 cells. Dose-response analysis of neutralization of the indicated variants by S309/Sotrovimab, VYD222/Pemivibart, AZD3152/Sipavibart, and SA55. Data are presented as mean ± standard deviation of 2-3 independent experiments. C. EC50 values (in ng/mL) for each mAb against the indicated viral strains in the 2 cell lines. “-”: no antiviral activity.

DISCUSSION

Altogether, our results indicate that AZD3152/Sipavibart totally lost antiviral activity against the prevalent strains circulating in mid-2024, most likely because of the presence of the F456L substitution in Spike protein. VYD222/Pemivibart remains active against JN.1.1, KP.1.1, LB.1, and KP.3.3, with however a strong increase in IC50. The loss of activity of VYD222/Pemivibart has been recently reported in a preprint, using VSV-based pseudotypes [19]. As of August 2024, the KP3.1.1 variant that combines the F456L and Q493E mutations found in KP.3 and KP.3.3 with the S31 deletion found in LB.1 has been on the rise [19]. Future work will help to assess the sensitivity of the rapidly diversifying JN.1 family to these mAbs.

CONCLUSION

Our *in vitro* results may not directly translate into clinical efficacy but raise concerns about the medical use of AZD3152/Sipavibart and warrant a close surveillance of Pemivibart, when most of the circulating strains totally or partially escape neutralization by the 2 antibodies. The mAb SA55 represents a promising alternative.

ACKNOWLEDGMENTS

The authors thank Timothée Bruel for critical reading of the manuscript, members of the Virus and Immunity Unit and other teams for discussion and help, Nathalie Aulner and the staff at the UtechS Photonic BioImaging (UPBI) core facility (Institut Pasteur), a member of the France Bio-Imaging network, for image acquisition and analysis, the members of the National Reference Center for Viruses of Respiratory Infections hosted at Institut Pasteur, the RELAB network, and the Mutualized Platform for Microbiology (P2M). This work has used the computational and storage services (Maestro cluster) provided by the IT department at Institut Pasteur, Paris. We gratefully acknowledge the originating laboratories responsible for obtaining the specimens, as well as the submitting laboratories where the genome data were generated and shared via GISAID.

FUNDING

Work in OS lab is funded by Institut Pasteur, Urgence COVID-19 Fundraising Campaign of Institut Pasteur, Fondation pour la Recherche Médicale (FRM), ANRS, the Vaccine Research Institute (VRI) (ANR-10-LABX-77), Labex IBEID (ANR-10-LABX-62-IBEID), the HERA Project DURABLE (grant 101102733) and Integrating Multi-Disciplinary Expertise in a Learning and Adaptive European Pandemic Preparedness System (LEAPS project), DP is supported by the VRI. The E.S.-L. laboratory is funded by Institut Pasteur, the INCEPTION program (Investissements d'Avenir grant ANR-16-CONV-0005), the Ixcore foundation for research, the French Government's Investissement d'Avenir programme, Laboratoire d'Excellence 'Integrative Biology of Emerging Infectious Diseases' (grant no. ANR-10-LABX-62-IBEID), the HERA Project DURABLE (grant no 101102733) and the NIH PICREID (grant no U01AI151758).

POTENTIAL CONFLICTS OF INTEREST

CP, HM, and OS have a patent application for anti-SARS-CoV-2 monoclonal antibodies not used in the present study (PCT/FR2021/070522, WO 2022/228827A1), and HM is a scientific consultant for SpikImm biotech. The remaining authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Experimental strategy design, experiments: DP, IS, CP, FG-B, ES-L, HM, M-ARW, OS. Vital materials: CP, EY, BJ, YR, VE, HM, M-ARW. Phylogenetic analysis: ES-L. Viral sequencing: EY, BJ, YR, VE, MP, FL, ES-L, M-ARW. Manuscript writing and editing: DP, ES-L, HM, M-ARW, OS.

REFERENCES

1. Planas D, Staropoli I, Michel V, Lemoine F, Donati F, Prot M, Porrot F, Guivel-Benhassine F, Jeyarajah B, Brisebarre A, Dehan O, Avon L, Bolland WH, Hubert M, Buchrieser J, Vanhoucke T, Rosenbaum P, Veyer D, Pere H, Lina B, Trouillet-Assant S, Hocqueloux L, Prazuck T, Simon-Loriere E, Schwartz O. Distinct evolution of SARS-CoV-2 Omicron XBB and BA.2.86/JN.1 lineages combining increased fitness and antibody evasion. *Nat Commun.* 2024;15(1):2254. doi: [10.1038/s41467-024-46490-7](https://doi.org/10.1038/s41467-024-46490-7). PubMed PMID: 38480689; PMCID: PMC10938001.
2. Kaku Y, Yo MS, Tolentino JE, Uriu K, Okumura K, Genotype to Phenotype Japan C, Ito J, Sato K. Virological characteristics of the SARS-CoV-2 KP.3, LB.1, and KP.2.3 variants. *Lancet Infect Dis.* 2024;24(8):e482-e3. doi: [10.1016/S1473-3099\(24\)00415-8](https://doi.org/10.1016/S1473-3099(24)00415-8). PubMed PMID: 38945150.
3. Kaku Y, Uriu K, Kosugi Y, Okumura K, Yamasoba D, Uwamino Y, Kuramochi J, Sadamasu K, Yoshimura K, Asakura H, Nagashima M, Genotype to Phenotype Japan C, Ito J, Sato K. Virological characteristics of the SARS-CoV-2 KP.2 variant. *Lancet Infect Dis.* 2024;24(7):e416. doi: [10.1016/S1473-3099\(24\)00298-6](https://doi.org/10.1016/S1473-3099(24)00298-6). PubMed PMID: 38782005.
4. Focosi D, Franchini M, Casadevall A, Maggi F. An update on the anti-spike monoclonal antibody pipeline for SARS-CoV-2. *Clin Microbiol Infect.* 2024;30(8):999-1006. doi: [10.1016/j.cmi.2024.04.012](https://doi.org/10.1016/j.cmi.2024.04.012). PubMed PMID: 38663655.
5. Cai Y, Diallo S, Rosenthal K, Ren K, Flores DJ, Dippel A, Oganessian V, van Dyk N, Chen X, Cantu E, Choudhary R, Sulikowski M, Adissu H, Chawla B, Kar S, Liu C, Djokaite-Guraliuc A, Mongkolsapaya J, Rajan S, Loo YM, Beavon R, Webber C, Chang LJ, Thomas S, Clegg L, Zhang H, Scream GR, Philbin N, Harre M, Selim A, Martinez-Alier N, Uriel A, Cohen TS, Perez JL, Esser MT, Blair W, Francica JR. AZD3152 neutralizes SARS-CoV-2 historical and contemporary variants and is protective in hamsters and well tolerated in adults. *Sci Transl Med.* 2024;16(753):eado2817. doi: [10.1126/scitranslmed.ado2817](https://doi.org/10.1126/scitranslmed.ado2817). PubMed PMID: 38924429.
6. AstraZeneca. Sipavibart EMA regulatory submission accepted under accelerated assessment for COVID-19 prevention. 2024.
7. Fact sheet for healthcare providers: Emergency Use Authorization of Pempgarda (pemi-vibart). Invivyd; 2024. Accessed March 25, 2024. In: Pempgarda, editor.: FDA.gov; 2024.
8. Cao Y, Jian F, Zhang Z, Yisimayi A, Hao X, Bao L, Yuan F, Yu Y, Du S, Wang J, Xiao T, Song W, Zhang Y, Liu P, An R, Wang P, Wang Y, Yang S, Niu X, Zhang Y, Gu Q, Shao F, Hu Y, Yin W, Zheng A, Wang Y, Qin C, Jin R, Xiao J, Xie XS. Rational identification of potent and broad sarbecovirus-neutralizing antibody cocktails from SARS convalescents. *Cell Rep.* 2022;41(12):111845. doi: [10.1016/j.celrep.2022.111845](https://doi.org/10.1016/j.celrep.2022.111845). PubMed PMID: 36493787; PMCID: PMC9712074.

9. Yang S, Yu Y, Xu Y, Jian F, Song W, Yisimayi A, Wang P, Wang J, Liu J, Yu L, Niu X, Wang J, Wang Y, Shao F, Jin R, Wang Y, Cao Y. Fast evolution of SARS-CoV-2 BA.2.86 to JN.1 under heavy immune pressure. *Lancet Infect Dis*. 2024;24(2):e70-e2. doi: [10.1016/S1473-3099\(23\)00744-2](https://doi.org/10.1016/S1473-3099(23)00744-2). PubMed PMID: 38109919.
10. Beijing Boren Hospital. Efficacy and Safety of the Anti-COVID-19 Antibody SA55 for Injection in Patients With Hematological Disorders Who Are Persistently Positive for COVID-19 [Internet]. ClinicalTrials.gov identifier: NCT123456. Available from: <https://clinicaltrials.gov/study/NCT05675943>.
11. Buchrieser J, Dufloo J, Hubert M, Monel B, Planas D, Rajah MM, Planchais C, Porrot F, Guivel-Benhassine F, Van der Werf S, Casartelli N, Mouquet H, Bruel T, Schwartz O. Syncytia formation by SARS-CoV-2-infected cells. *EMBO J*. 2020;39(23):e106267. doi: [10.15252/embj.2020106267](https://doi.org/10.15252/embj.2020106267). PubMed PMID: 33051876; PMCID: PMC7646020.
12. Planas D, Saunders N, Maes P, Guivel-Benhassine F, Planchais C, Buchrieser J, Bolland WH, Porrot F, Staropoli I, Lemoine F, Pere H, Veyer D, Puech J, Rodary J, Baele G, Dellicour S, Raymenants J, Gorissen S, Geenen C, Vanmechelen B, Wawina-Bokalanga T, Marti-Carreras J, Cuypers L, Seve A, Hocqueloux L, Prazuck T, Rey FA, Simon-Loriere E, Bruel T, Mouquet H, Andre E, Schwartz O. Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. *Nature*. 2022;602(7898):671-5. doi: [10.1038/s41586-021-04389-z](https://doi.org/10.1038/s41586-021-04389-z). PubMed PMID: 35016199.
13. Planas D, Bruel T, Grzelak L, Guivel-Benhassine F, Staropoli I, Porrot F, Planchais C, Buchrieser J, Rajah MM, Bishop E, Albert M, Donati F, Prot M, Behillil S, Enouf V, Maquart M, Smati-Lafarge M, Varon E, Schortgen F, Yahyaoui L, Gonzalez M, De Seze J, Pere H, Veyer D, Seve A, Simon-Loriere E, Fafi-Kremer S, Stefic K, Mouquet H, Hocqueloux L, van der Werf S, Prazuck T, Schwartz O. Sensitivity of infectious SARS-CoV-2 B.1.1.7 and B.1.351 variants to neutralizing antibodies. *Nat Med*. 2021;27(5):917-24. doi: [10.1038/s41591-021-01318-5](https://doi.org/10.1038/s41591-021-01318-5). PubMed PMID: 33772244.
14. Bruel T, Hadjadj J, Maes P, Planas D, Seve A, Staropoli I, Guivel-Benhassine F, Porrot F, Bolland WH, Nguyen Y, Casadevall M, Charre C, Pere H, Veyer D, Prot M, Baidaliuk A, Cuypers L, Planchais C, Mouquet H, Baele G, Mouthon L, Hocqueloux L, Simon-Loriere E, Andre E, Terrier B, Prazuck T, Schwartz O. Serum neutralization of SARS-CoV-2 Omicron sublineages BA.1 and BA.2 in patients receiving monoclonal antibodies. *Nat Med*. 2022;28(6):1297-302. doi: [10.1038/s41591-022-01792-5](https://doi.org/10.1038/s41591-022-01792-5). PubMed PMID: 35322239; PMCID: 35322239.
15. Planchais C, Fernandez I, Bruel T, de Melo GD, Prot M, Beretta M, Guardado-Calvo P, Dufloo J, Molinos-Albert LM, Backovic M, Chiaravalli J, Giraud E, Vesin B, Conquet L, Grzelak L, Planas D, Staropoli I, Guivel-Benhassine F, Hieu T, Boulle M, Cervantes-Gonzalez M, Ungeheuer MN, Charneau P, van der Werf S, Agou F, French CCSG, Group CS, Dimitrov JD, Simon-Loriere E, Bourhy H, Montagutelli X, Rey FA, Schwartz O, Mouquet H. Potent human broadly SARS-CoV-2-neutralizing IgA and IgG antibodies effective against Omicron BA.1 and BA.2. *J Exp Med*. 2022;219(7). doi: [10.1084/jem.20220638](https://doi.org/10.1084/jem.20220638). PubMed PMID: 35704748; PMCID: PMC9206116.

16. Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution to global health. *Glob Chall*. 2017;1(1):33-46. doi: [10.1002/gch2.1018](https://doi.org/10.1002/gch2.1018). PubMed PMID: 31565258; PMCID: PMC6607375.
17. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill*. 2017;22(13):30494. doi: [10.2807/1560-7917.ES.2017.22.13.30494](https://doi.org/10.2807/1560-7917.ES.2017.22.13.30494). PubMed PMID: 28382917; PMCID: PMC5388101.
18. Tsueng G, Mullen JL, Alkuzweny M, Cano M, Rush B, Haag E, Lin J, Welzel DJ, Zhou X, Qian Z, Latif AA, Hufbauer E, Zeller M, Andersen KG, Wu C, Su AI, Gangavarapu K, Hughes LD. Outbreak.info Research Library: a standardized, searchable platform to discover and explore COVID-19 resources. *Nat Methods*. 2023;20(4):536-40. doi: [10.1038/s41592-023-01770-w](https://doi.org/10.1038/s41592-023-01770-w). PubMed PMID: 36823331; PMCID: PMC10393269.
19. Wang Q, Guo Y, Ho J, Ho DD. Pemivibart is less active against recent SARS-CoV-2 JN.1 sublineages. *bioRxiv*. 2024:2024.08.12.607496. doi: [10.1101/2024.08.12.607496](https://doi.org/10.1101/2024.08.12.607496).

Footnotes

Submitted August 20, 2024 | Accepted September 4, 2024 | Published September 30, 2024

Copyright

Copyright © 2024 The Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License.